



A Journal of the Gesellschaft Deutscher Chemiker

Angewandte Chemie

GDCh

International Edition

www.angewandte.org

Accepted Article

Title: Reductive Molybdenum-Catalyzed Direct Amination of Boronic Acids with Nitro Compounds

Authors: Samuel Suárez-Pantiga, Raquel Hernández-Ruiz, Cintia Virumbrales, María R. Pedrosa, and Roberto Sanz

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: *Angew. Chem. Int. Ed.* 10.1002/anie.201812806
Angew. Chem. 10.1002/ange.201812806

Link to VoR: <http://dx.doi.org/10.1002/anie.201812806>
<http://dx.doi.org/10.1002/ange.201812806>

Reductive Molybdenum-Catalyzed Direct Amination of Boronic Acids with Nitro Compounds

Samuel Suárez-Pantiga,* Raquel Hernández-Ruiz, Cintia Virumbrales, María R. Pedrosa, and Roberto Sanz*

Dedicated to Prof. Dr. Francisco J. Fañanás on the occasion of his retirement.


Abstract: The synthesis of aromatic amines is of utmost importance in a wide range of chemical contexts. We report a direct amination of boronic acids with nitro compounds to yield (hetero)aryl amines. The novel combination of a dioxomolybdenum(VI) catalyst and triphenylphosphine as inexpensive reductant has revealed to be decisive to achieve this new C–N coupling. Our methodology has proven to be scalable, air and moisture tolerant, highly chemoselective and engages both aliphatic and aromatic nitro compounds. Moreover, this general and step-economical synthesis of aromatic secondary amines showcases orthogonality to other aromatic amine syntheses as it tolerates aryl halides and carbonyl compounds.

The prevalence of aromatic amines in the chemical, materials, agrochemical and pharmaceutical industries is a constant driver for the innovation of synthetic methods.^[1] The most general transformations to prepare (hetero)aryl amines include direct alkylation of amines with alkyl halides,^[2] Buchwald-Hartwig,^[3] Chan-Lam-Evans^[4] or Ullman-type^[4a,5] C–N cross-coupling reactions and reductive amination.^[6] These amination methodologies employ amines as the nitrogen source, particularly anilines, which are usually obtained by reduction of nitroarene derivatives. As an alternative, other nitrogen-containing compounds have also been used as starting materials in C–N bond forming aminations like nitrosoarenes,^[7] nitrenoids,^[8] or hydroxylamines,^[9] which are not commercially available and, again, must be obtained in a previous synthetic step by partial reduction of the nitro function. Therefore, the development of a general and efficient methodology for the synthesis of aromatic amines directly from abundant and readily available nitro(hetero)arenes is very attractive and remains an active research field. The use of nitro compounds as starting materials not only improves step-economy, saving time and cost, but may also allow tolerance of functional groups such as unprotected amine.^[10]


Although nitroaromatics are efficiently engaged as starting materials in C–N bond forming reductive cyclizations to synthesize indoles or carbazoles,^[11] the intermolecular amination to afford anilines in one single operational step has proved to be more challenging. The scarcely reported methodologies have in common that the nitro group evolves to a partially reduced

intermediate, which reacts with a proper alkylating or arylating reagent. Therefore, the key success factor lies in reducing the nitro group to the corresponding intermediate avoiding its strongly favored over-reduction or its evolution through side-reactions. In this field, Baran's and Hu's groups have described the Fe-catalyzed synthesis of *N*-alkylanilines through an *in situ* generated nitroso intermediate, which is subsequently trapped by highly reactive alkyl radicals derived from alkenes^[12] and alkyl halides,^[13] respectively (Scheme 1a). Conceptually different electrophilic aminations have also been reported.^[14] Within this framework, Knochel developed the construction of diarylamines by initial addition of arylmagnesium compounds to nitroarenes, involving the intermediacy of aryl nitroso and diarylhydroxylamine derivatives (Scheme 1b).^[15] More recently, Niggemann has described an elegant reductive coupling of zinc organyls with both aromatic and aliphatic nitro compounds. This is based in the partial reduction of the nitro group with B₂Pin₂ that yields a nitrenoid as the electrophilic amination reagent (Scheme 1c).^[16] Despite the potential of this transformation, it is currently underexploited since a fine tuning of the nitrenoid could allow the amination of less nucleophilic reagents such as boronic acids.


radical amination



electrophilic amination



d) our hypothesis:



[*] Dr. S. Suárez-Pantiga, R. Hernández-Ruiz, C. Virumbrales, Dr. M. R. Pedrosa, Prof. Dr. R. Sanz
 Department of Chemistry
 Universidad de Burgos
 Pza Misael s/n, 09001 Burgos, (Spain)
 E-mail: svsuarez@ubu.es; rsd@ubu.es

Supporting information for this article is given via a link at the end of the document.

Scheme 1. Direct amination with nitroarenes: state-of-the-art.

Boronic acids are a relevant class of intermediates as they exhibit wide functional group compatibility and synthetic versatility and are readily available from commercial sources or easily synthesized.^[17] Based on these features, achieving a direct amination of boronic acids with nitro compounds is a major milestone, which repurposes these two accessible and simple building blocks as coupling partners delivering an excellent platform to easily construct diverse libraries of valuable aromatic amines. Only very recently, during the preparation of this manuscript, a protocol for this reaction by using PhSiH₃ as reductant and phosphetane P(III)/O=P(V) catalysis was reported by Radosevich and co-workers.^[18] Ideally, the design of a moisture and air tolerant procedure and the utilization of an affordable reducing agent would be crucial to provide a general, simple and scalable methodology. Based on our experience in molybdenum catalyzed reactions,^[19] we hypothesized that the Mo-catalyzed formation of a partially-reduced intermediate from a nitro compound may allow the direct electrophilic amination of boronic acids (Scheme 1d).

Initially, model reaction between phenylboronic acid and 1-chloro-4-nitrobenzene was chosen (Table 1) and a screen of several reducing agents was performed.^[20] Pinacol proved to be unsuitable for this transformation as only direct reduction to the corresponding aniline **B** was observed (entry 1).^[19a] However, PPh₃ and P(OEt)₃ were identified as viable reductants leading to high yields of diarylamine **A** and only minor amounts of the aniline side-product **B** (entries 2 and 3). However, when nitrobenzene was tested as substrate the conversion decayed considerably (entries 4 and 5). This decrease was even clearer with 4-nitrotoluene as starting material (entry 6). After some reoptimization,^[20] best results were achieved by simple addition of 2,2'-bipyridine (bpy) as ligand and using PPh₃ as reductant (entries 7–9). When pre-formed MoO₂Cl₂(bpy)₂ was used instead, similar results were obtained although conversion was slightly lower (entry 10).


Table 1. Optimization of the Mo-catalyzed reductive coupling.^[a]

Entry	X	Catalyst	Reductant	Conversion [%] ^[b]	Yield ^[b] A:B [%]
1	Cl	MoO ₂ Cl ₂ (dmf) ₂	Pinacol	70	—:54
2	Cl	MoO ₂ Cl ₂ (dmf) ₂	PPh ₃	94	78:12
3	Cl	MoO ₂ Cl ₂ (dmf) ₂	P(OEt) ₃	95	72:16
4	H	MoO ₂ Cl ₂ (dmf) ₂	PPh ₃	70	55:8
5	H	MoO ₂ Cl ₂ (dmf) ₂	P(OEt) ₃	62	40:15
6	Me	MoO ₂ Cl ₂ (dmf) ₂	PPh ₃	66	35:15
7	Cl	MoO ₂ Cl ₂ (dmf) ₂ /bpy	PPh ₃	>95	>95:<5
8	H	MoO ₂ Cl ₂ (dmf) ₂ /bpy	PPh ₃	>95	88:6

9	Me	MoO ₂ Cl ₂ (dmf) ₂ /bpy	PPh ₃	>95	84:7
10	Me	MoO ₂ Cl ₂ (bpy) ₂	PPh ₃	88	82:<5


[a] Reaction conditions: nitroarene (0.2 mmol), PhB(OH)₂ (0.3 mmol), reductant (0.48 mmol), catalyst (5 mol%) in toluene (0.6 mL) at 100 °C for 20 h, under air. [b] Determined by ¹H NMR analysis with CH₂Br₂ as internal standard.

With an optimized set of conditions in hand, the scope of both the nitro compound and the boronic acid partners was extensively studied. The ability of the reaction to tolerate sensitive functional groups in the nitroaromatic component is exceptional (Scheme 2). Halides are well tolerated in different positions with excellent yields (**2–4**, **13**, **14**, **17–19**), allowing for downstream C–C, C–O, and C–N cross-coupling chemistry. Alkyl (**5**, **20**) and methoxy (**6**) electron-donating groups (EDG) were also tested and remain unscathed. However, starting from 4-nitroanisole MW irradiation was required to achieve full conversion. Particularly remarkable is that amination takes place selectively over the nitro group when free amine is also present without the need for protecting group chemistry (**25**). This example clearly shows that the reaction does not proceed through the amino group, which would allow unconventional disconnections in retrosynthetic analysis. Notable performance is accomplished with nitroarenes bearing strong electron-withdrawing groups (EWG) such as cyano (**7**, **21**), ester (**10**, **22**) or amide (**11**). Despite being a reductive process, ketones (**8**, **9**, **23**) and even aldehydes (**15**) are not reduced, as opposed to the classic reductive amination that would again require protection of these functional groups. Moreover, α,β-unsaturated carbonyls and alkenes emerged unaltered affording the corresponding diarylamines in good yields (**12**, **16**). The process has also revealed to be compatible with nitroarenes bearing several functional groups in the same ring (**24–27**) and with highly sterically demanding *ortho*-disubstituted nitroarenes (**30**), although with more moderate yields. At the same time, polycyclic nitroarenes are well tolerated (**28**, **29**). Remarkably, nitroheteroarenes can also be used to deliver medicinally relevant building blocks containing indole (**31**), pyridine (**32**) and quinoline (**33**) scaffolds. Interestingly, selective mono-amination could be achieved in molecules bearing more than one nitro group (**34**).




Scheme 2. Nitroarene scope (isolated yields). Conditions: nitroarene (0.5 mmol), in toluene (1.5 mL), under air. [a] Performed under MW irradiation (135 °C, 25–40 min).

Next, a screen of diverse boronic acids was evaluated (Scheme 3). The transformation took efficiently place with excellent yields in presence of EWG in the aromatic ring such as halides (**13**, **18**, **35**, **36**) and ester (**37**). Moreover, *o*-substituted boronic acids also proved to effectively promote the transformation (**20**, **38**). At the same time, electron-donating alkoxy substituents were evaluated in different positions (**6**, **39–42**) providing satisfactory results without the need of MW irradiation, thus ideally complementing the nitroarene scope.^[21] In addition, naphthyl (**43**), benzothienyl (**44**) and pyridinyl (**32**) boronic acids revealed also suitable.



Scheme 3. (Hetero)aryl boronic acid scope (isolated yields). Conditions: nitrobenzene (0.5 mmol), in toluene (1.5 mL), under air. [a] Performed under MW irradiation (135 °C, 25 min).


Afterwards, diarylamines, such as **45–49**, possessing both functionalized coupling partners were also synthesized in high yields (Scheme 4). The usefulness of this protocol in the synthesis of bioactive compounds is shown with the efficient preparation of flufenamic acid (**50**), an anthranilic acid derivative.^[22]



Scheme 4. Synthesis of highly functionalized diarylamines (isolated yields). Conditions: nitroarene (0.5 mmol), in toluene (1.5 mL), under air.


Remarkably, alkyl boronic acids were also compatible with this methodology as it was demonstrated with the synthesis of alkyl aryl amines **51–54**. Both primary and secondary alkyl boronic acids could undergo amination with selected

functionalized nitroarenes, although lower yields were obtained compared with the formation of C_{sp2}-N bonds (Scheme 5).



Scheme 5. Amination of alkyl boronic acids (isolated yields). Conditions: nitroarene (0.5–1 mmol), in toluene (1.5–3 mL), under air.

Finally, amination with the more challenging nitroalkanes was evaluated,^[23] in which only Niggemann's very recent approach was able to succeed.^[16b] Preliminary results under conventional heating at 100 °C showcased low conversion and so, MW irradiation was tested. To our delight, high conversion to the desired alkylated anilines was achieved. Several primary and more sterically demanding secondary nitroalkanes were converted in presence of different aromatic boronic acids leading to alkyl amines **53** and **55–60** in moderate to good yields (Scheme 6). Interestingly, when nitromethane is employed as substrate this procedure offers a facile access to selectively monomethylated *N*-methylanilines alternatively to the problematic mono *N*-methylation of anilines.




Scheme 6. Nitroalkane scope (isolated yields). Conditions: nitroalkane (1 mmol), in toluene (1.5 mL), under air.

The described procedure is amenable to scale up due to the ready availability of both nitro compounds and boronic acids, as well as, the use of inexpensive PPh₃ as reductant in combination with a dioxomolybdenum(VI) catalyst easily prepared from affordable Na₂MoO₄.^[24] Accordingly, several nitroarenes were submitted to the reaction conditions at 7 mmol scale to provide gram amounts of aromatic amines **10** (1.42 g, 92%), **15** (1.19 g, 86%), **46** (1.33 g, 88%), and **51** (10 mmol scale, 1.10 g, 74%).

A plausible mechanism is depicted in Scheme 7. It is well known that dioxomolybdenum(VI) catalyst **I** is easily reduced by PPh₃ to a Mo(IV) species,^[25] which upon coordination to the nitro

compound would afford **II**. Next, reoxidation of the Mo center by the coordinated nitro compound would deliver intermediate **III** bearing a nitroso ligand. At this point two different pathways could be considered. Path **A** would involve a second deoxygenation of **III** with PPh₃ giving rise, after reduction of the nitroso ligand and reoxidation of the metal center, to a new molybdooxaziridine (or η²-nitroso) species **IV**.^[26] The attack of the N atom to the boronic acid would afford nitrenoid intermediate **V**, whose subsequent rearrangement would generate aminoboronic acid **VI**. Alternatively, path **B** would involve the release of free nitroso compound **VII**,^[7c] which after reaction with a second equiv of PPh₃ would generate reduced adduct **VIII**. Its interaction with the boronic acid would deliver nitrenoid borate intermediate **IX**, which would eventually release O=PPh₃ and the same aminoboronic acid **VI** through 1,2-migration of a nucleophilic R² group.^[27] Final hydrolysis would provide the isolated secondary aromatic amine.



Scheme 7. Mechanistic proposal.

In summary, we have described a practical, efficient, scalable, air and moisture tolerant procedure for the general synthesis of secondary aromatic amines from nitro compounds in a single synthetic operation. As these results demonstrate, the covered chemical space of our methodology is much wider and diverse than that achieved with classical reactions and includes the synthesis of many highly substituted and functionalized amines. Not only nitroarenes but also nitroalkanes proved to be suitable substrates for this transformation. The wide availability of nitro compounds and boronic acids, in conjunction with the use of affordable PPh₃ as reductant and an easily available Mo(VI) catalyst, make this simple strategy advantageous to construct libraries of (hetero)aryl amines.

Acknowledgements

We are grateful to Junta de Castilla y León (BU022G18), Junta de Castilla y León and FEDER (BU076U16) and Ministerio de Economía y Competitividad (MINECO) (CTQ2016-75023-C2-1-


P) for financial support. R. H.-R. and C. V. thank Ministerio de Educación and Universidad de Burgos for pre-doctoral contracts, respectively. S. S.-P. thanks Junta de Castilla y León and FEDER for a post-doctoral contract.

Keywords: amination • boronic acids • molybdenum • nitro compounds • reduction

- [1] a) *Amino Group Chemistry*, (Ed.: A. Ricci), Wiley-VCH: Weinheim, **2008**; b) M. Liang, J. Chen, *Chem. Soc. Rev.* **2013**, *42*, 3453–3488; c) J. Wang, K. Liu, L. Ma, X. Zhan, *Chem. Rev.* **2016**, *116*, 14675–14725.
- [2] R. N. Salvatore, C. H. Yoon, K. W. Jung, *Tetrahedron* **2001**, *57*, 7785–7811.
- [3] For selected reviews, see: a) J. Bariwal, E. V. der Eicken, *Chem. Soc. Rev.* **2013**, *42*, 9283–9303; b) P. Ruiz-Castillo, S. L. Buchwald, *Chem. Rev.* **2016**, *116*, 12564–12649.
- [4] Reviews: a) J. X. Qiao, P. Y. S. Lam, *Synthesis* **2011**, 829–856; b) K. Sanjeeva Rao, T.-S. Wu, *Tetrahedron* **2012**, *68*, 7735–7754.
- [5] a) Y. Jiang, D. Ma, in *Copper²⁺ Mediated Cross²⁺ Coupling Reactions* (Eds.: G. Evano, N. Blanchard), John Wiley & Sons, Inc., **2014**, pp. 3–40; b) S. Bhunia, G. G. Pawar, S. V. Kumar, Y. Jiang, D. Ma, *Angew. Chem. Int. Ed.* **2017**, *56*, 16136–16179; *Angew. Chem.* **2017**, *129*, 16313–16327.
- [6] For a review, see: H. Alinezhad, H. Yavari, F. Salehian, *Curr. Org. Chem.* **2015**, *19*, 1021–1049.
- [7] a) Y. Yu, J. Srogl, L. S. Liebeskind, *Org. Lett.* **2004**, *6*, 2631–2634; b) V. Dhayalan, C. Sämman, P. Knochel, *Chem. Commun.* **2015**, *51*, 3239–3242; c) S. Roscales, A. G. Csáky, *Org. Lett.* **2018**, *20*, 1667–1671.
- [8] a) P. Starkov, T. F. Jamison, I. Marek, *Chem. Eur. J.* **2015**, *21*, 5278–5300; b) T. V. Nykaza, A. Ramirez, T. S. Harrison, M. R. Luzung, A. T. Radosevich, *J. Am. Chem. Soc.* **2018**, *140*, 3103–3113.
- [9] For selected examples, see: a) C. Zhu, G. Li, D. H. Ess, J. R. Falck, L. Kuerti, *J. Am. Chem. Soc.* **2012**, *134*, 18253–18256; b) N. Matsuda, K. Hirano, T. Satoh, M. Miura, *Angew. Chem. Int. Ed.* **2012**, *51*, 3642–3645; *Angew. Chem.* **2012**, *124*, 3702–3705; c) M. Mailig, R. P. Rucker, G. Lalic, *Chem. Commun.* **2015**, *51*, 11048–11051; d) H.-B. Sun, L. Gong, Y.-B. Tian, J.-G. Wu, X. Zhang, J. Liu, Z. Fu, D. Niu, *Angew. Chem. Int. Ed.* **2018**, *57*, 9456–9460; *Angew. Chem.* **2018**, *130*, 9600–9604.
- [10] For amide syntheses from nitroarenes, see: a) C. W. Cheung, M. L. Ploeger, X. Hu, *Nat. Commun.* **2017**, *8*, 14878; b) C. W. Cheung, M. L. Ploeger, X. Hu, *ACS Catal.* **2017**, *7*, 7092–7096; c) C. W. Cheung, J.-A. Ma, X. Hu, *J. Am. Chem. Soc.* **2018**, *140*, 6789–6792.
- [11] Cadogan reaction: a) J. I. G. Cadogan, M. Cameron-Wood, R. K. Mackie, R. J. G. Searle, *J. Chem. Soc.* **1965**, 4831–4837; b) R. Sanz, J. Escribano, M. R. Pedrosa, R. Aguado, F. J. Arnáiz, *Adv. Synth. Catal.* **2007**, *349*, 713–718; c) H. Gao, Q.-L. Xu, M. Yousufuddin, D. H. Ess, L. Kürti, *Angew. Chem. Int. Ed.* **2014**, *53*, 2701–2705; *Angew. Chem.* **2014**, *126*, 2739–2743; d) S. Tong, Z. Xu, M. Mamboury, Q. Wang, J. Zhu, *Angew. Chem. Int. Ed.* **2015**, *54*, 11809–11812; *Angew. Chem.* **2015**, *127*, 11975–11978; e) K. Yang, F. Zhou, Z. Kuang, G. Gao, T. G. Driver, Q. Song, *Org. Lett.* **2016**, *18*, 4088–4091. Bartoli reaction: f) G. Bartoli, R. Dalpozzo, M. Nardi, *Chem. Soc. Rev.* **2014**, *43*, 4728–4750;
- [12] a) J. Gui, C.-M. Pan, Y. Jin, T. Qin, J. C. Lo, B. J. Lee, S. H. Spengel, M. E. Mertzman, W. J. Pitts, T. E. La Cruz, M. A. Schmidt, N. Darvatkar, S. R. Natarajan, P. S. Baran, *Science* **2015**, *348*, 886–891; b) K. Zhu, M. P. Shaver, S. P. Thomas, *Chem. Sci.* **2016**, *7*, 3031–3035; c) K. Zhu, M. P. Shaver, S. P. Thomas, *Chem. Asian J.* **2016**, *11*, 977–980; d) J. Xiao, Y. He, F. Ye, S. Zhu, *Chem* **2018**, *4*, 1645–1657.
- [13] C. W. Cheung, X. Hu, *Nat. Commun.* **2016**, *7*, 12494.
- [14] M. Corpet, C. Gosmini, *Synthesis* **2014**, *46*, 2258–2271. See also ref. [8a].
- [15] a) I. Sapountzis, P. Knochel, *J. Am. Chem. Soc.* **2002**, *124*, 9390–9391; b) I. Sapountzis, P. Knochel, *Angew. Chem. Int. Ed.* **2004**, *43*, 897–900; *Angew. Chem.* **2004**, *116*, 915–918.
- [16] a) M. Rauser, C. Ascheberg, M. Niggemann, *Angew. Chem. Int. Ed.* **2017**, *56*, 11570–11574; *Angew. Chem.* **2017**, *129*, 11728–11732; b) M. Rauser, C. Ascheberg, M. Niggemann, *Chem. Eur. J.* **2018**, *24*, 3970–3974.
- [17] *Boronic Acids*, Vols. 1&2, (Ed.: D. G. Hall), Wiley-VCH: Weinheim, **2011**. For a highlight on the amination of boronic acids, see: V. Coeffard, X. Moreau, C. Thomassigny, C. Greck, *Angew. Chem. Int. Ed.* **2013**, *52*, 5684–5686; *Angew. Chem.* **2013**, *125*, 5794–5796.
- [18] T. V. Nykaza, J. C. Cooper, G. Li, N. Mahieu, A. Ramírez, M. R. Luzung, A. T. Radosevich, *J. Am. Chem. Soc.* **2018**, *140*, 15200–15205.
- [19] a) N. García, P. García-García, M. A. Fernández-Rodríguez, R. Rubio, M. R. Pedrosa, F. J. Arnáiz, R. Sanz, *Adv. Synth. Catal.* **2012**, *354*, 321–327; b) N. García, P. García-García, M. A. Fernández-Rodríguez, D. García, M. R. Pedrosa, F. J. Arnáiz, R. Sanz, *Green Chem.* **2013**, *15*, 999–1005; c) N. García, R. Rubio-Presa, P. García-García, M. A. Fernández-Rodríguez, M. R. Pedrosa, F. J. Arnáiz, R. Sanz, *Green Chem.* **2016**, *18*, 2335–2340; d) R. Rubio-Presa, M. A. Fernández-Rodríguez, M. R. Pedrosa, F. J. Arnáiz, R. Sanz, *Adv. Synth. Catal.* **2017**, *359*, 1752–1757; e) R. Rubio-Presa, M. R. Pedrosa, M. A. Fernández-Rodríguez, F. J. Arnáiz, R. Sanz, *Org. Lett.* **2017**, *19*, 5470–5473; f) R. Rubio-Presa, S. Suárez-Pantiga, M. R. Pedrosa, R. Sanz, *Adv. Synth. Catal.* **2018**, *360*, 2216–2220. For a review, see: g) R. Hernández-Ruiz, R. Sanz, *Synthesis* **2018**, *50*, 4019–4036.
- [20] See Supporting Information for further details.
- [21] This tendency resembles that observed by Radosevich (ref. [18]) in which more efficient couplings were obtained in the reactions of electron-deficient nitroarenes with electron-rich arylboronic acids.
- [22] See, for instance: H. Gögelein, D. Dahlem, H. C. Englert, H. J. Lang, *FEBS Lett.* **1990**, *268*, 79–82.
- [23] T. Stemmler, F. A. Westerhaus, A.-E. Surkus, M.-M. Pohl, K. Junge, M. Beller, *Green Chem.* **2014**, *16*, 4535–4540.
- [24] R. Sanz, J. Escribano, R. Aguado, M. R. Pedrosa, F. J. Arnáiz, *Synthesis* **2004**, 1629–1632.
- [25] a) R. Sanz, J. Escribano, Y. Fernández, M. R. Pedrosa, F. J. Arnáiz, *Synlett* **2005**, 1389–1392; b) R. Aguado, J. Escribano, M. R. Pedrosa, A. De Cian, R. Sanz, F. J. Arnáiz, *Polyhedron* **2007**, *26*, 3842–3848.
- [26] L. S. Liebeskind, K. B. Sharpless, R. D. Wilson, J. A. Ibers, *J. Am. Chem. Soc.* **1978**, *100*, 7061–7063.
- [27] The results with nitroalkanes would suggest that no free nitroso is generated, because a primary or secondary nitroso alkane would rapidly suffer tautomerization to oxime making inaccessible the subsequent C–N amination. However, nitrosoarene compounds efficiently afforded the corresponding diarylamines when subjected to the standard reaction. So, regarding the reaction with nitroaromatics, the participation of free nitrosoarenes is feasible. See Supporting Information for further details.

Entry for the Table of Contents (Please choose one layout)

COMMUNICATION



Samuel Suárez-Pantiga,* Raquel Hernández-Ruiz, Cintia Virumbrales, María R. Pedrosa and Roberto Sanz*

Page No. – Page No.

Reductive Molybdenum-Catalyzed Direct Amination of Boronic Acids with Nitro Compounds

C–N Coupling: In the present work a new electrophilic amination of boronic acids with simple and inexpensive nitro compounds has been developed allowing the preparation of a variety of highly substituted and functionalized aromatic secondary amines. The reported process employs affordable PPh_3 as reducing agent, in the presence of an easily available and highly stable dioxomolybdenum(VI) complex as catalyst.