

Journal Pre-proof

Straightforward and direct access to β -seleno- amines and sulfonylamides *via* the controlled addition of phenylselenomethylithium (LiCH_2SePh) to imines

Raffaele Senatore, Monika Malik, Saad Touqeer, Roberta Listro, Simona Collina, Wolfgang Holzer, Vittorio Pace

PII: S0040-4020(20)30355-0

DOI: <https://doi.org/10.1016/j.tet.2020.131220>

Reference: TET 131220

To appear in: *Tetrahedron*

Received Date: 25 March 2020


Revised Date: 16 April 2020

Accepted Date: 17 April 2020

Please cite this article as: Senatore R, Malik M, Touqeer S, Listro R, Collina S, Holzer W, Pace V, Straightforward and direct access to β -seleno- amines and sulfonylamides *via* the controlled addition of phenylselenomethylithium (LiCH_2SePh) to imines, *Tetrahedron* (2020), doi: <https://doi.org/10.1016/j.tet.2020.131220>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier Ltd.



Straightforward and Direct Access to β -Seleno- Amines and Sulfonylamides *via* the Controlled Addition of Phenylselenomethylithium (LiCH_2SePh) to Imines

Raffaele Senatore,^a Monika Malik,^a Saad Touqeer,^a Roberta Listro,^b Simona Collina,^b Wolfgang Holzer^a and Vittorio Pace^{a,c*}

^a Department of Pharmaceutical Chemistry, University of Vienna, Althanstrasse 14, 1090 Vienna, Austria.


^b Department of Drug Sciences, University of Pavia, Via Taramelli 6, 27100 Pavia, Italy.

^c Department of Chemistry, University of Turin, Via P. Giuria 7, 10125 Turin, Italy.

*E-mail: vittorio.pace@univie.ac.at; vittorio.pace@unito.it – Web page: <https://drugsynthesis.univie.ac.at/>

- Twitter: @LabPace

Dedicated to Prof. Dr. Nuno Maulide for his outstanding contributions to the art of Synthesis.



Abstract. The transfer of a α -methyl phenylseleno carbanion to variously functionalized *N*-aryl and *N*-sulfonyl imines is reported. The fast selenium-lithium exchange conducted on a diselenoacetal with *n*-BuLi enables the generation of the attacking homologative nucleophile under chemoselective conditions preserving concomitant potentially sensitive functionalities to the lithiating conditions. Uniformly high yields were observed, thus establishing a valuable and conceptually simple approach to the title compounds.

Introduction.

Embodying selenium containing groups into an organic skeleton profoundly influences the physical-chemical features of the resulting scaffold, thus complementing the reach and well-established portfolio of (stereoselective) transformations these organometallics offer.¹ Indeed, there is a continuous and growing interest towards the biological properties displayed by organoselenium analogues whose structures are frequently encountered in enzymes governing important metabolic cascades (*e.g.* selenoprotein P, glutathione peroxidase, *inter alia*).² Additionally, recent uses of selenium-based pharmacologically active substances emerged as valuable tools for treating important diseases, as exemplified by ebselen, thus further increasing the potential of such a chalcogen-containing motifs.³ Evidently, the simultaneous derivatization of organoselenium species with additional functionalities expands the chemical space and the overall versatility of the so obtained materials. In this sense, including a nitrogen-centered group in a relevant position may represent an effective strategy for tuning pivotal molecular properties (*e.g.* solubility, cell permeability, metabolic parameters *etc.*) or, by focusing on pure synthetic aspects, it enables selective chemistries *en route* to more complex architectures.⁴ In fact, the proper activation of these β -seleno amine-type manifolds results in a versatile technique for assembling different aza-cycles such as aziridines⁵ or diastereomerically enriched 2,4-disubstituted pyrrolidines.⁶ This applicative significance merged with the potential exhibited in different areas including material science and drug design,⁷ motivated the development of variously oriented tactics for the construction of the motif. In recent years, the amino- or amido- selenation of olefins – *i.e.* the simultaneous installation into the sp^2 -hybridized C-C bond of both the

selenium atom and the nitrogenated moiety - emerged as an attractive and practical solution. Mechanistically, the electrophilic organoselenium agent generates a seleniranium ion susceptible of nucleophilic attack by the *N*-containing partner. The protocol manifests strong dependence on the nature and level of substitution of the competent olefin, as well as, on the nucleophilicity of the attacking amine or amide groups. As a consequence, the adequate modulation – *via* ionic,⁸ electrochemical⁹ or radical¹⁰ pathway - of these events becomes critical for the successful outcome of the targeted strategy. A logically distinct approach involves the ring opening of an aziridine with an *in situ* generated selenium anion.¹¹ It appears conceivable that using these highly valuable synthons may pose severe limitations in terms of costs. Alternatively, Outurquin proposed a two-steps methodology paved on forming highly unstable and difficult to purify α -selenoimines (from the corresponding ketones) followed by the controlled reduction to the desired compounds.^{5a,12}


Inspired by our work on homologation of carbon electrophiles with α -substituted methyllithiums (LiCHXY),¹³ in 2018 we reported a general approach for introducing substituted seleniomethyl fragments – *i.e.* RSeCH₂ – onto Weinreb amides.^{13k,14} The strategy paved on the generation of the carbanion from a diselenoacetal¹⁵ by adapting seminal Reich's studies,¹⁶ documented high versatility and chemoselectivity. We reasoned that delivering this Se-centered carbanion¹⁷ to a given imine would produce the β -seleno nitrogenated adduct through a conceptually intuitive and novel approach. To the best of our knowledge, the employment of C-C bond forming operations for producing the title compounds is still unprecedented and, herein we disclose our findings.

Relevance of Organoseleniums in Med Chem


**Ebselen**

Prototypal Se-containing drug


- Cytoprotective
- Anti-inflammatory
- Anti-oxidant

Versatility of β -Seleno Nitrogenated MotifsThe Assembly of β -Seleno Nitrogenated Motifs



a) Oxidative Difunctionalization of Olefins



b) Ring Opening of Aziridines

c) Construction-Reduction of α -SelenoiminesThe Direct Transfer of LiCH_2SePh to C-Electrophiles

Weinreb Amides (Pace, 2018)

**This work: Addition to Imines**


- One-pot process
- High yielding
- Chemoselective
- 19 Examples

Scheme 1. General context of the presented work.**Results and Discussion.**

We selected the naphthyl-substituted imine **1** as the model compound for accomplishing the formal homologation with the lithiated α -methyl seleno carbanion (LiCH_2SePh). As anticipated, a convenient source of this species was individuated in the diselenoacetal^{15a} ($\text{PhSeCH}_2\text{SePh}$) which undergoes an extremely fast selenium-lithium exchange upon treatment with *n*-BuLi. Evidently, the genesis of the requested carbanion is accompanied by the formation of the interconversion product – the selenoether *n*-BuSePh (**2b**) – which guarantees a quantitative estimation of the process. The optimization study revealed

the adaptability of the conditions established for preparing α -selenoketones starting from Weinreb amides:^{14a} PhSeCH₂SePh (1.3 equiv), *n*-BuLi (1.25 equiv) in Et₂O at -78 °C under non-Barbier type conditions (Table 1). Although the initial experiment showed the formation of the expected β -selenoamine **2** in a reasonably good 70% yield, a noticeable amount of the aminic compound **2a** generated by the direct attack of *n*-BuLi to the azomethinic carbon was detected (entry 1). With the aim of improving the yield of **2**, we briefly screened the effect of additional parameters: *i*) the simple switching to THF as the reaction medium did not alter significantly the **2:2a** ratio (entry 2); *ii*) decreasing the lithiation time was detrimental in agreement with our previous study on Weinreb amides (entries 3-4); *iii*) the use of alternative ethereal-type solvents such as 2-MeTHF¹⁸ and CPME¹⁹ significantly decreased the efficiency (entries 5-6); *iv*) pleasingly, the progressive increase of the loading of Se-carbanion enables to maximize the yield up to 93%, being practically undetectable the formation of the side product **2a** (entries 7-8); *v*) using a smaller excess of diselenoacetal (2.0 equiv) compared to the lithiating agent (1.90 equiv) guarantees the full generation of LiCH₂SePh and, thus to skip the concomitant, undesired attack of *n*-BuLi to **1**.

Table 1. Model Reaction: Optimization





Entry	Solvent	LiCH ₂ SePh (equiv)	Lithiation time (min)	Ratio 2:2a	Yield of 2 (%) ^a
1 ^b	Et ₂ O	1.25	60	76:24	70
2	THF	1.25	60	78:22	73
3	THF	1.25	45	75:25	68
4	THF	1.25	30	77:23	64
5	2-MeTHF	1.25	60	65:45	50
6	CPME	1.25	60	71:29	58
7	THF	1.60	60	93:7	84
8	THF	1.90	60	>99:1	93
9 ^c	THF	1.90	60	95:5	90

^a Isolated yield. ^b **2a** was isolated in 19% yield. ^c PhSeCH₂SePh (1.90 equiv) and *n*-BuLi (1.90 equiv) were used.


With the optimal conditions in hand, we then explored the scope of the transformation. A series of variously functionalized *N*-aryl imines easily undergoes the transformation giving the expected β -phenylseleno adducts in high yields. The substitution on the aromatic ring of the starting *N*-phenyl-type aldimines was perfectly tolerated: the presence of electron-donating functionalities such as methoxy at different positions did not alter the efficiency of the methodology. Analogously, potentially sensitive halogens – including the reactive bromo-derivatives (**5** and **8**) – could be introduced on the nucleus yielding the corresponding seleno-amines. The chemoselectivity of the approach is further illustrated in the case of the nitrile-bearing compound **10**, in which this electrophilic group was preserved by the nucleophilic attack. Switching to heteroaromatic aldimines was also possible, as documented by the 2-furyl (**12**) derivative. The nature of the *N*-imine group can be conveniently shifted to a sulfonyl-analogue in order to prepare β -seleno

sulfonyl imides (**13-19**). Interestingly, our approach constitutes an effective solution for accessing **14** whose previous synthesis through the ring-opening of an aziridine was affected by limited regiocontrol.¹¹ A comparable chemoselective profile was observed: halogen-containing groups – such as chlorine, fluoro, trifluoromethyl and bromo – maintain unaltered the productivity of the homologative technique. Similarly, poly- (naphtyl, **13**) or heteroaromatic (2-furyl, **18** – 2-thienyl, **19**) benzenesulfonylimines acted as competent starting materials without affecting the chemical integrity of these systems.


Journal Pre-proof

*N-Aryl Imines*


2 (93 %)




3 (88 %)




4 (83 %)




5 (85 %)




6 (87 %)




7 (92 %)




8 (86 %)




9 (88 %)




10 (92 %)




11 (89 %)




12 (86 %)

N-Sulfonyl Imines


13 (89 %)




14 (91 %)




15 (85 %)




16 (90 %)



17 (92 %)



18 (87 %)



19 (90 %)

Scheme 2. Scope of the protocol.

Conclusions.

In conclusion, we have developed a straightforward and versatile methodology for the transfer of a selenium-centered methyl lithium carbanion to variously functionalized imines. The selective lithiation of a diselenoacetal (PhSeCH₂SePh) smoothly provides the homologating agent (LiCH₂SePh) which can be easily released to the electrophilic azomethinic partners. The high degree of chemoselectivity, as well as the very good to excellent yields observed, make this conceptually intuitive strategy of wide and general applicability.

Experimental Part

Instrumentation and General Analytical Methods

Melting points were determined on a Reichert–Kofler hot-stage microscope and are uncorrected. Mass spectra were obtained on a Shimadzu QP 1000 instrument (EI, 70 eV) and on a Bruker maXis 4G instrument (ESI-TOF, HRMS). ¹H, ¹³C, ¹⁹F, ¹⁵N and ⁷⁷Se NMR spectra were recorded with a Bruker Avance III 400 spectrometer (400 MHz for ¹H, 100 MHz for ¹³C, 376 MHz for ¹⁹F, 40 MHz for ¹⁵N and 76 MHz for ⁷⁷Se) at 297 K using a directly detecting broadband observe (BBFO) probe. The center of the (residual) solvent signal was used as an internal standard which was related to TMS with δ 7.26 ppm (¹H in CDCl₃), δ 77.0 ppm (¹³C in CDCl₃). ¹⁹F NMR spectra were referenced via the Ξ ratio (absolute referencing). ¹⁵N NMR spectra (gs-HMBC, gs-HSQC) were referenced against neat, external nitromethane. ⁷⁷Se spectra were referenced against diphenyldiselenane (δ Ph₂Se₂ 463 ppm). Spin-spin coupling constants (*J*) are given in Hz.


In nearly all cases, full and unambiguous assignment of all resonances was performed by combined application of standard NMR techniques, such as APT, HSQC, HMBC, HSQCTOCSY, COSY and NOESY experiments.

THF was distilled over Na / benzophenone. Chemicals were purchased from SigmaAldrich, Acros, Alfa Aesar and TCI Europe. Solutions were evaporated under reduced pressure with a rotary evaporator. TLC was carried out on aluminium sheets precoated with silica gel 60F254 (Merchery-Nagel, Merk); the spots were visualised under UV light ($\lambda = 254$ nm) and/or KMnO₄ (aq.) was used as revealing system.

General Procedure

To a solution of 1,1'-(methylenediselanyl)dibenzene (2.0 equiv) in dry THF (2 mL) a 2.5 M solution of *n*-BuLi in hexanes (1.9 equiv) was added dropwise at -78 °C under argon and the reaction mixture was stirred at -78 °C for 1 hour. Then, a solution of the starting imine (1.0 equiv) in dry THF (1 mL) was added and the mixture was stirred at -78 °C for 2.5 hours before quenching with sat. aqueous NH₄Cl. The mixture was exhaustively extracted with EtOAc. The combined organic extracts were washed with brine, dried over Na₂SO₄ and the solvent was removed under reduced pressure. The product was obtained after purification by column chromatography on silica gel (*n*-hexane:EtOAc).

4-methyl-*N*-[1-(2-naphthyl)-2-(phenylselanyl)ethyl]aniline (2)



By Following the general procedure, starting from 1,1'-(methylenediselanyl)dibenzene (0.652 g, 2.0 mmol, 2.0 equiv) in dry THF (2 mL), *n*-BuLi (2.5 M, 0.76 mL, 1.9 mmol, 1.9 equiv) and 4-methyl-*N*-(naphthalen-2-ylmethylene)aniline (0.245 g, 1.0 mmol, 1.0 equiv) in dry THF (1 mL), compound **2** was obtained in 93 % yield (0.387 g) as colourless oil after purification by column chromatography on silica gel (*n*-hexane:EtOAc 99:1).


¹H NMR (400 MHz, CDCl₃): δ 7.81 (m, 1H, Naph H-1), 7.80 (m, 2H, Naph), 7.78 (m, 1H, Naph), 7.54 (m, 2H, Ph 2 H-2,6), 7.47 (m, 1H, Naph H-3), 7.46 (m, 1H, Naph), 7.45 (m, 1H, Naph), 7.27 (m, 1H, Ph 2 H-4), 7.26 (m, 2H, Ph 2 H-2,6), 6.87 (m, 2H, Ph 1 H-3,5), 6.43 (m, 2H, Ph 1 H-2,6), 4.59 (dd, ³*J* = 8.9 Hz, ³*J* = 4.6 Hz, 1H, CH), 4.5 (brs, 1H, NH), 3.44 (dd, ²*J* = 12.6 Hz, ³*J* = 4.6 Hz, 1H, CH₂), 3.28 (dd, ²*J* = 12.6 Hz, ³*J* = 8.9 Hz, 1H, CH₂), 2.18 (s, 3H, CH₃).

¹³C NMR (100 MHz, CDCl₃): δ 144.7 (Ph 1 C-1), 140.3 (Naph C-2), 133.55 (Ph 2 C-2,6), 133.51 (Naph 8a), 133.0 (Naph 4a), 129.6 (Ph 1 C-3,5), 129.3 (Ph 2 C-1), 129.2 (Ph 2 C-3,5), 128.6 (Naph), 127.9 (Naph), 127.7 (Naph), 127.5 (Ph 2 C-4), 127.1 (Ph 1 C-4), 126.1 (Naph), 125.7 (Naph), 125.2 (Naph C-1), 124.4 (Naph C-3), 114.0 (Ph 1 C-2,6), 58.4 (CH), 36.4 (CH₂), 20.3 (CH₃).

⁷⁷Se NMR (76 MHz, CDCl₃): δ 267.3 (m).

HRMS (ESI), *m/z* [M + H]⁺ calcd. for C₂₅H₂₄NSe: 418.1068; found: 418.1057.

4-methyl-*N*-[1-(2-naphthyl)pentyl]aniline (**2a**)




¹H NMR (400 MHz, CDCl₃): δ 7.82 (m, 3H, Naph H-4,5,8), 7.81 (s, 1H, Naph H-1), 7.50 (m, 1H, Naph H-3), 7.47 (m, 1H, Naph H-6), 7.45 (m, 1H, Naph H-7), 6.89 (m, 2H, Ph H-3,5), 6.50 (m, 2H, Ph H-2,6), 4.44 (m, 1H, CH), 3.9 – 4.4 (brs, 1H, NH), 2.18 (s, 3H, CH₃), 1.88 (m, 2H, CH₂CH₂CH₂CH₃), 1.37 (m, 2H, CH₂CH₂CH₂CH₃), 1.36 (m, 2H, CH₂CH₂CH₂CH₃), 0.90 (t, ³*J* = 7.0 Hz, 3H, CH₂CH₂CH₂CH₃).

¹³C NMR (100 MHz, CDCl₃): δ 145.2 (Ph C-1), 142.0 (Naph C-2), 133.5 (Naph C-8a), 132.8 (Naph C-4a), 129.6 (Ph C-3,5), 128.3 (Naph), 127.8 (Naph), 127.6 (Naph), 126.4 (Ph C-4), 125.9 (Naph C-6), 125.4 (Naph C-7), 125.1 (Naph C-1), 124.8 (Naph C-3), 113.5 (Ph C-2,6), 58.8 (CH), 38.6 (CH₂CH₂CH₂CH₃), 28.5 (CH₂CH₂CH₂CH₃), 22.6 (CH₂CH₂CH₂CH₃), 20.3 (CH₃), 14.0 (CH₂CH₂CH₂CH₃).

^{15}N NMR (40 MHz, CDCl_3): δ -305.4.

HRMS (ESI), m/z $[\text{M} + \text{H}]^+$ calcd. for $\text{C}_{22}\text{H}_{26}\text{N}$: 304.2060; found: 304.2057.

Butyl phenyl selenide (2b)^{14a}




^1H NMR (400 MHz, CDCl_3): δ 7.48 (m, 2H, Ph H-2,6), 7.25 (m, 2H, Ph H-3,5), 7.23 (m, 1H, Ph H-4), 2.92 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.69 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.43 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 0.91 (t, $^3J = 7.4$ Hz, 3H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$).

^{13}C NMR (100 MHz, CDCl_3): δ 132.3 (Ph C-2,6), 130.7 (Ph C-1), 129.0 (Ph C-3,5), 126.6 (Ph C-4), 32.2 ($\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 27.6 ($\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 22.9 ($\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 13.5 ($\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$).

^{77}Se NMR (76 MHz, CDCl_3): δ 290.5.

HRMS (ESI), m/z $[\text{M} + \text{H}]^+$ calcd. for $\text{C}_{10}\text{H}_{15}\text{Se}$: 215.0333; found: 215.0336.

***N*-[1-(4-methoxyphenyl)-2-(phenylselanyl)ethyl]-4-methylaniline (3)**



By Following the general procedure, starting from 1,1'-(methylenediselanyl)dibenzene (0.652 g, 2.0 mmol, 2.0 equiv) in dry THF (2 mL), *n*-BuLi (2.5 M, 0.76 mL, 1.9 mmol, 1.9 equiv) and *N*-(4-methoxybenzylidene)-4-methylaniline (0.225 g, 1.0 mmol, 1.0 equiv) in dry THF (1 mL), compound **3** was obtained in 88 % yield (0.348 g) as colourless oil after purification by column chromatography on silica gel (*n*-hexane:EtOAc 9:1).


^1H NMR (400 MHz, CDCl_3): δ 7.50 (m, 2H, Ph 3 H-2,6), 7.26 (m, 3H, Ph 3 H-3,4,5), 7.25 (m, 2H, Ph 1 H-2,6), 6.88 (m, 2H, Ph 2 H-3,5), 6.83 (m, 2H, Ph 1 H-3,5), 6.40 (m, 2H, Ph 2 H-2,6), 4.39 (dd, $^3J = 8.4$ Hz, $^3J = 5.1$ Hz, 1H, CH), 4.0-4.8 (brs, 1H, NH), 3.78 (s, 3H, OCH_3), 3.33 (dd, $^2J = 12.5$ Hz, $^3J = 5.1$ Hz, 1H, CH_2), 3.22 (m, 1H, CH_2), 2.18 (s, 3H, CH_3).

^{13}C NMR (100 MHz, CDCl_3): δ 158.9 (Ph 1 C-4), 144.9 (Ph 2 C-1), 134.6 (Ph 1 C-1), 133.4 (Ph 3 C-2,6), 129.5 (Ph 2 C-3,5), 129.4 (Ph 3 C-1), 129.2 (Ph 3 C-3,5), 127.5 (Ph 1 C-2,6), 127.4 (Ph 3 C-4), 127.2 (Ph 2 C-4), 114.1 (Ph 1 C-3,5, Ph 2 C-2,6), 57.7 (CH), 55.2 (OCH_3), 36.4 (CH_2), 20.4 (CH_3).

^{77}Se NMR (76 MHz, CDCl_3): δ 265.0 (m).

HRMS (ESI), m/z $[\text{M} + \text{H}]^+$ calcd. for $\text{C}_{22}\text{H}_{24}\text{NOSe}$: 398.1018; found: 398.1016.

***N*-[1-(2,4-dimethoxyphenyl)-2-(phenylselanyl)ethyl]-4-methylaniline (4)**



By Following the general procedure, starting from 1,1'-(methylenediselanyl)dibenzene (0.652 g, 2.0 mmol, 2.0 equiv) in dry THF (2 mL), *n*-BuLi (2.5 M, 0.76 mL, 1.9 mmol, 1.9 equiv) and *N*-(2,4-dimethoxybenzylidene)-4-methylaniline (0.255 g, 1.0 mmol, 1.0 equiv) in dry THF (1 mL), compound **4** was obtained in 83 % yield (0.354 g) as brown oil after purification by column chromatography on silica gel (*n*-hexane:EtOAc 99:1).

^1H NMR (400 MHz, CDCl_3): δ 7.51 (m, 2H, Ph 3 H-2,6), 7.24 (m, 3H, Ph 3 H-3,4,5), 7.21 (d, $^3J = 8.4$ Hz, 1H, Ph 1 H-6), 6.89 (m, 2H, Ph 2 H-3,5), 6.43 (d, $^4J = 2.4$ Hz, 1H, Ph 1 H-3), 6.39 (m, 3H, Ph 1 H-5, Ph 2 H-2,6), 4.78 (dd, $^3J = 8.3$ Hz, $^3J = 4.8$ Hz, 1H, CH), 4.38 (brs, 1H, NH), 3.82 (s, 3H, 2- OCH_3), 3.77 (s, 3H, 4- OCH_3), 3.45 (dd, $^2J = 12.4$ Hz, $^3J = 4.8$ Hz, 1H, CH_2), 3.22 (dd, $^2J = 12.4$ Hz, $^3J = 8.3$ Hz, 1H, CH_2), 2.19 (s, 3H, CH_3).


^{13}C NMR (100 MHz, CDCl_3): δ 160.0 (Ph 1 C-4), 157.6 (Ph 1 C-2), 144.8 (Ph 2 C-1), 132.9 (Ph 3 C-2,6), 130.2 (Ph 3 C-1), 129.5 (Ph 2 C-3,5), 128.9 (Ph 3 C-3,5), 128.0 (Ph 1 C-6), 126.9 (Ph 3 C-4), 126.6 (Ph 2 C-4), 122.2 (Ph 1 C-1), 113.8 (Ph 2 C-2,6), 104.1 (Ph 1 C-5), 98.7 (Ph 1 C-3), 55.3 (4- OCH_3), 55.2 (2- OCH_3), 53.0 (CH), 34.2 (CH_2), 20.3 (CH_3).

^{77}Se NMR (76 MHz, CDCl_3): δ 263.3 (m).

^{15}N NMR (40 MHz, CDCl_3): δ -308.0.

HRMS (ESI), m/z $[\text{M} + \text{H}]^+$ calcd. for $\text{C}_{23}\text{H}_{26}\text{NO}_2\text{Se}$: 428.1123; found: 428.1120.

4-bromo-*N*-[1-(4-methoxyphenyl)-2-(phenylselanyl)ethyl]aniline (5)



By Following the general procedure, starting from 1,1'-(methylenediselanyl)dibenzene (0.652 g, 2.0 mmol, 2.0 equiv) in dry THF (2 mL), *n*-BuLi (2.5 M, 0.76 mL, 1.9 mmol, 1.9 equiv) and 4-bromo-*N*-(4-methoxybenzylidene)aniline (0.290 g, 1.0 mmol, 1.0 equiv) in dry THF (1 mL), compound **5** was obtained in 85 % yield (0.392 g) as yellow oil after purification by column chromatography on silica gel (*n*-hexane:EtOAc 98:2).

$^1\text{H NMR}$ (400 MHz, CDCl_3): δ 7.51 (m, 2H, Ph 3 H-2,6), 7.28 (m, 1H, Ph 3 H-4), 7.27 (m, 2H, Ph 3 H-3,5), 7.22 (m, 2H, Ph 1 H-2,6), 7.13 (m, 2H, Ph 2 H-3,5), 6.84 (m, 2H, Ph 1 H-3,5), 6.30 (m, 2H, Ph 2 H-2,6), 4.34 (dd, $^3J = 8.8$ Hz, $^3J = 4.6$ Hz, 1H, CH), 3.8-5.0 (brs, 1H, NH), 3.78 (s, 3H, OCH_3), 3.32 (dd, $^2J = 12.6$ Hz, $^3J = 4.6$ Hz, 1H, CH_2), 3.15 (dd, $^2J = 12.6$ Hz, $^3J = 8.8$ Hz, 1H, CH_2).


$^{13}\text{C NMR}$ (100 MHz, CDCl_3): δ 159.1 (Ph 1 C-4), 146.0 (Ph 2 C-1), 134.0 (Ph 1 C-1), 133.6 (Ph3 C-2,6), 131.7 (Ph 2 C-3,5), 129.3 (Ph 3 C-3,5), 129.0 (Ph 3 C-1), 127.6 (Ph 3 C-4), 127.3 (Ph 1 C-2,6), 115.3 (Ph 2 C-2,6), 114.2 (Ph 1 C-3,5), 109.5 (Ph 2 C-4), 57.2 (CH), 55.2 (OCH_3), 36.4 (CH_2).

$^{77}\text{Se NMR}$ (76 MHz, CDCl_3): δ 265.9 (m).

$^{15}\text{N NMR}$ (40 MHz, CDCl_3): δ -303.9.

HRMS (ESI), m/z $[\text{M} + \text{Na}]^+$ calcd. for $\text{C}_{21}\text{H}_{20}\text{BrNNaOSe}$: 483.9786; found: 483.9785.

N-[1-(4-chlorophenyl)-2-(phenylselanyl)ethyl]-4-methylaniline (**6**)



By Following the general procedure, starting from 1,1'-(methylenediselanyl)dibenzene (0.652 g, 2.0 mmol, 2.0 equiv) in dry THF (2 mL), *n*-BuLi (2.5 M, 0.76 mL, 1.9 mmol, 1.9 equiv) and *N*-(4-chlorobenzylidene)-4-methylaniline (0.229 g, 1.0 mmol, 1.0 equiv) in dry THF (1 mL), compound **6** was obtained in 87 % yield (0.349 g) as brown oil after purification by column chromatography on silica gel (*n*-hexane:EtOAc 99:1).

¹H NMR (400 MHz, CDCl₃): δ 7.51 (m, 2H, Ph 3 H-2,6), 7.27 (m, 7H, Ph 1 H-2,3,5,6, Ph 3 H-3,4,5), 6.90 (m, 2H, Ph 2 H-3,5), 6.35 (m, 2H, Ph 2 H-2,6), 4.40 (dd, ³J = 8.8 Hz, ³J = 4.6 Hz, 1H, CH), 4.36 (brs, 1H, NH), 3.32 (dd, ²J = 12.6 Hz, ³J = 4.5 Hz, 1H, CH₂), 3.16 (dd, ²J = 12.6 Hz, ³J = 8.8 Hz, 1H, CH₂), 2.20 (s, 3H, CH₃).


¹³C NMR (100 MHz, CDCl₃): δ 144.4 (Ph 2 C-1), 141.3 (Ph 1 C-1), 133.6 (Ph 3 C-2,6), 133.0 (Ph 1 C-4), 129.6 (Ph 2 C-3,5), 129.3 (Ph 3 C-3,5), 129.0 (Ph 3 C-1), 128.9 (Ph 1 C-3,5), 127.7 (Ph 1 C-2,6), 127.6 (Ph 3 C-4), 127.3 (Ph 2 C-4), 113.9 (Ph 2 C-2,6), 57.4 (CH), 36.4 (CH₂), 20.3 (CH₃).

⁷⁷Se NMR (76 MHz, CDCl₃): δ 266.0 (m).

¹⁵N NMR (40 MHz, CDCl₃): δ -307.1.

HRMS (ESI), *m/z* [M + H]⁺ calcd. for C₂₁H₂₁ClNSe: 402.0522; found: 402.0523.

N-[1-(4-fluorophenyl)-2-(phenylselanyl)ethyl]-4-methylaniline (**7**)



By Following the general procedure, starting from 1,1'-(methylenediselanyl)dibenzene (0.652 g, 2.0 mmol, 2.0 equiv) in dry THF (2 mL), *n*-BuLi (2.5 M, 0.76 mL, 1.9 mmol, 1.9 equiv) and *N*-(4-fluorobenzylidene)-4-methylaniline (0.213 g, 1.0 mmol, 1.0 equiv) in dry THF (1 mL), compound **7** was obtained in 92 % yield (0.353 g) as yellow oil after purification by column chromatography on silica gel (*n*-hexane:EtOAc 99:1).

¹H NMR (400 MHz, CDCl₃): δ 7.50 (m, 2H, Ph 3 H-2,6), 7.30 (m, 2H, Ph 1 H-2,6), 7.27 (m, 1H, Ph 3 H-4), 7.26 (m, 2H, Ph 3 H-3,5), 6.98 (m, 2H, Ph 1 H-3,5), 6.89 (m, 2H, Ph 2 H-3,5), 6.36 (m, 2H, Ph 2 H-2,6), 4.41 (dd, ³J = 8.7 Hz, ³J = 4.8 Hz, 1H, CH), 4.35 (brs, 1H, NH), 3.32 (dd, ²J = 12.6 Hz, ³J = 4.8 Hz, 1H, CH₂), 3.18 (dd, ²J = 12.6 Hz, ³J = 8.7 Hz, 1H, CH₂), 2.19 (s, 3H, CH₃).


¹³C NMR (100 MHz, CDCl₃): δ 162.1 (d, ¹J_{C,F} = 245.5 Hz, Ph 1 C-4), 144.4 (Ph 2 C-1), 138.3 (d, ⁴J_{C,F} = 3.1 Hz, Ph 1 C-1), 133.5 (Ph 3 C-2,6), 129.6 (Ph 2 C-3,5), 129.3 (Ph 3 C-3,5), 129.1 (Ph 3 C-1), 127.9 (d, ³J_{C,F} = 8.0 Hz, Ph 1 C-2,6), 127.5 (Ph 3 C-4), 127.4 (Ph 2 C-4), 115.6 (d, ²J_{C,F} = 21.4 Hz, Ph 1 C-3,5), 114.0 (Ph 2 C-2,6), 57.5 (CH), 36.5 (CH₂), 20.3 (CH₃).

⁷⁷Se NMR (76 MHz, CDCl₃): δ 265.6 (m).

¹⁹F NMR (376 MHz, CDCl₃): δ -115.2 (m).

¹⁵N NMR (40 MHz, CDCl₃): δ -312.9.

HRMS (ESI), *m/z* [M + H]⁺ calcd. for C₂₁H₂₁FNSe: 386.0818; found: 386.0812.

***N*-[1-[4-bromophenyl]-2-(phenylselanyl)ethyl]-4-methylaniline (8)**


By Following the general procedure, starting from 1,1'-(methylenediselanyl)dibenzene (0.652 g, 2.0 mmol, 2.0 equiv) in dry THF (2 mL), *n*-BuLi (2.5 M, 0.76 mL, 1.9 mmol, 1.9 equiv) and *N*-(4-bromobenzylidene)-4-methylaniline (0.274 g, 1.0 mmol, 1.0 equiv) in dry THF (1 mL), compound **8** was obtained in 86 % yield (0.383 g) as yellowish oil after purification by column chromatography on silica gel (*n*-hexane:EtOAc 98:2).

¹H NMR (400 MHz, CDCl₃): δ 7.49 (m, 2H, Ph 3 H-2,6), 7.40 (m, 2H, Ph 1 H-3,5), 7.27 (m, 1H, Ph 3 H-4), 7.26 (m, 2H, Ph 3 H-3,5), 7.21 (m, 2H, Ph 1 H-2,6), 6.89 (m, 2H, Ph 2 H-3,5), 6.36 (m, 2H, Ph 2 H-2,6), 4.37 (dd, ³*J* = 8.6 Hz, ³*J* = 4.8 Hz, 1H, CH), 4.0-5.0 (brs, 1H, NH), 3.32 (dd, ²*J* = 12.6 Hz, ³*J* = 4.8 Hz, 1H, CH₂), 3.18 (dd, ²*J* = 12.6 Hz, ³*J* = 8.6 Hz, 1H, CH₂), 2.19 (s, 3H, CH₃).

¹³C NMR (100 MHz, CDCl₃): 144.0 (Ph 2 C-1), 141.6 (Ph 1 C-1), 133.6 (Ph 3 C-2,6), 131.8 (Ph 1 C-3,5), 129.6 (Ph 2 C-3,5), 129.3 (Ph 3 C-3,5), 129.0 (Ph 3 C-1), 128.2 (Ph 1 C-2,6), 127.6 (Ph 2 C-4, Ph 3 C-4), 121.2 (Ph 1 C-4), 114.2 (Ph 2 C-2,6), 57.8 (CH), 36.1 (CH₂), 20.4 (CH₃).

⁷⁷Se NMR (76 MHz, CDCl₃): δ 266.5 (m).

HRMS (ESI), *m/z* [M + H]⁺ calcd. for C₂₁H₂₁BrNSe: 446.0017; found: 446.0021.

***N*-[1-(4-chlorophenyl)-2-(phenylselanyl)ethyl]-4-fluoroaniline (9)**

By Following the general procedure, starting from 1,1'-(methylenediselanyl)dibenzene (0.652 g, 2.0 mmol, 2.0 equiv) in dry THF (2 mL), *n*-BuLi (2.5 M, 0.76 mL, 1.9 mmol, 1.9 equiv) and *N*-(4-chlorobenzylidene)-4-fluoroaniline (0.233 g, 1.0 mmol, 1.0 equiv) in dry THF (1 mL), compound **9** was obtained in 88 % yield (0.356 g) as yellowish oil after purification by column chromatography on silica gel (*n*-hexane:EtOAc 98:2).

¹H NMR (400 MHz, CDCl₃): δ 7.50 (m, 2H, Ph 3 H-2,6), 7.27 (m, 7H, Ph 1 H-2,3,5,6, Ph 3 H-3,4,5), 6.78 (m, 2H, Ph 2 H-3,5), 6.35 (m, 2H, Ph 2 H-2,6), 4.31 (dd, ³J = 8.9 Hz, ³J = 4.5 Hz, 1H, CH), 4.0-4.9 (brs, 1H, NH), 3.32 (dd, ²J = 12.7 Hz, ³J = 4.5 Hz, 1H, CH₂), 3.14 (dd, ²J = 12.7 Hz, ³J = 8.9 Hz, 1H, CH₂).

¹³C NMR (100 MHz, CDCl₃): δ 156.2 (d, ¹J_{C,F} = 235.9 Hz, Ph 2 C-4), 142.9 (Ph 2 C-1), 140.8 (Ph 1 C-1), 133.7 (Ph 3 C-2,6), 133.3 (Ph 1 C-4), 129.3 (Ph 3 C-3,5), 129.0 (Ph 1 C-3,5), 128.7 (Ph 3 C-1), 127.7 (Ph 1 C-2,6, Ph 3 C-4), 115.6 (d, ²J_{C,F} = 22.4 Hz, Ph 2 C-3,5), 114.8 (m, Ph 2 C-2,6), 58.0 (CH), 36.2 (CH₂).


⁷⁷Se NMR (76 MHz, CDCl₃): δ 266.6 (m).

¹⁹F NMR (376 MHz, CDCl₃): δ -127.0 (m).

¹⁵N NMR (40 MHz, CDCl₃): δ -307.9.

HRMS (ESI), *m/z* [M + H]⁺ calcd. for C₂₀H₁₈ClFNSe: 406.0272; found: 406.0270.

4-{1-[(4-methylphenyl)amino]-2-(phenylselanyl)ethyl}benzonitrile (**10**)



By Following the general procedure, starting from 1,1'-(methylenediselanyl)dibenzene (0.652 g, 2.0 mmol, 2.0 equiv) in dry THF (2 mL), *n*-BuLi (2.5 M, 0.76 mL, 1.9 mmol, 1.9 equiv) and 4-((*p*-tolylimino)methyl)benzonitrile (0.220 g, 1.0 mmol, 1.0 equiv) in dry THF (1 mL), compound **10** was obtained in 92 % yield (0.360 g) as yellowish oil after purification by column chromatography on silica gel (*n*-hexane:EtOAc 99:1).

¹H NMR (400 MHz, CDCl₃): δ 7.57 (m, 2H, Ph 1 H-2,6), 7.49 (m, 2H, Ph 3 H-2,6), 7.45 (m, 2H, Ph 1 H-3,5), 7.28 (m, 1H, Ph 3 H-4), 7.26 (m, 2H, Ph 3 H-3,5), 6.89 (m, 2H, Ph 2 H-3,5), 6.32 (m, 2H, Ph 2 H-2,6), 4.44 (dd, ³J = 8.7 Hz, ³J = 4.5 Hz, 1H, CH), 4.1-4.9 (brs, 1H, NH), 3.32 (dd, ²J = 12.8 Hz, ³J = 4.5 Hz, 1H, CH₂), 3.16 (dd, ²J = 12.8 Hz, ³J = 8.7 Hz, 1H, CH₂), 2.19 (s, 3H, CH₃).


¹³C NMR (100 MHz, CDCl₃): δ 148.3 (Ph 1 C-4), 143.9 (Ph 2 C-1), 133.7 (Ph 3 C-2,6), 132.6 (Ph 1 C-2,6), 129.7 (Ph 2 C-3,5), 129.3 (Ph 3 C-3,5), 128.6 (Ph 3 C-1), 127.8 (Ph 2 C-4, Ph 3 C-4), 127.3 (Ph 1 C-3,5), 118.7 (C≡N), 114.0 (Ph 2 C-2,6), 111.3 (Ph 1 C-1), 57.9 (CH), 35.9 (CH₂), 20.3 (CH₃).

⁷⁷Se NMR (76 MHz, CDCl₃): δ 267.8

¹⁵N NMR (40 MHz, CDCl₃): δ -308.3

HRMS (ESI), m/z $[M + H]^+$ calcd. for $C_{22}H_{21}N_2Se$: 393.0864; found: 393.0866.

***N*-[1-phenyl-2-(phenylselanyl)ethyl]aniline (11)**



By Following the general procedure, starting from 1,1'-(methylenediselanyl)dibenzene (0.652 g, 2.0 mmol, 2.0 equiv) in dry THF (2 mL), *n*-BuLi (2.5 M, 0.76 mL, 1.9 mmol, 1.9 equiv) and *N*-benzylideneaniline (0.181 g, 1.0 mmol, 1.0 equiv) in dry THF (1 mL), compound **11** was obtained in 89 % yield (0.313 g) as yellow solid after purification by column chromatography on silica gel (*n*-hexane:EtOAc 99:1).

mp: 125-127 °C

1H NMR (400 MHz, $CDCl_3$): δ 7.52 (m, 2H, Ph 3 H-2,6), 7.35 (m, 2H, Ph 1 H-2,6), 7.31 (m, 2H, Ph 1 H-3,5), 7.27 (m, 3H, Ph 3 H-3,4,5), 7.25 (m, 1H, Ph 1 H-4), 7.07 (m, 2H, Ph 2 H-3,5), 6.67 (m, 1H, Ph 2 H-4), 6.45 (m, 2H, Ph 2 H-2,6), 4.46 (dd, $^3J = 8.8$ Hz, $^3J = 4.7$ Hz, 1H, CH), 4.3-4.7 (brs, 1H, NH), 3.37 (dd, $^2J = 12.6$ Hz, $^3J = 4.7$ Hz, 1H, CH_2), 3.22 (dd, $^2J = 12.6$ Hz, $^3J = 8.8$ Hz, 1H, CH_2).


^{13}C NMR (100 MHz, $CDCl_3$): δ 147.0 (Ph 2 C-1), 142.6 (Ph 1 C-1), 133.5 (Ph 3 C-2,6), 129.3 (Ph 3 C-1), 129.2 (Ph 3 C-3,5), 129.0 (Ph 2 C-3,5), 128.8 (Ph 1 C-3,5), 127.53 (Ph 3 C-4), 127.49 (Ph 1 C-4), 126.3 (Ph 1 C-2,6), 117.8 (Ph 2 C-4), 113.8 (Ph 2 C-2,6), 57.9 (CH), 36.4 (CH_2).

^{77}Se NMR (76 MHz, $CDCl_3$): δ 266.7 (m)

^{15}N NMR (40 MHz, $CDCl_3$): δ -305.1.

HRMS (ESI), m/z $[M + H]^+$ calcd. for $C_{20}H_{20}NSe$: 354.0755; found: 354.0751.

***N*-[1-(2-furyl)-2-(phenylselanyl)ethyl]-4-methylaniline (12)**



By Following the general procedure, starting from 1,1'-(methylenediselanyl)dibenzene (0.652 g, 2.0 mmol, 2.0 equiv) in dry THF (2 mL), *n*-BuLi (2.5 M, 0.76 mL, 1.9 mmol, 1.9 equiv) and *N*-(furan-2-ylmethylene)-4-methylaniline (0.185 g, 1.0 mmol, 1.0 equiv) in dry THF (1 mL), compound **12** was obtained in 86 % yield (0.306 g) as yellow oil after purification by column chromatography on silica gel (*n*-hexane:EtOAc 99:1).


¹H NMR (400 MHz, CDCl₃): δ 7.49 (m, 2H, Ph 2 H-2,6), 7.30 (dd, ³J = 1.8 Hz, ⁴J = 0.9 Hz, 1H, Furyl H-5), 7.25 (m, 3H, Ph 2 H-3,4,5), 6.93 (m, 2H, Ph 1 H-3,5), 6.45 (m, 2H, Ph 1 H-2,6), 6.27 (dd, ³J = 3.2 Hz, ³J = 1.8 Hz, 1H, Furyl H-4), 6.19 (m, 1H, Furyl H-3), 4.68 (m, 1H, CH), 4.03-4.30 (brs, 1H, NH), 3.40 (dd, ²J = 12.5 Hz, ³J = 6.5 Hz, 1H, CH₂), 3.36 (dd, ²J = 12.5 Hz, ³J = 6.3 Hz, 1H, CH₂), 2.21 (s, 3H, CH₃).

¹³C NMR (100 MHz, CDCl₃): δ 154.6 (Furyl C-2), 144.1 (Ph 1 C-1), 141.8 (Furyl C-5), 133.4 (Ph 2 C-2,6), 129.7 (Ph 1 C-3,5), 129.5 (Ph 2 C-1), 129.1 (Ph 2 C-3,5), 127.6 (Ph 1 C-4), 127.3 (Ph 2 C-4), 114.0 (Ph 1 C-2,6), 110.3 (Furyl C-4), 106.8 (Furyl C-3), 52.4 (CH), 32.9 (CH₂), 20.4 (CH₃).

⁷⁷Se NMR (76 MHz, CDCl₃): δ 262.2 (m).

HRMS (ESI), *m/z* [M + H]⁺ calcd. for C₁₉H₂₀NOSe: 358.0705; found: 358.0707.

N-(1-(naphthalen-2-yl)-2-(phenylselanyl)ethyl)benzenesulfonamide (**13**)



By Following the general procedure, starting from 1,1'-(methylenediselanyl)dibenzene (0.652 g, 2.0 mmol, 2.0 equiv) in dry THF (2 mL), *n*-BuLi (2.5 M, 0.76 mL, 1.9 mmol, 1.9 equiv) and *N*-(naphthalen-2-ylmethylene)benzenesulfonamide (0.295 g, 1.0 mmol, 1.0 equiv) in dry THF (1 mL), compound **13** was obtained in 89 % yield (0.414 g) as colourless oil after purification by column chromatography on silica gel (*n*-hexane:EtOAc 9:1).

¹H NMR (400 MHz, CDCl₃): δ 7.74 (m, 1H, Naph H-5), 7.67 (m, 1H, Naph H-8), 7.63 (m, 1H, Naph H-4), 7.61 (m, 2H, Ph 1 H-2,6), 7.46 (m, 1H, Naph H-1), 7.45 (m, 2H, Naph H-6, Naph H-7), 7.38 (m, 1H, Ph 1 H-4), 7.37 (m, 2H, Ph 2 H-2,6), 7.24 (m, 3H, Ph 1 H-3,5, Ph 2 H-4), 7.22 (m, 2H, Ph 2 H-3,5), 7.14 (m, 1H, Naph H-3), 5.43 (d, ³J = 5.5 Hz, 1H, NH), 4.58 (m, 1H, CH), 3.32 (dd, ²J = 12.8 Hz, ³J = 7.3 Hz, 1H, CH₂), 3.25 (dd, ²J = 12.8 Hz, ³J = 6.4 Hz, 1H, CH₂).


¹³C NMR (100 MHz, CDCl₃): δ 139.9 (Ph1 C-1), 136.4 (Naph C-2), 133.3 (Ph 2 C-2,6), 132.96 (Naph C-8a), 132.94 (Naph C-4a), 132.4 (Ph 1 C-4), 129.3 (Ph 2 C-3,5), 128.7 (Ph 1 C-3,5), 128.52 (Naph C-4), 128.51 (Ph 2 C-1), 127.9 (Naph C-8), 127.59 (Ph 2 C-4), 127.77 (Naph C-5), 127.1 (Ph 1 C-2,6), 126.3 (Naph C-7), 126.2 (Naph C-6), 126.1 (Naph C-1), 124.0 (Naph C-3), 57.3 (CH), 35.0 (CH₂).

⁷⁷Se NMR (76 MHz, CDCl₃): δ 257.8 (m).

^{15}N NMR (40 MHz, CDCl_3): δ -271.7.

HRMS (ESI), m/z $[\text{M} + \text{H}]^+$ calcd. for $\text{C}_{24}\text{H}_{22}\text{NO}_2\text{SSe}$: 468.0531; found: 468.0532.

4-methyl-*N*-[1-phenyl-2-(phenylselanyl)ethyl]benzenesulfonamide (14)¹¹



By Following the general procedure, starting from 1,1'-(methylenediselanyl)dibenzene (0.652 g, 2.0 mmol, 2.0 equiv) in dry THF (2 mL), *n*-BuLi (2.5 M, 0.76 mL, 1.9 mmol, 1.9 equiv) and *N*-benzylidene-4-methylbenzenesulfonamide (0.261 g, 1.0 mmol, 1.0 equiv) in dry THF (1 mL), compound **14** was obtained in 91 % yield (0.391 g) as white solid after purification by column chromatography on silica gel (*n*-hexane:EtOAc 99:1).

mp: 92-93 °C (lit. 92-93 °C).

^1H NMR (400 MHz, CDCl_3): δ 7.51 (m, 2H, Ph 2 H-2,6), 7.34 (m, 2H, Ph 3 H-2,6), 7.26 (m, 1H, Ph 3 H-4), 7.22 (m, 2H, Ph 3 H-3,5), 7.20 (m, 2H, Ph 1 H-3,5), 7.19 (m, 1H, Ph 1 H-4), 7.12 (m, 2H, Ph 2 H-3,5), 7.06 (m, 2H, Ph 1 H-2,6), 5.27 (m, 1H, NH), 4.37 (m, 1H, CH), 3.24 (dd, $^2J = 12.8\text{Hz}$, $^3J = 7.2\text{Hz}$, 1H, CH_2), 3.16 (dd, $^2J = 12.8\text{Hz}$, $^3J = 6.4\text{Hz}$, 1H, CH_2), 2.37 (s, 3H, CH_3).


^{13}C NMR (100 MHz, CDCl_3): δ 143.2 (Ph 2 C-4), 139.5 (Ph 1 C-1), 136.9 (Ph 2 C-1), 133.2 (Ph 3 C-2,6), 129.4 (Ph 2 C-3,5), 129.2 (Ph 3 C-3,5), 128.7 (Ph 3 C-1), 128.5 (Ph 1 C-3,5), 128.0 (Ph 1 C-4), 127.5 (Ph 3 C-4), 127.2 (Ph 2 C-2,6), 126.6 (Ph 1 C-2,6), 57.1 (CH), 35.1 (CH_2), 21.5 (CH_3).

^{77}Se NMR (76 MHz, CDCl_3): δ 256.8 (m)

^{15}N NMR (40 MHz, CDCl_3): δ -271.4.

HRMS (ESI), m/z $[\text{M} + \text{Na}]^+$ calcd. for $\text{C}_{21}\text{H}_{21}\text{NNaO}_2\text{SSe}$: 454.0350; found: 454.0342.

***N*-{1-[4-chloro-3-(trifluoromethyl)phenyl]-2-(phenylselanyl)ethyl} benzenesulfonamide (15)**



By Following the general procedure, starting from 1,1'-(methylenediselanyl)dibenzene (0.652 g, 2.0 mmol, 2.0 equiv) in dry THF (2 mL), *n*-BuLi (2.5 M, 0.76 mL, 1.9 mmol, 1.9 equiv) and *N*-(4-chloro-3-(trifluoromethyl)benzylidene)benzenesulfonamide (0.347 g, 1.0 mmol, 1.0 equiv) in dry THF (1 mL), compound **15** was obtained in 85 % yield (0.440 g) as yellow solid after purification by column chromatography on silica gel (*n*-hexane:EtOAc 99:1).

mp: 125-126 °C

¹H NMR (400 MHz, CDCl₃): δ 7.59 (m, 2H, Ph 2 H-2,6), 7.50 (m, 1H, Ph 2 H-4), 7.35 (m, 2H, Ph 2 H-3,5), 7.32 (m, 2H, Ph 3 H-2,6), 7.28 (m, 1H, Ph 3 H-4), 7.25 (m, 1H, Ph 1 H-5), 7.23 (m, 1H, Ph 1 H-2), 7.22 (m, 2H, Ph 3 H-3,5), 7.18 (m, 1H, Ph 1 H-6), 5.43 (d, ³J = 5.0 Hz, 1H, NH), 4.45 (m, 1H, CH), 3.11 (d, ³J = 6.6 Hz, 2H, CH₂).


¹³C NMR (100 MHz, CDCl₃): δ 139.7 (Ph 2 C-1), 138.5 (Ph 1 C-1), 133.7 (Ph 3 C-2,6), 132.9 (Ph 2 C-4), 131.8 (q, ³J_{C,F} = 1.9 Hz, Ph 1 C-4), 131.4 (Ph 1 C-5), 131.3 (Ph 1 C-6), 129.4 (Ph 3 C-3,5), 128.9 (Ph 2 C-3,5), 128.3 (q, ²J_{C,F} = 31.5 Hz, Ph 1 C-3), 128.1 (Ph 3 C-4), 127.6 (Ph 3 C-1), 127.1 (Ph 2 C-2,6), 125.9 (q, ³J_{C,F} = 5.4 Hz, Ph 1 C-2), 122.4 (q, ¹J_{C,F} = 273.7 Hz, CF₃), 56.2 (CH), 34.9 (CH₂).

¹⁹F NMR (376 MHz, CDCl₃): δ -62.7 (s, CF₃).

¹⁵N NMR (40 MHz, CDCl₃): δ -272.8.

HRMS (ESI), *m/z* [M + Na]⁺ calcd. for C₂₁H₁₇ClF₃NNaO₂SSe: 541.9678; found: 541.9691.

N-[1-(4-bromophenyl)-2-(phenylselanyl)ethyl]benzenesulfonamide (**16**)



By Following the general procedure, starting from 1,1'-(methylenediselanyl)dibenzene (0.652 g, 2.0 mmol, 2.0 equiv) in dry THF (2 mL), *n*-BuLi (2.5 M, 0.76 mL, 1.9 mmol, 1.9 equiv) and *N*-(4-bromobenzylidene)benzenesulfonamide (0.324 g, 1.0 mmol, 1.0 equiv) in dry THF (1 mL), compound **16** was obtained in 90 % yield (0.445 g) as white solid after purification by column chromatography on silica gel (*n*-hexane:EtOAc 99:1).

mp: 124-125 °C

¹H NMR (400 MHz, CDCl₃): δ 7.61 (m, 2H, Ph 2 H-2,6), 7.50 (m, 1H, Ph 2 H-4), 7.35 (m, 2H, Ph 2 H-3,5), 7.33 (m, 2H, Ph 3 H-2,6), 7.27 (m, 1H, Ph 3 H-4), 7.27 (m, 2H, Ph 1 H-3,5), 7.22 (m, 2H, Ph 3 H-3,5), 6.91 (m, 2H, Ph 1 H-2,6), 5.37 (d, ³J = 5.5 Hz, 1H, NH), 4.35 (m, 1H, CH), 3.16 (dd, ²J = 12.9 Hz, ³J = 7.4 Hz, 1H, CH₂), 3.11 (dd, ²J = 12.9 Hz, ³J = 6.2 Hz, 1H, CH₂).


¹³C NMR (100 MHz, CDCl₃): δ 139.8 (Ph 2 C-1), 138.4 (Ph 1 C-1), 133.4 (Ph 3 C-2,6), 132.6 (Ph 2 C-4), 131.6 (Ph 1 C-3,5), 129.4 (Ph 3 C-3,5), 128.9 (Ph 2 C-3,5), 128.4 (Ph 1 C-2,6), 128.2 (Ph 3 C-1), 127.8 (Ph 3 C-4), 127.1 (Ph 2 C-2,6), 122.0 (Ph 1 C-4), 56.6 (CH), 34.9 (CH₂).

⁷⁷Se NMR (76 MHz, CDCl₃): δ 257.9 (m).

¹⁵N NMR (40 MHz, CDCl₃): δ -272.2.

HRMS (ESI), *m/z* [M + H]⁺ calcd. for C₂₀H₁₉BrNO₂SSe: 495.9480; found: 495.9476.

***N*-[1-(4-fluorophenyl)-2-(phenylselanyl)ethyl]benzenesulfonamide (17)**



By Following the general procedure, starting from 1,1'-(methylenediselanyl)dibenzene (0.652 g, 2.0 mmol, 2.0 equiv) in dry THF (2 mL), *n*-BuLi (2.5 M, 0.76 mL, 1.9 mmol, 1.9 equiv) and *N*-(4-fluorobenzylidene)benzenesulfonamide (0.263 g, 1.0 mmol, 1.0 equiv) in dry THF (1 mL), compound **17** was obtained in 92 % yield (0.399 g) as yellowish solid after purification by column chromatography on silica gel (*n*-hexane:EtOAc 98:2).

mp: 99-102 °C

¹H NMR (400 MHz, CDCl₃): δ 7.61 (m, 2H, Ph 2 H-2,6), 7.48 (m, 1H, Ph 2 H-4), 7.35 (m, 2H, Ph 3 H-2,6), 7.34 (m, 2H, Ph 2 H-3,5), 7.27 (m, 1H, Ph 3 H-4), 7.23 (m, 2H, Ph 3 H-3,5), 7.00 (m, 2H, Ph 1 H-2,6), 6.83 (m, 2H, Ph 1 H-3,5), 5.38 (d, 1H, ³J = 5.4 Hz, 1H, NH), 4.38 (m, 1H, CH), 3.18 (dd, ²J = 12.9 Hz, ³J = 7.4 Hz, 1H, CH₂), 3.12 (dd, ²J = 12.9 Hz, ³J = 6.3 Hz, 1H, CH₂).

¹³C NMR (100 MHz, CDCl₃): δ 162.3 (d, ¹J_{C,F} = 247.0 Hz, Ph 1 C-4), 139.9 (Ph 2 C-1), 135.1 (d, ⁴J_{C,F} = 3.2 Hz, Ph 1 C-1), 133.4 (Ph 3 C-2,6), 132.6 (Ph 2 C-4), 129.3 (Ph 3 C-3,5), 128.8 (Ph 2 C-3,5), 128.4 (d, ³J_{C,F} = 8.3 Hz, Ph 1 C-2,6), 128.3 (Ph 3 C-1), 127.7 (Ph 3 C-4), 127.1 (Ph 2 C-2,6), 115.4 (d, ²J_{C,F} = 21.6 Hz, Ph 1 C-3,5), 56.5 (CH), 35.1 (CH₂).


⁷⁷Se NMR (76 MHz, CDCl₃): δ 258.0 (m).

^{19}F NMR (376 MHz, CDCl_3): δ -113.9 (m).

^{15}N NMR (40 MHz, CDCl_3): δ -271.6.

HRMS (ESI), m/z $[\text{M} + \text{Na}]^+$ calcd. for $\text{C}_{20}\text{H}_{18}\text{FNNaO}_2\text{SSe}$: 458.0100; found: 458.0108.

***N*-[1-(2-furyl)-2-(phenylselanyl)ethyl]benzenesulfonamide (18)**



By Following the general procedure, starting from 1,1'-(methylenediselanyl)dibenzene (0.652 g, 2.0 mmol, 2.0 equiv) in dry THF (2 mL), *n*-BuLi (2.5 M, 0.76 mL, 1.9 mmol, 1.9 equiv) and *N*-(furan-2-ylmethylene)benzenesulfonamide (0.235 g, 1.0 mmol, 1.0 equiv) in dry THF (1 mL), compound **18** was obtained in 87 % yield (0.353 g) as yellowish oil after purification by column chromatography on silica gel (*n*-hexane:EtOAc 9:1).

^1H NMR (400 MHz, CDCl_3): δ 7.70 (m, 2H, Ph 1 H-2,6), 7.50 (m, 1H, Ph 1 H-4), 7.39 (m, 2H, Ph 1 H-3,5), 7.39 (m, 2H, Ph2 H-2,6), 7.25 (m, 1H, Ph 2 H-4), 7.24 (m, 2H, Ph 2 H-3,5), 7.11 (dd, $^3J = 1.8$ Hz, $^4J = 0.8$ Hz, 1H, Furyl H-5), 6.12 (dd, $^3J = 3.3$ Hz, $^3J = 1.8$ Hz, 1H, Furyl H-4), 5.99 (m, 1H, Furyl H-3), 5.30 (d, $^3J = 8.0$ Hz, 1H, NH), 4.63 (m, 1H, CH), 3.36 (dd, $^2J = 12.7$ Hz, $^3J = 5.7$ Hz, 1H, CH_2), 3.17 (dd, $^2J = 12.7$ Hz, $^3J = 7.3$ Hz, 1H, CH_2).


^{13}C NMR (100 MHz, CDCl_3): δ 151.2 (Furyl C-2), 142.2 (Furyl C-5), 140.2 (Ph 1 C-1), 133.2 (Ph 2 C-2,6), 132.5 (Ph 1 C-4), 129.2 (Ph 2 C-3,5), 128.9 (Ph 1 C-3,5), 128.8 (Ph 2 C-1), 127.4 (Ph 2 C-4), 127.0 (Ph 1 C-2,6), 110.2 (Furyl C-4), 108.0 (Furyl C-3), 51.4 (CH), 32.5 (CH_2).

^{77}Se NMR (76 MHz, CDCl_3): δ 260.3 (m).

^{15}N NMR (40 MHz, CDCl_3): δ -274.5.

HRMS (ESI), m/z $[\text{M} + \text{Na}]^+$ calcd. for $\text{C}_{18}\text{H}_{17}\text{NNaO}_3\text{SSe}$: 429.9987; found: 429.9987.

***N*-[2-(phenylselanyl)-1-(2-thienyl)ethyl]benzenesulfonamide (19)**



By Following the general procedure, starting from 1,1'-(methylenediselanyl)dibenzene (0.652 g, 2.0 mmol, 2.0 equiv) in dry THF (2 mL), *n*-BuLi (2.5 M, 0.76 mL, 1.9 mmol, 1.9 equiv) and *N*-(thiophen-2-ylmethylene)benzenesulfonamide (0.251 g, 1.0 mmol, 1.0 equiv) in dry THF (1 mL), compound **19** was obtained in 90 % yield (0.379 g) as yellowish oil after purification by column chromatography on silica gel (*n*-hexane:EtOAc 9:1).

¹H NMR (400 MHz, CDCl₃): δ 7.68 (m, 2H, Ph 1 H-2,6), 7.50 (m, 1H, Ph 1 H-4), 7.42 (m, 2H, Ph 2 H-2,6), 7.37 (m, 2H, Ph 1 H-3,5), 7.27 (m, 1H, Ph 2 H-4), 7.26 (m, 2H, Ph 2 H-3,5), 7.12 (dd, ³J = 5.1 Hz, ⁴J = 1.2 Hz, 1H, Th H-5), 6.81 (dd, ³J = 5.1 Hz, ³J = 3.5 Hz, 1H, Th H-4), 6.74 (m, 1H, Th H-3), 5.31 (d, ³J = 6.8 Hz, 1H, NH), 4.79 (m, 1H, CH), 3.40 (dd, ²J = 12.8 Hz, ³J = 6.0 Hz, 1H, CH₂), 3.20 (dd, ²J = 12.8 Hz, ³J = 6.7 Hz, 1H, CH₂).

¹³C NMR (100 MHz, CDCl₃): δ 143.1 (Th C-2), 140.0 (Ph 1 C-1), 133.3 (Ph 2 C-2,6), 132.6 (Ph 1 C-4), 129.3 (Ph 2 C-3,5), 128.9 (Ph 1 C-3,5), 128.8 (Ph 2 C-1), 127.6 (Ph 2 C-4), 127.1 (Ph 1 C-2,6), 126.7 (Th C-4), 125.5 (Th C-3), 125.4 (Th C-5), 53.1 (CH), 35.7 (CH₂).

⁷⁷Se NMR (76 MHz, CDCl₃): δ 259.0 (m).

¹⁵N NMR (40 MHz, CDCl₃): δ -268.8.

HRMS (ESI), *m/z* [M + H]⁺ calcd. for C₁₈H₁₈NO₂S₂Se: 423.9939; found: 423.9936.

Acknowledgments. The Authors are grateful to the University of Vienna for generous support. S. T. thanks the OEAD for a Pakistan-Austria doctoral grant. R. L. thanks the University of Pavia for a visiting PhD fellowship.

References

- (1) For authoritative references, see: (a) *Organoselenium Chemistry: Modern Developments in Organic Synthesis in Top. Curr. Chem.*; Wirth, T., Ed.; Springer: Berlin, Heidelberg, 2000; Vol. 208. (b) Freudendahl, D. M.; Santoro, S.; Shahzad, S. A.; Santi, C.; Wirth, T. *Angew. Chem. Int. Ed.* **2009**, *48*, 8409-8411. (c) Freudendahl, D. M.; Shahzad, S. A.; Wirth, T. *Eur. J. Org. Chem.* **2009**, *2009*, 1649-1664. (d) Wirth, T. *Organoselenium Chemistry: Synthesis and Reactions*; Wiley-VCH: Weinheim, 2011. (e) Mukherjee, A. J.; Zade, S. S.; Singh, H. B.; Sunoj, R. B. *Chem. Rev.* **2010**, *110*, 4357-4416. (f) Paulmier, C. *Selenium Reagents and Intermediates in Organic Synthesis*; Pergamon: Oxford, 1986.
- (2) For excellent reviews, see: (a) Weekley, C. M.; Harris, H. H. *Chem. Soc. Rev.* **2013**, *42*, 8870-8894. (b) Nogueira, C. W.; Zeni, G.; Rocha, J. B. T. *Chem. Rev.* **2004**, *104*, 6255-6286.
- (3) (a) Sarma, B. K.; Mugesh, G. *J. Am. Chem. Soc.* **2005**, *127*, 11477-11485. (b) Mugesh, G.; du Mont, W.-W.; Sies, H. *Chem. Rev.* **2001**, *101*, 2125-2180. (c) Bhuyan, B. J.; Mugesh, G. Biological and Biochemical Aspects of Selenium Compounds. In *Organoselenium Chemistry*; Wiley-VCH Verlag GmbH & Co. KGaA: 2011; pp. 361-396. (d) Sands, K. N.; Back, T. G. *Tetrahedron* **2018**, *74*, 4959-4967.
- (4) (a) Mailyan, A. K.; Eickhoff, J. A.; Minakova, A. S.; Gu, Z.; Lu, P.; Zakarian, A. *Chem. Rev.* **2016**, *116*, 4441-4557. (b) Bariwal, J.; Van der Eycken, E. *Chem. Soc. Rev.* **2013**, *42*, 9283-9303.
- (5) (a) Miniejew, C.; Outurquin, F.; Pannecoucke, X. *Org. Biomol. Chem.* **2004**, *2*, 1575-1576. (b) Ward, V. R.; Cooper, M. A.; Ward, A. D. *J. Chem. Soc., Perkin Trans. 1* **2001**, 944-945. (c) Demarcus, M.; Filigheddu, S. N.; Mann, A.; Taddei, M. *Tetrahedron Lett.* **1999**, *40*, 4417-4420.
- (6) (a) Besev, M.; Engman, L. *Org. Lett.* **2000**, *2*, 1589-1592. (b) Shanks, D.; Berlin, S.; Beşev, M.; Ottosson, H.; Engman, L. *J. Org. Chem.* **2004**, *69*, 1487-1491.

- (7) For comprehensive overviews, see: (a) Manjare, S. T.; Kim, Y.; Churchill, D. G. *Acc. Chem. Res.* **2014**, *47*, 2985-2998. (b) *Selenium and Tellurium Chemistry. From Small Molecules to Biomolecules and Materials*; Woollins, J. D.; Laitinen, R., Eds.; Springer-Verlag: Berlin, 2011.
- (8) Tang, E.; Wang, W.; Zhao, Y.; Zhang, M.; Dai, X. *Org. Lett.* **2016**, *18*, 176-179.
- (9) Sun, L.; Yuan, Y.; Yao, M.; Wang, H.; Wang, D.; Gao, M.; Chen, Y.-H.; Lei, A. *Org. Lett.* **2019**, *21*, 1297-1300.
- (10) (a) Sun, K.; Wang, X.; Lv, Y.; Li, G.; Jiao, H.; Dai, C.; Li, Y.; Zhang, C.; Liu, L. *Chem. Commun.* **2016**, *52*, 8471-8474. (b) Liu, Y.; Li, C.; Mu, S.; Li, Y.; Feng, R.; Sun, K. *Asian J. Org. Chem.* **2018**, *7*, 720-723.
- (11) Ganesh, V.; Chandrasekaran, S. *Synthesis* **2009**, 3267-3278.
- (12) Miniejew, C.; Outurquin, F.; Pannecoucke, X. *Tetrahedron* **2005**, *61*, 447-456.
- (13) For precedents from our group, see: (a) Castoldi, L.; Monticelli, S.; Senatore, R.; Ielo, L.; Pace, V. *Chem. Commun.* **2018**, *54*, 6692-6704. (b) Pace, V. *Aust. J. Chem.* **2014**, *67*, 311-313. (c) Pace, V.; Holzer, W.; De Kimpe, N. *Chem. Rec.* **2016**, *16*, 2061-2076. (d) Pace, V.; Castoldi, L.; Monticelli, S.; Rui, M.; Collina, S. *Synlett* **2017**, *28*, 879-888. (e) Monticelli, S.; Urban, E.; Langer, T.; Holzer, W.; Pace, V. *Adv. Synth. Catal.* **2019**, *361*, 1001-1006. (f) Pace, V.; Castoldi, L.; Mamuye, A. D.; Langer, T.; Holzer, W. *Adv. Synth. Catal.* **2016**, *358*, 172-177. (g) Ielo, L.; Touqeer, S.; Roller, A.; Langer, T.; Holzer, W.; Pace, V. *Angew. Chem. Int. Ed.* **2019**, *58*, 2479-2484. (h) Pace, V.; Castoldi, L.; Mazzeo, E.; Rui, M.; Langer, T.; Holzer, W. *Angew. Chem. Int. Ed.* **2017**, *56*, 12677-12682. (i) Pace, V.; Castoldi, L.; Monticelli, S.; Safranek, S.; Roller, A.; Langer, T.; Holzer, W. *Chem. Eur. J.* **2015**, *21*, 18966-18970. (j) Castoldi, L.; Ielo, L.; Holzer, W.; Giester, G.; Roller, A.; Pace, V. *J. Org. Chem.* **2018**, *83*, 4336-4347. (k) Parisi, G.; Colella, M.; Monticelli, S.; Romanazzi, G.; Holzer, W.; Langer, T.; Degennaro, L.; Pace, V.; Luisi, R. *J. Am. Chem. Soc.* **2017**, *139*, 13648-13651. (l) Pace, V.; Castoldi, L.; Mamuye, A. D.; Holzer, W. *Synthesis* **2014**, *46*, 2897-2909.
- (14) (a) Senatore, R.; Castoldi, L.; Ielo, L.; Holzer, W.; Pace, V. *Org. Lett.* **2018**, *20*, 2685-2688. (b) Castoldi, L.; Holzer, W.; Langer, T.; Pace, V. *Chem. Commun.* **2017**, *53*, 9498-9501. (c) Pace, V.; Murgia, I.; Westermayer, S.; Langer, T.; Holzer, W. *Chem. Commun.* **2016**, *52*, 7584-7587. (d) Monticelli, S.; Holzer, W.; Langer, T.; Roller, A.; Olofsson, B.; Pace, V. *ChemSusChem* **2019**, doi:10.1002/cssc.201802815. (e) Senatore, R.; Ielo, L.; Urban, E.; Holzer, W.; Pace, V. *Eur. J. Org. Chem.* **2018**, 2466-2470. (f) Mamuye, A. D.; Castoldi, L.; Azzena, U.; Holzer, W.; Pace, V. *Org. Biomol. Chem.* **2015**, *13*, 1969-1973. (g) Miele, M.; Citarella, A.; Micale, N.; Holzer, W.; Pace, V. *Org. Lett.* **2019**, *21*, 8261-8265. (h) Senatore, R.; Ielo, L.; Monticelli, S.; Castoldi, L.; Pace, V. *Synthesis* **2019**, *51*, 2792-2808. (i) Castoldi, L.; Ielo, L.; Hoyos, P.; Hernáiz, M. J.; De Luca, L.; Alcántara, A. R.; Holzer, W.; Pace, V. *Tetrahedron* **2018**, *74*, 2211-2217.
- (15) (a) Pace, V.; Pelosi, A.; Antermite, D.; Rosati, O.; Curini, M.; Holzer, W. *Chem. Commun.* **2016**, *52*, 2639-2642. (b) Touqeer, S.; Castoldi, L.; Langer, T.; Holzer, W.; Pace, V. *Chem. Commun.* **2018**, *54*, 10112-10115.
- (16) For seminal studies, see: (a) Reich, H. J.; Reich, I. L.; Renga, J. M. *J. Am. Chem. Soc.* **1973**, *95*, 5813-5815. (b) Reich, H. J.; Renga, J. M.; Reich, I. L. *J. Am. Chem. Soc.* **1975**, *97*, 5434-5447. (c) Reich, H. J.; Chow, F.; Shah, S. K. *J. Am. Chem. Soc.* **1979**, *101*, 6638-6648. (d) Reich, H. J. *Acc. Chem. Res.* **1979**, *12*, 22-30. (e) Reich, H. J.; Jasperse, C. P.; Renga, J. M. *J. Org. Chem.* **1986**, *51*, 2981-2988.
- (17) For an excellent overview of seleno-stabilized carbanions, see: (a) Krief, A.; Kremer, A. 3.02 Organolithium Compounds Bearing a Phenyl-, a Vinyl-, and/or a Seleno Group on their Carbanionic Centers: Synthesis by Se/Li Exchange and Unusual Synthetic Applications A2 - Knochel, P. In *Comprehensive Organic Synthesis (Second Edition)*; Elsevier: Amsterdam, 2014; pp. 56-156. (b) Krief, A. Selenium Stabilization. In *Comprehensive Organic Synthesis*; Trost, B. M., Fleming, I., Eds.; Pergamon: Oxford, 1991; Vol. 1; pp. 629-728. (c) Krief, A. *Tetrahedron* **1980**, *36*, 2531-2640. (d) Comasseto, J. V.; Dos Santos, A. A.; Wendler, E. P. Selenium-Stabilized Carbanions. In *Organoselenium Chemistry: Synthesis and Reactions*; Wirth, T., Ed.; Wiley: Weinheim, 2011; pp. 147-189. (e) The addition of a selenium-containing titanium enolate to imines has been reported, see: Silveira, C. C.; Vieira, A. S.; Braga, A. L.; Russowsky, D. *Tetrahedron* **2005**, *61*, 9312-9318.
- (18) For recent reviews, see: (a) Monticelli, S.; Castoldi, L.; Murgia, I.; Senatore, R.; Mazzeo, E.; Wackerlig, J.; Urban, E.; Langer, T.; Pace, V. *Monat. Chem.* **2017**, *148*, 37-48. (b) Pace, V.; Hoyos, P.; Castoldi, L.; Domínguez de María, P.; Alcántara, A. R. *ChemSusChem* **2012**, *5*, 1369-1379.
- (19) For a recent review, see: Azzena, U.; Carraro, M.; Pisano, L.; Monticelli, S.; Bartolotta, R.; Pace, V. *ChemSusChem* **2019**, *12*, 40-70.