

Chemistry Europe

European Chemical

Societies Publishing

Chemistry A European Journal



Accepted Article

Title: Evaluation of the Synthetic Scope and the Reaction Pathways of Proton-Coupled Electron Transfer with Redox-Active Guanidines in C-H Activation Processes

Authors: Ute Wild, Petra Walter, Olaf Hübner, Elisabeth Kaifer, and Hans-Jörg Himmel

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: Chem. Eur. J. 10.1002/chem.202003424

Link to VoR: https://doi.org/10.1002/chem.202003424



WILEY-VCH

Evaluation of the Synthetic Scope and the Reaction Pathways of Proton-Coupled Electron Transfer with Redox-Active Guanidines in C-H Activation Processes

Ute Wild, Petra Walter, Olaf Hübner, Elisabeth Kaifer, Hans-Jörg Himmel*

[a] U. Wild, P. Walter, Dr. O. Hübner, Dr. E. Kaifer, Prof. H.-J. Himmel Institut für Anorganische Chemie Ruprecht-Karls-Universität Heidelberg Im Neuenheimer Feld 270, 69120 Heidelberg E-mail: hans-jorg.himmel@aci.uni-heidelberg.de

Supporting information for this article is given via a link at the end of the document.

Abstract: Proton-coupled electron transfer (PCET) is currently intensively studied due to its importance in synthetic chemistry and biology. In recent years it was shown that redox-active guanidines are capable PCET reagents for the selective oxidation of organic molecules. In this work the scope of their PCET reactivity regarding reactions that involve C–H activation is explored and kinetic studies carried out to disclose the reaction mechanisms. Organic molecules with potential up to 1.2 V vs. ferrocenium/ferrocene are efficiently oxidized. Reactions are initiated by electron transfer, followed by slow proton transfer from an electron-transfer equilibrium.

Introduction

Proton-coupled electron transfer is of key importance for the progress in modern synthetic chemistry,[1,2] as it allows e.g. selective and green oxidation of organic substrates, including C-H activation processes, carbon dioxide conversion or water splitting. In the last decades, important advancements were made concerning theory, concepts and synthetic applications of PCET, helping to decode biological processes and paving the grounds for new applications in synthetic chemistry.^[1-8] It was systematically explored how the leveling effect, diminishing the effect of derivatisations (by which the redox potential or the pKa value could be tuned) on the PCET reactivity, could be circumvented by the design of bidirectional PCET reactions, in which the electron and proton are transferred to different molecules or different sites of the same molecule.[9, 10] Applications of bidirectional PCET in synthesis were comprehensively reviewed by Knowles.[11-13] Detailed studies showed how environmental effects (e.g. hydrogen-bonding, the solvent polarity or the presence of acids) could be used to enable or speed up PCET reactions. The accumulation of charges in photoredox catalytic systems could be avoided by PCET, allowing the accumulation of oxidative and reductive equivalents instead of charges. ^[14,15] Conceptual work demonstrated how electron transfer could initiate proton movement over large distances. Hence, first proton shuttles were designed in which the transfer of several protons within hydrogen-bonds is triggered by electron transfer. ^[16,17] Hydrogen-bonding triggered by redox processes could be employed in sensor devices, and in hydrogen-bonded molecular shuttles that might be usable as a basis for artificial molecular machines.

Hence, in theory an extensive, rational and in some areas biomimetic use of organic PCET reagents in synthetic chemistry is now possible. However, progress does not only depend on the detailed knowledge of the mechanisms and the developments of advanced concepts. Due to the various areas of use of PCET reactions in modern synthetic chemistry, the availability of a larger number of PCET compound classes is required. Many applications still suffer from the liability of the known organic PCET reagents to side reactions, prohibiting the formation or diminishing the yield of the desired product. In stoichiometric reactions, an excess of the PCET reagent is not seldomly required to obtain good yields. High catalyst loadings are necessary in a number of catalytic reactions, making them unattractive for largescale processes. Some reactions (e.g. Scholl-type aryl-aryl coupling reactions) need to be carried out in highly acidic media, that could initiate degradation of the PCET reagent or the organic substrate. Photochemical applications are still limited due to the relatively small number of available photoactive PCET reagents. Finally, frequently employed quinones such as 2,3-dichloro-5,6dicyano-1,4-benzoquinone (DDQ) and tetrachloro-1,4benzoquinone (chloranil, CA), are highly toxic. Therefore, it is essential to develop new classes of PCET reagents to overcome limitations concerning stability, reactivity and toxicity of traditionally applied compounds. Numerous fields in synthetic

10.1002/chem.202003424

WILEY-VCH

FULL PAPER

chemistry could profit immensely from the developments of such new PCET reagents.

In the last years, we developed redox-active guanidines as a new class of capable PCET reagents.^[18] We already studied to some PCET extend the chemistry of 1,2,4,5tetrakis(tetramethylguanidino)-benzene (1). lt is readily accessible from commercially available 1,2,4,5-tetrakisaminobenzene-tetrahydrochloride,[19] and a relatively strong electron donor with a redox potential ($E_{1/2}$ value) of -0.73 V vs. ferrocenium/ferrocene for the redox couple $1^{2+}/1$ in CH₃CN solution.^[20] The compound looses two electrons at the same potential. The free radical monocation 1^{.+} is unstable towards disproportionation into 1 and 12+. Hence it is not formed in mixtures of 1 and 12+. On the other hand, the radical monocationic form is stable as bridging ligand in several dinuclear late-transition metal complexes.^[21,22] A number of salts of 1²⁺ were fully characterized, including $1(PF_6)_2^{[23]}$ and $1(BF_4)_2^{[24]}$ used in this work, and $1(I_3)_2$.^[19] Also, the nitrogen-rich $1[N(CN)_2]_2$ was prepared,^[25] that melts above 200 °C and decomposes smoothly at 220 °C, demonstrating its excellent thermal stability. In the dication 1^{2+} , the bond length between the carbons in 1 and 2 positions and that between the carbons in 4 and 5 positions (all directly bound to guanidino groups) are considerably elongated, in line with the Lewis structure in Figure 1. The twofold protonated, reduced form (1+2H)²⁺ could be re-oxidized to the dication 1²⁺ by dioxygen under mild conditions with a copper or cobalt catalyst,^[26,27] opening up the possibility to use 1 as redox-catalyst for the green aerobic oxidation of a variety of organic molecules.^[27] The scope for stoichiometric PCET reactions with 1²⁺ could be greatly extended in the presence of strong acids.^[28] leading in the first place to di-protonation of 12+ to give the tetracation (1+2H)⁴⁺ (see Lewis structure in Figure 1)^[29] with ca. 0.7 V higher oxidation potential.

In this work we elaborate on the PCET chemistry of the three redox-active guanidines 1,2,4,5-tetrakis(tetramethylguanidino)benzene (1), 1,4-bis(tetramethylguanidino)-benzene (2) and the new compound 1,4-bis(*N*,*N*-dimethylethylene)guanidinobenzene (3) shown in Figure 1, in PCET reactions that involve C-H activation. The conversion was followed by NMR or UV-vis spectroscopy, and the yields were estimated by NMR signal integration. Please note that the equations shown in the following do not account for protonation equilibria that are observed for the reduced, protonated guanidines arising as products in these reactions (see SI for NMR spectra of the protonated compounds). Also, in the presence of strong acids all guanidino groups become protonated.



Figure 1. The three redox-active guanidines studied in this work and equations to illustrate the PCET reactivity of 1²⁺.

(**1**+4H)⁴⁺

Results and Discussion

(**1**+2H)⁴⁺

1) Expansion of the scope of PCET chemistry with 12+.

We decided to test first the oxidative coupling of N-ethylcarbazole to N.N'-diethyl-3,3'-bicarbazole (Scheme 1). N-ethylcarbazole exhibits an Eox value of 1.12 V vs. SCE, [30] translating into a value of 0.66 V vs. Fc⁺/Fc.^[31] Since previous experiments showed that substrates with a redox potential of up to 0.77 V vs. Fc+/Fc could be oxidized.^[28] it was clear that the potential is not too high for a reaction with initial electron transfer. In our experiments, one equivalent of 1(BF₄)₂ or 1(ClO₄)₂ (see SI for synthesis and characterization of this new salt) was reacted with Nethylcarbazole in the presence of 16 equivalents of HBF₄·OEt₂. Acetonitrile was chosen as solvent, since the protonated, oxidized guanidine (1+2H)4+ that forms immediately in the presence of a strong acid, is not soluble in unpolar solvents such as CH₂Cl₂. Indeed, near quantitative conversion (94% and 95%, respectively) was obtained within 1 h reaction time. Recently, Venkatakrishnan et al. showed that reaction of DDQ or CA with N-ethylcarbazole gives quantitative yield (>99%) of the bicarbazole coupling product in very short time when carried out in CH₂Cl₂ solution with an excess of methanesulfonic acid.[32, 33] However, two equivalents of the guinone (fourfold excess) had to be used; the

yield decreased to 79% for CA and 87% for DDQ when only one equivalent was used (like in our experiments). Moreover, the yield decreased to 83% when the reaction was carried out in CH₃CN instead of CH₂Cl₂. Hence the results demonstrate that salts of 1^{2+} are valuable alternatives to toxic CA or DDQ in aryl-aryl coupling reactions.



Next we inspected the oxidative coupling of 3,3"-dimethoxy-3',4'dimethyl-o-terphenyl to 3,10-dimethoxy-6,7-dimethyltriphenylene with salts of 12+, again in the presence of excess (ca. 20 equivalents) of HBF₄·OEt₂ (Scheme 2). With 1(BF₄)₂, a yield of 76% was obtained after 130 min reaction time; use of $1(PF_6)_2$ resulted in a slightly lower yield of 71%. This reaction demonstrates that salts of 1^{2+} could oxidize compounds with redox potentials of 1.2 V vs. Fc+/Fc, being thus at least similar in their oxidizing capability to DDQ or CA. The strong acid leads to double-protonation of 1^{2+} to $(1+2H)^{4+,[29]}$ which is the oxidant in this reaction. Previous quantum-chemical calculations indicate that (1+2H)⁴⁺ is even a stronger oxidant in PCET reactions than DDQ or CA.^[28] The reaction requires the use of 1.5 equivalents of 1^{2+} . This can be explained by the considerably lower redox potential of the triphenylene product (0.72 V vs. Fc⁺/Fc in CH₂Cl₂) compared with the reactant (1.22 V vs. Fc+/Fc in CH2Cl2 for 3,3"dimethoxy-3',4'-dimethyl-o-terphenyl), leading to preferred oxidation of the product by $(1+2H)^{4+}$ to the radical monocation. The formation of this radical cation leads to broad signals in the NMR spectra, and therefore the reaction was guenched before analysis. Oxidative coupling of 3,3"-dimethoxy-3',4'-dimethyl-oterphenyl could also be carried out with 1.5 equivalent of DDQ and an excess of MeSO₃H in CH₂Cl₂, giving quantitatively the triphenylene coupling product in short time.[34,35] A detailed analysis showed that this reaction follows a cation-radical (electron transfer) mechanism rather than an arenium-ion (proton transfer) mechanism.^[35] Similarly, electron transfer between (1+2H)⁴⁺ and 3,3"-dimethoxy-3',4'-dimethyl-o-terphenyl is assumed to proceed proton transfer (see discussion in section 3). Please note that for both reactions the guanidine PCET reagent could be recycled from the reaction mixture. For this purpose it was first separated from the aryl-aryl coupling product by addition of ether (in which only the coupling product is soluble) and

filtration. Then the reduced, protonated $(1+2H)^{2+}$ was re-oxidized to 1^{2+} by catalytic oxidation with dioxygen (see SI for details).



Scheme 2. Intramolecular oxidative coupling of 3,3"-dimethoxy-3',4'-dimethylo-terphenyl to the corresponding dimethoxy-triphenylene. Conditions: acetonitrile, 1.5 eq. 1(BF₄)₂, 20 eq. HBF₄·OEt₂, 10 min at 0 °C, 120 min at r.t., 76% yield.

2) Further C-H bond cleavage reactions with salts of 1²⁺.

Having demonstrated the scope of oxidative aryl-aryl coupling reactions for substrates with potentials of up to 1.2 V vs. Fc+/Fc, we next focussed on the oxidation of 1-benzyl-1,4dihydronicotinamide (BNAH), 10-methyl-9,10-dihydroacridane (AcrH₂) and 9,10-dihydroanthracene (AnH₂) to systematically study PCET reactions without and with addition of a strong acid. 10-Methyl-9,10-dihydroacridane (AcrH₂) exhibits a bond dissociation energy (BDE) of 308 kJ mol-1 in CH₃CN solution,^[36] and an E_{ox} value of 0.492 V vs. Fc⁺/Fc in CH₃CN. With a reduction potential (E_{red}) of -0.77 V vs. Fc⁺/Fc for 1^{2+,[20]} the energy gap ΔG_{el} for electron transfer between AcrH₂ and 1²⁺ is 1.26 V or 122 kJ mol⁻¹, exceeding clearly the assumed limit for conventional electron transfer at standard conditions of ca. 1 V (96.5 kJ mol⁻¹).^[37] Hence, 1²⁺ is too weak to oxidize AcrH₂. Since electron transfer is supposed to be the first step of the reaction (see discussion in section 5), the expectation is that no reaction takes place. Indeed, only traces of N-methylacridinium (AcrH⁺) are observed if the reaction is carried out in the absence of an acid. On the other hand, a high product yield (90%) in 6.5 h reaction time is obtained upon addition of 7 equivalents of HBF₄·OEt₂ (Scheme 3). Again, the acid instantly protonates the guanidine to give the strong oxidant (1+2H)⁴⁺ (see Figure 1),^[24] decreasing the energy gap between the AcrH₂ electron donor and the acceptor to ca. 0.5 V, enabling electron-transfer as initial reaction step. The further excess of acid warrents fast conversion. A small but significant isotope effect was observed when AcrH₂ was replaced by AcrD₂ (see SI), implying that the rate of proton transfer enters into the overall reaction rate (see discussion below).



Scheme 3. Oxidation of 10-methyl-9,10-dihydroacridane (AcrH₂). Conditions: acetonitrile, 1 eq. $1(PF_6)_2$, 7 eq. HBF₄·OEt₂, 6.5 h at r.t., 90% yield.

3

The BDE value of 9,10-dihydroanthracene (ca. 326 kJ mol-1) is higher than that of AcrH₂.^[36] Although quantum-chemical calculations (B3LYP/def2-TZVP) predict the reaction of 12+ with 9,10-dihydroanthracene (AnH₂) to give anthracene (An) to be still exothermic (Gibbs free reaction enery of -104.5 kJ mol⁻¹ at ε_r = 38 and -87.7 kJ mol⁻¹ at ε_r = 1), a massive barrier is expected for the unfavored initial electron transfer step. Therefore, also reaction of 1^{2+} with 9.10-dihydroanthracene requires the presence of an excess of HBF₄·OEt₂. With 5 equivalents of HBF₄·OEt₂ and using the salt $1(PF_6)_2$, a yield of 78% anthracene is obtained within 3 h at room temperature (Scheme 4). The guanidine PCET reagent could be recycled (see SI). When more HBF₄·OEt₂ is applied, the rate increases, but the An yield decreases (e.g. 65% yield after 1.5 h with 9 equivalents of HBF₄·OEt₂). Interestingly, an induction period with low rate is visible in the conversion vs. time plot (see SI). A possible explanation is an autocatalytic effect by formation of the anthracene radical. Indeed, NMR spectra recorded after 2 h reaction time displayed broad signals indicating the presence of radicals in the reaction mixture. By contrast, the signals in the NMR spectra recorded after 3 h were again sharp, arguing for complete conversion of the radical intermediates. For comparison, kinetic measurements for the reaction between anthracene (An) and 9,10-dihydrophenanthrene (PhenH₂) to give eventually AnH₂ and Phen [38] showed that fast reaction between the produced AnH₂ and the reactant An generates the reactive radical AnH⁻.



 $\begin{array}{l} \mbox{Scheme 4. Oxidation of 9,10-dihydroanthracene. Conditions: acetonitrile, 1 eq. } \\ 1(PF_6)_2, 5 eq. \ HBF_4\cdot OEt_2, 3 \ h \ at r.t., 78\% \ yield. \end{array}$

Finally, the reaction of 1^{2+} with 1-benzyl-1,4-dihydronicotinamide (BNAH) was tested. The addition of a strong acid is not possible in this reaction due to acid-catalysed hydration of BNAH.^[39] The BDE value of BNAH is 284 kJ mol⁻¹,^[36] and the E_{ox} value is 0.259 V vs. Fc⁺/Fc in CH₃CN. The energy gap ΔG_{el} for electron transfer between 1^{2+} ($E_{red} = -0.77$ V vs. Fc⁺/Fc) and BNAH of 1.0 V is just at the assumed limit value for conventional electron transfer at standard conditions. Hence, reaction between 1^{2+} and BNAH might be possible at higher temperatures. Indeed, a slow reaction was observed at a temperature of 60 °C, yielding 27% of the pyridinium salt after 24 h (Scheme 5). Prolonged reaction times (2 d) lead to degradation of BNAH.



Scheme 5. Oxidation of 1-benzyl-1,4-dihydronicotinamide (BNAH). Conditions: acetonitrile, 1 eq. $1(\text{PF}_6)_2,$ 24 h at 60 °C, 27% yield.

3) Comparison between the PCET reactivity of 12+ and 22+.

The results assembled in the previous section demonstrate that salts of 12+ are capable PCET reagents. However, oxidation of higher potential substrates requires the addition of a strong acid. Since some substrates, e.g. BNAH, degrade in the presence of acids, we thought of ways to increase the oxidation power in the absence of acids. We previously showed that the redox potential of 1,4-bis(tetramethylguanidino)-benzene (2) is significantly higher than that of 1 ($E_{1/2}$ = -0.21 V vs. Fc⁺/Fc in CH₃CN for 2 ^[40] and -0.73 V for 1 [20]). Although 22+ is therefore a stronger oxidant in electron-transfer reactions, it is not neccessarily a stronger oxidant in PCET reactions, since the thermodynamics of PCET reactions depends not only on the potentials, but also on the pKa values of the reduced, protonated species. Generally, an increase of the potential is accompanied by a decrease of the pKa value of the protonated, reduced form. This "leveling effect" limits the possibility of tuning the PCET reactivity by derivatization. In this section we compare the PCET reactivity of 2²⁺ with that of 1²⁺ by direct reaction between the oxidized form of one compound with the reduced, doubly protonated form of the other compound.

Our experiments show that 2^{2+} reacts fast with $(1+2H)^{2+}$ to give $(2+2H)^{2+}$ and 1^{2+} . In an acetonitrile solution with concentrations of $1.25 \cdot 10^{-2}$ mol/l for both reactants, quantitative conversion (>99%) is obtained after 5 min at 25 °C (Scheme 6). On the other hand, no reaction takes place when 1^{2+} is mixed with $(2+2H)^{2+}$. From these experiments one could deduce that 2^{2+} is indeed a stronger PCET reagent than 1^{2+} , at least in the absence of a strong acid. This result is supported by quantum-chemical calculations predicting a Gibbs free reaction energy of -84.5 kJ mol⁻¹ at $\varepsilon_r = 38$ and -105.7 kJ mol⁻¹ at $\varepsilon_r = 1$.



Scheme 6. Oxidation of $(1+2H)^{2+}$. Conditions: acetonitrile, 1 eq. $2(PF_6)_2$, 5 min at r.t., >99% yield.

The fast rate of the reaction between 2^{2+} and $(1+2H)^{2+}$ gets evident from the appearance of the green colour characteristic for

FULL PAPER

1²⁺, motivating further analysis by UV-vis spectroscopy (Figure 2). For equal concentrations (5.6·10⁻⁵ mol/l) of the two reactants, the reaction is completed after 20 min. The growth of a strong band at 425 nm signals the formation of 1²⁺. The presence of an isosbestic point at ca. 357 nm indicates clean conversion, in line with the NMR experiments at higher concentrations. In further experiments, we added 0.25 equivalents of HBF₄·OEt₂ to see if the reaction rate is influenced by the presence of acid (see SI). A decrease of the reaction rate upon acid addition was observed. This could be explained by the protonation of the (1+2H)²⁺ reactant to give (1+4H)⁴⁺, which is less oxidizable. On the other hand, additional NMR experiments showed that the reaction could not be reversed by addition of larger quantities of HBF₄·OEt₂. Also, addition of HBF₄·OEt₂ to a 1²⁺ / (2+2H)²⁺ mixture only leads to the protonated, oxidized guanidine (1+2H)⁴⁺, but not to PCET.



Figure 2. UV-vis spectra recorded for the 1:1 reaction between 2^{2+} and $(1+2H)^{2+}$. The strong band at 426 nm signals formation of 1^{2+} as one reaction product.

To analyse the kinetics in more detail, stopped-flow measurements were conducted in which 2^{2+} was applied in excess (concentrations of 3.94·10⁻⁴ mol/l (ratio $2^{2+}:(1+2H)^{2+} = 10:1$), 7.87 10⁻⁴ mol/l (20:1), and 1.58 10⁻³ mol/l (40:1)). The results are in line with a pseudo-first order reaction only within the first seconds, hampering an unambigeous identification of the reaction order. On the other hand, the estimated pseudo-first order rate constants increase linearly with the concentration of 2^{2+} . Hence, the results might be in line with a first order in both 2^{2+} and $(1+2H)^{2+}$ in the initial reaction period (see SI for details), and the analysis yielded a second order rate constant $k_{\rm H} = 257 \pm 27 \,{\rm M}^{-1}{\rm s}^{-1}$ at room temperature.

The results in this section clearly show that 2^{2+} is a stronger PCET oxidant than 1^{2+} . Additional NMR experiments indicated degradation of 2^{2+} to so far unidentified products in the presence of excess HBF₄·OEt₂ (see SI). Therefore most reactions reported in the following were carried out in the absence of acid.

4) PCET reactions with 22+.

The reactions of 2^{2+} with 1-benzyl-1,4-dihydronicotinamide (BNAH), 10-methyl-9,10-dihydroacridane (AcrH₂) and 9,10dihydroanthracene (AnH₂) were carried out in the absence of a strong acid. Salts of 2^{2+} react very fast with BNAH; quantitative yield (>99%) is obtained instantly upon mixing BNAH with $2(PF_6)_2$ together at room temperature in acetonitrile solution (Scheme 7). With $E_{red} = -0.24$ V for 2^{2+} , the energy for electron transfer, ΔG_{el} , is 0.50 V (48 kJ mol⁻¹) and therefore well below 1 V. Hence the observation of fast reaction is in line with electron transfer in the first step.



Scheme 7. Oxidation of BNAH. Conditions: acetonitrile, 1 eq. $2(\mathsf{PF}_6)_2, \ r.t.,$ instantly, >99% yield.

As expected due to its higher potential, reaction of 2^{2+} (applied as $2(PF_6)_2$ salt) with 10-methyl-9,10-dihydroacridane is slower. NMR experiments showed that a yield of ca. 92% is obtained after 2.5 h at room temperature using 2 equivalents of $2(PF_6)_2$ (Scheme 8). The reaction is further slowed down if only one equivalent of $2(PF_6)_2$ is applied, yielding 73% ArcH⁺ in 3 h at room temperature. Again, this result is in line with electron transfer in the first step, since ΔG_{el} is 0.73 V (70 kJ mol⁻¹).



 $\label{eq:Scheme 8. Oxidation of N-methylacridane (AcrH_2)$. Conditions: acetonitrile, 2 eq. $2(PF_6)_2$, 160 min at r.t., 92% yield.}$

The reaction of 2^{2+} with 9,10-dihydroanthracene is again slower. A yield of only 6% of anthracene is obtained after 50 h at 60 °C. The yield could be increased to 29% by addition of ca. 2 equivalents of NH₄PF₆ (Scheme 9). In difference to the results obtained for the reaction of 1^{2+} with 9,10-dihydroanthracene in the presence of acid, there is no induction period, and the conversion increases almost linearly with time. Quantum chemical calculations (B3LYP/def2-TZVP) predict the reaction of 2^{2+} with 9,10-dihydroanthracene to be significantly exothermic (Gibbs free reaction energy of -189.0 kJ mol⁻¹ at $\varepsilon_r = 38$, and -193.4 kJ mol⁻¹ at $\varepsilon_r = 1$). The slow reaction rate could be explained by a

FULL PAPER

substantial barrier for the initial electron-transfer step. A similar strong correlation between the potential (E_{ox} or $E_{1/2}$ value) of the PCET reagent and the reaction yield was recently reported for oxidative C-H amination reactions of 9,10-dihydro-9-heteroanthracenes with quinones as PCET reagents.^[41]



 $\begin{array}{l} \mbox{Scheme 9. Oxidation of 9,10-dihydroanthracene. Conditions: acetonitrile, 1 eq. $$2(PF_6)_2, 2 eq. NH_4PF_6, 50 h at 60 °C, 29\% yield. $$ \end{array}$

Next, we tested the performance of 2^{2+} in an aryl-aryl coupling reaction. Reaction of 2²⁺ with 3,3"-4,4"-tetramethoxy-o-terphenyl (TMTP) at room temperature did not produce the product in the absence of an acid. The oxidation potential of TMTP is 0.74 V, and hence the energy for electron transfer, ΔG_{el} , is 0.98 V, close to the assumed limit value of 1.0 V. On the other hand, only small amounts of HBF₄·OEt₂ are required to initiate fast and clean reaction. With 12.5 mol% of HBF₄·OEt₂, a yield of 95% is obtained within 10 min reaction time at a temperature of 60 °C (Scheme 10). For comparison, the analogue reaction with 1²⁺ in place for 2^{2+} required the use of a large excess (21 eq.) of HBF₄·OEt₂ to be completed in relatively short time (45 min at room temperature, 99% yield).^[28] In the case of 2²⁺, small amounts of a strong acid could be applied and do not lead to acid-induced degradation of 2^{2+} . However, an excess of the acid (e.g. 4 equivalents and more) has to be avoided as it leads to quite fast degradation of 2^{2+} (see section 3 and SI).



Scheme 10. Oxidation of 3,3"-4,4"-tetramethoxy-o-terphenyl. Conditions: acetonitrile, 1 eq. 2(PF₆)₂, 12.5 mol% HBF₄·OEt₂, 10 min at 60 °C, 95% yield.

Finally, we inspected the reaction of 2^{2+} with *p*-dihydrobenzoquinone (Scheme 11). NMR experiments (c = $2.9 \cdot 10^{-2}$ mol·l⁻¹ for both reactants) showed complete conversion (>99%) within 12 min at 55 °C. At room temperature, 60 min are required (98% yield). Fits of the conversion vs. time curves, obtained for different concentrations of the two reactants, while keeping the 1:1 stoichiometric ratio, and for experiments at different temperatures (see SI), gave no satisfactorily results

when assuming second or first order rate equations. An analysis by UV-vis spectroscopy is hampered by the close proximity of the absorptions due to reactants and products. Quantum-chemical calculations (B3LYP/def2-TZVP) found a Gibbs free reaction energy of -96.1 kJ mol⁻¹ at $\varepsilon_r = 38$ and -107.8 kJ mol⁻¹ at $\varepsilon_r = 1$. For comparison, the analogue reaction with 1^{2+} in place for 2^{2+} is only slightly exergonic (-11.6 kJ mol⁻¹ at $\varepsilon_r = 38$, calculations with counterions).



Scheme 11. Oxidation of p-dihydrobenzoquinone. Conditions: acetonitrile, 1 eq. $2({\rm PF}_6)_2,\,12$ min at 55 °C, >99% yield.

5) Kinetic measurements and mechanistic considerations.

The reaction of 2²⁺ with 10-methyl-9,10-dihydroacridane was examined in more detail to derive information about the reaction mechanism. Preliminary NMR experiments point to a slower reaction with ArcD₂ in place for AcrH₂ (see SI); UV-vis experiments were conducted for further kinetic analysis. To obtain pseudo-first order conditions, a ten-, twenty- and forthy-fold excess of AcrH₂ was applied. The formation of Nmethylacridinium is clearly visible from the appearance of the typical sharp band at 357 nm and a broader one around 420 nm (with maxima at 395/415/440 nm). The band at 357 nm was chosen for the analysis. Figure 3a shows the spectra recorded with a tenfold excess of AcrH₂. In the first 5 min, the reaction nicely follows a pseudo first-order kinetics as seen from the In(A) vs. time plot in the inlet. The k_{obs} values derived from such fits were plotted as a function of the AcrH₂ concentration (see Figure 3b). The slope of this plot gives the second-order rate constant at room temperature, $k_{\rm H} = 3.32 \pm 0.16 \text{ M}^{-1}\text{s}^{-1}$. For comparison, a ca. 4.5 times larger rate constant ($k_{\rm H} = 15 \text{ M}^{-1}\text{s}^{-1}$) was previously published for the reaction between chloranil and AcrH₂ (also in acetonitrile).^[42] A similar analysis for AcrD₂ yielded a second-oder rate constant $k_D = 0.593 \pm 0.021 \text{ M}^{-1}\text{s}^{-1}$ (see plot of the k_{obs} values as a function of AcrD₂ concentration in Figure 3b and SI for further details). Thus a quite large kinetic isotope effect (KIE = $k_{\rm H}/k_{\rm D}$) of 5.6 results, being nevertheless still smaller than the KIE of 8.8 obtained for the reaction between CA and N-methyl-acridane (AcrH₂).^[42]



Figure 3. a) Selected UV-vis spectra for the reaction between $2(PF_6)_2$ and AcrH₂ (10 equivalents) in CH₃CN solution. The inlet shows the ln(A) vs. time plot in the first 4 min of the reaction, from which the pseudo first-order rate constant is determined. b) Plot of the pseudo first-order rate constants as a function of the acridane concentration (for an invariant concentration of $2(PF_6)_2$ of ca. 8.10⁻⁵ M). See SI for details.

The large KIE clearly shows that the rate constant for the protonation step is contributing to the overall rate constant. AcrH₂ is known to prefer a stepwise e-, H+, e- pathway for hydride transfer in many reactions. The energy for electron transfer, ΔG_{el} , for reaction between 2²⁺ and AcrH₂⁺ of 0.73 V (70 kJ mol⁻¹) is well within the region of conventional electron transfer under standard conditions, supporting electron transfer in the first step. Therefore a stepwise e-, H+, e- pathway, as sketched in Scheme 11, is likely to be operative here, too. The last step of the proposed mechanism is a fast second electron transfer, profiting from the low potential of the AcrH⁻ radical ($E_{ox} = -0.46$ V vs. SCE, ca. -0.06 V vs. Fc⁺/Fc).^[43] Hence this last step plays no role in the overall rate. According to the formula $k_{\rm H} = k_{\rm P} k_{\rm et} / (k_{\rm et} + k_{\rm P})$ ^[42] the observation of a large KIE value means that $k_{\rm P}$ (the rate constant for proton transfer) is much smaller than k_{-et} (the rate constant for back electron-transfer regenerating the reactants).



Scheme 11. Suggested mechanism for the oxidation of *N*-methylacridane with

In the UV-vis experiment, a small and broad absorption around 575 nm first increased in intensity, reaching a maximum after ca. 26 min, and then decreased (see difference spectrum in Figure 4 a), belonging to a reaction intermediate. According to the proposed reaction scheme, the obvious candidates for this intermediate are the radicals 2^{.+} and AcrH₂^{.+}, formed upon first electron transfer. A broad, weak absorption around 640 nm was previously assigned to the radical cation AcrH2.+. [42-44] We did not observe this band, most likely because the concentrations of AcrH2⁺⁺ are too low. To obtain more information on the other candidate, 2^{.+}, we prepared a 1:1 mixture of 2 (colourless) and 2(PF₆)₂ (yellow) in CH₃CN solution. The reaction turned instantly deep purple. In the UV-vis spectrum, bands at 575 nm (with a shoulder at 542 nm) and 404 nm appeared (Figure 4 b), vanishing within a few hours. TD-DFT calculations (B3LYP/def2-TZVP) predicted electronic transitions at 543/501 and 365/346 nm for 2^{.+}, in good agreement with the experimental values (see SI for details). The band at 575 nm could therefore be assigned to the radical 2⁺⁺. The other band of 2⁺⁺ in the visible region at 404 nm is obscured by the strong, broad bands of AcrH+ in this region (Figure 4 a). The detection of 2^{+} as a reaction intermediate is a valuable additional evidence for the reaction pathway sketched in Scheme 11.



Figure 4. a) Difference spectrum (26 minus 78 min reaction time) for the reaction between $2({\sf PF}_6)_2$ and AcrH_2 (10 equivalents) in CH_3CN solution. The band assigned to 2^+ is highlighted by an asterisk. b) Decay of the absorptions due to the radical monocation 2^+ , formed instantly upon mixing 2 and 2^{2+} in acetonitrile solution.

6) Reaction between two bisguanidines.

To test PCET between closely related redox-active guanidines, we synthesized the new bisguanidine **3** (77% yield) by reaction of *p*-phenylenediamine-dihydrochloride with 2-chloro-1,3-dimethyl-4,5-dihydro-1*H*-imidazolium chloride (prepared in situ from 1,3-dimethyl-2-imidazolidinone and oxalyl chloride, see SI for details). Compound **3** was then oxidized with FcPF₆ to give the salt **3**(PF₆)₂ (48% isolated yield), and protonated with NH₄PF₆ to give (**3**+2H)(PF₆)₂ (72% isolated yield), and these two states completely characterized. In addition, (**3**+H)PF₆ was synthesized from a 1:1 mixture of **3** and (**3**+2H)(PF₆)₂. The structures in the solid state of all relevant states are illustrated in Figure 5. Further experiments showed that **3**²⁺ does not degrade in the presence of larger amounts of a strong acid, in contrast to **2**²⁺.



Figure 5. Visualisation of the solid-state structures of the stable protonation and redox states of 3. C-H protons omitted. Displacement ellipsoids drawn at the 50% probability level. Colour code: N blue, C grey, P pink, F green. See SI for details.

According to cyclic voltammetry (Figure 6), the redox properties of **3** are very similar to those of **2**. In both cases two-electron waves are observed, leading reversibly to the dicationic state. An $E_{1/2}$ value of -0.21 V vs. Fc⁺/Fc ($E_{ox} = -0.18$ V, $E_{red} = -0.24$ V) for **2** compares with an $E_{1/2}$ value of -0.24 V vs. Fc⁺/Fc ($E_{ox} = -0.20$ V, $E_{red} = -0.27$ V) for **3**. Although the radical cation **3**⁺⁺ does not appear in the CV measurements, it could be generated (similar to **2**⁺⁺) by mixing neutral **3** and **3**(PF₆)₂ in CH₃CN solutions or by titration of solutions of **3** with **3**(PF₆)₂, and displays strong absorptions at 566 and 383 nm (see SI). Interestingly, degradation of **3**⁺⁺ is much slower than **2**⁺⁺ (see SI).



Figure 6. CV curves (CH₃CN, Ag/AgCl reference electrode, 0.1 M N(^{*n*}Bu)₄(PF₆) as supporting electrolyte, scan speed 100 mV s⁻¹) for the two compounds 2 and 3 measured in oxidation direction. Potentials measured vs. Fc⁺/Fc.

We then directly compared the PCET reactivity of the two compounds. Reaction between 2^{2+} and $(3+2H)^{2+}$ gave almost

anus

epte

quantitative yields of 3^{2+} . For example, reaction at 55 °C for a period of 45 min gave 99% yield (Scheme 12). On the other hand, no reaction was observed when 3^{2+} and $(2+2H)^{2+}$ were mixed together. These results are in line with quantum chemical calculations (B3LYP/def2-TZVP), predicting the reaction (at $\varepsilon_r =$ 1) in Scheme 12 to exhibit a reaction enthalpy at 0 K of -66.7 kJ mol⁻¹ and a Gibbs free energy at 298 K of -58.2 kJ mol⁻¹.



Scheme 12. Oxidation of $(3+2H)^{2+}.$ Conditions: acetonitrile, 1 eq. $2(\mathsf{PF}_6)_2,$ 45 min at 55 °C, 99% yield.

From the temperature dependence of the conversion vs. time plots and assuming a second-order rate law, the activation energy E_A could be estimated from an Arrhenius plot (see Figure 7 and SI for details). An E_A value of 54 ± 8 kJ mol⁻¹ was obtained in this way. This quite low activation energy might argue for a concerted e⁻, H⁺ transfer; a stepwise e⁻, H⁺ transfer would create a highly unfavorable intermediate (3+2H)³⁺, and a stepwise H⁺, e⁻ transfer an similarly unfavorable intermediate (2+H)3+. The basicity of GFAs with tetramethylguanidino groups is significantly higher than that of GFAs with N,N-dimethylethylene-guanidino groups.^[45] Thus, the redox potential and the pKa value, being the two decisive factors for the PCET thermodynamics, seem to operate in the same direction, making 32+ a weaker PCET reagent than 2^{2+} . Hence, the leveling effect oberved for quinones, that results from opposite trends of potential and pKa value, might not occur for redox-active guanidines.



Figure 7. Conversion vs. time plots for the reaction between 2^{2+} and $(3+2H)^{2+}$ at different temperatures. The second-order *k* values estimated from each curve were used in the ln*k* vs. 1/T plot in the inlet, allowing to estimate the Arrhenius activation energy.

Conclusion

Proton-coupled electron transfer (PCET) reactions between redox-active guanidines and a variety of organic compounds in which C-H bonds are cleaved were systematically evaluated. First it is demonstrated that substrates with oxidation potentials up to at least 1.2 V vs. $Fc^{+/0}$ could be oxidized in the presence of a strong acid. Then, the two redox-active guanidines 1,2,4,5tetrakis(tetramethylguanidino)-benzene (1) and 1,4bis(tetramethylguanidino)-benzene (2), both applied in their oxidized, dicationic state (12+ or 22+) are compared in their PCET reactivity. In the course of our analysis, the reactions with the three substrates 1-benzyl-1,4-dihydronicotinamide (BNAH), 10methyl-9,10-dihydroacridane (AcrH₂) and 9,10dihydroanthracene (AnH₂) were studied for both 1^{2+} and 2^{2+} . The results clearly show that 2^{2+} is a significantly stronger oxidant in PCET reactions. However, in contrast to 1²⁺, it degrades in the presence of larger quantities of a strong acid. In all PCET reactions discussed in this work, slow proton transfer is likely to initial proceed from electron-transfer equilibrium. an Consequently, a relatively large kinetic isotope effect (KIE) of 5.6 was obtained for the reaction between 2^{2+} and AcrH₂.

Finally, we synthesized a new redox-active guanidine, 1,4bis(N,N-dimethylethylene-guanidino)-benzene (**3**) and compared the PCET reactivity of **3**²⁺ and **2**²⁺. Compound **3**²⁺ has a slightly lower reduction potential and is a weaker oxidant in PCET reactions. The comparison between the two closely related bisguanidines indicates that a "leveling effect" (resulting from opposite trends of potential and pK_a) as observed for quinones, might not be an issue for redox-active guanidines, allowing the tuning of the PCET thermodynamics by derivatisations. However, more work is necessary to confirm this assumption, that would be very helpful for a directed approach to PCET reactions.

The results presented in this work clearly show that redox-active guanidinces are potent PCET reagents and real alternatives to toxic quinones like chloranil or DDQ.

Acknowledgements

The authors gratefully acknowledge financial support by the *Deutsche Forschungsgemeinschaft* (DFG).

Keywords: proton-coupled electron transfer • guanidine • oxidation • C-H activation • oxidative coupling

FULL PAPER

- [1] M. H. V. Huynh, T. J. Meyer, Chem. Rev. 2007, 107, 5004–5064.
- D. R. Weinberg, C. J. Gagliardi, J. F. Hull, C. F. Murphy, C. A. Kent, B.
 C. Westlake, A. Paul, D. H. Ess, D. G. McCafferty, T. J. Meyer, *Chem. Rev.* 2012, *112*, 4016–4093.
- [3] a) S. Hammes-Schiffer, A. V. Soudackov, J. Phys. Chem. B 2008, 112, 14108–14123; b) S. Hammes-Schiffer, A. A. Stuchebrukhov, Chem. Rev. 2010, 110, 6939–6960; c) S. Hammes-Schiffer, Acc. Chem. Res. 2018, 51, 1975–1983.
- [4] a) J. M. Mayer, Annu. Rev. Phys. Chem. 2004, 55, 363–390; b) J. J.
 Warren, T. A. Tronic, J. M. Mayer, Chem. Rev. 2010, 110, 6961–7001;
 c) J. M. Mayer, Acc. Chem. Res. 2011, 44, 36–46; d) J. W. Darcy, B.
 Koronkiewicz, G. A. Parada, J. M. Mayer, Acc. Chem. Res. 2018, 31, 2391–2399; e) J. W. Darcy, B. Koronkiewicz, G. A. Parada, J. M. Mayer, Acc. Chem. Res. 2018, 31, 2391–2399.
- C. Costentin, M. Robert, J.-M. Savéant, Chem. Rev. 2008, 108, 2145– 2179; Update 1: Chem. Rev. 2010, 110, PR1–PR40.
- [6] J. C. Lennox, D. A. Kurtz, T. Huang, J. L. Dempsey, ACS Energy Lett. 2017, 2, 1246–1256.
- [7] C. Costentin, M. Robert, J.-M. Savéant, Chem. Rev. 2008, 108, 2145– 2179; Chem. Rev. 2010, 110, PR1–PR40.
- [8] N. Elgrishi, B. D. McCarthy, E. S. Rountree, J. L. Dempsey, ACS Catal.
 2016, 6, 3644–3659.
- [9] S. Y. Reece, D. G. Nocera, Annu. Rev. Biochem. 2009, 78, 673–699.
- [10] S. Y. Reece, J. M. Hodgkiss, J. Stubbe, D. G. Nocera, *Phil. Trans. R.* Soc. B 2006, 361, 1351–1364.
- [11] E. C. Gentry, R. R. Knowles, Acc. Chem. Res. 2016, 49, 1546–1556.
- [12] D. C. Miller, K. T. Tarantino, R. R. Knowles, *Top. Curr. Chem. (Z)* 2016, 374:30.
- [13] H. G. Yayla, R. R. Knowles, Synlett 2014, 25, 2819–2826.
- [14] A. Pannwitz, O. S. Wenger, J. Am. Chem. Soc. 2017, 139, 13308–13311.
- [15] A. Pannwitz, O. S. Wenger, Chem. Commun. 2019, 55, 4004–4014.
- [16] C. Costentin, M. Robert, J.-M. Savéant, C. Tard, Angew. Chem. 2010, 122, 3891–3894; Angew. Chem. Int. Ed. 2010, 49, 3803–3806.
- E. Odella, S. J. Mora, B. L. Wadsworth, M. T. Huynh, J. J. Goings, P. A. Liddell, T. L. Groy, M. Gervaldo, L. E. Sereno, D. Gust, T. A. Moore, G. F. Moore, S. Hammes-Schiffer, A. L. Moore, *J. Am. Chem. Soc.* 2018, 140, 15450–15460.
- [18] H.-J. Himmel, Synlett 2018, 29, 1957–1977.
- [19] A. Peters, E. Kaifer, H.-J. Himmel, Eur. J. Org. Chem. 2008, 5907–5914.
- [20] B. Eberle, O. Hübner, A. Ziesak, E. Kaifer, H.-J. Himmel, *Chem. Eur. J.* 2015, *21*, 8578–8590.
- [21] C. Trumm, O. Hübner, P. Walter, S. Leingang, U. Wild, E. Kaifer, B. Eberle, H.-J. Himmel, *Eur. J. Inorg. Chem.* 2014, 6039–6050.

- [22] B. Eberle, M. Damjanovic, M. Enders, S. Leingang, J. Pfisterer, C. Krämer, O. Hübner, E. Kaifer, H.-J. Himmel, *Inorg. Chem.* 2016, 55, 1683–1696.
- [23] U. Wild, S. Federle, A. Wagner, E. Kaifer, H.-J. Himmel, Chem. Eur. J. 2016, 22, 11971–11976.
- [24] U. Wild, O. Hübner, L. Greb, M. Enders, E. Kaifer, H.-J. Himmel, Eur. J. Org. Chem. 2018, 5910–5915.
- [25] B. Eberle, H. Herrmann, E. Kaifer, H.-J. Himmel, Eur. J. Inorg. Chem. 2013, 3671–3679.
- [26] U. Wild, F. Schön, H.-J. Himmel, Angew. Chem. 2017, 129, 16630– 16633; Angew. Chem. Int. Ed. 2017, 56, 16410–16413.
- [27] F. Schön, E. Kaifer, H.-J. Himmel, *Chem. Eur. J.* **2019**, *25*, 8279–8288.
- [28] U. Wild, O. Hübner, H.-J. Himmel, Chem. Eur. J. 2019, 25, 15988–15992.
- [29] U. Wild, O. Hübner, L. Greb, M. Enders, E. Kaifer, H.-J. Himmel, *Eur. J. Org. Chem.* 2018, 5910–5915.
- [30] J. F. Ambrose, R. F. Nelson, J. Electrochem. Soc. 1968, 115, 1159– 1164.
- [31] N. G. Connelly, W. E. Geiger, Chem. Rev. 1996, 96, 877-810.
- [32] S. Mallick, S. Maddala, K. Kollimalayan, P. Venkatakrishnan, J. Org. Chem. 2019, 84, 73–93.
- [33] For a related work on aryl-aryl coupling reactions of arylamines, see: S. Maddala, S. Mallick, P. Venkatakrishnan, J. Org. Chem. 2017, 82, 8958– 8972.
- [34] L. Zhai, R. Shukla, R. Rathore, Org. Lett. 2009, 11, 3474–3477.
- [35] L. Zhai, R. Shukla, S. H. Wadumethrige, R. Rathore, J. Org. Chem. 2010, 75, 4748–4760.
- [36] X.-Q. Zhu, H.-R. Li, Q. Li, T. Ai, J.-Y. Lu, Y. Yang, J.-P. Cheng, *Chem. Eur. J.* 2003, *9*, 871–880.
- [37] J.-P. Cheng, Y. Lu, X. Zhu, L. Mu, J. Org. Chem. 1998, 63, 6108–6114.
- [38] R. Billmers, R. L. Brown, S. E. Stein, Int. J. Chem. Kinetics 1989, 21, 375–386.
- [39] C. C. Johnston, J. L. Gardner, C. H. Suelter, D. E. Metzler, *Biochem.* 1963, 2, 689–696.
- [40] J. Hornung, O. Hübner, E. Kaifer, H.-J. Himmel, RSC Adv. 2016, 6, 39323–39329.
- [41] N. P. van Leest, L. Grooten, J. I. van der Vlugt, B. de Bruin, *Chem. Eur. J.* 2019, 25, 5987–5993.
- [42] S. Fukuzumi, H. Kotani, Y.-M. Lee, W. Nam, J. Am. Chem. Soc. 2008, 130, 15134–15142.
- [43] S. Fukuzumi, Y. Tokuda, T. Kitano, T. Okamoto, J. Otera, J. Am. Chem. Soc. 1993, 115, 8960–8968.
- [44] J. Yuasa, S. Yamada, S. Fukuzumi, J. Am. Chem. Soc. 2008, 130, 5808– 5820.
- [45] A. Peters, H. Herrmann, M. Magg, E. Kaifer, H.-J. Himmel, *Eur. J. Inorg. Chem.* 2012, 1620–1631.

FULL PAPER

Entry for the Table of Contents



Redox-active guanidines are used in PCET reactions that involve C-H bond cleavage, disclosing the scope and the mechanism of PCET reactions with and without addition of a strong acid. Substrates with potentials of up to 1.2 V vs. ferrocenium/ferrocene could be oxidized in stepwise PCET processes initiated by electron transfer. Experiments with different guanidines show the possibility to tune the PCET reactivity.