This article was downloaded by: [The University Of Melbourne Libraries] On: 05 June 2013, At: 08:50 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Synthetic Communications: An International Journal for Rapid Communication of Synthetic Organic Chemistry

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/lsyc20

# An Improved Procedure for the Large Scale Preparation of 2-Chloro-4,6-dimethoxy-1,3,5triazine

Jason S. Cronin<sup>a</sup>, Francis O. Ginah<sup>a</sup>, Angela R. Murray<sup>a</sup> & James D. Copp<sup>a</sup>

<sup>a</sup> Lilly Research Laboratories, Chemical Process Research and Development, A Division of Eli Lilly & Company, Indianapolis, IN, 46285-4813 Published online: 21 Aug 2006.

To cite this article: Jason S. Cronin , Francis O. Ginah , Angela R. Murray & James D. Copp (1996): An Improved Procedure for the Large Scale Preparation of 2-Chloro-4,6-dimethoxy-1,3,5-triazine, Synthetic Communications: An International Journal for Rapid Communication of Synthetic Organic Chemistry, 26:18, 3491-3494

To link to this article: http://dx.doi.org/10.1080/00397919608003754

## PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <u>http://www.tandfonline.com/page/terms-and-conditions</u>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

### AN IMPROVED PROCEDURE FOR THE LARGE SCALE PREPARATION OF 2-CHLORO-4,6-DIMETHOXY-1,3,5-TRIAZINE

Jason S. Cronin, Francis O. Ginah, Angela R. Murray, James D. Copp\*

Chemical Process Research and Development, Lilly Research Laboratories, A Division of Eli Lilly & Company, Indianapolis, IN 46285-4813

Abstract: A robust process for the preparation of multikilogram quantities of 2chloro-4,6-dimethoxy-1,3,5-triazine (1) is described.

Recent advances in the preparation of esters, amides, anhydrides and peptides have utilized 2-chloro-4,6-dimethoxy-1,3,5-triazine 1 as a carboxylate activating agent<sup>1</sup>. Racemization of the position  $\alpha$  to the carboxylate functionality is minimized with the use of 1, thereby making it especially useful for peptide synthesis and other chiral syntheses<sup>1</sup>. The preparation of 1 from cyanuric chloride and methanol has been previously described in the literature<sup>2</sup>. The reaction utilizes methanol and water as the solvent system with 2 equivalents of sodium bicarbonate as the hydrogen chloride scavenger (Scheme 1)<sup>3</sup>. The reaction temperature is controlled at 30°C. In our hands, the application of this protocol to the production of kilogram quantities of 1 was not successful.

\*To whom correspondence should be addressed. Internet: Jim@ Lilly.Com

Copyright © 1996 by Marcel Dekker, Inc.



Scheme 1

Due to the requirement for kilogram quantities of 1, we required a robust synthesis. In our studies of the reaction, we observed the production of the trimethoxytriazine 2 as the reaction neared completion. Analysis using reaction calorimetry indicated a significant exotherm was associated with the conversion of the 1 to 2. Additional experimentation demonstrated that the conversion of 1 to 2 was facilitated by higher temperatures. Given the concern of temperature control ability at a larger scale, we investigated alternative reaction conditions which might provide for a more robust process.

We discovered that the addition of 3 equivalents of sodium bicarbonate to the reaction mixture significantly decreased the conversion of 1 to 2 and eliminated the observed exotherm at the end of the reaction in which only 2 equivalents of sodium bicarbonate were used. Table 1 is a comparison of the results obtained from the two reaction procedures<sup>4</sup>. The data in Table 1 suggest

entry	equivalents	reaction	% 2 30 minutes	time for
	of base	temperature	after endpoint <sup>5</sup>	conversion of 1
				to <b>2</b>
1	2.0	30° C	>10%	100% @(2 hr)
2	2.0	40° C	>20%	100% @(1 hr)
3	3.0	30° C	1%	3% @ (12 hr)
4	3.0	40° C	2%	9% @ 12 hr

 Table 1. Production of 2,4,6-Trimethoxy-1,3,5-triazine 2 as a Function of Reaction Conditions

that the use of an additional equivalent of base provides a good measure of robustness to the synthesis. Only 9% of the trimethoxy derivative 2 was formed at the higher temperature after 12 hours compared to quantitative conversion of 1 to 2 in 1 hour under the same conditions with 2 equivalents of base. Application of the protocol using 3 equivalents of sodium bicarbonate yields 1 in 65% yield. The procedure improves the yields of 1 and results in less conversion of 1 to 2 as the reaction approaches completion. This protocol has been run successfully on a scale of 20 Kg.

### Experimental

A mixture of 12.3 equivalents of methanol, 2.7 equivalents of water and 3 equivalents of sodium bicarbonate is cooled to 10 to  $15^{\circ}$  C. Cyanuric chloride is added and the resulting mixture is warmed to  $35^{\circ}$  C<sup>6</sup>. The reaction mixture is stirred until the intermediate 2,4-dichloro-6-methoxy-1,3,5-triazine is no longer present (about 12 hours as demonstrated by gas chromatographic analysis<sup>4</sup>). Water (5 volumes) is added and the resulting mixture stirred for 30 minutes. The product is isolated by filtration and the filter cake is washed with deionized water. The wet cake is dried under vacuum at 30° C to give a 65% yield of 1 as white crystals: mp 72 - 74.6 (lit<sup>2b</sup> 74.2 - 76.2); Anal. Calcd for C5H6N3O2Cl<sub>1</sub>: C, 34.21; H, 3.45; N, 23.93. Found: C, 34.47; H, 3.72; N, 23.68.

Acknowledgements: We express our appreciation to Tim Lonis for the reaction calorimetry data generated for this reaction.

#### **References and notes**

 Kaminski, Z.J. *Tetrahedron Lett.* 1985, *26*, 2901; Cook, G.K., Hornback, W.J., Jordan, C.L., McDonald III, J.H., Munroe, J.E. *J. Org. Chem.* 1989, *54*, 5828; Taylor, E.C., Gillespie, P. *J. Org. Chem.* 1992, *57*, 5757; Barnett, C.J., Wilson, T.M., Wendel, S.R., Winningham, M.J., Deeter, J.B. *J. Org. Chem.* 1994, *59*, 7038; Hipskind, P.A., Howbert, J.J., Cho, S., Cronin, J.S., Fort, S.L., Ginah, F.O., Hansen, G.J., Huff, B.E., Lobb, K.L., Martinelli, M.J., Murray, A.R., Nixon, J.A., Staszak, M.A., Copp, J.D. *J. Org. Chem.* 1995, *60*, 7033.

2. (a) Dudley, J.R., Thuyrston, J.T., Schaefer, F.C., Holm-Hansen, D., Hull, C.J., Adams, P. J. Am. Chem. Soc. 1951, 73, 2986; (b) Weber, A.J.M., Huysmans, W.G.B., Mijs, W.J., Bovee, W.M.M.J., Smidt, J., Vriend, J. Recl. Trav. Chim. *Pays-Bas* **1978**, *97*, 107, (c) Menicagli, R., Malanga, C., Peluso, P. Synth. *Commun.* **1994**, *24*, 2153.

3. Potassium bicarbonate functioned equally well in the reaction. Other bases such as sodium, potassium, and cesium carbonate gave increased levels of 2 and/or lower yields of 1.

4. The reaction mixture was analysed using a Hewitt Packard 5980 capillary GC with an DB-1 crosslinked methyl-silicon column,  $30m \times 0.25mm \times -.25\mu m$ , at a column temperature of  $250^{\circ}$  C and flow rate of 1.7 mL/min. The retention times are as follows: 2,4-dichloro-6-methoxy-1,3,5-triazine: 2.5 min; 2-chloro-4,6-dimethoxy-1,3,5-triazine: 3.0 min; 2,4,6-trimethoxy-1,3,5-triazine: 3.4 min.

5. The endpoint of this reaction is defined as the time at which >97% conversion of the intermediate 2,4-dichloro-6-methoxy-1,3,5-triazine to CDMT has occurred. There is no residual cyanuric chloride.

6. The conversion of cyanuric chloride to 2,4-dichloro-6-methoxy-1,3,5-triazine is accompanied by an exotherm. This appears to start at around 20° C and facilitates the warm-up to 35° C.

(Received in the USA 31 March 1996)