

Cu(I)-Catalyzed Intramolecular Hydroamination of Unactivated Alkenes Bearing a Primary or Secondary Amino Group in Alcoholic Solvents


Hirohisa Ohmiya, Toshimitsu Moriya, and Masaya Sawamura*

Department of Chemistry, Faculty of Science, Hokkaido University,
Sapporo 060-0810, Japan

sawamura@sci.hokudai.ac.jp

Received April 9, 2009

ABSTRACT



The Cu-Xantphos system [Cu(O-*t*-Bu)-Xantphos, 10–15 mol %] catalyzes the intramolecular hydroamination of unactivated terminal alkenes bearing an unprotected aminoalkyl substituent in alcoholic solvents, giving pyrrolidine and piperidine derivatives in excellent yields. This system is applicable to both primary and secondary amines and tolerates a variety of functional groups.

Metal-catalyzed intramolecular hydroamination of alkenes is one of the simplest methods for the preparation of nitrogen heterocycles.¹ Recent research in the intramolecular hydroamination of alkenes has pursued the establishment of a universal hydroamination methodology. However, the de-

velopment of a catalytic system applicable to the addition of primary or secondary amines to unactivated alkenes remains an interesting challenge. Various catalysts that involve different metals have been developed. Lanthanide and group 4 metal complexes catalyze the hydroamination with superb efficiency.² However, the poor functional group compatibility and high air- and moisture-sensitivity of these metal complexes have limited their synthetic utility. Recently, some complexes incorporating precious metals such as Rh,^{3a,e} Pt,^{3b,c} and Ir^{3d,e} have been introduced as catalysts for the intramolecular hydroamination of unactivated alkenes bearing an amino group. Although these catalysts exhibited excellent

(1) For reviews on the hydroamination of alkenes and alkynes, see: (a) Müller, T. E.; Beller, M. *Chem. Rev.* **1998**, *98*, 675–703. (b) Müller, T. E.; Hultsch, K. C.; Yus, M.; Foubelo, F.; Tada, M. *Chem. Rev.* **2008**, *108*, 3795–3892.

(2) For selected papers on intramolecular hydroaminations of alkenes catalyzed by lanthanide and group 4 metal complexes, see: (a) Hong, S.; Marks, T. J. *Acc. Chem. Res.* **2004**, *37*, 673–686. (b) Stubbert, B. D.; Marks, T. J. *J. Am. Chem. Soc.* **2007**, *129*, 4253–4271. (c) Bexrud, J. A.; Beard, J. D.; Leitch, D. C.; Schafer, L. L. *Org. Lett.* **2005**, *7*, 1959–1962. (d) Gribov, D. V.; Hultsch, K. C. *Angew. Chem. Int. Ed.* **2004**, *43*, 5542–5546. See also ref 1.

(3) For late transition metal catalyzed intramolecular hydroamination of unactivated alkenes with alkylamines, see the following. Rh: (a) Liu, Z.; Hartwig, J. F. *J. Am. Chem. Soc.* **2008**, *130*, 1570–1571. Pt: (b) Bender, C. F.; Widenhoefer, R. A. *J. Am. Chem. Soc.* **2005**, *127*, 1070–1071. (c) Bender, C. F.; Hudson, W. B.; Widenhoefer, R. A. *Organometallics* **2008**, *27*, 2356–2358. Ir: (d) Hesp, K. D.; Stradiotto, M. *Org. Lett.* **2009**, *11*, 1449–1452. Ir and Rh: (e) Bauer, E. B.; Andavan, G. T. S.; Hollis, T. K.; Rubio, R. J.; Cho, J.; Kuchenbeiser, G. R.; Helgert, T. R.; Letko, C. S.; Tham, F. S. *Org. Lett.* **2008**, *10*, 1175–1178.

(4) For hydroaminations and hydroamidations of (activated) alkenes catalyzed by Cu or Fe, see the following. Cu: (a) Taylor, J. G.; Whittall, N.; Hii, K. K. *Org. Lett.* **2006**, *8*, 3561–3564. (b) Munro-Leighton, C.; Delp, S. A.; Blue, E. D.; Gunnoe, T. B. *Organometallics* **2007**, *26*, 1483–1493. (c) Munro-Leighton, C.; Delp, S. A.; Alsop, N. M.; Blue, E. D.; Gunnoe, T. B. *Chem. Commun.* **2008**, 111–113. Fe: (d) Komeyama, K.; Morimoto, T.; Takaki, K. *Angew. Chem., Int. Ed.* **2006**, *45*, 2938–2941.


(5) For Pd, see: (a) Michael, F. E.; Cochran, B. M. *J. Am. Chem. Soc.* **2006**, *128*, 4246–4247. For Au, see: (b) Zhang, J.; Yang, C.-G.; Chuan, C. *J. Am. Chem. Soc.* **2006**, *128*, 1798–1799. See also ref 1.

functional group compatibility, there still remains room for further investigation. Specifically, the development of hydroamination using a cheap and environmentally benign metal catalyst is greatly anticipated.⁴ To date, most of the investigations on the intramolecular *hydroamination* of alkenes catalyzed by late transition metals have focused on the use of amides or carbamates bearing electron-withdrawing *N*-substituents such as Ts, Cbz, Boc, or Ac groups instead of free amines.⁵

Here we report the intramolecular hydroamination of unactivated alkenes catalyzed by a Cu-phosphine system.^{4,6–8} Several aspects of this transformation are noteworthy: (i) Cu, which is a relatively abundant metal in the earth crust and hence cheap and environmentally benign, is used as the metal; (ii) alcoholic solvents are used;⁷ (iii) both secondary and primary amines can be used; (iv) not only pyrrolidine but also piperidine derivatives are obtained; (v) a variety of functional groups are tolerated; and (vi) olefinic side products due to β -hydride elimination or alkene isomerization are not formed, and the yields are generally high.

The reaction of ω -alkenic secondary amine **1aa** in the presence of Cu(O-*t*-Bu) (10 mol %) and Xantphos (10 mol %) in MeOH (1 mL) proceeded at 60 °C and afforded the pyrrolidine derivative **2aa** in 80% yield (97% NMR yield) after 18 h reaction time (Scheme 1).

Scheme 1. Cu(I)-Catalyzed Intramolecular Hydroamination of Aminoalkene **1aa**



The employment of the alcoholic solvent MeOH had a significant impact on the yield of **2aa**.⁹ The use of other solvents such as toluene, hexane, THF, dioxane, CH₃CN, acetone, DMF, and DMSO resulted in lower yields of **2aa** (0–19% NMR yields). The effectiveness of the protic solvent would imply a tolerance of this reaction toward a wide variety of functional groups. Furthermore, its crucial role in the rate acceleration is of considerable interest from a mechanistic viewpoint.^{7,8,10}

(6) For Cu-catalyzed enantioselective intramolecular carboamination and aminoxygenation of unactivated alkenes, see: (a) Zeng, W.; Chemler, S. R. *J. Am. Chem. Soc.* **2007**, *129*, 12948–12949. (b) Fuller, P. H.; Kim, J.-W.; Chemler, S. R. *J. Am. Chem. Soc.* **2008**, *130*, 17638–17639.

(7) For Cu-catalyzed addition of terminal alkynes to aldehydes with alcoholic solvents, see: (a) Asano, Y.; Hara, K.; Ito, H.; Sawamura, M. *Org. Lett.* **2007**, *9*, 3901–3904. (b) Asano, Y.; Hara, K.; Ito, H.; Sawamura, M. *Organometallics* **2008**, *27*, 5984–5996.

(8) For related studies from our group, see: (a) Ito, H.; Watanabe, A.; Sawamura, M. *Org. Lett.* **2005**, *7*, 1869–1871. (b) Ito, H.; Kawakami, C.; Sawamura, M. *J. Am. Chem. Soc.* **2005**, *127*, 16034–16035. (c) Ito, H.; Ito, S.; Sasaki, Y.; Matsuura, K.; Sawamura, M. *J. Am. Chem. Soc.* **2007**, *129*, 14856–14857. (d) Ito, H.; Kosaka, Y.; Nonoyama, K.; Sasaki, Y.; Sawamura, M. *Angew. Chem., Int. Ed.* **2008**, *47*, 7424–7427. (e) Ito, H.; Sasaki, Y.; Sawamura, M. *J. Am. Chem. Soc.* **2008**, *130*, 15774–15775.

(9) For details of the solvent effect, see Supporting Information.

(10) Yamamoto, Y.; Kirai, N.; Harada, Y. *Chem. Commun.* **2008**, 2010–2012.

Our ligand screening has identified Xantphos, which features an extraordinary large bite-angle, to be the best, while the ligand effect was moderate.¹¹ Even without the ligand, the hydroamination of **1aa** proceeded in 49% NMR yield under essentially the same conditions. Nitrogen-based ligands such as 1,10-phenanthroline reduced the activity of the catalyst (20% NMR yield).

Analogues (**1ab–ag**, **1ba–be**) of aminopentene **1aa** bearing different *N*-substituents were transformed into the corresponding pyrrolidine derivatives (**2ab–ag**, **2ba–be**) smoothly at higher temperatures (100 or 140 °C) in a mixed solvent (MeOH/toluene or MeOH/*o*-xylene) (Table 1, entries

Table 1. Cu(I)-Catalyzed Hydroamination of Unactivated Alkenes Bearing Secondary Amino or Amido Groups^a

entry	aminoalkene	product	temp (°C)	time (h)	yield ^b (%)
1	R = Et (1ab)	2ab	100	48	87
2	R = <i>n</i> -Pr (1ac)	2ac	100	48	93
3	R = <i>i</i> -Bu (1ad)	2ad	100	48	88
4	R = (CH ₂) ₂ Ph (1ae)	2ae	100	48	87
5	R = COCH ₃ (1af)	2af	100	48	96 ^d
6	R = COPh (1ag)	2ag	140	48	94
7 ^c	R (FG)	2ba	100	48	99
8 ^c	R = H (1ba) FG = OMe (1bb)	2bb	100	48	94
9 ^c	FG = F (1bc)	2bc	100	48	98
10 ^c	FG = CN (1bd)	2bd	100	48	86
11 ^c	FG = CO ₂ Me (1be)	2be	100	48	86
12	1c	2c	100	4	91
13 ^c	1d	2d	140	48	89
14	1e	2e	100	24	92
15	1f	2f	100	24	94

^a Conditions: Cu(O-*t*-Bu)-Xantphos (10 mol %), **1** (0.5 mmol), MeOH/toluene (1:1, 1.0 mL), 100 °C or MeOH/*o*-xylene (1:1, 1.0 mL), 140 °C. ^b Isolated yield. ^c Conditions: Cu(O-*t*-Bu)-Xantphos (15 mol %), **1** (0.5 mmol), MeOH/toluene (1:1, 1.0 mL), 100 °C or MeOH/*o*-xylene (1:1, 1.0 mL), 140 °C. ^d The isolated product was contaminated with a small amount of **1af** (5%).

1–11). Thus, alkyl substituents such as Et, *n*-Pr, *i*-Bu, and 2-phenethyl groups were tolerated at the nitrogen atom (entries 1–4). Amide substrates **1af** and **1ag**, bearing electron-withdrawing *N*-substituents such as acetyl or benzoyl groups, also underwent hydroamidation in high yields (entries 5 and 6). Although the aminopentene (**1ba**) with an *N*-benzyl group was less reactive than **1aa–ae**, the reaction of this substrate also proceeded smoothly and led to completion with 15 mol % catalyst loading (entry 7). Notably, functionalities such as methoxy, fluoro, cyano, and ester were tolerated on the aromatic ring of the *N*-benzyl group (reactions of **1bb–be**, entries 8–11).

(11) DPPF was as effective as Xantphos at 60 °C, while slightly less effective at 40 °C. For details of the ligand effect, see Supporting Information.

The diphenylmethylene insert of the aminoalkene **1aa** could be replaced with a cyclohexane-1,1-diyl group without affecting the reactivity toward the hydroamination (reaction of **1c**, Table 1, entry 12). Unfortunately, however, the substrate without substituents in the linker chain, where the Thorpe–Ingold effect does not work, resisted the reaction.

The 1,1-disubstituted alkene **1d** also underwent hydroamination under more forcing conditions (15 mol % of Cu, 140 °C) (Table 1, entry 13). Conversely, the addition to a 1,2-disubstituted alkene did not proceed at all.

The extension of the Cu-catalyzed hydroamination from a five- to six-membered ring formation was feasible, as was the case with the other catalyst systems reported previously. Thus, the reaction of 1-amino-5-hexene derivatives **1e** and **1f** occurred within 24 h at 100 °C to give the corresponding piperidine derivatives (**2e** and **2f**) in excellent yields (Table 1, entries 14 and 15).¹

The Cu catalyst system was also effective for the reaction of ω -alkenic primary amines (**1g–j**) (Table 2). In fact, the

Table 2. Cu(I)-Catalyzed Hydroamination with Primary Aminoalkenes **1g–j**^a


entry	aminoalkene	Cu (mol %)	product	yield (%) ^b
1		15		92
2		15		79 ^c
3		15		87
4		10		91

^a Conditions: Cu(O-*t*-Bu) (0.05–0.075 mmol), Xantphos (1 equiv to Cu), **1** (0.5 mmol), MeOH/toluene (1:1, 1.0 mL), 100 °C, 72 h. ^b Isolated yield. ^c The isolated yield was reduced by the volatility of the product.

reaction of **1g–j** to form five- and six-membered rings in the presence of 10 or 15 mol % of Cu(I)-Xantphos catalyst proceeded at 100 °C, delivering the corresponding NH-free cyclic secondary amines (**2g–j**) in high yields. It should be noted that the late transition metal catalyzed intramolecular hydroamination of unactivated alkenes bearing a primary amino group has been achieved in only the cases with the Rh catalyst,^{3a} whereas various catalysts have been used for the additions of amides and carbamates, where the nitrogen moieties require installation and removal of the electron-withdrawing group.⁴

A possible mechanism for the Cu-catalyzed intramolecular hydroamination is proposed in Scheme 2.^{7,8} The catalytic

Scheme 2. Proposed Mechanism



cycle would be initiated by σ -bond metathesis between alkoxy copper **A** and the H–N bond of the substrate (**1**) to form copper amide **B**. Subsequently, the terminal alkene coordinates to the Cu atom to form Cu-alkene π -complex **C**. Next, the addition of the Cu–N bond across the C–C double bond affords alkylcopper intermediate **D** with a nitrogen heterocycle. Finally, protonolysis of the Cu–C bond of **D** with MeOH gives the final product (**2**). According to this mechanism, the protonation of **D** should be much faster than the β -hydride elimination to form enamine **3** and copper hydride **E**.

In summary, the Cu-Xantphos system catalyzed the intramolecular hydroamination of unactivated terminal alkenes bearing an unprotected aminoalkyl substituent in alcoholic solvents, giving pyrrolidine and piperidine derivatives in excellent yields. This system is applicable to both primary and secondary amines and tolerates a variety of functional groups. We believe that this is a significant step toward economically advantageous and environmentally benign alkene hydroaminations. Further studies to find more efficient catalyst systems based on copper are underway.

Acknowledgment. This work was supported by Grants-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology, Government of Japan.

Supporting Information Available: Experimental procedures and NMR spectra for new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL9007712