

Fig. 1 Examples of binaphthalene based chiroptical switches.

—a change of conjugation to the binaphthalene and therefore a significant change to the chromophore (causing a shift of the CD band maximum, e.g. **2**¹⁰).

—a combination of both: a minor change to the binaphthalene dihedral angle as well as a slight change to the chromophore attached to binaphthalene (resulting in change of intensity of the corresponding CD bands, such as the (*E/Z*)-isomerisation of **3**¹¹).

—a change in the supramolecular interactions between attached groups (a change of the ICD signal of chromophores in a chiral environment, e.g. **4**¹²).

In this work, we opted to explore possible modulation of the CD signal resulting from a significant chromophore change upon switching. For this purpose, we have designed, prepared and investigated the photochemical behaviour of a binaphthalene derivative with co-condensed pyran rings, **5** (consisting of two *2H*-naphtho[2,3-*b*]pyran units).

Results and discussion

Synthesis

Various synthetic routes for the preparation of diarylnaphthopyrans have been described.¹⁴ The most general and effective method involves the acid-catalysed reaction of propargyl alcohols with naphthols in the presence of a dehydrating agent (such as CH(OMe)₃,¹⁴ molecular sieves¹⁵ or activated silica¹⁶). However, none of them resulted in the formation of even traces of the desired product starting from BINOL (**6**). We also attempted to synthesise compound **5a** by a stepwise protocol: (i) we prepared the bispropargyl ether of BINOL in the presence of copper(i) iodide, potassium iodide and potassium carbonate in 29% yield (18 h at 90 °C), and (ii) we then tried to close the pyran ring in the presence of *N,N*-diethylaniline, in analogy with a previously described procedure.¹⁷ Even this attempt failed.

Therefore, we decided to introduce allyl alcohol **7** to positions 3 and 3' of BINOL prior to cyclisation of the pyran ring (Scheme 2). The 3,3'-diiodo precursor required for this approach, **8**, was prepared from (*R*)-BINOL (**6**) in three steps using previously described conditions: protection of the hydroxy groups as methyl ethers,¹⁸ bis-*ortho*-lithiation and iodination,¹⁹ and deprotection.²⁰ Allyl alcohol **7** was effectively (82% yield) introduced by Heck coupling using

palladium acetate and potassium carbonate in acetonitrile. The obtained open-chain intermediate, **9**, was then successfully (63% yield) cyclised using activated silica in boiling toluene. Thus, the desired product, **5a**, was prepared from (*R*)-BINOL (**9**) in five steps in a 31% overall yield.


Photochromism in solution

The electronic absorption spectrum of **5a** in deuterated benzene shows two intense high energy absorption bands at 298 and 308 nm, and two minor low energy absorption bands at 354 and 370 nm (Fig. 2). The photochromic behaviour of **5** was evaluated by the continuous irradiation of (*R*)-**5a** in C₆D₆ under argon with monochromatic light at 313 nm filtered from a 125 W high pressure mercury arc. After initial irradiation, a partial increase of the high energy band absorptions was observed. However, subsequent bleaching of these bands and the appearance of a new broad band at 336 nm (the formation of methyldene quinone chromophore) was detected upon prolonged irradiation as a result of electrocyclic pyran ring-opening (Fig. 2). Colour of the sample turned yellowish.

Photochemical pyran ring-opening was found to be relatively slow, as judged by HPLC monitoring of the irradiated mixture (50% conversion of (*R*)-**5a** was reached after 79 min at 26.5 °C). Due to the required long reaction time and the partial decomposition of the reaction mixture (as observed by HPLC), it became difficult to reach the photostationary state, but this was estimated to contain 30% of 'closed' form (*R*)-**5a**.

Pure 'open' form (*R*)-**5b** was isolated by flash chromatography from a mixture irradiated for 155 min, keeping the solution ice cold during the separation. The number and character of the signals in ¹H and ¹³C NMR spectra of **5b**, corresponding to C₁ symmetry, revealed that ring-opening of only one pyran ring took place (Scheme 3). The open chain propenylidene arm of **5b** has a prevailing *s-trans* conformation according to the spin-spin coupling constant between its inner protons (9.8 Hz). Due to overlap of the signals in the ¹H NMR spectrum, the (*E/Z*)-configuration of the exocyclic double bond could not be determined from NOE experiments.

The ring-closure activation barrier of 74.5 ± 5.8 kJ mol⁻¹ was calculated from the bleaching rate constants determined at various temperatures (Fig. 3) using the irradiated mixtures of



Scheme 2 Synthesis of (*R*)-5. Reagents and conditions: (i) CH_3I (9 equiv.), K_2CO_3 (3 equiv.), acetone, reflux, 36 h, 98% yield; (ii) (1) *n*-BuLi (3.8 equiv.), TMEDA (3.2 equiv.), diethyl ether, r.t., 4 h, (2) I_2 (3.8 equiv.), -78°C to r.t., 5 h, 63% yield; (iii) BBr_3 (5 equiv.), CH_2Cl_2 , 0°C , 4 h, 97% yield; (iv) **7** (5.2 equiv.), $\text{Pd}(\text{OAc})_2$ (8 mol%), K_2CO_3 (1 equiv.), CH_3CN , 100°C , 4 h, 82% yield; (v) silica, toluene, reflux, 24 h, 63% yield.




Fig. 2 The absorption spectra of (*R*)-5 in benzene (3.5×10^{-5} M) after irradiation with UV light (313 nm) for the given time. Concentration of **5** in C_6D_6 during irradiation: 1.52×10^{-2} M.





Fig. 3 Determination of the ring-closure activation barrier from the dependence of $\ln k$ (**5b** ring-closure rate constants) against $1/T$.



Scheme 3 Reversible switching of binaphthopyran (*R*)-5.

(*R*)-5a and (*R*)-5b. The time dependent composition of the mixtures was analysed by HPLC.

With the aim of designing conditions for the reversible switching of (*R*)-5 in d_6 -benzene, the solution was irradiated with UV light at 313 nm for only 5 min (6.2% conversion of **5a**) to avoid potential decomposition. Then, the solution was heated under reflux for 5 min. Each time, a small sample was taken and diluted with benzene to record electronic absorption spectra. As shown in Fig. 4, a certain level of switching was observed, but serious fatigue occurred within a few cycles.

The changes in the CD spectrum of (*R*)-5 in deuterated benzene upon irradiation and subsequent heating were examined (Fig. 5) under the conditions described above. The overlaid CD spectra do not contain clear isosbestic points (except for one at 386 nm).

This indicates a more complex character to the transformation, including the formation of by-products or more stereoisomers in a variable ratio. Moreover, the intensity




Fig. 4 Switching of the absorption intensity of (*R*)-5 (sample irradiated/heated in C_6D_6 at $c = 1.52 \times 10^{-2}$ M, diluted by benzene to $c = 3.5 \times 10^{-5}$ M for the measurement of absorbance) at 298 nm. The increase of the intensity after irradiation with UV light (313 nm) for 5 min and the decrease after heating (84°C) for 5 min.

changes of the major bands do not exhibit a reversible character, but their intensity gradually increases (the bisignate CD couplet shifted to a higher wavelength compared to simple binaphthalene derivatives,²¹ here with a negative branch at 278 nm and a positive branch at 307 nm) or diminishes (negative band at 392 nm).

Photochromism in polymer films

For their potential application in memory devices, molecular switches are expected to be used at solid surfaces. Therefore,




Fig. 5 The CD spectra of (*R*)-**5** in C_6D_6 ($c = 1.52 \times 10^{-2}$ M) before irradiation (black), after the 1st irradiation with UV light (313 nm) for 5 min (red), after the 1st heating (84 °C) for 5 min (blue), after the 2nd irradiation with UV light (313 nm) for 5 min (orange) and after the 2nd heating (84 °C) for 5 min (green).

we have prepared thin films of (*R*)-**5a** in poly(methyl methacrylate) (PMMA) and polystyrene (PS). After some experimentation, we succeeded in finding suitable conditions for irradiation and heating that kept the switching fully reversible (Fig. 6). An improved durability of switch (*R*)-**5a** after physical imprisonment in the polymer matrix could be caused by reduced side stereoconversions and interactions with oxygen and/or the recombination of activated species.

The changes in the absorbance spectrum of (*R*)-**5a** in PMMA and PS films upon switching did not exceed 9% of the intensity at the band's maximum (ESI, Fig. S22†). Interestingly, the CD spectrum of (*R*)-**5** in polymer films under the same conditions (Fig. 7) differed significantly in the intensity of the bisignate CD couplet (negative branch at 282 nm, positive branch at 308 nm). The higher amplitude CD signal in the PS film most probably results from π - π stacking interactions between switched (*R*)-**5** and the aromatic environment in the PS matrix. This interaction could lead to amplification of the chiroptical response²² due to reversibly-enhanced helicity in the film. As a result, over a six-fold increase in the CD signal intensity of the (*R*)-**5**/PS film at




Fig. 6 Switching of the absorbance intensity of (*R*)-**5** in a 50 μ m PMMA film (0.02 mol kg^{-1}) at 296 nm. The increase of the intensity after irradiation with UV light (313 nm) for 1 h and the decrease after heating (70 °C) for 8 min.




Fig. 7 The CD spectra of (*R*)-**5** in a 50 μ m PS film (solid lines) and a PMMA film (dashed lines), $c = 0.02 \text{ mol kg}^{-1}$: before irradiation (blue), after irradiation with UV light (313 nm) for 1 h (green) and after heating (70 °C) for 8 min (red).

308 nm could be maintained during multiple reversible switchings (Fig. 8). The detected amplitude of the CD signal intensity (108 ± 8 mdeg at 308 nm) is one of the largest observed for binaphthalene-based chiroptical switches so far seen.^{9–13}

Conclusions

In this paper, we reported the synthesis and characterisation of chiroptical molecular switch **5**, consisting of two 2*H*-naphtho[2,3-*b*]pyran units. Previously reported synthetic methods described for the preparation of naphthopyrans from naphthols did not yield the desired binaphthopyran, **5a**, from BINOL. The synthesis of **5a**, therefore, required the building up of the pyran rings through a Heck coupling of diiodo precursor **8** with allyl alcohol **7**, followed by a silica-assisted cyclisation.

The irradiation of **5a** with UV light at 313 nm caused a slow photochemical opening ($t_{1/2} = 79$ min) of only one of the pyran rings. Reverse ring-closure occurred spontaneously with an activation barrier of 74 kJ mol^{-1} . However, the progressive stereoisomerisation and/or fading of switch **5** resulted in an




Fig. 8 Switching of the CD signal intensity of (*R*)-**5** in a 50 μ m PS film (0.02 mol kg^{-1}) at 308 nm. The increase of the intensity after irradiation with UV light (313 nm) for 1 h and the decrease after heating (70 °C) for 8 min.

almost unidirectional increase/decrease of CD signal intensities upon switching in solution.

The imprisonment of **5** in a polymer film significantly improved its fatigue resistance, allowing multiple reversible switchings to be maintained. In addition, the interaction of **5** with a polystyrene film amplified its chiroptical response, as detected by increased CD signal amplitudes up to 108 mdeg during the switching (over 6:1 signal ratio of **5b/5a**).

Experimental

General

Flash column chromatography was performed on Merck silica gel (60H). Merck silica gel F254 plates were used for thin layer chromatography, and visualisation was effected using UV light (254 nm). Melting points were measured on a Electrothermal-IA9200 instrument and the values are uncorrected. Specific optical rotations were measured on a Perkin-Elmer 241 polarimeter and are given in $\text{deg cm}^2 \text{g}^{-1} \text{dm}^{-1}$. IR data (in cm^{-1}) were recorded on a SPECORD M 80 spectrophotometer. UV-vis spectra in solution were measured on a Hewlett Packard diode array 8245 spectrophotometer. Thin films were measured on a Shimadzu spectrometer UV 1650PC. CD spectra were measured on a Jasco J810 instrument. ^1H and ^{13}C NMR spectra were recorded on a Varian Gemini 300 instrument at 298 K. Chemical shifts are reported in ppm downfield of the internal standard Me_4Si (0.00 ppm), and the solvent was also used as a reference. The working frequency was 300 MHz for ^1H and 75.5 MHz for ^{13}C NMR. Coupling constants (J) are given in Hz. GC-MS spectra (70 eV, 150 μA , EI) were recorded on a Voyager GC-MS Finnigan instrument. Elemental analyses were determined using an Erba Science 1106 instrument. HPLC analyses were undertaken on a Chiralcel Daicel OD-H column (eluent: n-hexane-isopropyl alcohol 90/10, temperature: 21 $^\circ\text{C}$, flow: 0.75 mL min^{-1} , pressure: 27.5 bar, UV detector: $\lambda = 275 \text{ nm}$) and Waters Symmetry[®] C₁₈ 5 μm 3.9 \times 150 mm column (eluent: methanol-water 95/5, temperature: 21 $^\circ\text{C}$, flow: 0.25 mL min^{-1} , pressure: 27 bar, UV detector: $\lambda = 254 \text{ nm}$) using a UV-vis detector LCD 5000.

All chemicals were used as purchased if not stated otherwise. All reactions were carried out under an argon atmosphere using Schlenk techniques. Dichloromethane and acetonitrile were dried over calcium hydride and chloroform over calcium chloride. Toluene, diethyl ether and d_6 -benzene were dried over sodium/benzophenone and distilled over argon. The following compounds were prepared according to published procedures: $\text{CuCl}(\text{OH})\text{TMEDA}$,²³ (8*S*,9*R*)-(-)-*N*-benzylcinchonidinium chloride,²⁴ (*R*)-**6**,²⁵ (*R*)-2,2'-dimethoxy-1,1'-binaphthalene,¹⁸ (*R*)-3,3'-diiodo-2,2'-dimethoxy-1,1'-binaphthalene,¹⁹ **7**²⁶ and (*R*)-**8**.²⁰ All other chemicals were of analytical grade and were used without further purification.

Preparations

(*R*)-3,3'-Bis(3-hydroxy-3,3-diphenylprop-1-en-1-yl)-1,1'-binaphthalene-2,2'-diol (**9**). To a solution of (*R*)-**8** (200 mg, 0.37 mmol) in acetonitrile (1 mL) were added Cs_2CO_3

(121 mg, 0.37 mmol), $\text{Pd}(\text{OAc})_2$ (7 mg, 0.03 mmol) and **7** (391 mg, 1.9 mmol). The reaction mixture was stirred under argon at 100 $^\circ\text{C}$ for 4 h, diluted with dichloromethane and finally washed with brine. The organic layer was dried over Na_2SO_4 , concentrated under vacuum and the residue purified by chromatography on silica gel using hexanes/ethyl acetate (5:1) as the eluent to give product (*R*)-**9** ($R_F = 0.19$, 213 mg, 82%) as a yellow solid.

Found: C, 85.20; H, 5.47. $\text{C}_{50}\text{H}_{38}\text{O}_4$ requires C, 85.44; H, 5.45%. $\nu_{\text{max}}(\text{KBr})/\text{cm}^{-1}$ 3507 (OH), 3057 and 3025 (H_{arom}), 1716 ($\text{C}=\text{C}_{\text{alkene}}$), 1621 and 1598 ($\text{C}=\text{C}_{\text{arom}}$), 1211 (OH_{arom}), 1147 (OH_{tert}); $\delta_{\text{H}}(300 \text{ MHz; DMSO-}d_6; \text{Me}_4\text{Si})$ 6.16 (2H, s, $2 \times \text{OH}_{\text{tert}}$), 6.74 (2H, d, $J = 8.2 \text{ Hz}$, C(8)*H*, C(8')*H*), 7.07–7.49 (28H, m), 7.89 (2H, d, $J = 8.0 \text{ Hz}$, C(5)*H*, C(5')*H*), 8.23 (2H, s, C(4)*H*, C(4')*H*), 8.30 (2H, s, $2 \times \text{OH}_{\text{arom}}$); $\delta_{\text{C}}(75 \text{ MHz; DMSO-}d_6; \text{Me}_4\text{Si})$ 78.6, 114.7, 123.4, 124.0, 124.3, 126.4, 126.5, 127.1, 127.3, 127.6, 128.3, 128.5, 129.1, 134.0, 138.0, 148.1, 152.3; HRMS (ESI⁺) m/z 667.2637 [$\text{M} + \text{H}^+ - 2\text{H}_2\text{O}$]. $\text{C}_{50}\text{H}_{35}\text{O}_2^+$ requires 667.2637; $[\alpha]_{\text{D}}^{18} = +38.8$ ($c = 1$ in CHCl_3); mp 129–131 $^\circ\text{C}$.

(*R*)-2,2,2',2'-Tetraphenyl-10,10'-bi(2*H*-naphtho[2,3-*b*]pyran) (**5a**). A mixture of (*R*)-**9** (100 mg, 0.14 mmol) in dry toluene (1 mL) and silica gel (300 mg, freshly activated at 120 $^\circ\text{C}$) was heated under reflux for 24 h and monitored by thin layer chromatography. The hot mixture was filtered off and the silica washed with hot toluene ($2 \times 0.5 \text{ mL}$). The filtrate was concentrated under vacuum and the residue purified by chromatography on silica gel using hexanes/ethyl acetate (16:1) as the eluent to give product (*R*)-**5a** ($R_F = 0.34$, 60 mg, 63%) as a white solid.

Found: C, 90.29; H, 5.18. $\text{C}_{50}\text{H}_{34}\text{O}_2$ requires C, 90.06; H, 5.14%. $\nu_{\text{max}}(\text{KBr})/\text{cm}^{-1}$ 3057 and 3032 (H_{arom}), 1624 ($\text{C}=\text{C}_{\text{arom}}$), 1250 (C–O); $\delta_{\text{H}}(300 \text{ MHz; DMSO-}d_6; \text{Me}_4\text{Si})$ 6.77–7.40 (30H, m), 7.84 (2H, s, C(5)*H*, C(5')*H*), 7.90 (2H, d, $J = 8.1 \text{ Hz}$, C(6)*H*, C(6')*H*); $\delta_{\text{C}}(75 \text{ MHz; DMSO-}d_6; \text{Me}_4\text{Si})$ 81.7, 116.9, 121.8, 123.2, 123.8, 125.0, 125.6, 125.7, 125.9, 126.2, 126.9, 127.0, 127.6, 127.9, 128.1, 128.9, 132.1, 133.4, 144.7, 144.9, 148.4; $[\alpha]_{\text{D}}^{18} = +8.2$ ($c = 1$ in CHCl_3); λ_{max} (benzene)/nm ($\epsilon/\text{dm}^3 \text{ mol}^{-1} \text{ cm}^{-1}$) 298 (30460), 308 (29873), 354 (3626), 370 (3445); mp 123–126 $^\circ\text{C}$.

(*R*)-1-(2,2-Diphenyl-2*H*-naphtho[2,3-*b*]pyran-10-yl)-3-(3,3-diphenylprop-2-enylidene)naphthalen-2(3*H*)-one (**5b**). (*R*)-**5a** (223 mg, 0.335 mmol) was dissolved in UV grade benzene (220 mL, $c = 1.52 \times 10^{-3} \text{ M}$), placed into the reactor under an argon atmosphere and irradiated at 313 nm for 155 min (HPLC: $t_{(R)\text{-5a}} = 22.47 \text{ min}$, 52.5%; $t_{(R)\text{-5b}} = 23.95 \text{ min}$, 47.5%). The solvent was evaporated under reduced pressure and (*R*)-**5b** separated from the residue by flash chromatography on silica gel using hexanes/ethyl acetate (20:1) as the eluent ($R_F = 0.26$). The mixture and product were kept ice cold during the separation. This gave product (*R*)-**5b** (87 mg, 39%) as a yellowish coloured solid.

$\delta_{\text{H}}(300 \text{ MHz; CDCl}_3; \text{Me}_4\text{Si})$ 6.42 (1H, d, $J = 9.8 \text{ Hz}$, $=\text{CH}-\text{CH}=\text{CPh}_2$), 6.90–7.41 (25H, m), 7.50–7.54 (4H, m), 7.67 (1H, s, C(5')*H*), 7.79 (2H, d, $J = 8.1 \text{ Hz}$), 8.04 (1H, s, C(4)*H*); $\delta_{\text{C}}(75 \text{ MHz; CDCl}_3; \text{Me}_4\text{Si})$ 81.5, 91.9, 111.8, 114.3, 114.3, 120.7, 121.6, 122.4, 122.8, 123.5, 123.9, 124.1, 125.1,

125.1, 125.3, 125.7, 125.9, 126.1, 126.3, 126.4, 126.4, 126.7, 126.7, 126.8, 126.8, 127.1, 127.1, 127.4, 127.5, 127.5, 127.6, 127.8, 128.7, 131.6, 132.3, 132.8, 135.3, 143.3, 143.3, 148.2, 148.3, 208.0.

Switching in solution

A sample of (*R*)-**5a** (14.2 mg, 2.2×10^{-5} mol) in *d*₆-benzene (1.4 mL, 1.52×10^{-2} M) was placed into the NMR tube under argon and irradiated at 313 nm for 5 min. Reversible switching process was performed by heating of the sample at 84 °C in water bath for appropriate time. For UV-vis measurements, calculated volume of the sample was diluted with benzene to $c = 3.5 \times 10^{-5}$ M. CD spectra of the whole samples were recorded.

Kinetic measurements

A sample of (*R*)-**5a** (7.1 mg, 1.1×10^{-5} mol) in *d*₆-benzene (0.7 mL, 1.52×10^{-2} M) was placed into an NMR tube under argon and irradiated at 313 nm for 40 min. Small amounts of the sample were taken after regular time periods for HPLC analysis to determine the composition of the irradiated mixture. The ratios of **5a** and **5b**, calculated from peak areas at 254 nm, were corrected by molar ratio found from the ¹H NMR spectrum. The data corresponded to apparent first-order kinetics with a rate constant $k = (8.77 \pm 0.25) \times 10^{-3} \text{ min}^{-1}$ and a reaction half time $t_{1/2} = 79.0 \pm 2.3 \text{ min}$ at 26.5 °C.

Three samples of (*R*)-**5a** (7.1 mg, 1.1×10^{-5} mol) in *d*₆-benzene (0.7 mL, 1.52×10^{-2} M) were placed in tubes under argon and irradiated at 313 nm for 45 min at 26.5 °C. Each sample was, after irradiation, tempered for 15 min to a different temperature (14, 26.5 and 35 °C). From each sample was taken a small amount after regular time periods (1, 0.5 and 0.25 h, respectively) for HPLC analysis to determine the composition of the mixture. The found rate constants were: $(0.89 \pm 0.11) \times 10^{-3} \text{ min}^{-1}$ (14.0 °C), $(3.73 \pm 0.41) \times 10^{-3} \text{ min}^{-1}$ (26.5 °C) and $(7.30 \pm 0.66) \times 10^{-3} \text{ min}^{-1}$ (35.0 °C). The activation barrier for the ring-closure of (*R*)-**5b** was calculated from the dependence of the natural logarithm of the rate constant on reciprocal temperature as $74.5 \pm 5.8 \text{ kJ mol}^{-1}$.

Switching in polymer film

Films of poly(methyl methacrylate) (PMMA; Diacon, ICI, England) and polystyrene (PS; Chemische Werke Huels, F. R. G.) were prepared by casting a 1 mL chloroform solution of the polymer (5 g/100 mL) containing the respective amount of (*R*)-**5a** onto a glass plate (28 × 35 mm). The solvent was evaporated slowly in the dark. The final concentration of (*R*)-**5a** in the films was 0.02 mol kg⁻¹. The thickness of the films was 50 μm.

The photoisomerisation of (*R*)-**5** was examined in thin polymer films in a merry-go-round set up equipped with a 125 W medium pressure mercury arc lamp and a chromate liquid filter²⁷ transmitting monochromatic light $\lambda = 313 \text{ nm}$. Thermoisomerisation was performed in an oven at 70 °C for an appropriate time. Electronic absorption and CD spectra of

the films nestled at the standard cell wall were recorded in the ordinary transmittance mode.

Acknowledgements

This work was supported by the Slovak Research and Development Agency (grant APVV-0128-07) and the Slovak Grant Agency for Science (grant no. 1/0243/09).

References

- (a) V. Balzani, A. Credi and M. Venturi, *Molecular Devices and Machines—A Journey into the Nano World*, Wiley-VCH, Weinheim, 2003; (b) *Molecular Switches*, ed. B. L. Feringa, Wiley-VCH, Weinheim, 2001; (c) Photochromism: Memories and Switches, guest ed. M. Irie, *Chem. Rev.*, 2000, **100**, 1683–1890; (d) *Optical Sensors and Switches*, ed. V. Ramamurthy and K. S. Schanze, Marcel Dekker, New York, 2001; (e) J. Cusido, E. Deniz and F. M. Raymo, *Eur. J. Org. Chem.*, 2009, 2031; (f) H. Tian and Y. Feng, *J. Mater. Chem.*, 2008, **18**, 1617; (g) H. Mustroph, M. Stollenwerk and V. Bressau, *Angew. Chem., Int. Ed.*, 2006, **45**, 2016.
- (a) Y. S. Lee, *Self-Assembly and Nanotechnology: A Force Balance Approach*, John Wiley & Sons, Hoboken, New Jersey, 2008; (b) *Self-Assembled Nanomaterials*, ed. I. T. Shimizu, Springer-Verlag, Berlin, 2008; (c) T. J. White, R. L. Bricker, L. V. Natarajan, N. V. Tabiryan, L. Green, Q. Li and T. J. Bunning, *Adv. Funct. Mater.*, 2009, **19**, 3484; (d) M. Mathews and N. Tamaoki, *J. Am. Chem. Soc.*, 2008, **130**, 11409.
- (a) N. Chopra, G. Iftime and P. M. Kazmaier, *US Pat.* 2007/720206, 2007; (b) N. Menke, B. Yao, Y. Wang, Y. Zheng, M. Lei, L. Ren, G. Chen, Y. Chen, M. Fan and T. Li, *J. Opt. Soc. Am. A*, 2006, **23**, 267; (c) P. M. Kazmaier and G. Iftime, *US Pat.* 2009/7564030, 2009; (d) N. Menke, B. Yao, Y. Wang, Y. Zheng, M. Lei, Z. Ren, G. Chen, Y. Chen, M. Fan and T. Li, *Proc. SPIE*, 2005, **5642**, 152; (e) H. Takahashi, S. Hirano and I. Kawashima, *US Pat.* 2009/7474322, 2009.
- (a) M. Granstrom, K. Petritsch, A. C. Arias, A. Lux, M. R. Andersson and R. H. Friend, *Nature*, 1998, **395**, 257; (b) L. M. Garcia, *US Pat.* 2007/7273659, 2007.
- (a) *Organic Photochromic and Thermochromic Compounds, Main Photochromic Families*, ed. J. C. Crano and R. J. Guglielmetti, Kluwer Academic Publishers, Dordrecht, 2002, vol. 1; (b) *Photochromism: Molecules and Systems*, ed. H. Durr, H. Bouas-Laurent and H. Duerr, Elsevier, Amsterdam, 2003.
- B. L. Feringa, *Acc. Chem. Res.*, 2001, **34**, 504 and references therein.
- J. C. Crano, T. Flood, D. Knowles, A. Kumar and B. van Gemert, *Pure Appl. Chem.*, 1996, **68**, 1395.
- M.-J. R. P. Queiroz, P. M. S. Plasencia, R. Dubest, J. Aubard and R. Guglielmetti, *Tetrahedron*, 2003, **59**, 2567.
- M. Kawamoto, T. Aoki and T. Wada, *Chem. Commun.*, 2007, 930.
- Y. Zhou, D. Zhang, Y. Zhang, Y. Tang and D. Zhu, *J. Org. Chem.*, 2005, **70**, 6164.
- S. Pieraccini, G. Gottarelli, R. Labruto, S. Masiero, O. Pandoli and G. P. Spada, *Chem.-Eur. J.*, 2004, **10**, 5632.
- C. Wang, L. Zhu, Y. Yu, D. Zhang, Z. Shuai and D. Zhu, *J. Org. Chem.*, 2007, **72**, 4306.
- (a) M. B. Birau and Z. Y. Wang, *Tetrahedron Lett.*, 2000, **41**, 4025; (b) K. Takaiishi, M. Kawamoto, K. Tsubaki and T. Wada, *J. Org. Chem.*, 2009, **74**, 5723; (c) R. A. van Delden, T. Mecca, C. Rosini and B. L. Feringa, *Chem.-Eur. J.*, 2004, **10**, 61; (d) C. Wang, D. Zhang, G. Zhang, J. Xiang and D. Zhu, *Chem.-Eur. J.*, 2008, **14**, 5680; (e) R. Eelkema, M. M. Pollard, N. Katsonis, J. Vicario, D. J. Broer and B. L. Feringa, *J. Am. Chem. Soc.*, 2006, **128**, 14397; (f) Y. Yokoyama, S. Uchida, Y. Yokoyama, Y. Sugawara and Y. Kurita, *J. Am. Chem. Soc.*, 1996, **118**, 3100; (g) K. S. Burnham and G. B. Schuster, *J. Am. Chem. Soc.*, 1998, **120**, 12619.
- W. Zhao and E. M. Carreira, *Org. Lett.*, 2003, **5**, 4153 and references therein.
- W. Zhao and E. M. Carreira, *US Pat.* 2001/6281366, 2001.

- 16 K. Tanaka, H. Aoki, H. Hosomi and S. Ohba, *Org. Lett.*, 2000, **2**, 2133.
- 17 D. Bell, M. R. Davies, G. R. Geen and I. S. Mann, *Synthesis*, 1995, 707.
- 18 D. S. Lingenfelter, R. C. Helgeson and D. J. Cram, *J. Org. Chem.*, 1981, **46**, 393.
- 19 (a) L. Gobbi, P. Seiler, F. Diederich and V. Gramlich, *Helv. Chim. Acta*, 2000, **83**, 1711; (b) Y. Meng, W. T. Slaven, D. Wang, T. Liu, H. Chow and C. Li, *Tetrahedron: Asymmetry*, 1998, **9**, 3693.
- 20 (a) J. Hua and W. Lin, *Org. Lett.*, 2004, **6**, 861; (b) T. R. Wu, L. Shen and J. M. Chong, *Org. Lett.*, 2004, **6**, 2701; (c) L. Ma, P. S. White and W. Lin, *J. Org. Chem.*, 2002, **67**, 7577; (d) Y. Yamashita, H. Ishitani, H. Shimizu and H. Kobayashi, *J. Am. Chem. Soc.*, 2002, **124**, 3292.
- 21 (a) N. Harada and K. Nakanishi, *Circular Dichroic Spectroscopy—Exciton Coupling in Organic Stereochemistry*, Oxford University Press, Oxford, 1993; (b) N. Berova, N. Harada and K. Nakanishi, *Electronic Spectroscopy: Exciton Coupling, Theory and Applications*, in *Encyclopedia of Spectroscopy and Spectrometry*, ed. J. Lindon, G. Tranter and J. Holmes, Academic Press, New York, 1999; (c) K. Nakanishi and N. Berova, in *Circular Dichroism: Principles and Applications*, ed. K. Nakanishi, N. Berova and R. Woody, VCH Publishers, Inc., Deerfield Beach, FL, 1994, ch. 13, pp. 361–398; (d) N. Berova, B. Borhan, J. G. Dong, J. Guo, X. Huang, E. Karnaukhova, A. Kawamura, J. Lou, S. Matile, K. Nakanishi, B. Rickman, J. Su, Q. Tan and I. Zazne, *Pure Appl. Chem.*, 1998, **70**, 377.
- 22 (a) E. Yashima, K. Maeda and T. Nishimura, *Chem.–Eur. J.*, 2004, **10**, 42; (b) M. M. Green, J.-W. Park, T. Sato, A. Teramoto, S. Lifson, R. L. B. Selinger and J. V. Selinger, *Angew. Chem., Int. Ed.*, 1999, **38**, 3138.
- 23 C. R. H. I. de Jong, in *Organic Syntheses by Oxidation with Metal Compounds*, ed. W. J. Mijs and C. R. H. I. de Jong, Plenum Press, New York, 1986, pp. 423–443.
- 24 S. Colonna, A. Re and H. Wynberg, *J. Chem. Soc., Perkin Trans. 1*, 1981, 547.
- 25 (a) M. Noji, M. Nakajima and K. Koga, *Tetrahedron Lett.*, 1994, **35**, 7983; (b) D. Cai, D. L. Hughes, T. R. Verhoeven and P. J. Reider, *Org. Synth.*, 1999, **76**, 1; (c) Y. Wang, J. Sun and K. Ding, *Tetrahedron*, 2000, **56**, 4447; (d) Q. S. Hu, D. Vitharana and L. Pu, *Tetrahedron: Asymmetry*, 1995, **6**, 2123.
- 26 M. Hatano, S. Suzuki and K. Ishihara, *J. Am. Chem. Soc.*, 2006, **128**, 9998.
- 27 P. J. Wagner, *J. Am. Chem. Soc.*, 1967, **89**, 2820.