

This paper is published as part of a *Dalton Transactions* theme issue:

Dalton Discussion 11: The Renaissance of Main Group Chemistry

Guest Editor: John Arnold
University of California, Berkeley, CA, USA
23 - 25 June 2008

Published in [issue 33, 2008](#), of *Dalton Transactions*




Image reproduced with permission of Manfred Scheer

Papers published in this issue include:

[The coordination chemistry of group 15 element ligand complexes—a developing area](#)

Manfred Scheer, *Dalton Trans.*, 2008 DOI: [10.1039/b718179p](#)

[Formation, structure and bonding of metalloid Al and Ga clusters. A challenge for chemical efforts in nanosciences](#)

Hansgeorg Schnöckel, *Dalton Trans.*, 2008 DOI: [10.1039/b718784j](#)

[Polymeric materials based on main group elements: the recent development of ambient temperature and controlled routes to polyphosphazenes](#)

Vivienne Blackstone, Alejandro Presa Soto and Ian Manners, *Dalton Trans.*, 2008
DOI: [10.1039/b719361k](#)

[Recent developments in the chemistry of low valent Group 14 hydrides](#)

Eric Rivard and Philip P. Power, *Dalton Trans.*, 2008 DOI: [10.1039/b801400k](#)

[Chemistry and physics of silicon nanowire](#)

Peidong Yang, *Dalton Trans.*, 2008 DOI: [10.1039/b801440j](#)

Visit the *Dalton Transactions* website for more cutting-edge inorganic and organometallic research
www.rsc.org/dalton

6-Coordinate tungsten(vi) tris-*n*-isopropylanilide complexes: products of terminal oxo and nitrido transformations effected by main group electrophiles†‡

Christopher R. Clough, Peter Müller and Christopher C. Cummins*

Received 21st January 2008, Accepted 3rd March 2008

First published as an Advance Article on the web 1st July 2008

DOI: 10.1039/b801037d

The nitridotungsten(vi) complex $\text{NW}(\text{N}[i\text{-Pr}]\text{Ar})_3$ (**1-N**, Ar = 3,5-Me₂C₆H₃) reacts with (CF₃C(O))₂O followed by ClSiMe₃ to give the isolable trifluoroacetylido-chloride complex **1-(NC(O)CF₃)Cl**, with oxalyl chloride to give cyanate-dichloride **1-(OCN)(Cl)₂**, and with PCl₅ to give trichlorophosphinimide-dichloride **1-(NPCl₂)(Cl)₂**. The oxo-chloride complex **1-(O)Cl**, obtained from **1-N** upon treatment with pivaloyl chloride, reacts with PCl₅ to give trichloride **1-(Cl)₃**. Synthetic and structural details are reported for the new tungsten trisanilide derivatives.

Introduction

Recently we showed that the nitridotungsten(vi) complex $\text{NW}(\text{N}[i\text{-Pr}]\text{Ar})_3$ (**1-N**; Ar = 3,5-Me₂C₆H₃) can be used as a reagent for the transformation of acid chlorides into organic nitriles according to: **1-N** + R¹C(O)Cl → **1-(O)Cl** + R¹CN (see Scheme 1).¹ For R¹ = *t*-Bu or 1-Ad, the latter reaction proceeds quantitatively at 25 °C in less than 1 h and is an intriguing example of an isovalent N for (O)Cl exchange process. Acylimido-chloride complexes **1-(NC(O)R¹)Cl** are observable during the reaction as monitored by ¹H or ¹³C NMR spectroscopy, and they are kinetically competent to be intermediates.

The tungsten trifluoroacetylido trifluoroacetate complex **1-(NC(O)CF₃)(O₂CCF₃)**, was obtained previously by treatment of **1-N** with trifluoroacetic anhydride (TFAA), and was the subject of an X-ray diffraction study.¹ One possible explanation for the failure of **1-(NC(O)CF₃)(O₂CCF₃)** to thermally extrude CF₃CN was that bidentate trifluoroacetate coordination to W served to inhibit the formation of a metallacyclic acylimido complex similar to the proposed structure **2** (Scheme 1). This idea was rendered implausible by the observed structure of **1-(NC(O)CF₃)(O₂CCF₃)**,¹ in which monodentate trifluoroacetate was observed and the coordination geometry at tungsten (including ancillary N[*i*-Pr]Ar substituent conformation) was essentially identical to that found for oxo-chloride **1-(O)Cl**. It began to seem, therefore, that the electron-withdrawing nature of the CF₃ group in trifluoroacetylido **1-(NC(O)CF₃)(O₂CCF₃)** was principally responsible for the failure of this complex to mediate nitrile formation.

Results and discussion

Probing the stability of **1-(NC(O)CF₃)Cl**

To investigate the effect of electron withdrawing groups in the conversion of **1-(NC(O)CF₃)Cl** to **1-(O)Cl**, we treated **1-(NC(O)CF₃)(O₂CCF₃)** with excess ClSiMe₃ to provide trifluoroacetylido chloride **1-(NC(O)CF₃)Cl**. As was the case for its synthetic precursor we find **1-(NC(O)CF₃)Cl** to be thermally stable, and now report an X-ray structural study of this complex (Fig. 1). Like oxo-chloride **1-(O)Cl**, and like its precursor **1-(NC(O)CF₃)(O₂CCF₃)**, the trifluoroacetylido chloride complex **1-(NC(O)CF₃)(Cl)** incorporates trigonal-bipyramidal coordination at W, with an equatorial metal–ligand multiple bond.

If the mechanism of nitrile formation as effected by **1-N** indeed involves metallacycles such as **2** in Scheme 1, then either the specific choice of R¹ = CF₃ obviates metallacycle formation, or else it renders the reaction thermodynamically uphill.² Either way, CF₃CN is a nitrile not available when using the **1-N** reagent.

Since our utilization of sterically demanding ancillary anilide ligands (here, N[*i*-Pr]Ar) is motivated by a desire to foster low coordination number and low nuclearity, it was with some trepidation that we proposed the intermediacy of 6-coordinate metallacycles **2**. For this reason, we were fascinated to find examples of *bona fide* 6-coordinate, octahedral complexes supported by platform **1**. This occurred in the course of surveying the reactivity of **1-N** with a variety of acid chlorides.


Isolation of 6-coordinate complexes

Treatment of **1-N** with 0.5 equiv of oxalyl chloride was intended to provide cyanogen. Instead, only 0.5 equiv of the initial **1-N** was consumed, indicating that a 1 : 1 reaction was preferred. Accordingly, treatment of **1-N** with 1.0 equiv of oxalyl chloride was found to provide, with effervescence attributed to CO liberation, the cyanato-dichloride complex **1-(OCN)(Cl)₂**. The latter has interesting NMR spectroscopic properties. It is C₁ chiral as indicated by (i) the presence of three distinct N[*i*-Pr]Ar ligand environments in a 1 : 1 : 1 ratio, and (ii) the diastereotopic nature of each of the three N[*i*-Pr]Ar ligand environments. From this combination

Room 6-435, Department of Chemistry, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA, 02139-4307, United States of America

† Based on the presentation given at Dalton Discussion No. 11, 23–25 June 2008, University of California, Berkeley, USA.

‡ Electronic supplementary information (ESI) available: Details of crystal structures of all prepared complexes. CCDC reference numbers 675710–675713. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/b801037d



of facts, we can infer that the combination of a *fac* and C_3 arrangement of the $(\text{OCN})(\text{Cl})_2$ substituents, combined with a *fac* and frozen-out C_3 three-bladed propellor of *N*-isopropylanilide residues, leads to a chiral metal environment with overall C_1 symmetry. This is consistent with an X-ray structural study of **1**-(OCN)(Cl)₂ (Fig. 2). It is from the X-ray study that we tentatively assign **1**-(OCN)(Cl)₂ as equipped with an O-coordinated cyanate ligand, the IR data ($\nu_{\text{NCO}} = 2200 \text{ cm}^{-1}$, vs) being insufficient information to make the distinction.³

Normally, molecules with three *N*(*i*-Pr)Ar ligands, *e.g.* **1**-N or **1**-(O)Cl, are not at 25 °C frozen out into a static C_3 configuration. Such complexes typically evince a single, non-diastereotopic ligand environment. That **1**-(OCN)(Cl)₂ is frozen out at room temperature signifies substantial steric crowding.

Another example of a C_1 -symmetric derivative with an octahedral coordination environment at tungsten is that obtained by reaction of **1**-N with PCl_5 . It is known that PCl_5 has a propensity to react with N-containing compounds to give products with 4-coordinate P and P–N multiple bonding.⁴ As in the case of the oxalyl chloride reaction, tungsten accepts two chloride




Fig. 1 Selected interatomic distances (Å) and angles (°) for **1**-(NC(O)CF₃)Cl: W1–N4, 1.791(3); N4–C41, 1.329(5); O1–C41, 1.199(5); C41–N4–W1, 156.8(3).




Fig. 2 View of **1**-(OCN)(Cl)₂ normal to the 001 plane (space group $R\bar{3}$). The OCN and two Cl ligands are positionally disordered about the crystallographic C_3 axis passing through W1.

ligands while expanding to 6-coordination. The product molecule, **1**-(N=PCl₃)(Cl)₂, incorporating a rare trichlorophosphinimide ligand (³¹P NMR: $\delta = -49.8 \text{ ppm}$, $^2J_{\text{WP}} = 85 \text{ Hz}$), is an orange-red compound soluble in THF or benzene, but of limited ether or pentane solubility. An X-ray structural study of the complex revealed a moderately bent phosphinimide nitrogen, together with overall conformational attributes very much reminiscent of **1**-(OCN)(Cl)₂ (Fig. 3). The related trichlorophosphinimide complex Cl₃P(N=PCl₃) has been prepared by treatment of WCl_6 with Cl₃P=NSiMe₃.^{5,6} Also, $\text{Ph}_4\text{P}[\text{Cl}_5\text{Mo}(\text{N}=\text{PCl}_3)]$ has been prepared by treatment of nitride $\text{Ph}_4\text{P}[\text{Cl}_4\text{MoN}]$ with $\text{PCl}_3/\text{PCl}_5$, in what appears to be the closest precedent for our synthesis of **1**-(N=PCl₃)(Cl)₂.⁷

Since oxo-chloride **1**-(O)Cl is the ultimate product in the reaction of **1**-N with acid chlorides, we are interested in methods




Fig. 3 Selected interatomic distances (Å) and angles (°) for $1-(\text{NPCl}_3)(\text{Cl})_2$: W1–N4, 2.047(4); P1–N4, 1.449(4); P1–N4–W1, 155.9(2).

for the recycling of $1-(\text{O})\text{Cl}$ back to the nitride reagent $1-\text{N}$. This objective is similar to another realized recently, namely the activation of terminal oxo product $\text{ONb}(\text{N}[\text{Np}]\text{Ar})_3$ by reaction with triflic anhydride to give bistriflate $(\text{TfO})_2\text{Nb}(\text{N}[\text{Np}]\text{Ar})_3$; the latter is then reduced to its P_4 -activating niobaziridine-hydride form in our synthesis of phosphalkyne ($\text{RC}\equiv\text{P}$) molecules.⁸ Activation of the oxo in $1-(\text{O})\text{Cl}$ with triflic anhydride was not successful, this reaction giving oxo triflate $1-(\text{O})(\text{OTf})$ instead. On the other hand, we find that PCl_5 serves smoothly to transform oxo chloride $1-(\text{O})\text{Cl}$ into trichloride $1-(\text{Cl})_3$, with POCl_3 as the sole byproduct.⁹ We had expected that a *fac* arrangement of three chloride ligands together with a *fac* and C_3 frozen out arrangement of three $\text{N}[i\text{-Pr}]\text{Ar}$ ligands would provide trichloride $1-(\text{Cl})_3$ with a single yet diastereotopic set of $\text{N}[i\text{-Pr}]\text{Ar}$ ligand ^1H NMR resonances at 25 °C. That expectation was borne out in full: the ^1H NMR spectrum of $1-(\text{Cl})_3$ has a pair of aryl methyl resonances, three aryl proton signals in a 1:1:1 ratio, and a pair of isopropyl methyl doublets. The other example of a *fac* trisamide tungsten trichloride complex that is potentially C_3 with diastereotopic ligand environments (based on its crystal structure) is $\text{WCl}_3(\text{NET}_2)_3$.¹⁰ This molecule shows a single ethyl group triplet and quartet in its ^1H NMR spectrum, indicative of free rotation about its W–N linkages.

An X-ray structural study of trichloride $1-(\text{Cl})_3$ (space-filling diagram in Fig. 4) validates our formulation of this molecule while illustrating the severe inter-ligand steric interactions present in 6-coordinate tungsten systems based on the tris(anilide) platform **1**. Such interactions are expected to be a destabilizing influence on proposed metallacycles **2**.

Experimental

General

Unless stated otherwise, all operations were performed in a Vacuum Atmospheres drybox under an atmosphere of purified nitrogen or using Schlenk techniques under an argon atmosphere. $\text{N}\equiv\text{W}(\text{N}[i\text{-Pr}]\text{Ar})_3$ (**1-N**, Ar = 3,5-Me₂C₆H₃), $(\text{Ar}[i\text{-Pr}]\text{N})_3\text{W}(\text{O})\text{Cl}$ (**1-(O)Cl**), and $(\text{Ar}[i\text{-Pr}]\text{N})_3\text{W}(\text{NC}(\text{O})\text{CF}_3)(\text{O}_2\text{CCF}_3)$ (**1-(NC(O)-CF₃)(O₂CCF₃)**) were prepared as previously published.¹ Oxalyl




Fig. 4 Space filling model of $1-(\text{Cl})_3$.

chloride and *t*-BuC(O)Cl were purchased from Aldrich and distilled under N_2 . PCl_5 was purchased from Aldrich and used as received. Diethyl ether, *n*-pentane, and toluene were dried and deoxygenated by the method of Grubbs.¹¹ THF was distilled from purple Na/benzophenone and collected under nitrogen. C_6D_6 was degassed and dried over 4 Å molecular sieves. Other chemicals were purified and dried by standard procedures or were used as received. Celite® 545, alumina and 4 Å molecular sieves were dried *in vacuo* overnight at a temperature above 200 °C. ^1H , ^{13}C , ^{19}F , and ^{31}P NMR spectra were recorded on Varian Mercury-300, Varian INOVA-500, or Bruker AVANCE-400 spectrometers. ^1H and ^{13}C chemical shifts are reported with respect to internal solvent (C_6D_6 , 7.16 and 128.39 ppm, respectively). ^{19}F and ^{31}P chemical shifts are reported with respect to external reference (CFCl_3 , 0.0 ppm and 85% H_3PO_4 , 0.0 ppm, respectively). Infrared spectra were recorded on a Bio-Rad 135 Series FTIR spectrometer.

Crystallography

X-ray data collections were carried out on a Siemens Platform three-circle diffractometer equipped with a Bruker-AXS Apex CCD detector and an Oxford Cryosystems CryoStream 700 low-temperature device. Graphite-monochromated Mo- K_α radiation ($\lambda = 0.71073$ Å) was used in all cases. All software for diffraction data processing and crystal-structure solution and refinement are contained in the SHELXTL (v6.14) program suite (G. Sheldrick, Bruker AXS, Madison, WI).¹² Details of crystallographic data and refinement are given in Table 1 and in the ESI.†

Syntheses

Synthesis of $(\text{Ar}[i\text{-Pr}]\text{N})_3\text{W}(\text{NC}(\text{O})\text{CF}_3)\text{Cl}$ (1-(NC(O)CF₃)(Cl)**).** **1-(C(O)CF₃)(O₂CCF₃)** (525 mg, 0.767 mmol) was dissolved in minimal Me_3SiCl (~5 mL) and the resulting red solution was stirred for ~5 min, filtered through a bed of Celite® 545 and the filtrate cooled to –35 °C overnight. **1-(NC(O)CF₃)(Cl)** was obtained as a red precipitate which was washed with cold pentane, and dried *in vacuo* (160 mg, 0.196 mmol, 25.5%). X-Ray quality crystals can also be grown by following the same procedure using smaller amounts of **1-(C(O)CF₃)(O₂CCF₃)** (*ca.* 100 mg) in more dilute solutions of Me_3SiCl . ^1H NMR (500 MHz, C_6D_6): δ = 6.57 (s, 3H, *para*), 6.52 (s, 6H, *ortho*), 5.33 (septet, 3H, *i*-Pr methine), 2.05 (s, 18H, ArCH_3), 1.14 (d, 18H, *i*-Pr methyl) ppm. ^{13}C NMR (100 MHz, C_6D_6): δ = 149.3 (*ipso*), 138.2 (*meta*), 129.1 (*para*), 126.3 (*ortho*), 65.6 (*i*-Pr methine), 23.0 (methyl), 21.6 (methyl)

Table 1 Crystallographic data for **1**-(NC(O)CF₃)Cl, **1**-(OCN)(Cl)₂, **1**-(NPCl₃)(Cl)₂ and **1**-(Cl)₃

	1 -(NC(O)CF ₃)Cl	1 -(OCN)(Cl) ₂	1 -(NPCl ₃)(Cl) ₂	1 -(Cl) ₃
Empirical formula	C ₃₅ H ₄₈ ClF ₃ N ₄ OW	C ₄₂ H ₆₄ Cl ₂ N ₄ O ₃ W ^a	C ₃₇ H ₅₆ Cl ₅ N ₄ OPW ^b	C _{33.50} H ₄₉ Cl ₄ N ₃ W ^c
Formula weight	817.07	927.72	964.93	819.41
Color	Dark red	Yellow	Orange	Yellow
Morphology	Plate	Plate	Shard	Shard
Crystal size/mm	0.14 × 0.08 × 0.02	0.27 × 0.23 × 0.04	0.10 × 0.09 × 0.06	0.13 × 0.10 × 0.09
<i>T</i> /K	100(2)	100(2)	100(2)	100(2)
Crystal system	Monoclinic	Rhombohedral	Monoclinic	Triclinic
Space group	<i>P</i> 2 ₁ / <i>n</i>	<i>R</i> $\bar{3}$	<i>P</i> 2 ₁ / <i>c</i>	<i>P</i> $\bar{1}$
Unit cell dimensions:				
<i>a</i> /Å	18.1298(6)	13.3600(12)	15.9389(6)	10.7751(3)
<i>b</i> /Å	10.4918(3)	13.3600(12)	13.7357(5)	13.3987(3)
<i>c</i> /Å	19.4084(5)	39.556(8)	18.9527(6)	13.5032(4)
<i>a</i> /°	90	90	90	88.8520(10)
<i>β</i> /°	107.0410(10)	90	97.4620(10)	71.9870(10)
<i>γ</i> /°	90	120	90	83.6570(10)
<i>V</i> /Å ³	3529.67(18)	6114.5(14)	4114.2(3)	1842.38(9)
<i>Z</i>	4	6	4	2
Density calc./Mg m ⁻³	1.538	1.512	1.558	1.477
Absorption coefficient/mm ⁻¹	3.397	3.008	3.206	3.451
<i>F</i> (000)	1648	2856	1952	826
Theta range for data collection/°	1.83 to 28.28	1.54 to 25.14	1.84 to 27.88	1.53 to 28.28
Reflections collected	73315	10803	81238	38272
Independent reflections, <i>R</i> _{int}	8736 (0.0907)	2442 (0.0634)	9806 (0.0533)	9143 (0.0321)
Completeness to theta (%)	100.0	100.0	100.0	99.9
Max. and min. transmission	0.9352 and 0.6477	0.8891 and 0.4972	0.8309 and 0.7399	0.7465 and 0.6626
Data/restraints/parameters	8736/0/412	2442/345/288	9806/74/452	9143/85/406
Goodness-of-fit on <i>F</i> ²	1.075	1.152	1.070	1.103
Final <i>R</i> indices [<i>I</i> > 2σ(<i>I</i>)]	<i>R</i> ₁ = 0.0359 <i>wR</i> ₂ = 0.0727	<i>R</i> ₁ = 0.0450 <i>wR</i> ₂ = 0.1157	<i>R</i> ₁ = 0.0332 <i>wR</i> ₂ = 0.0839	<i>R</i> ₁ = 0.0293 <i>wR</i> ₂ = 0.0826
<i>R</i> indices (all data)	<i>R</i> ₁ = 0.0522 <i>wR</i> ₂ = 0.0791	<i>R</i> ₁ = 0.0650 <i>wR</i> ₂ = 0.1394	<i>R</i> ₁ = 0.0436 <i>wR</i> ₂ = 0.0895	<i>R</i> ₁ = 0.0328 <i>wR</i> ₂ = 0.0847
Largest diff. peak and hole/e Å ⁻³	1.302 and -1.122	2.017 and -1.728	2.006 and -0.720	2.061 and -0.416

^a Two heavily disordered molecules of tetrahydrofuran are present in the asymmetric unit. ^b A disordered molecule of tetrahydrofuran is present in the asymmetric unit. ^c One half of a heavily disordered molecule of methylene chloride is present in the asymmetric unit.

ppm. ¹⁹F NMR (376 MHz, C₆D₆): δ = -72.7 (s, 3F, CF₃) ppm. Anal. calcd for C₃₅H₄₈ClF₃N₄OW: C, 51.45; H, 5.92; N, 6.86. Found: C, 50.95; H, 6.16; N, 6.73.

Synthesis of (Ar[*i*-Pr]N)₃W(OCN)(Cl)₂ (1**-(OCN)(Cl)₂).** Oxalyl chloride (39.0 μL, 0.447 mmol) was added to a colorless solution of **1**-N (304 mg, 0.444 mmol) in Et₂O (10 mL) using a microliter syringe. The reaction mixture turned blood-red upon addition and shortly changed color to brown. The reaction mixture was stirred for 20 min at which point the volatiles were removed *in vacuo* giving a brownish-yellow solid. The crude material was scraped onto a fritted glass filter and washed with Et₂O revealing a bright yellow solid. The bright yellow solid was dissolved in minimal THF, the solution filtered through a plug of Celite® 545 and the filtrate cooled to -35 °C overnight. From the THF solution, small yellow crystals were harvested, washed with pentane and dried *in vacuo* (87 mg, 0.11 mmol, 25%). Recrystallized samples of **1**-(OCN)(Cl)₂ contain ~0.5 equiv of THF that persists even after drying *in vacuo* as evidenced by the ¹H and ¹³C NMR spectra. X-Ray quality crystals of **1**-(OCN)(Cl)₂ can be grown from a saturated THF solution layered with pentane and stored at -35 °C. ¹H NMR (500 MHz, C₆D₆): δ = 7.65 (s, 1H, *para*), 7.61 (s, 1H, *para*), 7.16 (s, 1H, *para*), 6.56 (s, 3H, *ortho*), 6.43 (septet, 1H, *i*-Pr methine), 6.36 (s, 3H, *ortho*), 6.26 (septet, 1H, *i*-Pr methine), 6.01 (septet, 1H, *i*-Pr methine), 2.15 (s, 9H, ArCH₃), 2.054 (s, 3H, ArCH₃), 2.049 (s, 3H, ArCH₃), 2.02 (s, 3H, ArCH₃), 1.56 (d, 3H, *i*-Pr methyl),

1.52 (d, 3H, *i*-Pr methyl), 1.39 (d, 3H, *i*-Pr methyl), -0.27 (d, 3H, *i*-Pr methyl), -0.30 (d, 3H, *i*-Pr methyl), -0.31 (d, 3H, *i*-Pr methyl) ppm. ¹³C NMR (125 MHz, C₆D₆): δ = 152.1 (*ipso*), 151.9 (*ipso*), 151.7 (*ipso*), 138.51 (*meta*), 138.50 (*meta*), 138.4 (*meta*), 137.3 (*meta*), 137.1 (*meta*), 137.0 (*meta*), 136.8 (OCN), 129.04 (*ortho*), 129.00 (*ortho*), 128.8 (*ortho*), 124.2 (*ortho*), 124.01 (*ortho*), 123.92 (*para*), 123.85 (*para*), 123.82 (*ortho*), 123.6 (*para*), 68.0 (*i*-Pr methine), 67.7 (*i*-Pr methine), 67.1 (*i*-Pr methine), 23.4 (*i*-Pr methyl), 23.3 (*i*-Pr methyl), 23.2 (*i*-Pr methyl), 23.0 (*i*-Pr methyl), 22.5 (*i*-Pr methyl), 22.08 (2C, ArCH₃), 22.06 (ArCH₃), 21.74 (ArCH₃), 21.72 (ArCH₃), 21.70 (ArCH₃), 21.5 (*i*-Pr methyl) ppm. FTIR (C₆D₆, KBr): ν_{NCO} = 2200 cm⁻¹ (vs). Anal. calcd for C₃₄H₄₈Cl₂N₄OW: C, 52.12; H, 6.17; N, 7.15. Found: C, 51.75; H, 6.18; N, 6.93.

Synthesis of (Ar[*i*-Pr]N)₃W(N=PCl₃)(Cl)₂ (1**-(N=PCl₃)(Cl)₂).** A thawing, colorless solution of **1**-N (505 mg, 0.738 mmol) in Et₂O (5 mL) was added to a thawing suspension of PCl₅ (154 mg, 0.740 mmol) in Et₂O (5 mL) resulting in an orange-red reaction mixture upon addition. The reaction mixture was allowed to warm to room temperature and stirred for 1 h after which time the volatiles were removed *in vacuo* leaving an orange-yellow solid. The solid was collected on a fritted glass filter, washed with pentane, and dried under vacuum (505 mg, 0.566 mmol, 76.6%). The sample used to collect the NMR spectra was recrystallized from THF layered with Et₂O at -35 °C. The sample contains ~1 equiv of

THF of co-crystallization as seen in the ^1H and ^{13}C NMR spectra. The THF remains in the sample even after prolonged periods of drying *in vacuo*. ^1H NMR (400 MHz, C_6D_6): $\delta = 7.74$ (s, 1H, *para*), 7.72 (s, 1H, *para*), 7.32 (s, 1H, *para*), 6.61 to 6.47 (7H, *ortho* and *i*-Pr methine), 6.19 (septet, 1H, *i*-Pr methine), 5.88 (septet, 1H, *i*-Pr methine), 2.20 (s, 9H, ArCH_3), 2.16 (s, 3H, ArCH_3), 2.09 (s, 3H, ArCH_3), 2.08 (s, 3H, ArCH_3), 1.65 (d, 3H, *i*-Pr methyl), 1.55 (d, 3H, *i*-Pr methyl), 1.44 (d, 3H, *i*-Pr methyl), -0.22 to -0.25 (9H, *i*-Pr methyl) ppm. ^{13}C NMR (100 MHz, C_6D_6): $\delta = 153.0$ (*ipso*), 152.5 (*ipso*), 151.9 (*ipso*), 138.1 (*meta*), 137.9 (*meta*), 137.8 (*meta*), 137.1 (*meta*), 136.6 (*meta*), 136.4 (*meta*), 125.4 (2C, *ortho*), 125.2 (2C, *ortho*), 124.5 (2C, *ortho*), 124.3 (*para*), 124.2 (*para*), 124.1 (*para*), 68.9 (*i*-Pr methine), 68.1 (*i*-Pr methine), 66.1 (*i*-Pr methine), 23.7 (methyl), 23.4 (methyl), 23.2 (methyl), 23.1 (methyl), 23.0 (methyl), 22.7 (methyl), 22.02 (methyl), 21.95 (methyl), 21.70 (methyl), 21.66 (methyl) ppm. ^{31}P NMR (162 MHz, C_6D_6): $\delta = -49.8$ ($^2J_{\text{WP}} = 85$ Hz) ppm. Anal. calcd for $\text{C}_{33}\text{H}_{48}\text{Cl}_5\text{N}_4\text{PW}$: C, 44.39; H, 5.42; N, 6.27. Calcd for $\text{C}_{37}\text{H}_{56}\text{Cl}_5\text{N}_4\text{OPW}$ (1.0 equiv THF—as seen in the crystal structure): C, 46.05; H, 5.85; N, 5.81. Found: C, 45.95; H, 5.84; N, 5.72.

Synthesis of $(\text{Ar}[i\text{-Pr}]\text{N})_3\text{W}(\text{Cl})_3$ (1**-(Cl) $_3$).** A thawing, red solution of **1**-(O)Cl (794 mg, 1.10 mmol) in Et_2O (5 mL) was added to a thawing suspension of PCl_5 (228 mg, 1.10 mmol) in Et_2O (3 mL) resulting in an orange reaction mixture upon addition. The reaction mixture was allowed to warm to room temperature and was stirred for 0.5 h. A canary yellow solid precipitated out of solution and was collected on a fritted glass filter. The solids were washed with pentane (3×20 mL) and dried under vacuum (595 mg, 0.766 mmol, 69.6%). Samples contain ~ 1 equiv of Et_2O as evinced by ^1H and ^{13}C NMR. The Et_2O remains even after prolonged periods of drying *in vacuo*. X-Ray quality crystals of **1**-(Cl) $_3$ can be grown from a saturated methylene chloride solution layered with diethyl ether and stored at -35°C . ^1H NMR (300 MHz, C_6D_6): $\delta = 7.77$ (s, 3H, *para*), 6.55 (s, 3H, *ortho*), 6.44 (s coincident with septet, 6H, *ortho* and *i*-Pr methine, respectively), 2.15 (s, 9H, ArCH_3), 2.05 (s, 9H, ArCH_3), 1.16 (d, 9H, *i*-Pr methyl), -0.27 (d, 9H, *i*-Pr methyl). ^{13}C NMR (100 MHz, C_6D_6): $\delta = 152.3$ (*ipso*), 138.2 (*meta*), 136.8 (*meta*), 124.0 (2C, *ortho*), 123.9 (*para*), 68.5 (*i*-Pr methine), 23.4 (methyl), 22.1 (methyl), 22.0 (methyl), 21.6 (methyl) ppm. Anal. calcd for $\text{C}_{33}\text{H}_{48}\text{Cl}_3\text{N}_3\text{W}$: C, 51.01; H, 6.23; N, 5.41. Calcd for $\text{C}_{33.5}\text{H}_{49}\text{Cl}_4\text{N}_3\text{W}$ (0.5 equiv methylene chloride—as seen in the crystal structure): C, 49.10; H, 6.04; N, 5.13. Found: C, 48.58; H, 6.38; N, 4.32.

Density functional calculations

All calculations were carried out using ADF 2004.01 from Scientific Computing and Modeling (<http://www.scm.com>).^{13,14} In all cases the LDA functional employed was that of Vosko, Wilk, and Nusair (VWN)¹⁵ while the GGA part was handled using the functionals of Becke and Perdew (BP86).^{16,17} In addition, all calculations were carried out using the Zero Order Regular Approximation (ZORA) for relativistic effects.^{18,19} In all cases the basis sets were triple-zeta with two polarization functions (TZ2P) as supplied with ADF. Frozen core approximations were utilized according to the following atom types: F, N, C, and O: 1 s frozen; Cl: core frozen through and including 2p; W: core frozen through and including 4f. Calculations were carried out on a four- or an eight-processor Quantum Cube workstation from

Table 2 Total bonding energies for molecules involved in the nitrile elimination reactions

Molecule	Total energy/kcal mol ^{-1a}
CH_3CN	-837.37
CF_3CN	-851.07
$\text{CH}_3\text{C}(\text{O})\text{NW}(\text{NH}_2)_3\text{Cl}$	-2417.45
$\text{CF}_3\text{C}(\text{O})\text{NW}(\text{NH}_2)_3\text{Cl}$	-2442.01
$\text{OW}(\text{NH}_2)_3\text{Cl}$	-1579.16

^a With respect to spherical atomic fragments.

Parallel Quantum Solutions (<http://www.pqs-chem.com>). All results reported are with reference to fully optimized geometries with no imaginary frequencies.^{20,21}

From the above total bonding energies (Table 2) we can compute ΔH_{rxn} for the two nitrile elimination reactions as follows:



Based on calculations with the above model complexes, extrusion of acetonitrile from $\text{CH}_3\text{C}(\text{O})\text{NW}(\text{NH}_2)_3$ is essentially thermoneutral while formation of trifluoroacetonitrile from the analogous tungsten complex is significantly uphill.

Conclusions

Synthesis and characterization of 6-coordinate tungsten complexes **1**-(N= PCl_3)(Cl) $_2$, **1**-(OCN)(Cl) $_2$ and **1**-(Cl) $_3$ lends credence to the proposed intermediate **2**. Similar metallacycles have been proposed both by us²² and others,²³ but there had been some doubt as to whether the tungsten trisanilide platform **1** could adopt a pseudo-octahedral structure. Additionally, we have discovered a new mode of reactivity for **1**-N and **1**-(O)Cl that awaits further exploitation.

Acknowledgements

For stimulating discussions the authors thank Richard R. Schrock. We gratefully acknowledge for funding this work the U. S. National Science Foundation (CHE-0316823).

References

- 1 C. R. Clough, J. B. Greco, J. S. Figueroa, P. L. Diaconescu, W. M. Davis and C. C. Cummins, *J. Am. Chem. Soc.*, 2004, **126**, 7742–7743.
- 2 Density functional theory calculations (ADF 2004.01, ZORA TZ2P, BP86) have been carried out for the nitrile extrusion reaction: $(\text{H}_2\text{N})_3\text{W}(\text{NC}(\text{O})\text{CR}_3)\text{Cl} \rightarrow (\text{H}_2\text{N})_3\text{W}(\text{O})\text{Cl} + \text{CR}_3\text{CN}$. For, R = H, $\Delta H_{\text{rxn}} = +1$ kcal mol⁻¹, while for R = F, $\Delta H_{\text{rxn}} = +12$ kcal mol⁻¹.
- 3 R. A. Bailey, S. L. Kozak, T. W. Michelsen and W. N. Mills, *Coord. Chem. Rev.*, 1971, **6**, 407–445.
- 4 M. Becke-Goehring, *Fortschr. Chem. Forsch.*, 1968, **10**, 207–237.
- 5 C. H. Honeyman, A. J. Lough and I. Manners, *Inorg. Chem.*, 1994, **33**, 2988–2993.
- 6 E. Rivard, C. H. Honeyman, A. R. McWilliams, A. J. Lough and I. Manners, *Inorg. Chem.*, 2001, **40**, 1489–1495.
- 7 I. Schmidt, U. Kynast, J. Hanich and K. Dehnicke, *Z. Naturforsch., B: Anorg. Chem. Org. Chem.*, 1984, **39**, 1248–1251.
- 8 J. S. Figueroa and C. C. Cummins, *J. Am. Chem. Soc.*, 2004, **126**, 13916–13917.
- 9 K. R. Seddon and V. H. Thomas, *Inorg. Chem.*, 1978, **17**, 749–751.

-
- 10 S. Dietz, V. Allured and M. R. Dubois, *Inorg. Chem.*, 1993, **32**, 5418–5420.
- 11 A. B. Pangborn, M. A. Giardello, R. H. Grubbs, R. K. Rosen and F. J. Timmers, *Organometallics*, 1996, **15**, 1518–1520.
- 12 G. M. Sheldrick, *Acta Crystallogr., Sect. A: Found. Crystallogr.*, 2008, **64**, 112–122.
- 13 G. te Velde, F. M. Bickelhaupt, E. J. Baerends, C. F. Guerra, S. J. A. van Gisbergen, J. G. Snijders and T. Ziegler, *J. Comput. Chem.*, 2001, **22**, 931–967.
- 14 C. F. Guerra, J. G. Snijders, G. te Velde and E. J. Baerends, *Theor. Chem. Acc.*, 1998, **99**, 391–403.
- 15 S. H. Vosko, L. Wilk and M. Nusair, *Can. J. Phys.*, 1980, **58**, 1200–1211.
- 16 A. D. Becke, *Phys. Rev. A*, 1988, **38**, 3098–3100.
- 17 J. P. Perdew, *Phys. Rev. B: Condens. Matter Mater. Phys.*, 1986, **34**, 7406–7406.
- 18 E. van Lenthe, E. J. Baerends and J. G. Snijders, *J. Chem. Phys.*, 1993, **99**, 4597–4610.
- 19 E. van Lenthe, A. Ehlers and E. J. Baerends, *J. Chem. Phys.*, 1999, **110**, 8943–8953.
- 20 L. Y. Fan and T. Ziegler, *J. Chem. Phys.*, 1992, **96**, 9005–9012.
- 21 L. Y. Fan and T. Ziegler, *J. Phys. Chem.*, 1992, **96**, 6937–6941.
- 22 J. K. Brask, V. Dura-Vila, P. L. Diaconescu and C. C. Cummins, *Chem. Commun.*, 2002, 902–903.
- 23 M. H. Chisholm, E. E. Delbridge, A. R. Kidwell and K. B. Quinlan, *Chem. Commun.*, 2003, 126–127.