

Catalytic Asymmetric Synthesis of α -Alkylidene- β -hydroxy Esters via Dynamic Kinetic Asymmetric Transformation Involving Ba-Catalyzed Direct Aldol Reaction

Akitake Yamaguchi, Shigeki Matsunaga,* and Masakatsu Shibasaki*

Graduate School of Pharmaceutical Sciences, The University of Tokyo, Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

Received June 5, 2009; E-mail: smatsuna@mol.f.u-tokyo.ac.jp; mshibasa@mol.f.u-tokyo.ac.jp

The direct catalytic asymmetric aldol reaction is a powerful and atom-economical method for synthesizing chiral β -hydroxycarbonyl compounds.¹ To date, many metal and organocatalysts for reactions of ketone and aldehyde donors have been developed.^{1,2} These catalysts kinetically control aldol reactions via a simple proton-transfer process, avoiding undesirable retro-aldol reactions that would cause racemization of the products. In contrast, direct catalytic asymmetric aldol reactions with ester donors are limited to reactions with α -isocyanoacetate,^{3a} α -cyanoacetate,^{3b} α -diazoacetate,⁴ and glycinate Schiff base donors.⁵ Although the utility of several α -alkyl-substituted ester-equivalent donors in direct aldol reactions has been nicely demonstrated,^{6,7} esters without α -heteroatoms and/or electron-withdrawing substituents have not been utilized, possibly because of the low acidity of α -protons in esters and/or undesirable retro-aldol reactions that occur under basic reaction conditions. Herein, we report the use of ester donors in dynamic kinetic asymmetric transformation (DYKAT)⁸ for catalytic asymmetric synthesis of β -hydroxy esters. DYKAT involving Ba(O-*i*Pr)₂/**1a**-catalyzed direct aldol/retro-aldol reactions of β,γ -unsaturated esters **2** (Figure 1) afforded α -alkylidene- β -hydroxy esters in up to 99% ee.

A retro-aldol reaction is generally considered an undesirable pathway in kinetically controlled direct asymmetric aldol reactions. We planned to utilize the direct aldol/retro-aldol process in DYKAT to obtain chiral α -alkylidene- β -hydroxy esters. Our working hypothesis is summarized in Scheme 1. Dienolates generated in situ from β,γ -unsaturated ester **2** by a chiral catalyst would react with aldehyde **3** at the α - and/or γ -position. If the chiral catalyst promotes rapid retro-aldol reaction of the α -adduct **4**, isomerization of **4** to the more thermodynamically stable α -alkylidene- β -hydroxy ester **5** can be a DYKAT.

On the basis of this hypothesis, we screened chiral catalysts for the reaction of ester **2a** and aldehyde **3a**, and 1:1 Ba(O-*i*Pr)₂/**1a** mixture gave promising results (Table 1).^{9,10} The desired aldol reaction/isomerization sequence was promoted by 10 mol % (*S*)-Ba-**1a** in THF at 0 °C, giving **5aa** in 41% yield and 91% ee (entry 1). To improve the α/γ selectivity, we optimized the reaction conditions (entries 2–7). Performing the reaction in THF/DME drastically improved the α/γ selectivity to 11:1 (entry 4). Other metal sources, such as alkali metal (entry 5, Li-**1a**) and rare-earth metal (entry 6, La-**1a**), resulted in poor reactivity and selectivity. The best result was obtained in DME alone with Ba-**1a**, giving predominantly the *E* adduct **5aa** with $\alpha/\gamma > 20:1$ in 85% yield and 99% ee after 24 h at 0 °C (entry 7). Ethyl ester **2b** also gave the desired product in 99% ee, but both the α/γ selectivity and yield of the desired product decreased (entry 8). With sterically hindered *t*-butyl ester **2c**, the reactivity was poor (entry 9).

The substrate scope of the reaction is summarized in Table 2. The reactions of some aldehydes in Table 2 gave a better yield using (*S*)-Ba-**1b** rather than (*S*)-Ba-**1a**. Thus, we examined both




Figure 1. Structures of (*S*)-BINOL (**1a**), (*S*)-6,6'-(MeO)₂-BINOL (**1b**), and β,γ -unsaturated ester **2**.

Scheme 1. Working Hypothesis of DYKAT Involving Direct Aldol/Retrosaldol Reaction for Catalytic Asymmetric Synthesis of α -Alkylidene- β -hydroxy Esters





Table 1. Optimization of Reaction Conditions



entry	M	x	2	solvent	5/6 ^a	% yield of 5	% ee of 5
1	Ba(O- <i>i</i> Pr) ₂	10	2a	THF	1.8:1	41	91
2	Ba(O- <i>i</i> Pr) ₂	10	2a	1:9 THF/toluene	—	trace	—
3	Ba(O- <i>i</i> Pr) ₂	10	2a	1:9 THF/EtOAc	0.8:1	4	69
4	Ba(O- <i>i</i> Pr) ₂	10	2a	1:9 THF/DME	11:1	79	98
5	Li(O- <i>i</i> Pr)	5	2a	1:9 THF/DME	1.6:1	8	9 ^b
6	La(O- <i>i</i> Pr) ₃	15	2a	1:9 THF/DME	—	0	—
7	Ba(O- <i>i</i> Pr) ₂	10	2a	DME	>20:1	85	99
8	Ba(O- <i>i</i> Pr) ₂	10	2b	DME	5.6:1	69	99
9	Ba(O- <i>i</i> Pr) ₂	10	2c	DME	—	trace	—

^a Determined by ¹H NMR analysis of the crude mixture. ^b *ent*-**5** was obtained as the major isomer.


(*S*)-Ba-**1a** and (*S*)-Ba-**1b** for each aldehyde, and the best results are shown in Table 2.¹¹ (*S*)-Ba-**1** catalysts were applicable to a broad range of aldehydes, giving predominantly *E* adducts **5** in all entries. Aryl aldehydes **3b–d** with either an electron-donating group (entries 2 and 3) or an electron-withdrawing group (entry 4) gave the desired products **5ba–5da** in 99–96% ee. High enantioselectivity was also achieved with heteroaryl and alkenyl aldehydes **3e–h** (entries 5–8, 99–97% ee). Readily enolizable alkyl aldehydes, including linear aldehyde **3j**, were applicable as well, although the enantioselectivity was slightly lower than for other aldehydes

Table 2. Ba-Catalyzed Asymmetric Synthesis of α -Alkylidene- β -hydroxy Esters via DYKAT^a


entry	R	3	1 (x)	temp (°C)	time (h)	5	5/6 ^b	% yield of 5 ^c	% ee
1	Ph	3a	1a (10)	0	24	5aa	>20:1	85	99
2	4-MeC ₆ H ₄	3b	1a (10)	0	42	5ba	17:1	77	98
3	3-MeOC ₆ H ₄	3c	1b (10)	-20	24	5ca	15:1	81	99
4	3-BrC ₆ H ₄	3d	1a (10)	0	42	5da	>20:1	78	96
5	2-thienyl	3e	1a (10)	0	34	5ea	>20:1	80	98
6	3-thienyl	3f	1b (10)	-20	40	5fa	>20:1	85	97
7	3-furyl	3g	1a (10)	0	34	5ga	>20:1	82	98
8	(E)-PhCH=CH	3h	1b (10)	-20	28	5ha	>20:1	63	99
9	iBu	3i	1b (10)	0	42	5ia	>20:1	76	91
10	nPr	3j	1a (10)	0	48	5ja	>20:1	53	87
11	Ph	3a	1a (5)	0	55	5aa	>20:1	84	99

^a In entries 1 and 11, 2 equiv of **2a** were used; in entries 2–10, 3 equiv of **2a** were used. ^b Determined by ¹H NMR analysis of the crude mixture. ^c Isolated yield after purification by column chromatography.

(entries 9 and 10, 91–87% ee). The catalyst loading was successfully reduced to 5 mol % without decreasing the yield or enantioselectivity, but a longer reaction time was required (entry 11, 55 h). (*S*)-Ba-**1a** also promoted the aldol/isomerization sequence of ester **2d** with an allenyl moiety at room temperature, giving (*E*)-**5ad** in 79% yield and 97% ee (eq 1).



To gain preliminary insight into the enantiodiscriminating step in the present reaction, several experiments were performed (Scheme 2). When the reaction of aldehyde **3a** with ester **2a** catalyzed by (*S*)-Ba-**1a** was analyzed at the initial stage (0.5 h), only trace amounts of **5aa** and **6aa** were observed. Instead, α -adduct **4aa** was obtained in 13% yield, but with poor dr and ee (Scheme 2a, dr = 1.6:1, major/minor = 0/15% ee). The poor ee for **4aa** indicated that the isomerization step from **4** to **5** was highly enantioselective. In fact, when diastereomixtures of racemic **4aa** were treated with (*S*)-Ba-**1a** in the presence of 1 equiv of **2a**, (*E*)-**5aa** was obtained in 69% yield and 99% ee after 24 h (Scheme 2b). The yield and ee of **5aa** in Scheme 2b suggested the

Scheme 2. Mechanistic Studies of Ba-Catalyzed DYKAT

presence of a retro-aldol reaction under the reaction conditions, and this was further confirmed by the crossover experiment shown in Scheme 2c. Treatment of aldehyde **3g** and racemic **4aa** with (*S*)-Ba-**1a** in DME gave both **5aa** (24% yield, 87% ee) and **5ga** (35% yield, 96% ee) after 24 h. These results confirmed that the present system is a Ba-**1**-catalyzed DYKAT involving a direct aldol/retro-aldol process.

In summary, we have developed a Ba-catalyzed DYKAT involving a direct aldol/retro-aldol reaction of β,γ -unsaturated ester donors. α -Alkylidene- β -hydroxy esters were obtained from aryl, heteroaryl, alkenyl, and alkyl aldehydes under simple proton-transfer conditions in 99–87% ee and >20:1 to 15:1 α/γ selectivity. Further studies to improve the reaction rate and catalyst loading are ongoing.

Acknowledgment. This work was supported by Grants-in-Aid for Scientific Research (S) (to M.S.) and Scientific Research on Priority Areas (20037010, Chemistry of Concerto Catalysis, to S.M.). A.Y. thanks JSPS for a fellowship.

Supporting Information Available: Experimental procedures, spectral data for new compounds, and determination of stereochemistry. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) For a general review of asymmetric aldol reactions, see: Geary, L. M.; Hultin, P. G. *Tetrahedron: Asymmetry* **2009**, *20*, 131.
- (2) Reviews of direct aldol reactions: (a) *Modern Aldol Reactions*; Mahrwald, R., Ed.; Wiley-VCH: Weinheim, Germany, 2004. For organocatalytic direct aldol reactions, also see: (b) Notz, W.; Tanaka, F.; Barbas, C. F., III. *Acc. Chem. Res.* **2004**, *37*, 580. (c) Mukherjee, S.; Yang, J. W.; Hoffmann, S.; List, B. *Chem. Rev.* **2007**, *107*, 5471.
- (3) (a) Ito, Y.; Sawamura, M.; Hayashi, T. *J. Am. Chem. Soc.* **1986**, *108*, 6405. (b) Kuwano, R.; Miyazaki, H.; Ito, Y. *Chem. Commun.* **1998**, 71.
- (4) (a) Trost, B. M.; Malhotra, S.; Fried, B. A. *J. Am. Chem. Soc.* **2009**, *131*, 1674. For early works, also see: (b) Yao, W.; Wang, J. *Org. Lett.* **2003**, *5*, 1527. (c) Hasegawa, K.; Arai, S.; Nishida, A. *Tetrahedron* **2006**, *62*, 1390.
- (5) (a) Ooi, T.; Kameda, M.; Taniguchi, M.; Maruoka, K. *J. Am. Chem. Soc.* **2004**, *126*, 9685. For early works, also see: (b) Gasparski, C. M.; Miller, M. J. *Tetrahedron* **1991**, *47*, 5367. (c) Yoshikawa, N.; Shibasaki, M. *Tetrahedron* **2002**, *58*, 8289.
- (6) Thiazolidinedione: (a) Evans, D. A.; Downey, C. W.; Hubbs, J. L. *J. Am. Chem. Soc.* **2003**, *125*, 8706. Malonic acid half thioester: (b) Magdziak, D.; Lalic, G.; Lee, H. M.; Fortner, K. C.; Aloise, A. D.; Shair, M. D. *J. Am. Chem. Soc.* **2005**, *127*, 7284. (c) Fortner, K. C.; Shair, M. D. *J. Am. Chem. Soc.* **2007**, *129*, 1032. α -Isothiocyanato imide: (d) Willis, M. C.; Cutting, G. A.; Piccio, V. J.-D.; Durbin, M. J.; John, M. P. *Angew. Chem., Int. Ed.* **2005**, *44*, 1543. (e) Li, L.; Klauber, E. G.; Seidel, D. *J. Am. Chem. Soc.* **2008**, *130*, 12248. Alkyl nitrile: (f) Suto, Y.; Tsuji, R.; Kanai, M.; Shibasaki, M. *Org. Lett.* **2005**, *7*, 3757.
- (7) For alternative catalytic asymmetric C–C bond-forming reactions for chiral β -hydroxy ester synthesis, see the following reviews: (a) Catalytic asymmetric reductive aldol reactions: Garner, S. A.; Kricheldorf, M. J. In *Modern Reduction Methods*; Andersson, P. G.; Munslow, I. J., Eds.; Wiley-VCH: Weinheim, Germany, 2008; p 387. (b) Catalytic asymmetric Morita–Baylis–Hillman reactions: Masson, G.; Housseman, C.; Zhu, J. *Angew. Chem., Int. Ed.* **2007**, *46*, 4614.
- (8) For a review of DYKAT, see: (a) Steinreiber, J.; Faber, K.; Griengl, H. *Chem.–Eur. J.* **2008**, *14*, 8060. For DYKAT involving a aldol/retro-aldol sequence, see: (b) Mascarenhas, C. M.; Miller, S. P.; White, P. S.; Morken, J. P. *Angew. Chem., Int. Ed.* **2001**, *40*, 601. (c) Gnanadesikan, V.; Horiuchi, Y.; Ohshima, T.; Shibasaki, M. *J. Am. Chem. Soc.* **2004**, *126*, 7782. (d) Córdova, A.; Ibrahim, I.; Casas, J.; Sundén, H.; Engqvist, M.; Reyes, E. *Chem.–Eur. J.* **2005**, *11*, 4772. (e) Steinreiber, J.; Schürmann, M.; Wolberg, M.; van Assema, F.; Reisinger, C.; Fesko, K.; Mink, D.; Griengl, H. *Angew. Chem., Int. Ed.* **2007**, *46*, 1624.
- (9) For a review of alkaline-earth metal catalysts for direct C–C bond-forming reactions, see: (a) Kazmaier, U. *Angew. Chem., Int. Ed.* **2009**, *48*, available online (DOI: 10.1002/anie.200901261). For examples of chiral Ba aryloxide catalysts, see: (b) Yamada, Y. M. A.; Shibasaki, M. *Tetrahedron Lett.* **1998**, *39*, 5561. (c) Saito, S.; Kobayashi, S. *J. Am. Chem. Soc.* **2006**, *128*, 8704. (d) Yamatsugu, K.; Yin, L.; Kamijo, S.; Kimura, Y.; Kanai, M.; Shibasaki, M. *Angew. Chem., Int. Ed.* **2009**, *48*, 1070.
- (10) β,γ -Unsaturated esters have been used for direct catalytic asymmetric Mannich-type reactions (see: Yamaguchi, A.; Aoyama, N.; Matsunaga, S.; Shibasaki, M. *Org. Lett.* **2007**, *9*, 3387). Benzyl crotonate (α,β -unsaturated ester) was not applicable for the present system because the γ -proton in crotonate is less acidic than the α -proton in β,γ -unsaturated esters.
- (11) Detailed results comparing (*S*)-Ba-**1a** with (*S*)-Ba-**1b** are shown in the Supporting Information.

JA904575E