FULL PAPER

Synthesis of carbido and related derivatives from calcium carbide and ruthenium carbonyl clusters

Michael I. Bruce,*^a Natasha N. Zaitseva,^a Brian W. Skelton^b and Allan H. White^b

- ^a Department of Chemistry, University of Adelaide, Adelaide, South Australia 5005, Australia. E-mail: michael.bruce@adelaide.edu.au
- ^b Department of Chemistry, University of Western Australia, Crawley, Western Australia 6009, Australia

Received 6th March 2002, Accepted 25th July 2002 First published as an Advance Article on the web 13th September 2002

Reactions between a carbide anion $(C_2^{2^-})$ source and ruthenium cluster carbonyls are reported for the first time. The reaction between CaC₂ and Ru₃(CO)₁₂, carried out in thf, affords putative Ca[Ru₁₀(C)₂(CO)₂₄], which can be converted into [ppn]₂[Ru₁₀(C)₂(CO)₂₄] (1) in overall 60% yield. The X-ray structure of 1 as its 2CH₂Cl₂ solvate is described, together with those of the minor products Ru₄(µ₄-HC₂H)₂(µ-CO)₂(CO)₉ (3), Ru₆C(µ₃-HC₂Me)(CO)₁₅ (4) and Ru₄(µ₄-CHCHCO)(µ-CO)(CO)₁₀(tmeda) (5). The latter contains an Ru-spiked Ru₃ core carrying a CHCHCO ligand; the tmeda chelates the spike Ru atom. The only product from Ru₃(µ-dppm)(CO)₁₀ and CaC₂ is the known compound Ru₆(µ₄-CCH₂)₂(µ-dppm)₂(CO)₁₂ (6), formed in 24% yield.

Introduction

The chemistry of metal cluster carbonyls containing carbido ligands has a long history, the first complex, $Fe_5C(CO)_{15}$, being discovered *via* an X-ray structural determination in 1962.¹ These interesting compounds have usually been made by pyrolysis of a simpler precursor carbonyl, the source of the carbido atom being shown to be one of the carbonyl groups.^{2,3} In the ruthenium system, extensive studies have revealed several intermediate clusters are formed *en route* from $Ru_3(CO)_{12}$ to $Ru_6C(CO)_{17}$.⁴ The chemistry of carbido cluster complexes has been reviewed.⁵

Other synthetic routes have been used to make cluster carbides. For example, condensation of anionic Group 9 cluster carbonyls with halocarbon derivatives results in formation of complexes such as $[Rh_6C(CO)_{15}]^{2-}$, from CCl_4 and $[Rh(CO)_4]^{-,6}$ or $[Co_6C(CO)_{15}]^{2-}$, from $Co_3(\mu_3$ -CCl)(CO)₉ and $[Co(CO)_4]^{-,7}$ Extensive hydrogenation of a ruthenium cluster-bonded ethynyl group has resulted in the addition of up to three H atoms to one carbon of the C₂ fragment, with a consequent decrease in the C–C bond order. In the last stage, cleavage of the C–C bond afforded a carbido complex, the second carbon atom being fully reduced to a methyl group.⁸

Although several experimental procedures have now been developed to produce carbido clusters in reasonable yield,⁹ to the best of our knowledge, there are no reported uses of main group carbides as sources of the carbido carbon atoms. This may be because of (i) the relative difficulties attendant on the production and manipulation of alkali metal acetylides, even though Li₂C₂·en is available commercially, or (ii) reports that in reactions with metal carbonyl halides, the acetylides tend to act as one-electron reducing agents. For example, the reaction of MnBr(CO)₅ with CaC₂ is reported to give Mn₂(CO)₁₀, while Fe₃(CO)₁₂ and Co₄(CO)₁₂ give [Fe₃(μ -H)(CO)₁₁]⁻ and [Co₆(CO)₁₄]⁴⁻, respectively.¹⁰ In addition, {Re(CO)₅ 2(μ -C=C) cannot be prepared from Re(FBF₃)(CO)₅ and Li₂C₂ or Na₂C₂.¹¹ Also relevant is the use of calcium carbide to form transition metal carbides *via* metathesis reactions with halides or oxides in the solid state, which has been reported recently.¹²

We have found that calcium carbide is a useful reagent for the preparation of the anionic dicarbido cluster $[Ru_{10}(C)_2$ -

 $(CO)_{24}]^{2-}$, which was first described in 1982 as the product from thermolysis (210–230 °C, 80 h) of $[Ru_6C(CO)_{16}]^{2-}$ in refluxing tetraglyme.¹³ Several other products have been isolated from the reaction mixture, but these are formed in low yields and probably result from reactions of various cluster complexes with ethyne formed from the calcium carbide reacting with adventitious water.

Results and discussion

When a 30-fold excess of solid calcium carbide (which is approximately 80% pure) is added to a solution of $Ru_3(CO)_{12}$ in thf and the mixture heated overnight, a dark-coloured solution containing solid material is obtained. Removal of solvent and filtration of an acetone extract of the residue into a solution of [ppn]Cl in CH₂Cl₂ produces a dark purple solution, from which the salt [ppn]₂[Ru₁₀(C)₂(CO)₂₄] (1) can be isolated in 60% yield. We have not yet characterised any complexes from the solid; based on the formulation Ca[Ru₁₀C₂(CO)₂₄], the yield is *ca*. 70%. The ν (CO) band frequencies in the IR spectrum are similar to those reported earlier, while the ¹³C resonances of the CO groups are found between δ 185 and 205 as five singlets, the interstitial carbide giving a singlet at δ 456.89 (*cf.* δ 456 in the [NEt₄]⁺ salt). In the electrospray (ES) mass spectrum, the doubly-charged parent anion is found at *m*/*z* 855.



(1) CO groups on Ru(3,6) omitted for clarity

While the formation of **1** can result from thermolysis of smaller ruthenium cluster carbides, we showed independently that heating $Ru_3(CO)_{12}$ in the overnight does not result in any cluster condensation. Consequently, the formation of **1** is a direct result of the interaction of CaC_2 with the carbonyl, an interesting and unusual cleavage of the C_2^{2-} anion taking place. Complex **1** was originally obtained by extended thermolysis of the mono-carbido cluster $[NEt_4]_2[Ru_6C(CO)_{16}]$; it was noted that the yield was dependent on the counter cation, only decomposition resulting when Na, $[ppn]^+$ or $[NEt_3(CH_2Ph)]^+$ were used.¹³ Fusion of two Ru_6 clusters with concomitant elimination of two $Ru(CO)_n$ fragments [or one $Ru_2(CO)_y$ fragment] had occurred.

With a similar experiment carried out with only a 10-fold excess of CaC_2 , the mother liquor from the separation of 1 was evaporated and the residue extracted with dichloromethane. Preparative t.l.c. enabled separation of several minor components, three of which have been purified and characterised initially by single-crystal X-ray crystallography as Ru₄(µ₄-HC₂-H)(CO)₁₂ (2), $Ru_4(\mu_4-HC_2H)_2(\mu-CO)_2(CO)_9$ (3) and $Ru_6C(\mu_3-\mu_4)$ $HC_2Me)(CO)_{15}$ (4). Subsequent assignments of their spectra (IR, NMR, ES MS) on the bases of their solid-state structures are discussed below. Complex 2 has been described before 14 and 3 is the prototype for a series of substituted complexes $M_4(\mu_4$ - $RC_{2}R')_{2}(CO)_{11}$ (M = Fe, R = H, R' = Et; M = Ru, R = Ph, R' = Me, Ph) which have been reported earlier and which contain two alkyne molecules attached to opposite sides of a buckled M₄ rhomb.¹⁵ It was characterised only by a single-crystal X-ray structural determination, while the pattern of v(CO) peaks in its IR spectrum is similar to those reported for the substituted analogues.



(3) CO groups on Ru(1,1') omitted for clarity



Complex **4** is a simple μ_3 -propyne complex of the hexanuclear carbido carbonyl Ru₆C(CO)₁₇. A range of analogous complexes, including the but-2-yne complex, has been described by others.^{16,17} In **4**, the propyne ligand is attached in the usual

3880 J. Chem. Soc., Dalton Trans., 2002, 3879–3885

 μ_3 - η^1 : η^1 : η^2 mode to one triangular face of the octahedral Ru₆C cluster, with bond parameters similar to those found for the but-2-yne analogue.¹⁶ All the CO ligands are terminal, as found in the solid-state structure, and give rise to ν (CO) bands between 2088 and 1982 cm⁻¹ in the IR spectrum. In the ¹H NMR spectrum, the Me and H protons of the alkyne resonate at δ 1.67 and 6.97, respectively. We have described in an earlier account the formation of Ru₆(μ -H)(μ_4 -C)(μ_4 -CCMe)(μ -CO)-(CO)₁₆ (5) during the reaction between Ru₃(CO)₁₀(NCMe)₂ and ethyne.¹⁸ Disproportionation of two molecules of C₂H₂ to give a carbide and propyne ligands occurred, and we suggest that a similar reaction takes place here.



In an effort to facilitate the initial reaction between Ru_3 -(CO)₁₂ and CaC₂, we added some tetramethyldiaminoethane (tmeda) to the reaction mixture, hoping that the calcium cations would be complexed by the diamine. As described in the Experimental section, little change was observed in the formation of **1**, but among the minor products, the cluster $Ru_4(\mu_4$ -CHCHCO)(μ -CO)(CO)₁₀(tmeda) (6) was characterised. We believe this to be the first example of a ruthenium cluster containing the tmeda ligand.



The molecular structure of 6 is based on an Ru-spiked Ru₃ triangle, which carries a CHCHCO ligand, formed by combination of a CO ligand with an ethyne molecule (again, presumably formed from the CaC₂). The chelating tmeda ligand is attached to the Ru spike. Complex 6 contains the familiar allylic ligand CHCHCO⁻ which is closely related to that found in $Ru_3(\mu\text{-}H)\{\mu_3\text{-}CHCHC(OH)\}(CO)_8(PPh_3).^{19}$ In the reactions which lead to 6, deprotonation of the hydroxy group has occurred and the resulting anion has become attached to the fourth Ru atom. The origin of the latter is not clear at this stage-one possibility is that an intermediate mononuclear species, such as Ru(CO)₃(tmeda), may be formed. The proximity of the fourth Ru atom results in Ru-Ru bond formation, giving the spiked cluster. The new bond is bridged by one CO group, which gives rise to the v(CO) band found at 1864 cm⁻¹ in the IR spectrum. In the ¹H NMR spectrum, the two protons on the CHCHCO ligand give two doublets at δ 6.28 and 7.85, while those of the tmeda ligand occur at δ 3.14 (CH₂), and 2.37 and 2.60 (Me). Surprisingly, the oxo anion has not attacked a CO, as found in the formation of Au₃Ru₃{µ₃-OC- $(O)CPh_2CHC\}(CO)_8(PPh_3)_3.^{20}$

We have also examined the reaction between CaC_2 and $Ru_3(\mu$ -dppm)(CO)₁₀, which often affords products different in structure from those obtained from the parent carbonyl. In this

instance, the sole product that we have been able to characterise is the hexanuclear vinylidene cluster $Ru_6(\mu_4$ -CCH₂)₂(μ -dppm)₂-(CO)₁₂ (7), isolated in only 18–24% yield, which we earlier obtained serendipitously and characterised as the benzene solvate from the reaction of $Ru_3(\mu$ -H)(μ_3 -C₂SiMe₃)(μ -dppm)-(CO)₇ with KF in MeOH.²¹ The present structure determination was carried out with a mixed CHCl₃–MeCN solvate, which showed considerable disorder (see Experimental section).



Molecular structures

Fig. 1–4 contain plots of the cation of 1 and molecules of 3, 4 and 6, respectively; significant bond parameters are given in Tables 1–4.



Fig. 1 Plot of the anion in $[ppn]_2[Ru_{10}(C)_2(CO)_{24}]$ (1), showing atom numbering scheme.



Fig. 2 Plot of a molecule of $Ru_4(\mu_4-HC_2H)_2(\mu-CO)_2(CO)_9$ (3). A crystallographic mirror plane passes through Ru(2,3) and carbonyls 21, 31, 32 and the C(1,2) alkyne group.



Fig. 3 Plot of a molecule of $Ru_6C(\mu_3-HC_2Me)(CO)_{15}$ (4).



Fig. 4 Plot of molecule 1 of $Ru_4(\mu_4$ -CHCHCO)(μ -CO)(CO)₁₀(tmeda) (6).

Table 1 Selected bond distances (Å) for $[ppn]_2[Ru_{10}(C)_2(\mu\text{-CO})_4\text{-}(CO)_{20}]\,(1)^{\,\alpha}$

Ru(1)-Ru(2) Ru(4)-Ru(5) Ru(3)-Ru(6) Ru(1)-Ru(6) Ru(2)-Ru(3) Ru(3)-Ru(4) Ru(5)-Ru(6) Ru(1)-Ru(7)	2.9575(6) 2.9659(5) 2.8670(5) 2.8962(5) 2.8962(5) 2.8813(5) 2.9023(5) 2.9537(6)	Ru(4)-Ru(10) Ru(5)-Ru(9) Ru(3)-Ru(7) Ru(3)-Ru(10) Ru(6)-Ru(8) Ru(6)-Ru(9) Ru(3)-Ru(8) Ru(3)-Ru(9)	2.8117(6) 2.8126(6) 2.8350(5) 2.8511(6) 2.8246(6) 2.8231(6) 3.1228(6) 3.1152(6)
Ru(1)-Ru(6)	2.9047(5)	Ru(3)-Ru(10)	2.8511(6)
Ru(2) - Ru(3) Ru(3) - Ru(4)	2.8962(5)	Ru(6) - Ru(8) Ru(6) - Ru(9)	2.8240(6)
Ru(5)-Ru(6)	2.9023(5)	Ru(3)- $Ru(8)$	3.1228(6)
$\operatorname{Ru}(1)$ – $\operatorname{Ru}(7)$	2.9537(6)	Ru(3)– $Ru(9)$	3.1152(6)
$\operatorname{Ru}(2) - \operatorname{Ru}(8)$	3.0081(6)	Ru(6) - Ru(7)	3.0888(6)
Ru(4) = Ru(9) Ru(5) = Ru(10)	2.9712(0)	Ru(0) - Ru(10) Ru(7) - Ru(9)	3.2279(5)
Ru(1)– $Ru(8)$	2.8198(6)	Ru(8) - Ru(10)	3.1020(5)
Ru(2)–Ru(7)	2.8353(6)		

^{*a*} Ru–Ru bond distances have been grouped by type. Ru–C distances 2.025–2.102(5); C(1)–C(2) 3.015(5) Å.

In the structure of the anion in $[ppn]_2[1]$, very similar to but more precisely defined than that found in $[NEt_4]_2[Ru_{10}(C)_2-(CO)_{24}]$ ($[NEt_4]_2[1]$),¹³ the core comprises two octahedra sharing an edge, one carbido atom being encapsulated in each. The Ru–Ru separations form several sets. The shortest are those bridged by CO ligands [2.8117–2.8353(6) Å], while Ru(3)– Ru(6), at 2.8670(5) Å, is shared between the octahedra. Of the non-bridged Ru–Ru vectors, those on the outside of the cluster are longer [2.9526–3.0081(6) Å] than the inner Ru(eq)–Ru(eq) separations [2.8813–2.9047(6) Å]. Several Ru–Ru distances are longer than 3 Å, ranging from Ru(2)–Ru(8) [3.0081(6) Å] to Ru(6)–Ru(10) [3.1472(6) Å], while the separations between the two pairs of apical Ru atoms are of the same order [Ru(7)–Ru(9) 3.2279(5), Ru(8)–Ru(10) 3.1020(5) Å]. Of some interest are the diagonal Ru(ap)–Ru(eq) distances involving atoms Ru(3,6); both atoms are involved in short [2.8231–2.8511(6) Å] and long interactions [3.1152–3.1472(6) Å], resulting in a twist being imparted to the bioctahedral cluster [torsion angles Ru(1,2,4,5) and Ru(2,1,5,4) –8.13 and –8.09(2)°]. The C(1) \cdots C(2) separation is 3.015(5) Å. The two [ppn]⁺ cations are bent, with N–P distances of 1.569–1.586(5) Å; P–N–P 141.7, 143.1(3)°.

Several reactions of **1** with unsaturated hydrocarbons have given derivatives in which the apical Ru atoms are considered to be either bonded or non-bonded.^{13,22} This feature has been discussed previously in terms of the cluster valence electron (c.v.e.) count for this anion (138 c.v.e.) and the occupation of both bonding and anti-bonding MOs as the HOMOs.^{13,22} This c.v.e. count arises from the presence of two edge-sharing octahedra, for which the Mingos procedure²³ gives 2 × 86 (Ru₆ octahedra) - 34 (Ru₂ edge) = 138 c.v.e. It seems to us that there is a continuum of Ru–Ru bonding interactions in all these complexes, the relative separations of the bonding and

Table 2 Selected bond distances (Å) and angles (°) for $Ru_4(\mu_4-HC_2-H)_2(\mu-CO)_2(CO)_9$ (3)^{*a*}

Ru(1)-Ru(2)	2.834(1)	Ru(2)–C(1)	2.11(2)
Ru(1)-Ru(3)	2.773(1)	Ru(2)–C(3)	2.31(1)
Ru(1)-Ru(1')	3.838(1)	Ru(3)–C(2)	2.14(2)
Ru(2)-Ru(3)	3.843(1)	Ru(3)–C(3)	2.28(1)
Ru(1)-C(1)	2.31(1)	C(1) - C(2)	1.39(3)
Ru(1)-C(2)	2.27(1)	C(3)–C(3')	1.37(2)
Ru(1)-C(3)	2.14(1)		
$\operatorname{Ru}(2)$ - $\operatorname{Ru}(1)$ - $\operatorname{Ru}(3)$	86.53(4)	Ru(2)-C(1)-C(2)	125(1)
Ru(1)-Ru(2)-Ru(1')	85.25(5)	Ru(3)-C(2)-C(1)	126(1)
Ru(1)- $Ru(3)$ - $Ru(1')$	87.57(5)	Ru(1)-C(3)-C(3')	125.2(9)
		1. 1. 1	

^a Primed atoms are related by the intramolecular plane.

Table 3 Selected bond distances (Å) for $Ru_6C(\mu_3-HC_2Me)(CO)_{15}$ (4)^{*a*}

Ru(1)–Ru(2)	2.8056(4)	Ru(2)–Ru(3)	2.9699(4)
Ru(1)-Ru(5)	2.8532(4)	Ru(1)-Ru(4)	2.9340(4)
Ru(2)-Ru(5)	2.7718(4)		
Ru(1)-Ru(6)	2.9423(4)	Ru(1)-C(2)	2.059(4)
Ru(2)-Ru(6)	2.9293(4)	Ru(2)-C(1)	2.189(4)
Ru(3)-Ru(5)	2.9088(5)	Ru(2)-C(2)	2.217(4)
Ru(4)-Ru(5)	2.9905(4)	Ru(5)-C(1)	2.046(4)
Ru(3)-Ru(4)	2.8943(4)		
Ru(3)-Ru(6)	2.8547(4)	C(1) - C(2)	1.400(6)
Ru(4)-Ru(6)	2.8882(4)	C(2)–C(3)	1.505(6)
Ru–C(carbide) 24.0(3)°.	2.028–2.069(4),	av. 2.05(1) Å.	C(1)-C(2)-C(3)

anti-bonding orbitals being determined by the electron density within the cluster, which is, in turn, determined by the nature of the other ligands present.

Of interest is the presence of two C atoms within the cluster, rather than the initial C₂ species present in the CaC₂ precursor. In closely related Group 9 carbido clusters, the presence of C or C₂ ligands has been related to the size of the cluster, the latter required a (larger) trigonal prismatic cavity.²⁴ However, the final geometry is not a simple consequence of electron count. In the isoelectronic $[Rh_{12}(C)_2(CO)_{24}]^{2-25}$ and $Rh_{12}(C_2)(CO)_{25}$ clusters, different cluster geometries are adopted as a result of 2e in the former being replaced by the sterically more demanding CO ligand in the neutral complex. In the present case, the more electron-rich Ru atoms result in octahedral cavities being formed, with resulting cleavage of the C–C bond. Further discussion of the electronic structures of these and related complexes is deferred until DFT studies have been completed.

The molecule of **3** has a crystallographic plane of symmetry passing through Ru(2,3)C(21,31,32) and the midpoint of the C(3)–C(3') vector. The four Ru atoms form a rhomb bent along the Ru(1) · · · Ru(1') vector, with two different bonded Ru–Ru distances: Ru(1)–Ru(2) [2.834(1) Å] and Ru(1)–Ru(3), bridged by CO(13) [2.773(1) Å]. Angles subtended at one Ru atom by the others are 85.25–87.57(5)°. The two ethyne molecules lie on opposite sides of the Ru₄ core, each being attached by two σ [Ru(1)–C(3) 2.14(2), Ru(3)–C(2) 2.14(2) Å] and two π [Ru–C 2.27–2.31(1) Å] bonds, the differences arising because only Ru(2) is not attached to a μ -CO ligand. The C(1)–C(2) separation is 1.39(3) Å. Unlike previous structurally characterised examples which contain only one,¹⁵ 3 contains two bridging CO ligands. This cluster contains 62 c.v.e.

For **4**, the octahedral Ru₆C cluster has Ru–Ru separations [2.9088–2.9905(4) Å] in the non-alkyne bridged faces and internal Ru–C(carbide) distances [2.028–2.069(4), av. 2.05(1) Å] which are similar to those found in the parent cluster²⁷ and the but-2-yne analogue.¹⁶ The Ru₃ face [Ru(1,2,5)] supporting the alkyne is slightly contracted [Ru–Ru 2.7718–2.8532(4) Å], as is the opposite face [Ru(3,4,6): Ru–Ru 2.8547–2.8943(4) Å]. The Ru–C(alkyne) distances [Ru(1)–C(2) 2.059(4), Ru(5)–C(1) 2.046(4); Ru(2)–C(1,2) 2.189, 2.217(4) Å] are similar to those found in Ru₃(µ₃-HC₂Me)(µ-CO)(CO)₉ and related Ru₆C clusters containing µ₃-alkynes. The C(1)–C(2) separation is 1.400(6) Å and the bend-back angle for the Me substituent is 56.0(3)°. All CO groups are terminal, a configuration that is preserved in solution.

Comparison of the structure of **6** with that of $Ru_3(\mu-H)-{\mu_3-CHCHC(OH)}(CO)_8(PPh_3)$ (**8**)¹⁸ shows similar Ru–Ru separations in the Ru₃ triangle, with the exception of Ru(2)–Ru(3) [2.8910, 2.8627(2) Å; values for molecules 1, 2 given], which here is bridged by CO rather than by H as in **8**. The Ru(3)–Ru(4) distance is 2.8259, 2.8111(3) Å, and this vector is bridged both by an η^1 -CO ligand and by the $\eta^2(C,O)$ -CO portion of the CHCHCO ligand [Ru(3)–C(3) 2.030, 2.027(2), Ru(4)–O(4) 2.075, 2.073(2) Å]. The tmeda ligand chelates

Table 4 Selected bond parameters for $Ru_4(\mu_4$ -CHCHCO)(μ -CO)(CO)_{10}(tmeda) (6; values for molecules 1, 2)

$\mu = 0$ ($\mu = 0$)(μ				
Ru(1)–Ru(2)	2.7699, 2.7656(3)	Ru(4)–O(4)	2.075, 2.073(2)	
Ru(1)-Ru(3)	2.8613, 2.8702(3)	Ru(4) - N(5)	2.232, 2.230(2)	
Ru(2)-Ru(3)	2.8910, 2.8627(2)	Ru(4)–N(6)	2.247, 2.250(2)	
Ru(3)-Ru(4)	2.8259, 2.8111(3)		· · · · · ·	
	· · · · · ·	C(1)-C(2)	1.401, 1.401(3)	
Ru(1)-C(1)	2.224, 2.235(2)	C(2) - C(3)	1.431, 1.435(3)	
Ru(1) - C(2)	2.273, 2.286(2)	C(3) - O(4)	1.319, 1.312(2)	
Ru(1) - C(3)	2.387, 2.383(2)	C(5) - N(5)	1.490, 1.487(3)	
Ru(2)-C(1)	2.069, 2.089(3)	C(5)-C(6)	1.503, 1.499(3)	
Ru(3) - C(3)	2.030, 2.027(3)	C(6) - N(6)	1.493, 1.483(3)	
	· · · · ·		· · · · · ·	
C(1)-C(2)-C(3)	120.4, 119.8(3)	C(3) - O(4) - Ru(4)	107.5, 107.9(1)	
C(2)–C(3)–O(4)	119.2, 119.8(2)	N(5)-Ru(4)-N(6)	81.11, 81.68(7)	

Ru(4), with Ru(4)–N(5,6) 2.232, 2.230; 2.247, 2.250(2) Å and an N(5)–Ru(4)–N(6) bite angle of 81.11, 81.68(7)°. The N atoms are *trans* to the Ru–Ru bond and to CO(41) and the tmeda takes up an envelope conformation. The geometry of the CHCHCO fragment is similar to that found in **8**, with C(1)– C(2) 1.401, 1.401(3), C(2)–C(3) 1.431, 1.435(3) and C(3)–O(4) 1.319, 1.312(2) Å; the angles C(1)–C(2)–C(3) and C(2)–C(3)– O(4) are 120.4, 119.8(3) and 119.2, 119.8(2)°, respectively. Normal σ [Ru(2)–C(1) 2.069, 2.089(3); Ru(3)–C(3) 2.030, 2.027(2) Å] and π bonds [Ru(1)–C(1,2,3) 2.224–2.387(2) Å] are found to the Ru₃ triangle.

Conclusion

We have shown for the first time that it is possible to use calcium carbide as a source of carbido carbons in ruthenium clusters, the major product from its reaction with $Ru_3(CO)_{12}$ being the decanuclear dicarbido anion $[Ru_{10}(C)_2(CO)_{24}]^{2-}$. An unusual feature of this reaction is the cleavage of the C_2^{2-} anion present in crystalline CaC₂. Some minor products have also been characterised: these contain ethyne or derived ligands and we are inclined to suggest that these complexes are formed from traces of ethyne present in the reaction mixture, possibly generated by the reaction of CaC₂ with water or another proton source.

Experimental

General reaction conditions

Reactions were carried out under an atmosphere of nitrogen, but no special precautions were taken to exclude oxygen during work-up. Common solvents were dried and distilled under nitrogen before use. Elemental analyses were performed by Chemical and Micro-analytical Services p/l, Melbourne. Preparative t.l.c. was carried out on glass plates (20×20 cm) coated with silica gel (Merck 60 GF₂₅₄, 0.5 mm thickness).

Instrumentation

IR: Perkin-Elmer 1720X FT IR. NMR: Bruker CXP300 orACP300 (¹H at 300.13 MHz, ¹³C at 75.47 MHz) or Varian Gemini 200 (¹H at 199.8 MHz, ¹³C at 50.29 MHz) spectrometers. Unless otherwise stated, spectra were recorded using solutions in CDC1₃ in 5 mm sample tubes. ES mass spectra: VG Platform 2 or Finnigan LCQ. Solutions were directly infused into the instrument. Chemical aids to ionisation were used as required.²⁸

Reagents

 $Ru_3(CO)_{12}^{29}$ and $Ru_3(\mu$ -dppm)(CO)_{10}^{30} were prepared according to the cited methods. CaC_2 (Aldrich) was a commercial sample, used as received. It appeared to be about 80% pure, the major impurity being Ca(OH)₂.

Reactions of CaC₂

(a) With Ru₃(CO)₁₂. A suspension of Ru₃(CO)₁₂ (100 mg, 0.16 mmol) and CaC₂ (100 mg, 1.6 mmol) in thf (20 ml) was heated at reflux point overnight to give a brown solution above a dark coloured precipitate. After removal of solvent (rotary evaporator), the residue was extracted separately with CH₂Cl₂ (fraction A) and acetone (fraction B). Fraction A was separated by preparative t.l.c. (hexane–acetone 7:3) to give several coloured bands: (a) a brown band ($R_{\rm f}$ 0.74) contained Ru₆C-(μ_3 -HC₂Me)(CO)₁₅ (4) (2.3 mg, 2.7%), isolated as dark red crystals (from CH₂Cl₂–MeOH). IR (cyclohexane): ν (CO) 2088w, 2062vw, 2043s, 2057m, 2022m, 2015m, 1994vw, 1982vw cm⁻¹. ¹H NMR (CDCl₃): δ 1.67 (s, 3H, Me), 6.97 (s, 1H, CH); (b) a red band ($R_{\rm f}$ 0.71) yielded Ru₄(μ_4 -HC₂H)(CO)₁₂ (2)

(4.8 mg, 5.3%), identified by comparison [IR ν (CO)] with an authentic sample; (c) a yellow band (R_f 0.68) contained Ru₄-(μ_4 -HC₂H)₂(μ -CO)₂(CO)₉ (**3**) (1.1 mg, 1.0%), obtained as dark yellow crystals (from CH₂Cl₂-pentane), identified by single-crystal X-ray diffractometry. IR (cyclohexane): ν (CO) 2093w, 2063m, 2040s, 2030vs, 2019 (sh), 1985m, 1954vw, 1848w (br) cm⁻¹.

Fraction B was also purified by t.l.c. (acetone-hexane 2:1) to give a dark purple band ($R_{\rm f}$ 0.33), probably containing $[Ru_{10}(C)_2(CO)_{24}]^{2-}$ as its solvated Ca^{2+} salt (43 mg). IR (thf): v(CO) 2049vw, 2004vs, 1960w, 1930w cm⁻¹. The compound is moderately stable in air, both in solution and as a solid. It was further characterised by conversion to the [ppn]⁺ salt by dissolving 20 mg in acetone (5 ml) and adding [ppn]Cl (1.4 mg, 0.025 mmol). After stirring at r.t. for 30 min, evaporation and extraction of the residue with CH₂Cl₂ and further purification by preparative t.l.c. (acetone-hexane 2:1) gave [ppn]₂[Ru₁₀(C)₂- $(CO)_{24}$] (1) (27 mg, 84%) as dark purple crystals (from CH_2Cl_2 -MeOH). Anal. found: C, 42.30; H, 2.04; N, 0.97; C₉₈H₆₀N₂O₂₄-P₄Ru₁₀ calc.: C, 42.28; H, 2.17; N, 1.01%. IR (CH₂Cl₂): v(CO) 2055vw, 2003vs, 1960w, 1927w, 1783m (br) cm⁻¹ [*cf*. 2046w, 2003vs, 1962w (sh), 1922w (sh), 1798m cm⁻¹ (thf)¹³⁴]. ¹H NMR (CDCl₃): δ 7.24–7.623 (m, Ph). ¹³C NMR (CDCl₃): δ 125.91– 133.59 (m, Ph), 185.40, 193.97, 202.36, 203.38, 205.29 (5 × s, CO), 456.89 (s, C). ³¹P NMR (CDCl₃): δ 22.33 (s, PPh₃). ES MS $(C_2H_4Cl_2, \text{ positive ion, } m/z)$: 538, $[ppn]^+$; (negative ion, m/z): 855, $[Ru_{10}C_2(CO)_{24}]^{2-}$ (isotopic peaks separated by 0.5 units).

A later experiment was carried out using $Ru_3(CO)_{12}$ (100 mg, 0.16 mmol) and CaC_2 (300 mg, 4.7 mmol) in thf (20 ml) as above, to give a dark brown solution over a dark coloured precipitate. The thf solution gave $Ca[Ru_{10}(C)_2(CO)_{24}]$ (9 mg), while extraction of the precipitate with acetone gave the same compound (46 mg). The combined product was dissolved in acetone (5 ml) and [ppn]Cl (43 mg, 0.075 mmol) was added. After 30 min at r.t., evaporation and crystallisation of the residue (CH₂Cl₂–MeOH) gave [ppn]₂[Ru₁₀(C)₂(CO)₂₄] (1) (75 mg, 57%) as dark purple crystals.

(b) With $Ru_3(CO)_{12}$ and tmeda. A similar reaction was carried out between $Ru_3(CO)_{12}$ (150 mg, 0.24 mmol) and CaC_2 (150 mg, 2.4 mmol) in thf (20 ml) containing tmeda (6 drops, ~40 mg). After removal of solvent, the residue was extracted with CH_2Cl_2 (fraction A) and acetone (fraction B). Separation of fraction A by preparative t.l.c. (acetone–hexane 2:1) gave a major red band (R_f 0.63) from which dark red crystals of $Ru_4(\mu_4$ -CHCHCO)-(CO)₁₁(tmeda) (6) (26 mg, 16.7%) were isolated. Anal. found: C, 27.11; H, 2.09; N, 3.25; $C_{20}H_{18}N_2O_{12}Ru_4$ calc.: C, 27.75; H, 2.06; N, 3.17%. IR (cyclohexane): v(CO) 2075m, 2032s, 2022vs, 2007w, 1998w (br), 1993w (br), 1972w, 1959w, 1949w, 1864w (br) cm⁻¹. ¹H NMR (CDCl₃): δ 2.37, 2.60 (2 × s, 2 × 6H, Me), 3.14 (m, 4H, CH₂), 6.28 [d, J(HH) 6.9, 1H, CH], 7.85 [d, J(HH) 6.9 Hz, 1 H, CH]. Other bands were separated but not identified.

Fraction B was purified by preparative t.l.c. (as above) to give putative $Ca[Ru_{10}(C)_2(CO)_{24}]$ (41 mg, 33%).

(c) With Ru₃(μ -dppm)(CO)₁₀. Tmeda (6 drops) was added to a suspension of Ru₃(μ -dppm)(CO)₁₀ (100 mg, 0.1 mmol) and CaC₂ (64 mg, 1 mmol) in thf (10 ml). The mixture was heated at reflux point overnight, after which the solvent was removed and the residue was extracted with CH₂Cl₂. Separation by preparative t.l.c. (acetone–hexane 3:7) afforded dark red Ru₆-(μ_4 -CCH₂)₂(μ -dppm)₂(CO)₁₂ (7) (22 mg, 24%), identified by comparison (IR, NMR) with an authentic sample²¹ and also by single-crystal X-ray structure determination of a disordered CHCl₃–MeCN solvate (the previous determination was carried out with a C₆H₆ solvate). Several minor products were also present but not characterised.

A similar reaction was carried out in the presence of Me_3NO instead of tmeda and produced 7 in 18% yield.

Structure determinations

Full spheres of data were measured at *ca*. 153 K using a Bruker AXS CCD area-detector instrument. N_{total} data merging after "empirical" /multiscan corrections (proprietary software) to N unique (R_{int} quoted), N_o with $F > 4\sigma(F)$ being used in the refinements. All data were measured using monochromatic Mo-K α radiation, $\lambda = 0.71073$ Å. In the refinements, anisotropic thermal parameter forms were used for the non-hydrogen atoms, (U_{iso})_H being constrained at estimated values. Conventional residuals R, R_w on |F| are quoted, statistical weights being employed. Neutral atom complex scattering factors were used; computation used the XTAL 3.7 program system.³¹ Pertinent results are given in the figures (which show nonhydrogen atoms with 50% probability amplitude displacement ellipsoids) and tables.

Crystal and refinement data

(1) $[ppn]_2[Ru_{10}(C)_2(CO)_{24}] \cdot 2CH_2Cl_2 \equiv C_{98}H_{60}N_2O_{24}P_4Ru_{10} \cdot 2CH_2Cl_2, M = 2954.0.$ Triclinic, space group $P\overline{1}$, a = 12.313(1), b = 18.211(2), c = 23.540(2) Å, a = 101.622(3), $\beta = 94.226(3)$, $\gamma = 94.918(2)^\circ$, V = 5129 Å³, D_c (Z = 2) = 1.913 g cm⁻³. Crystal $0.32 \times 0.30 \times 0.13$ mm, μ (Mo-K α) = 1.7 mm⁻¹, $T_{min,max} = 0.56$, $0.73. 2\theta_{max} = 65^\circ$; $N_{total} = 103731$, N = 37560 ($R_{int} = 0.044$), $N_o = 27340$, R = 0.052, $R_w = 0.073$.

(3) $\operatorname{Ru}_4(\mu_4\operatorname{-HC}_2\operatorname{H})_2(\mu\operatorname{-CO})_2(\operatorname{CO})_9 \equiv \operatorname{C}_{15}\operatorname{H}_4\operatorname{O}_{11}\operatorname{Ru}_4, M = 764.5.$ Orthorhombic, space group $Cmc2_1, a = 11.337(2), b = 13.456(2), c = 12.805(2)$ Å, V = 1953 Å³, $D_c (Z = 4) = 2.599$ g cm⁻³. Crystal $0.30 \times 0.10 \times 0.08$ mm, $\mu(\operatorname{Mo-K}\alpha) = 3.1$ mm⁻¹, $T_{\min,\max} = 0.69, 0.86. 2\theta_{\max} = 62.5^\circ; N_{total} = 14802, N = 1733 (R_{int} = 0.051), N_o = 1631, R = 0.050, R_w = 0.107. x_{abs} = 0.07(18).$ (4) $\operatorname{Ru}_6\operatorname{C}(\mu_3\operatorname{-HC}_2\operatorname{Me})(\operatorname{CO})_{15} \equiv \operatorname{C}_{19}\operatorname{H}_4\operatorname{O}_{15}\operatorname{Ru}_6, M = 1078.7.$

(4) $\operatorname{Ru}_6C(\mu_3-\operatorname{HC}_2\operatorname{Me})(\operatorname{CO})_{15} \equiv \operatorname{C}_{19}\operatorname{H}_4\operatorname{O}_{15}\operatorname{Ru}_6$, M = 1078.7. Monoclinic, space group C2/c, a = 16.9560(9), b = 9.4025(5), c = 32.929(2) Å, $\beta = 93.079(1)^\circ$, V = 5242 Å³, $D_c(Z=8) = 2.733$ g cm⁻³. Crystal 0.28 × 0.24 × 0.13 mm, $\mu(\operatorname{Mo-K}\alpha) = 3.4 \text{ mm}^{-1}$, $T_{\min,\max} = 0.58$, 0.84. $2\theta_{\max} = 75^\circ$; $N_{\text{total}} = 52942$, N = 13654 ($R_{\text{int}} = 0.042$), $N_o = 11414$, R = 0.038, $R_w = 0.071$.

(6) $\operatorname{Ru}_4(\mu_4\text{-CHCHCO})(\mu\text{-CO})(\operatorname{CO})_{10}(\operatorname{tmeda})\cdot 0.5\operatorname{CH}_2\operatorname{Cl}_2 = C_{20}\operatorname{H}_{18}\operatorname{N}_2\operatorname{O}_{12}\operatorname{Ru}_4\cdot 0.5\operatorname{CH}_2\operatorname{Cl}_2, M = 925.1.$ Triclinic, space group P1, a = 13.2174(6), b = 14.5234(9), c = 16.5710(7) Å, $a = 73.859(1), \beta = 82.855(1), \gamma = 68.495(1)^\circ, V = 2842$ Å³, D_c (Z = 4) = 2.162 g cm⁻³. Crystal 0.26 × 0.14 × 0.06 mm, μ (Mo-Ka) = 2.2 mm⁻¹, $T_{\min,\max} = 0.60, 0.78. 2\theta_{\max} = 75^\circ$; $N_{\text{total}} = 59267, N = 29321$ ($R_{\text{int}} = 0.031$), $N_o = 20247, R = 0.031, R_w = 0.026.$ (x, y, z, U_{iso})_H were refined.

(7) $\operatorname{Ru}_6(\mu_4\text{-}\operatorname{CCH}_2)_2(\mu\text{-}\operatorname{dppm})_2(\operatorname{CO})_{12}\cdot 0.5\operatorname{CHCl}_3\cdot 2\operatorname{MeCN} \equiv C_{66}H_{48}O_{12}P_4\operatorname{Ru}_6\cdot 0.5\operatorname{CHCl}_3\cdot 2\operatorname{CH}_3\operatorname{CN}, M = 1905.2.$ Triclinic, space group $P\overline{1}, a = 13.079(1), b = 15.254(1), c = 19.114(2)$ Å, $a = 90.636(2), \beta = 99.454(2), \gamma = 105.195(2)^\circ, V = 3624$ Å³, $D_c(Z = 2) = 1.746$ g cm⁻³. Crystal $0.49 \times 0.36 \times 0.30$ mm, $\mu(\operatorname{Mo-Ka}) = 1.42 \text{ mm}^{-1}, T_{\min,\max} = 0.57, 0.80.2\theta_{\max} = 65^\circ$. $N_{\text{total}} = 72818, N = 25430$ ($R_{\text{int}} = 0.035$), $N_o = 19208, R = 0.061, R_w = 0.113$.

Variata

(4) $(x, y, z, U_{iso})_{\rm H}$ were refined. (6) The dichloromethane was modelled with the chlorine atoms disordered over two sets of sites, occupancies set at 0.7 and complement after trial refinement; $(x, y, z, U_{iso})_{\rm H}$ were refined. (7) Molecular parameters determined from this structure are less precise than those of the original determination, which was carried out on a tris-benzene solvate, as a result of the presence of a disordered component, which can be described in terms of the heavy atom component being modelled as distributed over a pair of sites, population ca 1:9, displaced on average by ca 0.5 Å. It is not possible to define the light atom array associated with the minor component, so that we cannot say whether the two molecules are the same, one possibly displaced in concert with some component of solvent disorder, or whether there is a second component. However, we do not find any evidence for the latter in the NMR spectra. The chloroform solvent is modelled in terms of residues of occupancy 0.318(3), 1/2 - 0.318(3).

CCDC reference numbers 181260–181264.

See http://www.rsc.org/suppdata/dt/b2/b202364b/ for crystallographic data in CIF or other electronic format.

Acknowledgements

We thank the Australian Research Council for support of this work and Johnson Matthey plc, Reading, UK, for a generous loan of $RuCl_3 \cdot nH_2O$. We thank Professor B. K. Nicholson (University of Waikato, Hamilton, New Zealand) for some mass spectra.

References

- 1 E. H. Braye, L. F. Dahl, W. Hübel and D. L. Wampler, J. Am. Chem. Soc., 1962, 84, 4633.
- 2 (a) C. R. Eady, B. F. G. Johnson and J. Lewis, J. Organomet. Chem., 1972, **39**, 329; (b) B. F. G. Johnson, J. Lewis, K. Wong and M. McPartlin, J. Organomet. Chem., 1980, **185**, C17.
- 3 (a) J. S. Bradley, G. B. Ansell and E. W. Hill, J. Am. Chem. Soc., 1979, **101**, 7417; (b) J. S. Bradley, G. B. Ansell and E. W. Hill, J. Organomet. Chem., 1980, **184**, C33.
- 4 (a) C. E. Anson, P. J. Bailey, G. Conole, B. F. G. Johnson, J. Lewis, M. McPartlin and H. R. Powell, *J. Chem. Soc., Chem. Commun.*, 1989, 442; (b) P. J. Bailey, M. J. Duer, B. F. G. Johnson, J. Lewis, G. Conole, M. McPartlin, H. R. Powell and C. E. Anson, *J. Organomet. Chem.*, 1990, **383**, 441.
- 5 (a) M. Tachikawa and E. L. Muetterties, Prog. Inorg. Chem., 1981,
 28, 203; (b) J. S. Bradley, Adv. Organomet. Chem., 1983, 22, 1;
 (c) P. J. Dyson, Adv. Organomet. Chem., 1998, 43, 43.
- 6 V. G. Albano, P. Chini, S. Martinengo, D. J. A. McCaffrey, D. Strumolo and B. T. Heaton, J. Am. Chem. Soc., 1974, 96, 8106.
- 7 V. G. Albano, P. Chini, S. Martinengo, M. Sansoni and D. Strumolo, J. Chem. Soc., Chem. Commun., 1974, 299.
- 8 (a) C. J. Adams, M. I. Bruce, B. W. Skelton and A. H. White, J. Chem. Soc., Chem. Commun., 1985, 744; (b) C. J. Adams, M. I. Bruce, B. W. Skelton and A. H. White, J. Organomet. Chem., 1989, 369, 393.
- 9 See, for example, Fe: (a) J. W. Kolis, M. A. Drezdzon and D. F. Shriver, *Inorg. Synth.*, 1989, **26**, 246; (b) E. W. Hill and J. S. Bradley, *Inorg. Synth.*, 1990, **27**, 182; Ru: (c) J. N. Nicholls and M. D. Vargas, *Inorg. Synth.*, 1989, **26**, 280; (d) M. P. Cifuentes and M. G. Humphrey, *Inorg. Synth.*, 1998, **32**, 287; Rh: (e) S. Martinengo, D. Strumolo and P. Chini, *Inorg. Synth.*, 1980, **20**, 212.
- 10 M. C. Manning and W. C. Trogler, *Inorg. Chim. Acta*, 1981, **50**, 247. 11 J. Heidrich, M. Steimann, M. Appel, W. Beck, J. R. Phillips and
- W. C. Trogler, Organometallics, 1990, 9, 1296.
- 12 A. M. Nartowski, I. P. Parkin, M. Mackenzie and A. J. Craven, J. Mater. Chem., 2001, 11, 3116.
- 13 (a) C.-M. T. Hayward, J. R. Shapley, M. R. Churchill, C. Bueno and A. L. Rheingold, J. Am. Chem. Soc., 1982, 104, 7347; (b) M. R. Churchill, C. Bueno and A. L. Rheingold, J. Organomet. Chem., 1990, 395, 85.
- 14 M. I. Bruce, B. W. Skelton, A. H. White and N. N. Zaitseva, J. Chem. Soc., Dalton Trans., 1999, 1445.
- 15 M/R/R' = (a) Fe/H/Et: E. Sappa, A. Tiripicchio and M. Tiripicchio Camellini, J. Chem. Soc., Dalton Trans., 1978, 419; (b) Ru/Me/Ph: S. Aime, G. Nicola, D. Osella, A. M. Manotti Lanfredi and A. Tiripicchio, Inorg. Chim. Acta, 1984, 85, 161; (c) Ru/Ph/Ph: J. E. Davies, B. F. G. Johnson, C. M. Martion, R. H. H. Pearson and P. J. Dyson, J. Organomet. Chem., 1998, 550, 431.
- 16 (a) S. R. Drake, B. F. G. Johnson and J. Lewis, J. Chem. Soc., Dalton Trans., 1989, 243; (b) S. R. Drake, B. F. G. Johnson, J. Lewis, G. Conole and M. McPartlin, J. Chem. Soc., Dalton Trans., 1990, 995; (c) R. L. Mallors, A. J. Blake, P. J. Dyson, B. F. G. Johnson and S. Parsons, Organometallics, 1997, 16, 1668.
- 17 R. D. Adams and W. Wu, Organometallics, 1993, 12, 1238.
- 18 M. I. Bruce, B. W. Skelton, A. H. White and N. N. Zaitseva, J. Chem. Soc., Dalton Trans., 1999, 13.
- 19 M. I. Bruce, N. N. Zaitseva, B. W. Skelton and A. H. White, J. Chem. Soc., Dalton Trans., 1999, 2777.
- 20 M. I. Bruce, B. W. Skelton, A. H. White and N. N. Zaitseva, *Inorg. Chem. Commun.*, 1999, 2, 453.
- 21 M. I. Bruce, S. M. Pyke, N. N. Zaitseva, B. W. Skelton and A. H. White, *Helv. Chim. Acta*, 2001, **84**, 3197.
- 22 (a) L. Ma, D. P. S. Rodgers, S. R. Wilson and J. R. Shapley, Inorg.

Chem., 1991, **30**, 3591; (b) J. W. Benson, T. Ishida, K. Lee, S. R. Wilson and J. R. Shapley, *Organometallics*, 1997, **16**, 4929; (c) K. Lee and J. R. Shapley, *Organometallics*, 1998, **17**, 4030; (d) K. Lee and J. R. Shapley, *Organometallics*, 1998, **17**, 4113; (e) K. Lee and J. R. Shapley, *Organometallics*, 1998, **17**, 4368.

- 23 (a) D. M. P. Mingos, J. Chem. Soc., Chem. Commun., 1983, 706; (b) D. M. P. Mingos and A. S. May, in *The Chemistry of Metal Cluster Complexes*, ed. D. F. Shriver, H. D. Kaesz, R. D. Adams, VCH, New York, 1990, p. 89.
- 24 J.-F. Halet, in *Topics in Physical Organometallic Chemistry*, ed. M. Gielen, Freund, London, 1992, vol. 4, pp. 221–288.
- 25 V. G. Albano, P. Chini, S. Martinengo, M. Sansoni and D. Strumolo, J. Chem. Soc., Dalton Trans., 1978, 459.

- 26 V. G. Albano, D. Braga, P. Chini, D. Strumolo and S. Martinengo, J. Chem. Soc., Dalton Trans., 1983, 249.
- 27 (a) D. Braga, F. Grepioni, P. J. Dyson, B. F. G. Johnson, P. Frediani, M. Bianchi and F. Piacenti, J. Chem. Soc., Dalton Trans., 1992, 2565; (b) A. Sirigu, M. Bianchi and E. Benedetti, J. Chem. Soc., Chem. Commun., 1969, 596.
- 28 W. Henderson, J. S. McIndoe, B. K. Nicholson and P. J. Dyson, J. Chem. Soc., Dalton Trans., 1998, 519.
- 29 M. I. Bruce, C. M. Jensen and N. L. Jones, *Inorg. Synth.*, 1989, 26, 259.
- 30 M. I. Bruce, B. K. Nicholson and M. L. Williams, *Inorg. Synth.*, 1989, 26, 271.
- 31 S. R. Hall, D. J. du Boulay and R. Olthof-Hazekamp (eds), *The Xtal* 3.7 System, University of Western Australia, Crawley, 2002.