

REDUCTIVE COUPLING OF AROMATIC ALDEHYDES
BY OCTACARBONYL DIFERRATE

Keiji ITO,* Saburo NAKANISHI,** and Yoshio OTSUJI**


*Department of Industrial Chemistry, Osaka Prefectural Technical
College, Saiwaicho, Neyagawa, Osaka 572

**Department of Applied Chemistry, College of Engineering,
University of Osaka Prefecture, Mozu-Umemachi, Sakai, Osaka 591

The reaction of aromatic aldehydes with $\text{Fe}(\text{CO})_5$ or $\text{Fe}_3(\text{CO})_{12}$ in pyridine gives the corresponding 1,2-diaryl-1,2-ethanediols as major products in good yields. A reactive species of this reaction is octacarbonyl diferrate $[\text{Fe}_2(\text{CO})_8]^{2-}$.

Recently, mononuclear carbonyl ferrates such as $[\text{Fe}(\text{CO})_4]^{2-}$ and $[\text{HFe}(\text{CO})_4]^-$ have been proved to be useful reagents in organic synthesis.^{1,2)} However, the synthetic utility of polynuclear carbonyl ferrates have little been developed.³⁾ Previously, we reported that a polynuclear ferrate, which was prepared from $\text{Fe}_3(\text{CO})_{12}$ and pyridine N-oxide, effectively promotes the reductive coupling of benzylic and allylic halides.⁴⁾ We now report that a dinuclear carbonyl ferrate, which is prepared by treating iron carbonyls with pyridine,⁵⁾ induces the reductive coupling of aromatic aldehydes to give the corresponding 1,2-diaryl-1,2-ethanediols.⁶⁾

A solution of $\text{Fe}_3(\text{CO})_{12}$ (2.0 mmol) and benzaldehyde [1a] (20 mmol) in pyridine (10 ml) was heated at 80°C under N_2 atmosphere for 1.5 h. Water (10 mmol) was added, and the mixture was heated again at 80°C for 18.5 h and then treated with 6 M hydrochloric acid (10 ml) at room temperature for 30 min.⁷⁾ The resulting mixture was extracted with benzene (10 ml). The extract was washed with water and dried over Na_2SO_4 . Upon adding petroleum ether to the extract, 1,2-diphenyl-1,2-ethanediol [2a] (5.1 mmol) precipitated as crystals which were separated by filtration. The filtrate was concentrated and chromatographed on silica gel. Elution with benzene gave recovered [1a] (7.1 mmol). Subsequent elution with benzene-ether afforded [2a] (0.72 mmol) and benzyl alcohol [3a] (0.78 mmol). The total yield of [2a] based on the reacted [1a] was 91%, and that based on $\text{Fe}_3(\text{CO})_{12}$ used was 294%.



a: R=H, b: R=o- CH_3 , c: R=m- CH_3 , d: R=p- CH_3 , e: R=p-Cl, f: R=p- CH_3O

Table 1. Reaction of Aromatic Aldehydes and Ketones
 with Iron Carbonyls in Pyridine^{a)}

Run	Aromatic Substrate	Iron Carbonyl	Time (h)	Products, Yield % ^{b)}	
				[2]	[3]
1	[1a]	Fe(CO) ₅	14	88 (88)	11 (5.5)
2	[1a]	Fe ₃ (CO) ₁₂	20 ^{c)}	294 (91)	39 (6.0)
3	[1b]	Fe(CO) ₅	20 ^{c)}	82 (81)	22 (11)
4	[1b]	Fe ₃ (CO) ₁₂	20 ^{c)}	274 (80)	89 (14)
5	[1c]	Fe(CO) ₅	20 ^{c)}	76 (76)	25 (13)
6	[1c]	Fe ₃ (CO) ₁₂	20 ^{c)}	233 (72)	90 (13)
7	[1d]	Fe(CO) ₅	20 ^{c)}	85 (79)	34 (16)
8	[1d]	Fe ₃ (CO) ₁₂	20 ^{c)}	252 (78)	122 (19)
9	[1e]	Fe(CO) ₅	20	77 (74)	26 (13)
10	[1e]	Fe ₃ (CO) ₁₂	20	262 (84)	63 (10)
11	[1f]	Fe(CO) ₅	20 ^{c)}	31 (95)	—
12	[1f]	Fe ₃ (CO) ₁₂	20 ^{c)}	134 (96)	—
13	C ₆ H ₅ COCH ₃	Fe ₃ (CO) ₁₂	20	no reaction	
14	C ₆ H ₅ COC ₆ H ₅	Fe ₃ (CO) ₁₂	20	no reaction	

a) A solution of Fe(CO)₅ (6.0 mmol) or Fe₃(CO)₁₂ (2.0 mmol) and a substrate (20 mmol) in pyridine (10 ml) was heated at 80°C for 20 h except Run 1. b) The yields based on the iron carbonyl used, and those in parentheses based on aldehydes reacted.

c) A small amount of water (10 mmol) was added after heating the solution at 80°C for the following periods of time; 1.5 h in Runs 2 and 8, and 5 h in the other Runs.

Similar treatment of substituted benzaldehydes [1b-f] with Fe₃(CO)₁₂ in pyridine gave the corresponding 1,2-diaryl-1,2-ethanediols [2b-f] as major products and the corresponding benzyl alcohols [3b-f] as minor products. The reaction of [1a-f] with Fe(CO)₅ in pyridine also yielded the same products, [2a-f] and [3a-f], but with lower efficiency. However, no reaction occurred when acetophenone and benzophenone were treated with Fe₃(CO)₁₂ in pyridine under similar conditions. The results are summarized in Table 1.

In order to elucidate a mechanistic feature of these reactions, the reactions of [1a] with Fe(CO)₅ and Fe₃(CO)₁₂ in pyridine were conducted under the similar conditions as above. Then, amounts of [2a] produced were determined at appropriate time intervals by VPC analysis of the reaction mixture. The results are shown in Fig. 1. The induction period in the reaction using Fe₃(CO)₁₂ was much shorter than that in the reaction using Fe(CO)₅. However, amounts of [2a] produced per one Fe atom of each iron carbonyl were approximately equal for both Fe₃(CO)₁₂ and Fe(CO)₅; that is, one molar amount of [2a] was produced from one atom equiv of Fe in both cases.

Evolution of CO was observed even when pyridine solutions of Fe(CO)₅ and




Fig.1 Rates of formation of [2a].
A mixture of [1a] (20 mmol) and Fe(CO)₅ (6 mmol) or Fe₃(CO)₁₂ (2 mmol) in pyridine (10 ml) was heated at 80°C for 5 h in the case of Fe(CO)₅ and for 1.5 h in the case of Fe₃(CO)₁₂, and water (10 mmol) was added. Then, the mixture was heated at 80°C. ○ : Fe(CO)₅, ● : Fe₃(CO)₁₂.





Fig.2 Amounts of CO on the reaction time.
Fe(CO)₅=0.48 mmol, Fe₃(CO)₁₂=0.16 mmol, Pyridine=5 ml, [1a]=10 mmol.
○ : Fe(CO)₅, ● : Fe₃(CO)₁₂.

Fe₃(CO)₁₂ were heated at 80°C in the absence of [1a]. After evolution of CO ceased, further evolution of CO commenced as soon as [1a] was added to the above mixtures. Amounts of CO evolved during these reactions are shown in Fig.2. In the case of Fe(CO)₅, approximately 7/3 molar equiv of CO evolved from one molar amount of Fe(CO)₅. While, in the reaction of Fe₃(CO)₁₂, approximately 4 molar equiv of CO evolved from one molar amount of Fe₃(CO)₁₂. On the other hand, amounts of CO evolved after the addition of [1a] and its evolution rates were approximately equal for Fe(CO)₅ and Fe₃(CO)₁₂.

To isolate a key intermediate of the reaction, Fe₃(CO)₁₂ (2.0 mmol) was heated in pyridine (10 ml) at 80°C for 1 h, and then the mixture was treated with a CH₂Cl₂ solution of bis(triphenylphosphine)iminium chloride [(Ph₃P)₂N]Cl (2.44 mmol). Work-up of this mixture gave the dinuclear ferrate complex [(Ph₃P)₂N]₂[Fe₂(CO)₈].⁸⁾

From the above results, we presumed the reaction pathways of Scheme 1 for the reactions reported in this paper. The reactions of Fe(CO)₅ and Fe₃(CO)₁₂ with pyridine give the octacarbonyl diferrate [4]. The rate of formation of [4]



Scheme 1

is faster for $\text{Fe}_3(\text{CO})_{12}$ than for $\text{Fe}(\text{CO})_5$ (see Fig.2). Complex [4] induces the reductive coupling of aromatic aldehydes with concomitant oxidation of [4] to $\text{Fe}(\text{II})$ ions and evolution of CO. In the presence of water, [4] induces the reduction of aldehydes, giving benzyl alcohols as by-products. An active species of this reaction may be a hydrido complex of [4].⁹⁾

On the contrary, disodium tetracarbonyl ferrate $\text{Na}_2\text{Fe}(\text{CO})_4$, a mononuclear ferrate, was shown to react with aromatic aldehydes to give the corresponding benzyl benzoates.¹⁰⁾ The reactivity of the dinuclear ferrate is apparently different from that of the mononuclear ferrate.

References and Notes

- 1) J. P. Collman, *Acc. Chem. Res.*, **9**, 209 (1976).
- 2) Y. Watanabe and Y. Takegami, *Kagaku.*, **31**, 678 (1976); Y. Watanabe and Y. Takegami, *Yuki Gosei Kagaku Kyokai Shi.*, **35**, 585 (1977), and references cited therein.
- 3) J. P. Collman, R. G. Finke, P. L. Matlock, R. Wahren, R. G. Komoto, and J. I. Brauman, *J. Am. Chem. Soc.*, **100**, 1119 (1978), and references cited therein.
- 4) S. Nakanishi, T. Oda, T. Ueda, and Y. Otsuji, *Chem. Lett.*, **1978**, 1309.
- 5) W. Hieber and R. Werner, *Chem. Ber.*, **90**, 286 (1957); W. Hieber and N. Kahren, *ibid.*, **91**, 2234 (1958).
- 6) Recently, it has been demonstrated that the reductive coupling of aldehydes can also be effected by low-valent titanium reagents: E. J. Corey, R. L. Danheiser, and S. Chandrasekaran, *J. Org. Chem.*, **41**, 260 (1976), and references cited therein.
- 7) When [1a] was treated with $\text{Fe}_3(\text{CO})_{12}$ in pyridine at 80°C for a longer period of time (20 h) in the absence of water, benzoin and [2a] were obtained in a 3:1 ratio in the total yield of 270% based on $\text{Fe}_3(\text{CO})_{12}$ used, together with a small amount of [3a].
- 8) The following analytical and spectral data were obtained for $[(\text{Ph}_3\text{P})_2\text{N}]_2[\text{Fe}_2(\text{CO})_8]$. Found: C, 67.78; H, 4.27; Fe, 7.99; N, 1.83%. Calcd. for $\text{C}_{80}\text{H}_{60}\text{Fe}_2\text{N}_2\text{O}_8\text{P}_4$: C, 68.00; H, 4.28; Fe, 7.91; N, 1.98%. IR (KBr), 1930 cm^{-1} , 1910 cm^{-1} , 1870 cm^{-1} .
- 9) Aromatic aldehydes are reduced by hydrido ferrates to give the corresponding alcohols: H. W. Sternberg, R. Markby, and I. Wender, *J. Am. Chem. Soc.*, **78**, 5704 (1956); H. W. Sternberg, R. Markby, and I. Wender, *ibid.*, **79**, 6116 (1957).
- 10) M. Yamashita, Y. Watanabe, T. Mitsudo, and Y. Takegami, *Bull. Chem. Soc. Jpn.*, **49**, 3597 (1976).

(Received June 13, 1980)