Journal Pre-proofs

Unexpected migration of a benzoyl group in the intramolecular Wittig reaction of *o*-acyloxybenzylidenephosphoranes with benzoyl chlorides: one-pot synthesis of isomeric 3-benzoyl-2-phenylbenzofurans

Michela Begala, Michele Mancinelli, Giovanna Lucia Delogu

PII:	\$0040-4039(20)30047-2
DOI:	https://doi.org/10.1016/j.tetlet.2020.151634
Reference:	TETL 151634
To appear in:	Tetrahedron Letters
Received Date:	18 November 2019
Revised Date:	10 January 2020
Accepted Date:	14 January 2020



Please cite this article as: Begala, M., Mancinelli, M., Lucia Delogu, G., Unexpected migration of a benzoyl group in the intramolecular Wittig reaction of *o*-acyloxybenzylidenephosphoranes with benzoyl chlorides: one-pot synthesis of isomeric 3-benzoyl-2-phenylbenzofurans, *Tetrahedron Letters* (2020), doi: https://doi.org/10.1016/j.tetlet.2020.151634

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier Ltd.

Graphical Abstract

To create your abstract, type over the instructions in the template box below. Fonts or abstract dimensions should not be changed or altered.





Tetrahedron Letters journal homepage: www.elsevier.com

Unexpected migration of a benzoyl group in the intramolecular Wittig reaction of *o*-acyloxybenzylidenephosphoranes with benzoyl chlorides: one-pot synthesis of isomeric 3-benzoyl-2-phenylbenzofurans

Michela Begala^{a,1*,†}, Michele Mancinelli^b and Giovanna Lucia Delogu^{a,†}

^aDepartment of Life and Environmental Sciences, University of Cagliari, Via Ospedale 72, 09124 Cagliari, Italy

^bDepartment of Industrial Chemistry "Toso Montanari", University of Bologna, Viale del Risorgimento, 4, 40136 Bologna, Italy

ARTICLE INFO * Corresponding author.	ABSTRACT
e-mail address: michelabegala@unica.it (M. Bega	ıla).
ArTibushismsthors contributed equally to the work.	The reaction of <i>o</i> -acyloxybenzylidenetriphenylphosphoranes with substituted benzoyl chlorides in an aprotic solvent led, together with the expected 2-phenylbenzofuran, to isomeric 3-benzoyl- 2-phenylbenzofuran derivatives. This result formally corresponds to intramolecular migration of the benzoyl group from the <i>ortho</i> oxygen atom to the ylide carbanion <i>via</i> cyclization and ring
Received in revised form	
Accepted	
	opening of the starting o-acyloxybenzylidenetriphenylphosphoranes.
Keywords:	
Wittig reaction	
Phosphoranes	
2-phenylbenzofurans	

Introduction

3-aroylbenzofurans

3-Aroyl[b]benzofurans are present in the structural cores of a large number of bioactive molecules in current pharmaceutical use and development (Fig. 1) [1-4]. As a result, numerous approaches towards the synthesis of 3-acylbenzofurans have been disclosed in the literature [5,6]; however, most of them are only suitable for the preparation of 2-aroyl-3-benzoylbenzofuran derivatives bearing electron-donating groups [7].



Figure 1. Representative 3-aroyl[b]benzofurans of pharmacological interest.

In our previous work, we found that the reaction of 2hydroxybenzyltriphenylphosphonium bromide with benzoyl chlorides and triethylamine in an aprotic solvent (*i.e.* toluene), leads to 3-benzoyl-2-phenylbenzo[*b*]furans *via* ylide acylation in addition to the expected 2-phenylbenzofurans (Scheme 1) [8,9].



Scheme 1. Synthetic route towards 2-phenylbenzofurans and 3-benzoyl-2-phenylbenzofurans.

Hercouet and Le Corre [10] reported that the intramolecular Wittig reaction of *o*-acyloxybenzylidenetriphenylphosphoranes leads to the corresponding benzofurans in aprotic medium (toluene) or to acylated products in protic medium (*t*-BuOH). Therefore, we became interested in verifying whether the *o*-[(benzoyloxy)benzyl]-triphenyl-phosphorane was the key intermediate leading to the 3-benzoyl derivatives. Our results showed that benzoyl chloride reacts with phosphorane I to give a mixture of 2-phenylbenzofuran and 3-benzoyl-2-phenylbenzofuran, thus confirming this hypothesis (Scheme 2).



Scheme 2. Formation of 2-phenylbenzofuran and 3-benzoyl-2-phenylbenzofuran from I.

3-Benzoyl-2-phenylbenzofuran derivatives with strongly electron-withdrawing groups (*e.g.* NO_2 and CN) on the 2-phenyl or 3-benzoyl ring, could provide convenient intermediates in the

Journal Pre-proofs

preparation of more complete compounds [11]. However, no methods for the synthesis of such deactivated benzofurans have been described. For instance, during the Friedel-Crafts acylation of 2-phenylbenzofuran with nitrobenzoyl chloride, other positions of the benzofuran ring were also acylated, leading to a complex mixture of regioisomers where the expected derivative was formed as a minor product [12]. Thus, the development of new synthetic routes, especially those that allow access to deactivated analogues, is of considerable interest.

In this context, we envisaged that the reaction of *o*-acyloxybenzylidenetriphenylphosphoranes with substituted benzoyl chlorides would provide access to the corresponding 3-benzoyl-2-phenylbenzofurans bearing different substituents on the 2-phenyl and/or the 3-benzoyl ring.

In the present work, we describe our efforts toward the synthesis of diversely 2,3-disubstituted benzofurans with particular attention given to the preparation of 3-benzoyl-2-phenylbenzofuran derivatives bearing electron-withdrawing groups. Moreover, the unexpected formation of the 3-acyl isomers, prompted us to further investigate the Wittig reaction.

Results and Discussion

o-[(Benzoyloxy)benzyl]-triphenyl-phosphoranes **I**, prepared from *ortho*-cresol [8,13,14] (Scheme 3), were reacted with benzoyl chlorides substituted with electron withdrawing or donating groups.



Scheme 3. Synthetic route towards o-[(benzoyloxy)benzyl]-triphenyl-phosphoranes I (R = H, NO₂ OCH₃).

When I (R = H) was reacted with 4-nitrobenzoyl chloride in the presence of triethylamine in toluene, the reaction mixture showed the formation of two 3-acyl isomers (Scheme 4), *i.e.* the expected 3-(4-nitrobenzoyl)-2-phenylbenzofuran **2b** and 3benzoyl-2-(4-nitrophenyl)benzofuran **3b**, as identified by NMR spectroscopy and GC/MS analysis.



Scheme 4. Synthesis of isomeric compounds 2b and 3b.

The same behaviour was also observed when a variety of 4substituted benzoyl chlorides were used (*vide infra*). These results clearly suggest that benzoyl group migration occurred to some extent. Interestingly, Hercouet and Le Corre reported such a transposition only under protic condition (*t*-BuOH) and using a strong base (*t*-BuOK) [10]. An analogous 1,5-transposition of an ester acyl functional group was also observed by Gorgues and coworkers for the σ -bonded organochromium(III) complex resulting from the reduction of *ortho*-bromomethylphenyl benzoate with CrCl₂ [15].

A plausible mechanism is depicted in Scheme 5. Carbanion I reacts with the corresponding acyl chloride to give the isomer **2b** (Route B), as previously proposed by us [8]. Alternatively, the O-acylated phosphorane I can also cyclize directly into intermediate II (Route A) to give either the 2-phenylbenzofuran **1a** or undergo ring-opening to the C-acylated phenolate III. Intermediate III is O-acylated to give the phenol ester IV, that leads to the isomeric 3-benzoyl-2-phenylbenzofuran **3b** *via* a ylide intermediate.



Scheme 5. Proposed mechanism of benzoyl group migration from ylide I and the formation of isomeric compounds 2b and 3b.



Table 1. Synthetic route towards 2-phenylbenzofurans 1a-c, and 3-benzoyl-2-phenylbenzofurans 2-9a-e, 10 and 11.ª

^aI (1 equiv.), benzoyl chlorides (2.5 equiv.), Et₃N (2.5 equiv.), toluene 110 °C, 2 h.

ournal Pre-proo

Interestingly, the isomer arising from benzoyl migration 3b (45%) was the main product; therefore Route A was more favored than Route B. These data clearly indicate that the O-acylated phosphorane I spontaneously cyclizes into II, presumably because intramolecular attack of the ylide carbanion to the *ortho* carbonyl group is favoured by their proximity (Scheme 5 and Table 1, entry 1).

This behaviour was even more pronounced when 4nitrobenzoyl chloride was replaced by 4-methoxybenzoyl chloride (Table 1, entry 2). In the latter case, the reaction of **I** with the less electrophilic 4-methoxybenzoyl chloride (Route B) was suppressed (4c, 3%) in favour of Route A (1a, 20%; 5c, 15%). These results clearly indicate that deactivated benzoyl chlorides allow the formation of both isomeric 3-acyl derivatives, most likely because **I** and **III** rapidly attack the highly electrophilic carbonyl of the 4-nitrobenzoyl chloride. Accordingly, analogous trends were also observed with 4-bromo and 4-cyanobenzoyl chlorides (Table 1, entries 3 and 4).

When substituted o-[(benzoyloxy)benzyl]-triphenylphosphoranes were used (R = NO₂, OCH₃, entries 5 and 6), we observed that the electron withdrawing group further suppressed Route B in favour of cyclization to **II** (Route A), as reflected by the higher yield of 2-(4-nitrophenyl)benzofuran **1b** (42%) and the absence of 3-benzoyl-2-(4-nitrophenyl)benzofuran **3b** (Table 1, entry 5). This behaviour could be rationalized by the fact that the 4-nitro group increases the electrophilicity of the carbonyl group of **I** (R = NO₂), making it easier for the ylide carbanion to attack this electrophilic centre. Accordingly, o-[(benzoyloxy)benzyl]triphenyl-phosphorane bearing an electron donating group (R = OCH₃) still resulted in the formation of isomer **5c**; albeit in low yield (4%) (Table 1, entry 6).

The reaction of substituted o-[(benzoyloxy)benzyl]-triphenylphosphoranes I (R = OCH₃) with 4-nitrobenzoyl chloride (Table 1, entry 7) again leads predominantly to 2-(4methoxyphenyl)benzofuran 1c (12%) and isomer 11 (15%) *via* Route A.

Conclusion

Our previous results demonstrated that under aprotic conditions 3-acylbenzofurans can be obtained from the reaction of *o*-[(benzoyloxy)benzyl]-triphenyl-phosphoranes with benzoyl chlorides. We have also demonstrated that under these conditions phosphoranes preferentially undergo intramolecular migration of the benzoyl group *via* cyclization and ring opening. This behaviour clarifies the unexpected formation of two sets of 3-acylbenzofuran isomers when phosphoranes react with benzoyl chlorides.

In particular, we found that the isomer arising from benzoyl migration is further favoured by the use of deactivated benzoyl chlorides or by phosphoranes I with a 4-methoxy group. This finding is of considerable interest as it allows preparation of a wide variety of 3-acylbenzofuran derivatives which are difficult to obtain using alternative methods.

Acknowledgments

The present work was partially supported by FIR (Fondo Integrativo per la Ricerca – annualità 2018) - University of Cagliari.

References and notes

- Singh SN, Fletcher RD, Fisher SG, Singh BN, Lewis HD, Deedwania PC, Massie BM, Colling C, Layyeri D. N. Eng. J. Med. 1995;333:77-82; (b) Kálai T, Várbió G, Bognár Z, Pálfi A, Hantó K, Bognár B, Ősz E, Smegi B, Hideg K. Bioorg. Med. Chem. 2005;13:2629-2636; (c) Hartong R, Wiersinga WM, Plomp TA. Horm. Metab. Res. 1990;22:85-89; (d) Basaria S, Cooper DS. Am. J. Med. 2005;118:706-714.
- [2] Felder CC, Joyce KE, Briley EM, Glass M, Mackie KP, Fahey KJ, Cullian GJ, Hunden DC, Johnson DW, Chaney MO, Koppel GA, Brownstein M J. Pharmacol Exp Ther. 1998;284(1):291-297.
- [3] Heel RC, Brogden RN, Speight TM, Avery GS. Drugs 1977;14:349-366.
- [4] (a) Brenner LM, Brush CK. US Pat. 4.001.426 (1997); (b) Howlett DR, Perry AE, Godfrey F, Swatton JE, Jennings KH, Spitzfaden C, Wadsworth H, Wood SE, Markwell RE. *Biochem. J.* 1999;340:283-289.
- [5] Gong W, Liu D, Li F, Gao J, Li H, Lang J. Tetrahedron 2015;71:1269-1275.
- [6] (a) Nakamura M, Ilies L, Otsubo S, Nakamura E. Org Lett. 2006;8:2803-2805; (b) Zhao J, Fang H, Xie C, et al. Org Chem. 2013;2:1044-1047; (c) Farrar MW, Levine R. J Am Chem Soc. 1950;72:4433-4436.
- [7] (a) Oda K, Machida M. Chem. Pharm. Bull. 1993;41:1299-1301; (b) Yuan H, Bi K, Li B, Yue R, Ye J, Shen Y, Shan L, Jin H, Sun Q, Zhang W. Org. Lett. 2013;15:4742-4745; (c) Arcadi A, Cacchi S, Del Rosario M, Fabrizi G, Marinelli F. J. Org. Chem. 1996;61:9280-9288. (d) Jason H, Chaplin JH, Flynn BL. Chem. Commun. 2001;1594-1595; (e) Mothe SR, Susanti D, Hong Chan PW. Tetrahedron Lett. 2010;51:2136-2140.
- [8] Begala M, Caboni P, Matos MJ, Delogu GL. Tetrahedron Lett. 2018;59:1711-1714.
- [9] Begala M, Delogu GL. J. Mass Spectrom. 2019;54(9):750-760.
- [10] (a) Hercouet A, Le Corre M. *Tetrahedron Lett.* 1979;23:2145-2148; (b) Hercouet A, Le Corre M. *Tetrahedron* 1981;37(16):2867-2873.
- [11] (a) Kabalka GW, Varma RS. Comprehensive Organic Synthesis; Trost BM, Fleming I, eds, 8. Oxford: Pergamon Press; 1991:363–379; (b) Rappoport Z, ed. Chemistry of the Cyano Group. London: John Wiley & Sons; 1970.
- [12] (a) Twyman LJ, Allsop D. *Tetrahedron Lett.* **1999**;40:9383-9384; (b) Thévenin M, Thoret S, Grellier P, Dubois J. *Bioorg. Med. Chem.* **2013**;21(17):4885-4892.
- [13] Ghosh S, Das J. Tetrahedron Lett. 2011;52:1112-1116.
- [14] Routasalo T, Helaja J, Kavakka J, Koskinen AMP. Eur. J. Org. Chem. 2008;18:3190-3199.
- [15] Ledoussal B, Gorgues A, Le Coq A. Tetrahedron 1987;43:5841-5852.

Supporting Information

Details of the experimental procedures and spectroscopic data for new compounds. This material is available free of charge in the online version [http://].

Declaration of interests

 \boxtimes The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper. interests/personal relationships which may be considered as potential competing interests:

- benzoyl-2-phenylbenzofurans
- Access to 3-benzoyl-2-phenylbenzofurans with strongly deactivated functional groups

SEP

Graphical Abstract

To create your abstract, type over the instructions in the template box below. Fonts or abstract dimensions should not be changed or altered.

Highlights

1,5-transposition of benzoyl group in *o*benzoyloxybenzylidenetriphenylphosphoran es