

# Communication

# Total Synthesis of Salimabromide, a Tetracyclic Polyketide from a Marine Myxobacterium

Matthias Schmid, Adriana Grossmann, Klaus Wurst, and Thomas Magauer

J. Am. Chem. Soc., Just Accepted Manuscript • DOI: 10.1021/jacs.8b06228 • Publication Date (Web): 28 Jun 2018

Downloaded from http://pubs.acs.org on June 28, 2018

# Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036

Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

# Total Synthesis of Salimabromide, a Tetracyclic Polyketide from a **Marine Myxobacterium**

Matthias Schmid,<sup>†,‡</sup> Adriana S. Grossmann,<sup>†</sup> Klaus Wurst<sup>§</sup> and Thomas Magauer<sup>\*,‡</sup>

<sup>†</sup>Department of Chemistry, Ludwig-Maximilians-University Munich, Butenandtstrasse 5–13, 81377, Munich, Germany, <sup>‡</sup>Institute of Organic Chemistry and Center for Molecular Biosciences, University of Innsbruck, Innrain 80–82, 6020 Innsbruck. Austria. <sup>§</sup>Institute of General, Inorganic & Theoretical Chemistry, University of Innsbruck, Innrain 80–82, 6020 Innsbruck, Austria.

Supporting Information Placeholder

**ABSTRACT:** Salimabromide is an antibiotic polyketide that was previously isolated from the obligate marine myxobacterium Enhygromyxa salina, and its densely functionalized and conformationally rigid tetracyclic framework is unprecedented in nature. Herein we report the first chemical synthesis of the target structure by employing a series of well-orchestrated, robust transformations, highlighted by an acid-promoted, powerful Wagner-Meerwein rearrangement/Friedel-Crafts cyclization sequence to forge the two adjacent quaternary carbon centers embedded in the tetrahydronaphthalene. A high-yielding ketiminium mediated [2+2]-cycloaddition was also utilized for the simultaneous construction of the remaining three stereocenters.

Myxobacteria of terrestrial origin produce an abundance of structurally complex secondary metabolites with notable biological activities. Prominent examples include epothilone, corallopyronin and soraphen.<sup>1</sup> Marine myxobacteria, on the other hand, have eluded their cultivation and isolation on many occasions and constitute a largely unexplored treasure trove of bioactive molecules.<sup>2</sup> Haliangincin (2), which was reported in seminal work by Fudou in 2001,<sup>3</sup> represents the first natural product isolated from a strictly marine myxobacterium (Scheme 1a). Following this early report, only a handful of new marine myxobacterium molecules have appeared in the literature,<sup>2</sup> and in 2013, König disclosed the isolation of salimabromide (1) in minute quantities (0.5 mg from 64 L of)culture) from the obligate marine bacterium Enhygromxya salina.<sup>4</sup> Although a broad biological screening campaign was impossible at this stage, 1 was shown to possess inhibitory activity against Arthrobacter *cristallopoietes* (16  $\mu$ g mL<sup>-1</sup>).

From a structural point of view, salimabromide (1) contains an unprecedented tetracyclic ring-architecture that contains four consecutive stereocenters, one of which is quaternary. Additionally, the brominated tetrahydronaphthalene core is bridged at C12 and C15 to form a highly

substituted and synthetically challenging seven-membered carbocycle.<sup>5</sup> The conformational flexibility of this subunit is further reduced by fusion to a five-membered lactone.

Scheme 1. a) Structural diversity produced by marine myxobacteria and b) key retrosynthetic bond disconnections for salimabromide (1).



60

The structure for **1** was exclusively deduced from extensive NMR measurements. While this only confirmed the connectivity, the absolute stereochemistry was determined by comparison of calculated and measured CD spectra.

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22 23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57 58 59

60

Despite considerable progress of the Menche group,<sup>5</sup> no total synthesis of salimabromide (1) has been accomplished to date. In light of these considerations, we set out to develop a practical synthetic route to enable rapid access to 1 and fully synthetic derivatives thereof. Herein we describe a streamlined synthesis of 1 employing a series of robust chemical transformations. The successful realization of this route allowed us to produce 1.9 g of a highly advanced intermediate in a single batch from which salimabromide (1) was prepared in only three steps.

Our retrosynthetic bond disconnections were guided by our desire to rapidly generate molecular complexity and to install the individual stereocenters via a minimum number of synthetic operations (Scheme 1b). We planned to first install the pivotal C12 quaternary carbon center and utilize this handle for the subsequent one-step construction of the stereotriad along C11, C13 and C15. For this purpose and in analogy to the logic of two-phase terpene (bio)synthesis,<sup>6</sup> salimabromide (1) was first reduced to tetracycle 5. This intermediate was designed to eliminate steric hindrance and cross-reactivity of the bromine substituents en route to the tetracyclic carbon framework of 1. Compound 5 contains the retron for a powerful ketiminium mediated [2+2]-cycloaddition<sup>7</sup> to produce the simplified dihydronaphthalene 6. Further C-C bond disconnections involving truncation of the sidechain and a retro-Friedel-Crafts cyclization/Wagner-Meerwein<sup>8</sup> rearrangement sequence provided epoxide 7.

We commenced our synthesis with the Claisen-Schmidt condensation of commercially available 3-methoxybenzaldehyde 8 with pinacolone 9 (Scheme 2). While slow reactions (48 h) and moderate yields (48%) were observed for standard bases such as sodium or potassium hydroxide, the use of activated barium hydroxide  $[Ba(OH)_2 \cdot H_2O, C-200]^9$  led to rapid consumption of the reactants and clean conversion to the condensation product 10. On large scale, 10 was best obtained by simple filtration of the reaction mixture using a pad of Celite<sup>®</sup> and evaporation of excess ethanol and pinacolone 9. The following heterogeneous hydrogenation (1 atm  $H_2$ , Pd/C, EtOAc) was conducted on large-scale and reproducibly provided 11 in good yield (76-83% over two steps, 32 g). Exposure of 11 to standard Corey-Chaykovsky conditions (NaDMSO, Me<sub>3</sub>SI, 0 to 23 °C)<sup>10</sup> effected clean conversion to epoxide 7. With 7 in hand, we investigated the crucial rearrangement-cyclization sequence. We found that exposure of 7 to hexanes/sulfuric acid or dichloromethane/titanium(IV) chloride was low-vielding for 12 (~30%) and afforded substantial amounts of the undesired ortho-product 13 (~10%). Further screening revealed

1,1,1,3,3,3-hexafluoro-2-propanol (HFIP, 0.1 M), а strong hydrogen-bond donor and weak nucleophile,<sup>11</sup> as the solvent of choice and concentrated sulfuric acid as the optimal catalyst (10 mol%). Under these conditions, the para-product 12 was formed with very good regioselectivity (12:13 = 8:1). The remainder of the mass balance corresponds to a complex mixture of byproducts.<sup>12</sup> Similar results were obtained when the reaction was conducted in a hexameric resorcinarene capsule.<sup>13</sup> In agreement with related Lewis-acid catalyzed semi-pinacol and Wagner-Meerwein rearrangements,<sup>14</sup> a high level of stereoselectivity should be possible for the initial rearrangement step. Evidence was obtained when a solution of enantiomerically enriched 7 (82% ee) in dichloromethane was exposed to titanium(IV) chloride at -78 °C. Under these conditions, an enantiomeric excess of 70% was obtained for 12 (see Supporting Information for details). This result provides an opportunity to access 1 in an asymmetric fashion.





<sup>a</sup>Reagents and conditions: (a) [Ba(OH)<sub>2</sub>·H<sub>2</sub>O, C-200], pinacolone **9**, EtOH, 78 °C; (b) Pd/C, H<sub>2</sub>, EtOAc, 23 °C, 76–82% over two steps; (c) NaH, DMSO, 70 °C, then Me<sub>3</sub>SI, 0 to 23 °C; (d) H<sub>2</sub>SO<sub>4</sub>, HFIP, 23 °C, 41% for **12**, 5% for **13** over two steps.

Having successfully installed the crucial C12 quaternary stereocenter in only four steps, we turned our attention to the remaining functionalization of **12** (Scheme 3). To begin, the primary alcohol was protected as its *tert*butyldiphenylsilyl (TBDPS) ether, and styrene **14** was then formed via exposure to 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ, 2 equiv).<sup>15</sup> The use of substochiometric amounts of DDQ (30 mol%) in combination with manganese(IV) oxide (6 equiv) or alternative oxidation agents such as chloranil, Pd/C or electrochemical methods (4 V, chloranil, Pt/Pt, 0.1 M LiClO<sub>4</sub>, MeCN) were ineffective. 



<sup>a</sup>Reagents and conditions: (a) TBDPSCl, imidazole, DMAP, DMF, 23°C; (b) DDQ, dioxane, 93 °C, 91% over two steps; (c) LiPPh<sub>2</sub>, THF, 60 °C; (d) PhNTf<sub>2</sub>, NEt<sub>3</sub>, THF, 23 °C, 99% over two steps; (e) ZnEt<sub>2</sub>, Pd(dppf)Cl<sub>2</sub>, dioxane, 70 °C, 92%; (f) TBAF, THF, 23 °C, 99%; (g) DMP, K<sub>2</sub>CO<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>, 0 °C to 23 °C, 96%; (h) NaHMDS, **17**, THF, -78 °C to 23 °C, 85%; (i) pyrrolidine, 100 °C, 99%; (j) Tf<sub>2</sub>O, 2,4,6-collidine, ClCH<sub>2</sub>CH<sub>2</sub>Cl, 80 °C, 89%; (k) SeO<sub>2</sub>, SiO<sub>2</sub>, dioxane, 120 °C, 47% over five cycles; (l) *t*-BuCHO, Cu(OAc)<sub>2</sub>, O<sub>2</sub> (1 atm), ClCH<sub>2</sub>CH<sub>2</sub>Cl, 23 °C, then DMP, NaHCO<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>, 79%, **23**:**24** = 1:3.2; (m) AgTFA, Br<sub>2</sub>, CF<sub>3</sub>COOH, 50%.

The TBDPS group was crucial for the stability of the silyl ether under the reaction conditions, and the presence of the electron-donating methoxy substituent was beneficial for the oxidation.<sup>16</sup>

For the installation of the missing ethyl substituent, we initially investigated the direct coupling of **14** using nickel (e.g. Ni(acac)<sub>2</sub>, EtMgBr, MgI<sub>2</sub>, PhMe, 100 °C; Ni(cod)<sub>2</sub>, EtMgI, MgI<sub>2</sub>, PhMe, 100 °C).<sup>17</sup> Since our substrate proved to be remarkably unreactive under these conditions, we decided to replace the methoxy substituent with a more reactive triflate. For the removal of the methyl ether, freshly prepared lithium diphenylphosphide (Ph<sub>2</sub>PH, *n*-BuLi),<sup>17</sup> was found to be optimal. The free phenol was then converted to the triflate upon exposure

to the Hendrickson–McMurray reagent (PhNTf<sub>2</sub>)<sup>19</sup> to afford **15** in quantitative yield. Standard Negishi cross-coupling (Pd(dppf)Cl<sub>2</sub>, ZnEt<sub>2</sub>, dioxane, 70 °C)<sup>20</sup> enabled clean installation of the missing ethyl substituent (92%), and cleavage of the silyl ether (TBAF, THF, 23 °C) gave alcohol **16** (99%). The remaining carbon-chain was introduced in a three-step sequence beginning with the oxidation of **16** using Dess–Martin periodinane<sup>21</sup> (96%), followed by a high-yielding, *Z*-selective Wittig olefination and amide formation (pyrrolidine, 100 °C) to provide **6** (85%).

With robust access to the dihydronaphthalene **6**, we proceeded to investigate the key-step of our synthesis: a ketiminium ion mediated [2+2]-cycloaddition<sup>7</sup> to construct the seven-membered carbocycle and complete the

tetracyclic carbon framework. Under optimized conditions, a solution of amide **6** and *sym*-collidine (1.2 equiv) in dichloroethane was slowly added to a refluxing solution of freshly distilled trifluoromethanesulfonic anhydride (1.2 equiv) in dichloroethane (0.1 M). The cycloaddition produced the tetracycle **5** with almost perfect regioand diastereoselectivity in excellent yield (89%, 1.9 g). The exact mechanism, concerted (synchronous/asynchronous) or stepwise, has been a matter of debate for many decades.<sup>7</sup> Depending on the substitution pattern on both the alkene and the ketiminium salt, either of the two pathways might be operational.<sup>22</sup> The greater resonance stabilization of the benzylic cation **21** versus **19** was envisioned to govern the regioselectivity favoring formation of **5**.

1

2

3

4

5

6

7

8

9

10

11

12

13

14

57

58 59

60

15 Having installed the crucial stereocenters, we were 16 poised to tackle the remaining challenges: regioselective 17 oxidation of the carbon-framework and bromination of 18 the arene subunit. When 5 (500 mg) was treated with se-19 lenium dioxide (dioxane, 120 °C, 6 h) in the presence of 20 silicon dioxide (>230 mesh) to prevent agglomeration, 21 the diastereomerically pure allylic alcohol 22 was formed 22 together with unreacted 5. Extended reaction times were 23 detrimental as overoxidation and decomposition started 24 to prevail. Subjection of recovered 5 (76%) to the reaction 25 conditions enabled us to prepare 250 mg of 22 after five 26 cycles (47%, ~15% for the first cycle).<sup>23</sup> Subsequent Bae-27 ver-Villiger oxidation using standard conditions (m-28 CPBA, NaHCO<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>) gave two regioisomeric lac-29 tones, which were directly oxidized (DMP, NaHCO<sub>3</sub>, 30 CH<sub>2</sub>Cl<sub>2</sub>) to afford 23 and 24 in a ratio of 1.4:1 in 84% 31 combined yield. Separation of 23 and 24 was readily ac-32 complished by flash column chromatography. To improve 33 this undesired outcome, further optimization of the oxi-34 dant was performed. Interestingly, exposure of 22 to t-Bu-35 CHO (5 equiv) in the presence of molecular oxygen (1 36 atm) and copper(II) acetate (1 equiv) gave 24 as the major 37 product (79%, 23:24 = 1:3.2).<sup>24</sup> The directing effect of the 38 free hydroxy group was crucial as the corresponding me-39 thyl ether led to lower regioselectivity (1:1.1) only 40 slightly favoring the desired regioisomer (compare 24). It 41 is also noteworthy that replacement of t-BuCHO with m-42 CPBA in the Baeyer-Villiger step under otherwise iden-43 tical conditions was even less efficient and only poor re-44 gioselectivity (23:24 = 1.2:1) was obtained. For the intro-45 duction of the missing bromine substituents, 24 was ex-46 posed to a panel of brominating agents (e.g. Br<sub>2</sub>, CHCl<sub>3</sub>; 47 NBS, HOAc; BnMe<sub>3</sub>NBr<sub>3</sub>, ZnBr<sub>2</sub>, HOAc). Under these 48 conditions, formation of 1 was only observed in trace 49 amounts if at all. Finally, we found that treating a solution 50 of 24 in trifluoroacetic acid (0.1 M) with silver(I) tri-51 fluoroacetate (3 equiv) and elemental bromine (3 equiv) 52 enabled the desired bromination (50%) and thus com-53 pleted the synthesis of salimabromide (1, 50 mg).<sup>25</sup> The 54 analytical data for 1 (<sup>1</sup>H-NMR, <sup>13</sup>C-NMR, HRMS) fully 55 56

matched those reported for the natural compound. Additionally, the structure of **1** was unambiguously validated by single-crystal X-ray diffraction analysis.

In summary, we have completed the first total synthesis of salimabromide, a unique tetracyclic polyketide. The highlights of the developed route are: 1) A powerful Wagner–Meerwein rearrangement/Friedel–Crafts cyclization sequence to forge the tetrahydronaphthalene skeleton and 2) a high-yielding ketiminium mediated (2+2)-cycloaddition to set the remaining three stereocenters. The overall sequence benefits from a series of practical transformations that can be also conducted on large scale. The robustness of the developed synthesis is evident from the fact that more than 1.9 g of a highly advanced intermediate were prepared in a single batch.

### ASSOCIATED CONTENT

#### **Supporting Information**

The Supporting Information is available free of charge on the ACS Publications website at http://pubs.acs.org.

Experimental details and spectroscopic data (PDF) X-ray crystallographic data for 1, 5, 23 and 24 (CIF)

## AUTHOR INFORMATION

### **Corresponding Author**

thomas.magauer@uibk.ac.at

#### Notes

The authors declare no competing financial interests.

## ACKNOWLEDGMENT

T.M. acknowledges the European Research Council under the European Union's Horizon 2020 research and innovation program (grant agreement No 714049) and the Center for Molecular Biosciences (CMBI). M.S. gratefully acknowledges financial support from the German National Academic Foundation. We thank Dr. Kevin Mellem (Revolution Medicines) for helpful discussions, Prof. Albrecht Berkessel (University of Cologne) for a screen of epoxidation catalysts and Prof. Konrad Tiefenbacher (University of Basel/ETH) for the screen of supramolecular catalysts.

#### REFERENCES

- (1) Hermann, J.; Fayad, A. A.; Müller, R. Nat. Prod. Rep. 2017, 34, 135.
- (2) (a) Dávila-Céspedes, A.; Hufendiek, P.; Crüsemann, M.; Schäberle, T. F.; König, G. M. *Beilstein J. Org. Chem.* 2016, *12*, 969.
  (b) Felder, S.; Kehraus, S.; Neu, E.; Bierbaum, G.; Schäberle, T. F.; König, G. M. *ChemBioChem* 2013, *14*, 1363.
- (3) Fudou, R.; Iizuka, T.; Sato, S.; Ando, T.; Shimba, N.; Yamanaka, S.; J. Antibiotics, 2001, 54, 153.
- (4) Felder, S.; Dreisigacker, S.; Kehraus, S.; Neu, E.; Bierbaum, G.; Wrigth, P. R.; Menche, D.; Schäberle, T. F.; König, G. M. Chem. Eur. J. 2013, 19, 9319.
- (5) Schmalzbauer, B.; Menche, D. Org. Lett. 2015, 17, 2956.
- (6) Chen, K.; Baran, P. S. *Nature* **2009**, *459*, 824.
- (7) (a) Snider, B. B. Chem. Rev. 1988, 88, 793. (b) Madelaine, C.; Valerio, V.; Maulide, N. Chem. Asian J. 2011, 6, 2224.
- (8) (a) Knölker, H.-J.; Baum, E.; Graf, R.; Jones, P. G.; Spieß, O. Angew. Chem. Int. Ed. 1999, 38, 2583. (b) Khalaf, A. A.; Albar, H. A.; El-Fouty, K. O. *Ind. J. Chem.* 2010, 49B, 203.
- (9) Sinisterra, J. V.; Garcia-Raso, A. Synthesis 1984, 502.
- (10) Corey, E. J.; Chaykovsky, M. J. Am. Chem. Soc. 1965, 87, 1353.
- (11) Bégué, J.-P.; Bonnet-Delpon, D.; Crousse, B. Synlett 2004, 1, 18.

- (12) The mixture of byproducts could not be separated, however, careful analysis of the <sup>1</sup>H NMR revealed minor traces of the Meinwald rearrangement product (Meinwald, J.; Labana, S. S.; Chadha, M. S. J. *Am. Chem. Soc.* **1963**, *85*, 582). An indane, resulting from the direct Friedel-Crafts reaction, was not detected.
- (13) Zhang, Q.; Tiefenbacher, K. J. Am. Chem. Soc. 2013, 135, 16213.
- (14) (a) Cram, D. J.; J. Am. Chem. Soc. 1949, 71, 3863. (b) Suzuki, K.; Katayama, E.; Tsuchihashi, G. Tetrahedron Lett. 1984, 25, 1817.
  (c) Maruoka, K.; Hasegawa, M.; Yamamoto, H. J. Am. Chem. Soc. 1986, 108, 3827. (d) Matsubara, S.; Yamamoto, H.; Oshima, K. Angew. Chem. Int. Ed. 2002, 41, 2837.
- (15) Trost, B. M. J. Am. Chem. Soc. 1967, 89, 1847.
- (16) Replacement of the methoxy with a triflate group gave low yields in the subsequent oxidation.
- (17) Tobisu, M.; Takahira, T.; Morioka, T.; Chatani, N. J. Am. Chem. Soc. 2016, 138, 6711.
- (18) (a) Mann, F. G.; Pragnell, M. J. J. Chem. Soc. 1965, 4120. (b) Ireland, R. E.; Welch, S. J. Am. Chem. Soc. 1970, 92, 7232.

- (19) (a) Hendrickson, J. B.; Bergeron, R. *Tetrahedron Lett.* 1973, *46*, 4607. (b) Mc Murry, J. E.; Scott, W. J. *Tetrahedron Lett.* 1983, *24*, 979.
- (20) (a) Hayashi, T.; Konishi, M.; Kobori, Y., Kumada, M.; Higuchi, T.; Hirotsu, K. J. Am. Chem. Soc. 1984, 106, 158. (b) Negishi, E. Acc. Chem. Res. 1982, 15, 340.
- (21) Dess, D. B.; Martin, J. C. J. Am. Chem. Soc. 1991, 113, 7277.
- (22) Ding, W. J.; Fang, D.-C. J. Org. Chem. 2001, 66, 6673.
- (23) For an alternative Schönecker C-H oxidation, see Supplementary Information and See, Y. Y.; Hermann, A. T.; Aihara, Y.; Baran, P. S. J. Am. Chem. Soc. 2015, 137, 13776.
- (24) (a) Bolm, C.; Schlingloff, G.; Weickhardt, K. *Tetrahedron Lett.* 1993, 34, 3405. (b) Bolm, C.; Schlingloff, G.; Weickhardt, K. *Tetrahedron Lett.* 1993, 34, 3405.
- (25) (a) Haszeldine, R. N. J. Chem. Soc. 1951, 584. (b) Haszeldine, R. N.; Sharpe, A. G. J. Chem. Soc. 1952, 993.

