

Mild and Efficient Synthesis of Benzoxazoles, Benzothiazoles, Benzimidazoles, and Oxazolo[4,5-*b*]pyridines Catalyzed by Bi(III) Salts Under Solvent-Free Conditions

Iraj Mohammadpoor-Baltork*, Ahmad R. Khosropour, and Seyedeh F. Hojati

Department of Chemistry, Isfahan University, Isfahan, Iran

Received December 22, 2006; accepted January 22, 2007; published online May 11, 2007

© Springer-Verlag 2007

Summary. A series of benzoxazoles, benzothiazoles, benzimidazoles, and oxazolo[4,5-*b*]pyridines was efficiently synthesized from the reactions of *o*-aminophenols, *o*-aminothiophenol, *o*-phenylenediamines, and 2-amino-3-hydroxypyridine with orthoesters in the presence of catalytic amounts of Bi(III) salts, such as Bi(*TFA*)₃, Bi(*OTf*)₃, and BiOCIO₄ · *x*H₂O under solvent-free conditions. The remarkable features of this new protocol are high conversion, very short reaction times, cleaner reaction profiles under solvent-free conditions, straightforward procedure, and use of relatively non-toxic catalysts.

Keywords. Benzoxazoles; Benzothiazoles; Benzimidazoles; Oxazolo[4,5-*b*]pyridines; Bi(III) salts.

Introduction

Benzoxazole, benzothiazole, benzimidazole, and oxazolo[4,5-*b*]pyridine moieties have attracted a great deal of interest in diverse fields of chemistry [1]. Some of these nuclei are found in various naturally occurring compounds [2]. These heterocycles have achieved significance in pharmacology as antibacterial [3], antiviral [4], antifungal [5], anticancer [6], antibiotic [7], anticonsultant [8], and immunosuppressant [9] agents. They have also been used as ligands in asymmetric syntheses [10].


Several methods have been developed for the preparation of these heterocycles including the con-

densation of carboxylic acids [11], acid chlorides [12], nitriles [13], orthoesters [14], amides [15], esters [16], and aldehydes [17] with *o*-aminophenols, *o*-aminothiophenols, and *o*-phenylenediamines, dehydration of *o*-acylaminophenols [18], the reaction of *o*-quinones with amines [19], *Beckmann* rearrangement of *o*-acylphenol oximes [20], and photocyclization of phenolic *Schiff* bases [21]. However, all of these procedures suffer from one or more of the following disadvantages such as long reaction times, low yields of the products, harsh reaction conditions, the use of excess amounts of reagents, tedious workup procedures, and co-occurrence of several side reactions. In addition, some of the catalysts and reagents are expensive, toxic, and air sensitive. Therefore, there is still a need to search for better catalysts that could be superior to the existing ones with regards to toxicity, handling, and operational simplicity.

Results and Discussion

Recently there has been growing interest in the use of Bi(III) salts as catalysts in various organic reactions due to their low toxicity, ease of handling, and relative insensitivity to air and moisture [22]. Bismuth has been heralded as a green element, and the low toxicity of many bismuth compounds is evident from their *LD*₅₀ (rat, oral *LD*₅₀ = 2–22 g/kg [23]). In continuation of our research on the use of Bi(III)

* Corresponding author. E-mail: imbaltork@sci.ui.ac.ir



Scheme 1

salts in organic transformations [24], we now report a simple and environmentally benign methodology for the synthesis of benzoxazole, benzothiazole, benzimidazole, and oxazolo[4,5-*b*]pyridine derivatives from orthoesters under solvent-free conditions using Bi(III) salts as catalysts.

At first, the solvent effect was examined by the reaction of 1 equiv. *o*-aminophenol (**1a**) with 1.1 equiv. triethyl orthoacetate (**2b**) in the presence of 0.015 equiv. Bi(*TFA*)₃ (Scheme 1) in different solvents, such as ClCH₂CH₂Cl, CHCl₃, CH₃OH, CH₃CN, and also under solvent-free conditions at room temperature (Table 1). The best result was obtained


under solvent-free conditions, which is also economically and environmentally favourable (Table 1, entry 5). On the other hand, further examination illustrated that Bi(*TFA*)₃ is essential for this transformation whereas in the absence of the catalyst, only 5% of the product was obtained from the reaction mixture (Table 1, entry 6).

The applicability of this method was then examined for the preparation of benzoxazole, benzothiazole, benzimidazole, and oxazolo[4,5-*b*]pyridine derivatives from various orthoesters under solvent-free conditions. To this aim, different orthoesters were reacted with 2-substituted anilines and 2-amino-3-hydroxypyridine in the presence of catalytic amounts of Bi(*TFA*)₃, Bi(*OTf*)₃, or BiOClO₄ · *x*H₂O (Scheme 2) and the corresponding products were obtained in high to excellent yields (Table 2).

The results showed that 2-substituted benzoxazoles were efficiently and rapidly formed under mild reaction conditions at room temperature in the presence of the above mentioned catalysts in excellent yields (Table 2, entries 1–6), whereas synthesis of benzoxazoles unsubstituted at 2-position needs heating at 85°C (Table 2, entries 7–9). 2-Aminothiophe-

Table 1. Investigation of solvent and catalyst effects on the synthesis of 2-methylbenzoxazole (**3d**)

Entry	Catalyst	Solvent	Time/min	Yield/%
1	Bi(<i>TFA</i>) ₃	ClCH ₂ CH ₂ Cl	15	83
2	Bi(<i>TFA</i>) ₃	CHCl ₃	10	87
3	Bi(<i>TFA</i>) ₃	CH ₃ OH	20	50
4	Bi(<i>TFA</i>) ₃	CH ₃ CN	10	85
5	Bi(<i>TFA</i>) ₃	–	2	96
6	–	–	20	5



Scheme 2

Table 2. Synthesis of benzoxazoles, benzothiazoles, benzimidazoles, and oxazolo[4,5-*b*]pyridines catalyzed by Bi(III) salts under solvent-free conditions

Entry	1	2	3	Catalyst (eq.)	Time/min (Yield/% ^{a,b})			M.P. °C	Ref.
					Bi(TFA) ₃	Bi(OTf) ₃	BiOCIO ₄ · xH ₂ O		
1	1a	2a	3a	0.01	4 (97)	5 (93)	10 (92)	Oil	[14c]
2	1b	2a	3b	0.005	0.5 (98)	1 (95)	2 (95)	59–61	[20]
3	1c	2a	3c	0.01	2.5 (95)	4 (93)	10 (90)	29–31	[20]
4	1a	2b	3d	0.015	2 (96)	3 (93)	10 (85)	Oil	[14c]
5	1b	2b	3e	0.01	0.5 (95)	1 (90)	5 (90)	53–55	[20]
6	1c	2b	3f	0.02	5 (95)	10 (90)	15 (85)	Oil	[20]
7 ^c	1a	2c	3g	0.005	12 (88)	10 (93)	10 (90)	Oil	[14c]
8 ^c	1b	2c	3h	0.005	8 (88)	7 (92)	8 (90)	36–38	[25a]
9 ^c	1c	2c	3i	0.005	15 (87)	10 (90)	15 (88)	42–44	[25b]
10	1d	2a	3j	0.01	2 (86)	4 (81)	4 (80)	Oil	[14c]
11	1d	2b	3k	0.01	3 (81)	5 (80)	5 (80)	Oil	[14a]
12	1d	2c	3l	0.01	5 (83)	5 (81)	7 (83)	Oil	[14a]
13 ^c	1e	2a	3m	0.03	11 (87)	9 (93)	8 (88)	163–165	[25c]
14 ^c	1f	2a	3n	0.04	5 (84)	6 (91)	5 (88)	160–163	[25d]
15 ^c	1e	2b	3o	0.03	5 (82)	4 (90)	4 (82)	172–174	[25e]
16 ^c	1f	2b	3p	0.04	5 (90)	4 (85)	4 (85)	198–200	[25d]
17 ^c	1e	2c	3q	0.015	9 (92)	7 (85)	6 (80)	170–172	[14a]
18 ^c	1f	2c	3r	0.04	8 (85)	8 (83)	10 (80)	112–114	[25f]
19 ^c	1g	2a	3s	0.01	10 (83)	6 (85)	8 (83)	51–53	[14b]
20 ^c	1g	2b	3t	0.01	12 (82)	8 (85)	10 (83)	70–72	[25g]
21 ^c	1g	2c	3u	0.03	12 (52)	10 (55)	12 (53)	69–72	[14b]

^a Products were identified by comparison of their physical and spectral data with those of authentic samples. ^b Isolated yields.^c The reaction was performed at 85°C

nol was also rapidly reacted with orthoesters at room temperature to afford the corresponding benzothiazoles in high yields (Table 2, entries 10–12). The reactions of *o*-phenylenediamines with orthoesters were also investigated. We found that *o*-phenylenediamines reacted rapidly with orthoesters in the presence of Bi(III) salts at 85°C and the corresponding benzimidazoles were isolated in high yields (Table 2, entries 13–18). 2-Amino-3-hydroxypyridine was also reacted with orthoesters at 85°C and the corresponding oxazolo[4,5-*b*]pyridines were isolated in good to high yields (Table 2, entries 19–21).

In conclusion, a very simple and convenient protocol for the synthesis of 2-substituted benzoxazoles, benzothiazoles, benzimidazoles, and oxazolo[4,5-*b*]pyridines using catalytic amounts of Bi(III) salts under solvent-free conditions was demonstrated. Furthermore, availability, stability, and non-toxicity of the catalysts are other noteworthy advantages which make this method eco-friendly and environmentally benign. In addition, mild reaction conditions, absence of solvent, very short reaction times, high

yields of the products, and easy workup make this methodology a valid contribution to the existing processes in the field of these heterocycle syntheses.

Experimental

Chemicals were purchased from Merck or Aldrich chemical companies. Melting points were determined using a Stuart scientific apparatus, the accuracy of which was checked using standard compounds of known melting points. IR spectra were recorded on a Shimadzu IR-435 spectrophotometer. ¹H NMR spectra were recorded in CDCl₃ solvent on a Bruker 500 MHz.

General Procedure for the Synthesis of Benzoxazoles, Benzothiazoles, Benzimidazoles and Oxazolo[4,5-*b*]pyridines Under Solvent-Free Conditions

To a mixture of orthoester (1.1 mmol) and *o*-aminophenol, *o*-aminothiophenol, *o*-phenylenediamine, or 2-amino-3-hydroxypyridine (1 mmol) was added the catalyst (0.005–0.04 mmol Bi(TFA)₃, Bi(OTf)₃, or BiOCIO₄ · xH₂O). The mixture was stirred at room temperature or at 85°C for the appropriate time according to Table 2. The progress of the reaction was monitored by TLC (*n*-hexane:EtOAc, 2:1). After completion of the reaction, the crude product was purified by col-

umn chromatography or recrystallization to afford the pure product (Table 2).

Acknowledgements

We are grateful to the center of excellence (Chemistry) and Office of Graduate Studies of Isfahan University for their financial support.

References

- [1] a) Denny WA, Rewcastle GW, Baguley B (1990) *J Med Chem* **33**: 814; b) Kondo J, Suzuki N, Imaoka T, Kawasaki T, Nakanishi A, Kawahara Y (1994) *Anal Sci* **10**: 17; c) Zhao M, Samoc M, Prasad PN, Reinhardt BA, Sinky M (1992) *Chem Mater* **2**: 670; d) Cassidy PE (1980) *Thermally Stable Polymers*, Marcel Dekker, New York; e) Lin S, Yang L (2005) *Tetrahedron Lett* **46**: 4315; f) Mortimer CG, Wells G, Crochard J-P, Stone EL, Bradshaw TD, Stevens MFG, Westwell AD (2006) *J Med Chem* **49**: 179; g) Chen C, Chen Y-J (2004) *Tetrahedron Lett* **45**: 113; h) Frachy G, Crestini C, Berini R, Salidino R, Micione E (1994) *Heterocycles* **38**: 2621
- [2] a) Kobayashi J, Madono T, Shigemori H (1995) *Tetrahedron Lett* **36**: 5589; b) Rodriguez AD, Ramirez C, Rodriguez II, Gonzales E (1999) *Org Lett* **1**: 527; c) Rodriguez II, Rodriguez AD (2003) *J Nat Prod* **66**: 855; d) Ueki M, Ueno K, Miyadoh S, Abe K, Shibata K, Taniguchi M, Oi S (1993) *J Antibiot* **46**: 1089; e) Paramashivappa R, Kumar PP, Subba Rao PV, Srinivasa Rao A (2003) *Bioorg Med Chem Lett* **13**: 657; f) Koči J, Klimešová V, Waisser K, Kaustová J, Dahse H-M, Möllmann U (2002) *Bioorg Med Chem Lett* **12**: 3275; g) Mann J, Baron A, Opoku-Boahen Y, Johansson E, Parkinson G, Kelland LR, Neidle S (2001) *J Med Chem* **44**: 138; h) Craig WA, LeSueur BW, Skibo EB (1999) *J Med Chem* **42**: 3324
- [3] Yildiz-Oren I, Yalcin I, Aki-Sener E, Ucarturk N (2004) *Eur J Med Chem* **39**: 291
- [4] Song X, Vig BS, Lorenzi PL, Drach JC, Townsend LB, Amidon GL (2005) *J Med Chem* **48**: 1274
- [5] Yamato M (1992) *J Pharm Soc Jpn* **112**: 81
- [6] a) Kumar D, Jacob MR, Reynolds MB, Kerwin SM (2002) *Bioorg Med Chem* **10**: 3997; b) DeLuca MR, Kerwin SM (1997) *Tetrahedron Lett* **38**: 199; c) Sato S, Kajiura T, Noguchi M, Takehana K, Kobayasho T, Tsuji T (2001) *J Antibiot* **54**: 102; d) Ueki M, Shibata K, Taniguchi M (1998) *J Antibiot* **51**: 883
- [7] Evans DA, Sacks CE, Kleschick WA, Taber TR (1979) *J Am Chem Soc* **101**: 6789
- [8] a) Bywater WG, Coleman WR, Kamm O, Merritt HH (1945) *J Am Chem Soc* **67**: 905; b) Benazzouz A, Boraud T, Dubédat P, Boireau A, Stutzmann J-M, Gross C (1995) *Eur J Pharmacol* **284**: 299
- [9] a) Grimmett MR (1996) In: Katritzky AR, Rees CW, Scriven EFV (eds) *Comprehensive Heterocyclic Chemistry II*, vol. 3, Elsevier Science Ltd, Oxford; b) Grimmett MR (1997) *Imidazole and Benzimidazole Synthesis*, Academic Press, San Diego
- [10] Figge A, Altenbach HJ, Brauer DJ, Tielmann P (2002) *Tetrahedron: Asymmetry* **13**: 137
- [11] a) Alcalde E, Dinarés I, Pérez-García L, Roca T (1992) *Synthesis* 395; b) So Y-H, Heeschen JP (1997) *J Org Chem* **62**: 3552; c) Heuser S, Keenan M, Weichert AG (2005) *Tetrahedron Lett* **46**: 9001
- [12] a) Pottorf RS, Chadha NK, Katkevics M, Ozola V, Suna E, Ghane H, Regberg T, Player MR (2003) *Tetrahedron Lett* **44**: 175; b) Nadaf RN, Siddiqui SA, Daniel T, Lahoti RJ, Srinivasan KV (2004) *J Mol Catal A: Chem* **214**: 155
- [13] Hein DW, Alheim RJ, Leavitt JJ (1957) *J Am Chem Soc* **79**: 427
- [14] a) Villemin D, Hammadi M, Martin B (1996) *Synth Commun* **26**: 2895; b) Doise M, Dennin F, Blondeau D, Sliwa H (1990) *Tetrahedron Lett* **31**: 1155; c) Jenkins GL, Knevel AM, Davis CS (1961) *J Org Chem* **26**: 274
- [15] Terashima M, Ishii M (1982) *Synthesis* 484
- [16] a) Chakraborti AK, Rudrawar S, Kaur G, Sharma L (2004) *Synlett* 1533; b) Dolbier Jr WR, Burkholder CR, Médebielle M (1999) *J Fluorine Chem* **95**: 127
- [17] a) Hari A, Karan C, Rodrigues WC, Miller BL (2001) *J Org Chem* **66**: 991; b) Beaulieu PL, Haché B, Von Moos E (2003) *Synthesis* 1683; c) Curini M, Epifano F, Montanari F, Rosati O, Taccone S (2004) *Synlett* 1832; d) Gogoi P, Konwar D (2006) *Tetrahedron Lett* **47**: 79; e) Trivedi R, De SK, Gibbs RA (2006) *J Mol Catal A: Chem* **245**: 8; f) Navarrete-Vázquez G, Moreno-Díaz H, Aguirre-Crespo F, León-Rivera I, Villalobos-Molina R, Muñoz-Muñoz O, Estrada-Soto S (2006) *Bioorg Med Chem Lett* **16**: 4169; g) Lee CL, Lam Y, Lee S-Y (2001) *Tetrahedron Lett* **42**: 109
- [18] Boyd GV (1984) In: Katritzky AR, Rees CW, Potts KT (eds) *Comprehensive Heterocyclic Chemistry*, Pergamon Press, New York, pp 177
- [19] Bu X-Z, Huang Z-S, Zhang M, Ma L, Xiao G-W, Gu L-Q (2001) *Tetrahedron Lett* **42**: 5737
- [20] Bhawal BM, Mayabhate SP, Likhite AP, Deshmukh ARAS (1995) *Synth Commun* **25**: 3315
- [21] Chen Y, Zeng DX (2004) *J Org Chem* **69**: 5037
- [22] a) Suzuki H, Ikegami T, Matano Y (1997) *Synthesis* 249; b) Reglinski J (1998) In: Norman NC (ed) *Chemistry of Arsenic, Antimony and Bismuth*, Balackie Academic and Professional, New York; c) Anderson AM, Blazek JM, Garg P, Payne BJ, Mohan RS (2000) *Tetrahedron Lett* **41**: 1527; d) Leonard NM, Wieland LC, Mohan RS (2002) *Tetrahedron* **58**: 8373; e) Chakraborti AK, Shivani GR (2003) *Synlett* 1805; f) Sabitha G, Venkata Reddy E, Swapna R, Mallikarjun Reddy N, Yadav JS (2004) *Synlett* 1276; g) Yadav JS, Reddy BVS, Premalatha K (2004) *Synlett* 963; h) Gosh R, Maiti S, Chakraborty A (2005) *Synlett* 115; i) Sreedhar B, Swapna V, Sridhar C (2005) *Catal Commun* **6**: 293; j) Ollevier T, Mwene-Mbeja TM (2006) *Tetrahedron Lett* **47**: 4051; k) Fu H, Zhang L, Yao W, Zhu Y (2006)

- Appl Catal B: Env **66**: 100; l) Van Well WJM, Le MT, Schiødt NC, Hoste S, Stoltze P (2006) J Mol Catal A: Chem **256**: 1; m) Sparks DE, Patterson PM, Jacobs G, Dogimont N, Tackett A, Crocker M (2006) Appl Catal B: Env **65**: 44
- [23] a) Irving Sax N, Lewis RJ (1988) *Dangerous Properties of Industrial Materials*, Van Nostrand Reinhold, New York, p 522; b) Wormser U, Nir I (1994) In: Patai S (ed) *The Chemistry of Organic Arsenic, Antimony and Bismuth Compounds*, Wiley, New York, p 715
- [24] a) Firouzabadi H, Mohammadpoor-Baltork I, Kolagar S (2001) Synth Commun **31**: 905; b) Mohammadpoor-Baltork I, Khosropour AR (2001) Molecules **996**; c) Mohammadpoor-Baltork I, Khosropour AR, Aliyan H (2001) Synth Commun **31**: 3411; d) Mohammadpoor-Baltork I, Khosropour AR, Aliyan H (2001) J Chem Res (S) **280**; e) Mohammadpoor-Baltork I, Aliyan H, Khosropour AR (2001) Tetrahedron **57**: 5851; f) Mohammadpoor-Baltork I, Khosropour AR (2002) Monatsh Chem **133**: 189; g) Mohammadpoor-Baltork I, Kharamesh B, Kolagar S (2002) Synth Commun **32**: 1633; h) Mohammadpoor-Baltork I, Khodaei MM, Nikoofar K (2003) Tetrahedron Lett **44**: 591; i) Mohammadpoor-Baltork I, Khosropour AR, Hojati SF (2005) Synlett **2747**; j) Khosropour AR, Mohammadpoor-Baltork I, Ghorbankhani H (2006) Catal Commun **7**: 713; k) Khosropour AR, Mohammadpoor-Baltork I, Ghorbankhani H (2006) Tetrahedron Lett **47**: 3561; l) Mohammadpoor-Baltork I, Memarian HR, Khosropour AR, Nikoofar K (2006) Heterocycles **68**: 1837
- [25] a) Sam J, Plampin JN (1964) J Pharm Sci **53**: 538; b) Roussos M, Lecomte J (1962) Ger **1** **124**: 499, (1963) Chem Abstr **57**: 9858g; c) Raban M, Chang H, Craine L, Hortelano E (1985) J Org Chem **50**: 2205; d) Wang H, Partch RE, Li Y (1997) J Org Chem **62**: 5222; e) Aldrich Catalogue Handbook of Fine Chemicals (2000–2001); f) Kato H, Shiba T, Kitajima E, Kiyosawa T (1976) J Chem Soc Perkin Trans I **863**; g) Takahashi T, Yoneda F (1957) Pharm Bull **5**: 350