

Syntheses of ruthenium-containing heterometallic complexes by use of tridentate phosphine ligands

Nagwa Nawar^a, Anthony K. Smith^{b,*}

^a Department of Chemistry, Mansoura University, Faculty of Science, Mansoura 35516, Egypt

^b Department of Chemistry, University of Liverpool, P.O. Box 147, Liverpool L69 3BX, UK

Received 11 November 1994

Abstract

Treatment of $[\text{RuCl}(\text{dppee})(\eta\text{-C}_5\text{H}_5)]$ [$\text{dppee} = 1,1\text{-bis}(\text{diphenylphosphino})\text{ethene}$] with HPPH_2 gives the addition product $[\text{RuCl}(\text{PPh}_2)_2\text{CHCH}_2\text{PPh}_2](\text{C}_5\text{H}_5)$, **1**. The uncoordinated phosphine group of complex **1** reacts with $[\text{IrCl}(\text{CO})_2(p\text{-toluidine})]$ to give the heterobimetallic complex $[\text{RuCl}(\text{C}_5\text{H}_5)(\text{PPh}_2)_2\text{CHCH}_2\text{PPh}_2]\text{IrCl}(\text{CO})_2$, **2**. Complex **1** reacts with $[\text{Fe}(\text{CO})_5]$ to yield $[\text{RuCl}(\text{C}_5\text{H}_5)(\text{PPh}_2)_2\text{CHCH}_2\text{PPh}_2]\text{Fe}(\text{CO})_4$, **3**, and with $[\text{Mo}(\text{CO})_6]$ to give the heterometallic complex $[\text{RuCl}(\text{C}_5\text{H}_5)(\text{PPh}_2)_2\text{CHCH}_2\text{PPh}_2]\text{Mo}(\text{CO})_5$, **4**. Complex **1** also reacts with $[\text{RuCl}_2(p\text{-cymene})]_2$, $[\text{Ru}_3(\text{CO})_{12}]$ and $[\text{Ru}_3(\text{CO})_{10}(\text{dppee})]$ to give the homometallic complexes $[\text{RuCl}(\text{C}_5\text{H}_5)(\text{PPh}_2)_2\text{CHCH}_2\text{PPh}_2]\text{RuCl}_2(p\text{-cymene})$, **5**, $[\text{RuCl}(\text{C}_5\text{H}_5)(\text{PPh}_2)_2\text{CHCH}_2\text{PPh}_2]\text{Ru}_3(\text{CO})_{11}$, **6**, and $[\text{RuCl}(\text{C}_5\text{H}_5)(\text{PPh}_2)_2\text{CHCH}_2\text{PPh}_2]\text{Ru}_3(\text{CO})_9(\text{dppee})$, **7**, respectively.

Keywords: Ruthenium; Iron; Iridium; Molybdenum; Heterometallic complexes; Tridentate phosphines

1. Introduction

We have previously reported the ring-opening reaction of $[\text{RuCl}(\text{dppee})(\eta\text{-C}_5\text{H}_5)]$ with $[\text{RhCl}(\text{CO})_2]_2$ which leads to formation of the heterobimetallic complex $[(\text{C}_5\text{H}_5)\text{Ru}(\mu\text{-CO})_2\{\mu\text{-PPh}_2\text{C}(\text{=CH}_2)\text{PPh}_2\}\text{RhCl}_2]$ [**1**]. It has been reported that the dppee ligand readily undergoes a Michael-type addition reaction with HPPH_2 to give the tridentate phosphine ligand $(\text{PPh}_2)_2\text{-CHCH}_2\text{PPh}_2$. This addition reaction occurs both on the uncomplexed dppee [**2**] and, more readily, on complexed dppee [**3**]. We have now employed this reaction to synthesise $[\text{RuCl}\{(\text{PPh}_2)_2\text{CHCH}_2\text{PPh}_2\}(\text{C}_5\text{H}_5)]$ **1**, and made use of the dangling phosphine created in this way to synthesise heterometallic complexes. We report below the results of the treatment of complex **1** with $[\text{IrCl}(\text{CO})_2(p\text{-toluidine})]$, $[\text{Fe}(\text{CO})_5]$, $[\text{Mo}(\text{CO})_6]$, $[\text{RuCl}_2(p\text{-cymene})]_2$, $[\text{Ru}_3(\text{CO})_{12}]$ and $[\text{Ru}_3(\text{CO})_{10}(\text{dppee})]$.

2. Results and discussion

The complex $[\text{RuCl}(\text{dppee})(\text{C}_5\text{H}_5)]$ is prepared in high yield by treatment of $[\text{RuCl}(\text{PPh}_3)_2(\text{C}_5\text{H}_5)]$ with dppee [**1**]. The complex $[\text{RuCl}\{(\text{PPh}_2)_2\text{CHCH}_2\text{PPh}_2\}(\text{C}_5\text{H}_5)]$, **1**, is formed in quantitative yield by the base (KO^tBu) catalysed addition of diphenylphosphine to $[\text{RuCl}(\text{dppee})(\text{C}_5\text{H}_5)]$ solution in tetrahydrofuran. The ^{31}P NMR spectrum of **1** (Table 1) consists of a doublet at δ 36.8 ppm [$^3J(\text{PP}) = 9.8$ Hz] as a result of the two coordinated phosphorus atoms and a triplet at δ -22.0 because of the uncoordinated phosphorus atom.

The presence of a “dangling” phosphine in complex **1** provides an opportunity for further reactions with different metal centres, and reactions of this type reported here are summarised in Scheme 1. Thus, complex **1** reacts with $[\text{IrCl}(\text{CO})_2(p\text{-toluidine})]$ at 50°C to give the heterometallic complex $[\text{RuCl}(\text{C}_5\text{H}_5)(\text{PPh}_2)_2\text{-CHCH}_2\text{PPh}_2]\text{IrCl}(\text{CO})_2$, **2**, in good yield. On coordination to Ir, the ^{31}P NMR signal of the dangling phosphine moves from δ -22.0 to δ 17.2, and the PP coupling constant is reduced to 3.7 Hz (see Table 1). The IR spectrum of **2** shows two $\nu(\text{CO})$ bands (2015,

* Corresponding author.

Table 1
Spectroscopic data

Compound	^{31}P NMR ^a	IR (cm ⁻¹) ^b
(1)	36.8 [d, $J(\text{PP})$ 9.8, P_A], -22.0 [t, P_X]	
(2)	43.5 [d, $J(\text{PP})$ 3.7, P_A], 17.2 [t, P_X]	2015s, 1965s
(3)	66.8 [t, $J(\text{PP})$ 4.9, P_X], 42.4 [d, P_A]	2070m, 2020s, 1990vs, 1940m
(4)	41.1 [d, $J(\text{PP})$ 4.9, P_A], 32.3 [t, P_X]	2072m, 1980s, 1940vs
(5)	39.9 [d, $J(\text{PP})$ 4.9, P_A], 21.7 [t, P_X]	
(6)	41.6 [d, $J(\text{PP})$ 4.9, P_A], 31.5 [t, P_X]	2100w, 2060s, 2050sh, 2025s, 2010s, 1988sh, 1950sh
(7)	41.1 [d, $J(\text{PP})$ 3.7, P_A], 32.7 [m, P_X , dppee]	2080w, 2050w, 2000sh, 1990s, 1972s, 1940sh, 1925sh


^a In THF/C₆D₆; coupling constants in Hz; P_A refers to two equivalent P atoms bound to Ru, P_X refers to the unique P atom

^b $\nu(\text{CO})$, in CH₂Cl₂

1965 cm⁻¹), indicating the expected *cis* arrangement of the CO ligands at the square-planar Ir atom.

Complex (1) also reacts with Fe(CO)₅ to give [RuCl(C₅H₅)(PPh₂)₂CHCH₂PPh₂Fe(CO)₄] 3, and with Mo(CO)₆ on irradiation to give [RuCl(C₅H₅)(PPh₂)₂CHCH₂PPh₂Mo(CO)₅] 4. The ruthenium–iron complex 3 was identified from its IR spectrum, which shows $\nu(\text{CO})$ bands typical of a Fe(CO)₄ moiety [4]

(Table 1). The ^{31}P NMR spectrum of 3 shows a triplet at δ 66.8 because of the phosphorus atom coordinated to Fe(CO)₄, similar to that previously reported for the di-iron complex [Fe₂(CO)₇{PPh₂CH}] which also contains a Fe(CO)₄P fragment [5]. The complex 4 is also characterised by IR and NMR spectroscopy (Table 1). Thus, the IR spectrum of 4 in the $\nu(\text{CO})$ region is virtually identical to that reported for complexes of the



Scheme 1.

type $[\text{Mo}(\text{CO})_5(\text{PR}_3)]$ [6], and the ^{31}P NMR chemical shift of the P atom bound to Mo (δ 32.3) is similar to that reported for $[\text{Mo}(\text{CO})_5(\text{PPh}_3)]$.

The reactions of complex **1** with other ruthenium complexes was also investigated. Thus, complex **1** reacts with $[\text{RuCl}_2(p\text{-cymene})]_2$ to form the diruthenium complex $[\text{RuCl}(\text{C}_5\text{H}_5)\{(\text{PPh}_2)_2\text{CHCH}_2\text{PPh}_2\}\text{RuCl}_2(p\text{-cymene})]$, **5**, with $[\text{Ru}_3(\text{CO})_{12}]$ to give $[\text{RuCl}(\text{C}_5\text{H}_5)\{(\text{PPh}_2)_2\text{CHCH}_2\text{PPh}_2\}\text{Ru}_3(\text{CO})_{11}]$ **6**, and with $[\text{Ru}_3(\text{CO})_{10}\{(\text{PPh}_2)_2\text{C}=\text{CH}_2\}]$ to give $[\text{RuCl}(\text{C}_5\text{H}_5)\{(\text{PPh}_2)_2\text{CHCH}_2\text{PPh}_2\}\text{Ru}_3(\text{CO})_9\{(\text{PPh}_2)_2\text{C}=\text{CH}_2\}]$, **7**. These complexes were characterised by IR and NMR spectroscopy (Table 1). The ruthenium cluster complex **6** has an IR spectrum which is virtually identical in the $\nu(\text{CO})$ region to that of $[\text{Ru}_3(\text{CO})_{11}\text{L}]$ [7], while the IR spectrum of **7** is very similar to that of the previously reported complexes $[\text{Ru}_3(\text{CO})_9(\text{dppm})\text{L}]$ [$\text{L}=\text{PPh}_3$ or $\text{PPh}_2\text{CH}_2\text{CH}_2\text{Si}(\text{OEt})_3$] [8] and $[(\text{CO})_3\text{Fe}(\text{PPh}_2)_2\text{CHCH}_2\text{PPh}_2]\text{Ru}_3(\text{CO})_9\{(\text{PPh}_2)_2\text{C}=\text{CH}_2\}$ [4]. The ^{31}P NMR spectra of **6** and **7** also support their formulation as cluster complexes of triruthenium in which an equatorial carbonyl ligand has been substituted by the dangling phosphine group of complex **1**, although in the case of complex **7** the spectrum is complicated by the overlap of the resonances from the $(\text{PPh}_2)_2\text{C}=\text{CH}_2$ ligand with that of the unique phosphorus on the $(\text{PPh}_2)_2\text{CHCH}_2\text{PPh}_2$ ligand.

Complex **5** is characterised by its ^{31}P NMR spectrum, which indicates that the expected halide bridge cleavage reaction has occurred on reaction with the dangling phosphine group of complex **1**.

3. Experimental details

All reactions were carried out under nitrogen unless otherwise stated, using dry, degassed solvents and standard Schlenk-line techniques. IR spectra were recorded as dichloromethane solutions in 0.5 mm NaCl cells on a Perkin-Elmer 681 spectrophotometer; NMR spectra were recorded on Jeol FX-60 or Bruker WM250 instruments. Chemical shifts are relative to 85% H_3PO_4 for ^{31}P NMR spectra. Microanalyses were carried out in the Department of Chemistry, Liverpool University. The compounds $(\text{PPh}_2)_2\text{C}=\text{CH}_2$ [9], $(\text{PPh}_2)_2\text{CHCH}_2\text{PPh}_2$ [2], $[\text{IrCl}(\text{CO})_2(p\text{-toluidene})]$ [10], $[\text{RuCl}_2(p\text{-cymene})]_2$ [11], $[\text{Ru}_3(\text{CO})_{12}]$ [12] and $[\text{Ru}_3(\text{CO})_{10}\{(\text{PPh}_2)_2\text{C}=\text{CH}_2\}]$ [13] were prepared by published procedures.

3.1. Preparation of $[\text{RuCl}\{(\text{PPh}_2)_2\text{C}=\text{CH}_2\}(\text{C}_5\text{H}_5)]$

A solution of $[\text{RuCl}(\text{PPh}_3)_2(\text{C}_5\text{H}_5)]$ (0.362 g, 0.48 mmol) and $(\text{PPh}_2)_2\text{C}=\text{CH}_2$ (0.203 g, 0.52 mmol) in benzene (100 cm^3) was refluxed for 5 h. The volume of the solution was reduced to 15 cm^3 by evaporation under vacuum, and hexane (50 cm^3) was added. On

standing for 24 h at -20°C the solution gave dark red crystals of $[\text{RuCl}\{(\text{PPh}_2)_2\text{C}=\text{CH}_2\}(\text{C}_5\text{H}_5)]$ (0.24 g, 85%). Anal. Found: C, 62.0; H, 4.4. $\text{C}_{31}\text{H}_{27}\text{ClP}_2\text{Ru}$ calc.: C, 62.2; H, 4.5%; M^+ at m/z 598. M^+ based on ^{101}Ru and ^{35}Cl , 598.

3.2. Preparation of $[\text{RuCl}\{(\text{PPh}_2)_2\text{CHCH}_2\text{PPh}_2\}(\text{C}_5\text{H}_5)]$ **1**

A mixture of diphenylphosphine (0.012 g, 0.064 mmol) and $[\text{RuCl}\{(\text{PPh}_2)_2\text{C}=\text{CH}_2\}(\text{C}_5\text{H}_5)]$ (0.038 g, 0.064 mmol) in THF (30 cm^3) was stirred at room temperature for 0.5 h in the presence of a catalytic amount of KO^tBu. The resulting pale-yellow solution was evaporated to dryness and the yellow residue was washed with pentane (3×10 cm^3 portions) and recrystallised from THF, to give complex **1** as a yellow solid (0.035 g, 70%). Anal. Found: C, 65.1; H, 4.7. $\text{C}_{43}\text{H}_{38}\text{ClP}_3\text{Ru}$ calc.: C, 65.9; H, 4.9%.

3.3. Preparation of $[\text{RuCl}(\text{C}_5\text{H}_5)\{(\text{PPh}_2)_2\text{CHCH}_2\text{PPh}_2\}\text{IrCl}(\text{CO})_2]$ **2**

A mixture of $[\text{IrCl}(\text{CO})_2(p\text{-toluidene})]$ (0.025 g, 0.064 mmol) and complex **1** (0.050 g, 0.064 mmol) in THF (30 cm^3) was stirred at 50°C for 24 h. The resulting solution was evaporated to dryness and the residue was recrystallised from THF/benzene to give complex **2** as a yellow solid (0.049 g, 70%). Anal. Found: C, 50.1; H, 3.2. $\text{C}_{45}\text{H}_{38}\text{Cl}_2\text{O}_2\text{P}_3\text{IrRu}$ calc.: C, 50.6; H, 3.6%.

3.4. Preparation of $[\text{RuCl}(\text{C}_5\text{H}_5)\{(\text{PPh}_2)_2\text{CHCH}_2\text{PPh}_2\}\text{Fe}(\text{CO})_4]$ **3**

A mixture of complex **1** (0.050 g, 0.064 mmol) with an equimolar amount of $[\text{Fe}(\text{CO})_5]$ (0.013 g, 0.064 mmol) in THF (10 cm^3) was stirred at 40°C for 4 h. The resulting red solution was evaporated to dryness, and the residue was recrystallised from THF/benzene to yield small red crystals of complex **3** in 80% yield. Anal. Found: C, 59.0; H, 4.0. $\text{C}_{47}\text{H}_{38}\text{ClO}_4\text{P}_3\text{FeRu}$ calc.: C, 59.3; H, 4.0%.

3.5. Preparation of $[\text{RuCl}(\text{C}_5\text{H}_5)\{(\text{PPh}_2)_2\text{CHCH}_2\text{PPh}_2\}\text{Mo}(\text{CO})_5]$ **4**

A mixture of $[\text{Mo}(\text{CO})_6]$ (0.0176 g, 0.067 mmol) and complex **1** (0.052 g, 0.067 mmol) in THF (30 cm^3) was heated at 40°C for 2 h. ^{31}P NMR spectroscopy showed that no reaction had occurred. The mixture was placed in a quartz tube and irradiated with light from a mercury UV lamp for 2 h. Slow evaporation of the solution produced orange crystals of complex **4** (0.039 g, 57%). Anal. Found: C, 56.0; H, 3.3. $\text{C}_{48}\text{H}_{38}\text{ClO}_5\text{P}_3\text{MoRu}$ calc.: C, 56.5; H, 3.7%.

3.6. Preparation of $[RuCl(C_5H_5)\{(PPh_2)_2CHCH_2-PPh_2\}RuCl_2(p\text{-cymene})]$ **5**

A mixture of $[RuCl_2(p\text{-cymene})]_2$ (0.027 g, 0.044 mmol) in $CHCl_3$ (3 cm³) and complex **1** (0.07 g, 0.089 mmol) in THF (30 cm³) was heated at 50°C for 2 h. The resulting red-orange solution was evaporated to dryness and the residue recrystallised from THF/benzene, by slow evaporation, to yield orange crystals of complex **5** (0.074 g, 77%). Anal. Found: C, 58.3; H, 4.3. $C_{53}H_{52}Cl_3P_3Ru_2$ calc.: C, 58.4; H, 4.8%.

3.7. Preparation of $[RuCl(C_5H_5)\{(PPh_2)_2CHCH_2-PPh_2\}Ru_3(CO)_{11}]$ **6**

A mixture of $[Ru_3(CO)_{12}]$ (0.044 g, 0.069 mmol) and complex **1** (0.054 g, 0.069 mmol) in THF (30 cm³) was stirred at 50°C for 3 h, during which the colour of the solution gradually changed from yellow-orange to red-orange. The solution was evaporated to dryness and the residue recrystallised from THF/benzene to yield red-orange crystals of complex **6** (0.063 g, 65%). Anal. Found: C, 46.1; H, 2.6. $C_{54}H_{38}ClO_{11}P_3Ru_4$ calc.: C, 46.5; H, 2.7%.

3.8. Preparation of $[RuCl(C_5H_5)\{(PPh_2)_2CHCH_2-PPh_2\}Ru_3(CO)_9\{(PPh_2)_2C=CH_2\}]$ **7**

A mixture of $[Ru_3(CO)_{10}\{(PPh_2)_2C=CH_2\}]$ (0.051 g, 0.052 mmol) and complex **1** (0.041 g, 0.052 mmol) in THF (30 cm³) was stirred at 50°C for 2 h. The resulting red-orange solution was evaporated to dryness and the residue recrystallised from THF/benzene to yield deep-orange crystals of complex **7** (0.07 g, 79%). Anal.

Found: C, 53.8; H, 4.0. $C_{78}H_{60}ClO_9P_5Ru_4$ calc.: C, 54.0; H, 3.5%.

Acknowledgement

We thank the British Council for a Fellowship (to NN).

References

- [1] M.P. Brown, D. Burns, R. Das, P.A. Dolby, M.M. Harding, R.W. Jones, E.J. Robinson and A.K. Smith, *J. Chem. Soc., Dalton Trans.*, (1991) 351.
- [2] J.L. Bookham, W. McFarlane and I.J. Colquhoun, *J. Chem. Soc., Chem. Commun.*, (1986) 1041.
- [3] G.R. Cooper, D.M. McEwan and B.L. Shaw, *Inorg. Chim. Acta*, 122 (1986) 207.
- [4] P.A. Dolby, M.M. Harding, N. Nawar and A.K. Smith, *J. Chem. Soc., Dalton Trans.*, (1992) 2939.
- [5] J.T. Mague and S.E. Dessens, *J. Organomet. Chem.*, 262 (1984) 347.
- [6] S.O. Grim, D.A. Wheatland and W. McFarlane, *J. Am. Chem. Soc.*, 89 (1967) 5573.
- [7] M.I. Bruce, G. Shaw and F.G.A. Stone, *J. Chem. Soc., Dalton Trans.*, (1972) 2094.
- [8] D.F. Foster, J. Harrison, B.S. Nicholls and A.K. Smith, *J. Organomet. Chem.*, 295 (1985) 99.
- [9] H. Schmidbaur, R. Herr and J. Riede, *Chem. Ber.*, 117 (1984) 2322.
- [10] U. Klabunde, *Inorg. Synth.*, 15 (1974) 82.
- [11] M.A. Bennett and A.K. Smith, *J. Chem. Soc., Dalton Trans.*, (1974) 233.
- [12] M.I. Bruce, J.G. Matison, R.C. Wallis, J.M. Patrick, B.W. Skelton and A.H. White, *J. Chem. Soc., Dalton Trans.*, (1983) 2365.
- [13] J.A. Clucas, R.H. Dawson, P.A. Dolby, M.M. Harding, K. Pearson and A.K. Smith, *J. Organomet. Chem.*, 311 (1986) 153.