

## KF/Al<sub>2</sub>O<sub>3</sub> Catalyzed One-Pot Three-Component Process for the Synthesis of Some 2-Thioxoquinazolin-4(1*H*)-one Derivatives

FARHAD HATAMJAFARI, FARZAD ALJANICHAKOLI\* and MOHAMMADJAVAD MOHAMMADMOHTASHAM

Department of Chemistry, Faculty of Science, Tonekabon Branch, Islamic Azad University, Tonekabon, Iran

\*Corresponding author: E-mail: farzadaljanichakoli@yahoo.com

(Received: 19 June 2012;

Accepted: 22 April 2013)

AJC-13381

Three component one-pot reaction for synthesis of 2-thioxoquinazolin-4(1*H*)-ones are given by the condensation of isatoic anhydride, primary amine and phenyl isothiocyanate by using KF/Al<sub>2</sub>O<sub>3</sub> as an environmentally friendly catalyst.


**Key Words:** 2-Thioxoquinazolin-4(1*H*)-one, One-pot, Three component, KF/Al<sub>2</sub>O<sub>3</sub>, Microwave.

### INTRODUCTION

Today multicomponent reactions (MCRs) emphasizes the development of organic compounds as drug precursor on environmentally friendly when synthesized compounds without solvent and catalyst or water solvent and non-toxic catalyst. Multicomponent reactions aid to play a key role in the development of synthetic compounds because of their possible generation of an adduct in a single step from multi reactants usually used by bond-forming efficiency<sup>1</sup>. Application of multicomponent reaction is on the high chemoselectivity in the presence of all the reactants<sup>2</sup>.

Quinazolinones are very important structure in the field of drug chemistry with wide range of biological and pharmacological activity<sup>3</sup>. The quinazolinones scaffold are important class of fused heterocycles with a show promise such as anti-inflammatory<sup>4</sup>, antimalarial<sup>5</sup>, anti HIV<sup>6</sup> and anti cancer<sup>7</sup> and dihydrofolate reductase inhibitors<sup>8</sup>. Moreover, few methods are reported for the synthesis of 2-thioxoquinazolin-4-ones, as most of the methods reported are for quinazolin-2,4(1*H*,3*H*)-diones<sup>9,10</sup>.

Several methods have been proposed for quinazolinones over the years, provide a simple method with the environmentally friendly catalyst for synthesis of 2-thioxoquinazolinones. As part of our ongoing interest in developing new methods for the synthesis of various heterocyclic scaffolds<sup>11-17</sup>, therefore in this study, we have synthesized derivatives of 2-thioxoquinazolin-4(1*H*)-one with the condensation of isatoic anhydride, primary amine and phenyl isothiocyanate by using KF/Al<sub>2</sub>O<sub>3</sub> as a catalyst (Scheme-I).



Scheme-I

IR spectral data of 3-benzyl-2,3-dihydro-2-thioxoquinazoline-4(1*H*)-one (**4b**) shows the characteristic single broad N-H peak in the range 3280 cm<sup>-1</sup>, a sharp C=O peak at 1698 cm<sup>-1</sup> and C=S peak at 1185 cm<sup>-1</sup>. <sup>1</sup>H NMR shows characteristic broad singlet for N-H at δ 12.01 in addition to those due to aromatic protons at δ 7.33-7.86 ppm<sup>13</sup>. In <sup>13</sup>C NMR C=S appears at δ 170 C=O appears at δ 161 ppm. Therefore in the present investigation we have used KF/Al<sub>2</sub>O<sub>3</sub> is a widely used solid supported reagent which is cheap and easy for catalysis of a variety of reactions. Due to its strongly basic nature it has been used as a replacement for organic bases in a number of organic reactions<sup>18</sup>. Various substituted isothiocyanates were condensed isatoic anhydride, primary amine and phenyl isothiocyanate in water by using KF/Al<sub>2</sub>O<sub>3</sub> as an environmentally friendly catalyst to afford the desired 2-thioxoquinazolin-4(1*H*)-one derivatives in 60-79 % yields and was completed within 90-140 min (Table-1).

All chemicals were obtained from Merck or Fluka without further purification. Melting points were determined on an Electrothermal 9100 melting point apparatus and are uncorrected. IR spectra were measured on a Shimadzu IR-470 spectrophotometer. <sup>1</sup>H and <sup>13</sup>C NMR spectra were determined on Bruker 500 DRX AVANCE instrument at 500 and 125 MHz, respectively. The element analyses (C, H, N) were obtained from a Carlo ERBA Model EA 1108 analyzer carried out on Perkin-Elmer 240c analyzer.

TABLE-1  
SYNTHESIS OF VARIOUS 2-THIOXOQUINAZOLIN-4(1H)-ONES FROM DIFFERENT  
ISATOIC ANHYDRIDE, PRIMARY AMINE AND ISOTHIOCYANATE

| Entry | R <sub>1</sub> | R <sub>2</sub> -NH <sub>2</sub> | m.p. (°C) | Yields (%) | Reaction time (min) |
|-------|----------------|---------------------------------|-----------|------------|---------------------|
| 4a    | H              |                                 | 307-309   | 76         | 100                 |
| 4b    | H              |                                 | 253-256   | 70         | 120                 |
| 4c    | meta-Cl        |                                 | 324-325   | 65         | 120                 |
| 4d    | H              |                                 | 264-266   | 60         | 140                 |
| 4e    | H              |                                 | 317-319   | 76         | 90                  |

**Typical synthesis of compounds (4a-e):** A mixture of KF/Al<sub>2</sub>O<sub>3</sub> (0.2 g), isatoic anhydride (1 mmol), primary amine (2 mmol) and phenyl isothiocyanate (1 mmol) was refluxed in ethanol (5 mL) for 90-140 min. The progress of reaction was monitored by TLC. The mixture was extracted with 3 cm × 30 cm CH<sub>2</sub>Cl<sub>2</sub>, filtered and dried with anh. Na<sub>2</sub>SO<sub>4</sub> sulphate. The solution was dried under high vacuum and the resulting solid residue recrystallized from ethanol to give the pure crystalline solid (4a-e).

**2,3-Dihydro-3-phenyl-2-thioxoquinazolin-4(1H)-one (4a):** White powder, m.p. 307-309 °C; IR (KBr,  $\nu_{\max}$ , cm<sup>-1</sup>): 3239 (N-H), 3091 (C-H<sub>arom</sub>), 1676 (C=O), 1189 (C=S); <sup>1</sup>H NMR (500 MHz, DMSO):  $\delta$  = 7.29-7.96 (m, 9H, CH<sub>arom</sub>), 12.03 (s, 1H, NH); <sup>13</sup>C NMR (125 MHz, DMSO): 118.6, 119.1, 121.3, 124.6, 125.0, 128.8, 129.9, 132.8, 137.1, 138.6, 158.8, 177.0 ppm. MS (m/z, %): 254 (M<sup>+</sup>). Anal. calcd. (%) for C<sub>14</sub>H<sub>10</sub>N<sub>2</sub>OS: C, 66.12; H, 3.96; N, 11.02. Found (%): C, 65.71; H, 3.81; N, 10.74.

**3-Benzyl-2,3-dihydro-2-thioxoquinazolin-4(1H)-one (4b):** White powder, m.p. 253-256 °C; IR (KBr,  $\nu_{\max}$ , cm<sup>-1</sup>): 3280 (N-H), 3080 (C-H<sub>arom</sub>), 1698 (C=O), 1185 (C=S); <sup>1</sup>H NMR (500 MHz, DMSO):  $\delta$  = 5.59 (s, 2H), 7.33-7.86 (m, 8H, CH<sub>arom</sub>), 12.01 (s, 1H, NH); <sup>13</sup>C NMR (125 MHz, DMSO): 44.8, 111.5, 114.4, 122.8, 125.1, 126.7, 127.8, 128.9, 134.1, 135.7, 138.1, 161.0, 170.0 ppm. MS (m/z, %): 268 (M<sup>+</sup>). Anal. calcd. (%) for C<sub>15</sub>H<sub>12</sub>N<sub>2</sub>OS: C, 67.14; H, 4.51; N, 10.44. Found (%): C, 66.89; H, 4.21; N, 10.26.

**7-Chloro-2,3-dihydro-3-phenyl-2-thioxoquinazolin-4(1H)-one (4c):** White powder, m.p. 324-326 °C; IR (KBr,  $\nu_{\max}$ , cm<sup>-1</sup>): 3188 (N-H), 3010 (C-H<sub>arom</sub>), 1700 (C=O), 1200 (C=S); <sup>1</sup>H NMR (500 MHz, DMSO):  $\delta$  = 7.30-7.92 (m, 8H, CH<sub>arom</sub>), 12.06 (s, 1H, NH); <sup>13</sup>C NMR (125 MHz, DMSO): 20.1, 55.5, 113.1, 115.2, 123.9, 124.2, 125.7, 126.5, 129.2, 131.9, 135.6, 138.8, 153.7, 169.3 ppm. MS (m/z, %): 288 (M<sup>+</sup>). Anal. calcd. (%) for C<sub>14</sub>H<sub>9</sub>N<sub>2</sub>OSCl: C, 58.23; H, 3.14; N, 9.70. Found (%): C, 57.95; H, 2.85; N, 9.58.

**2,3-Dihydro-3-(1-phenylethyl)-2-thioxoquinazolin-4(1H)-one (4d):** White powder, m.p. 264-266 °C; IR (KBr,

$\nu_{\max}$ , cm<sup>-1</sup>): 3291 (N-H), 3036 (C-H<sub>arom</sub>), 1677 (C=O), 1179 (C=S); <sup>1</sup>H NMR (500 MHz, DMSO):  $\delta$  = 1.79 (d, *J* = 7.1 Hz, 3H), 2.02 (q, *J* = 7.1 Hz, H), 7.15-7.82 (m, 9H, CH<sub>arom</sub>), 11.96 (s, 1H, NH); <sup>13</sup>C NMR (125 MHz, DMSO): 27.7, 46.5, 58.2, 113.8, 114.5, 115.8, 118.6, 121.9, 125.6, 128.5, 132.4, 134.9, 138.0, 145.4, 147.3, 160.1, 171.9, ppm. MS (m/z, %): 282 (M<sup>+</sup>). Anal. calcd. (%) for C<sub>16</sub>H<sub>14</sub>N<sub>2</sub>OS: C, 68.06; H, 5.00; N, 9.92. Found (%): C, 67.81; H, 4.76; N, 9.71.

**3-(4-Chlorobenzyl)-2,3-dihydro-2-thioxoquinazolin-4(1H)-one (4e):** White powder, m.p. 317-319 °C; IR (KBr,  $\nu_{\max}$ , cm<sup>-1</sup>): 3122 (N-H), 3032 (C-H<sub>arom</sub>), 1681 (C=O), 1209 (C=S); <sup>1</sup>H NMR (500 MHz, DMSO):  $\delta$  = 5.16 (s, 2H), 7.16-7.79 (m, 8H, CH<sub>arom</sub>), 12.02 (s, 1H, NH); <sup>13</sup>C NMR (125 MHz, DMSO): 46.4, 115.8, 122.4, 125.0, 126.7, 127.7, 130.6, 133.2, 134.5, 140.8, 145.8, 157.1, 172.3 ppm. MS (m/z, %): 302 (M<sup>+</sup>). Anal. calcd. (%) for C<sub>15</sub>H<sub>11</sub>N<sub>2</sub>OSCl: C, 59.50; H, 3.66; N, 9.25. Found (%): C, 59.33; H, 3.50; N, 9.18.

## Conclusion

We have developed an efficient, one-pot synthesis of 2-thioxoquinazolin-4(1H)-ones by the condensation of isatoic anhydride, primary amine and phenyl isothiocyanate by using KF/Al<sub>2</sub>O<sub>3</sub> in providing increased yields. KF/Al<sub>2</sub>O<sub>3</sub> can act as heterogeneous catalyst which is low cost and environmentally friendly.

## ACKNOWLEDGEMENTS

The authors gratefully acknowledged the financial support from the Research Council of Tonekabon Branch Islamic Azad University, Iran.

## REFERENCES

- K.C. Nicolaou, D.J. Edmonds and P.G. Bulger, *Angew. Chem. Int. Ed.*, **45**, 7134 (2006).
- F. Hatamjafari and N. Montazeri, *Turkish J. Chem.*, **33**, 797 (2009).
- G.P. Ellis, *Synthesis of Fused Heterocycles.*, John Wiley & Sons, Ltd.: Chichester, 47 (1987).
- K. Tereshima, H. Shimamura, A. Kawase, Y. Tanaka, T. Tanimura, T. Kamisaki, Y. Ishizuka and M. Sato, *Chem. Pharm. Bull.*, **45**, 2021 (1995).

5. D. Gueyrard, V. Gurnel, O. Leoni, S. Palmieri and P. Rollin, *Heterocycles*, **32**, 827 (2000).
6. S.N. Pandeya, D. Sriram, G. Nath and E. Declerca, *Pharm. Acta Helv.*, **74**, 11 (1999).
7. J.F. Wolfe, T.L. Rathman, M.C. Sleevi, J.S.A. Campbell and T.D. Greenwood, *J. Med. Chem.*, **33**, 161 (1990).
8. A. Rosowsky, C.E. Mota, J.E. Wright and S.F. Queener, *J. Med. Chem.*, **37**, 4522 (1994).
9. N.D. Moirangthem and W.S. Laitonjam, *Beilstein J. Org. Chem.*, **6**, 1056 (2010).
10. A. Saeed and U. Flörke, *Crystals*, **1**, 254 (2011).
11. J. Azizian, F. Hatamjafari, A.R. Karimi and M. Shaabanzadeh, *Synthesis*, 765 (2006).
12. J. Azizian, M. Shaabanzadeh, F. Hatamjafari and M.R. Mohammadizadeh, *Arkivoc*, 47 (2006).
13. F. Hatamjafari, *Synth. Commun.*, **36**, 3563 (2006).
14. J. Azizian, F. Hatamjafari and A.R. Karimi, *J. Heterocycl. Chem.*, **43**, 1349 (2006).
15. F. Hatamjafari, N. Montazeri and K. Pourshamsian, *Asian J. Chem.*, **23**, 3287 (2011).
16. F. Hatamjafari, *Orient. J. Chem.*, **28**, 141 (2012).
17. F. Hatamjafari, *Asian J. Chem.*, **25**, 2339 (2013).
18. B.E. Blass, *Tetrahedron*, **58**, 9301 (2002).