

Di- and Linear Tri-nuclear Carbonyl Ruthenium Clusters containing Asymmetrically Bridging 2,7-Disubstituted Naphthyridines†




Markus Mintert and William S. Sheldrick*

Lehrstuhl für Analytische Chemie, Ruhr-Universität Bochum, D-44780 Bochum, Germany

The diruthenium(I) complex *cis*-[Ru₂(bmnapy)₂(CO)₄] **1** was prepared by reaction of 2-benzylamino-7-methyl-1,8-naphthyridine (Hbmnapy) with [Ru₃(CO)₁₂] in benzene. The N¹,N² co-ordination mode of the bridging ligands is supplemented by N⁸ as an axial donor atom. In contrast, the analogous reaction with 7-methyl-1,8-naphthyridin-2-one (Hmonap) or the corresponding 5-methyl-7-phenyl derivative (Hmopnapy) afforded the respective trinuclear clusters *cis*-[Ru₃(monap)₂(CO)₆] **2** and *cis*-[Ru₃(mopnapy)₂(CO)₆] **3**, which exhibit three-atom chains with 46 valence electrons. X-Ray structural analyses demonstrate that the Ru–Ru bond distance remains effectively unchanged on going from the Ru₂ unit in **1** [2.707(1) Å] to the Ru₃ sequence in **3** [average 2.701(2) Å], for which an Ru–Ru–Ru angle of 168.6(3)° is observed.

A number of complexes containing 2,7-disubstituted 1,8-naphthyridines as dinucleating ligands have been reported in the past decade. These include dirhodium(II) species of the type [Rh₂(O₂CMe)₃L]PF₆, which contain symmetrically substituted crescent-shaped ligands such as dpnapy [2,7-bis(2-pyridyl)-1,8-naphthyridine].¹ This neutral compound was also employed in the preparation of [Ru₂Cl₂(bipy)₂(μ-dpnapy)]·[PF₆]₂ (bipy = 2,2'-bipyridine) and [Ru₂(O₂CMe)₃(μ-dpnapy)]PF₆, both of which contain a Ru₂⁴⁺ core and axially co-ordinating dpnapy pyridine rings.² The latter complex and the structurally analogous Ru₂⁵⁺ species [Ru₂(O₂CMe)₃(μ-dcnapy)] (dcnapy = 1,8-naphthyridine-2,7-dicarboxylate) were characterized by X-ray structural analyses.^{2,3} We have also recently described⁴ the quadruply bridged dinuclear complexes [Mo₂(monap)₄] and [Ru₂(monap)₄], in which the asymmetrically substituted ligands monap (Hmonap = 7-methyl-1,8-naphthyridin-2-one) display respectively N,O and N,N co-ordination modes. An extension of this approach led us to study the reaction of [Ru₂Cl(O₂CMe)₄] with Hmopnapy (5-methyl-7-phenyl-1,8-naphthyridin-2-one) in methanol at reflux, which leads to the successive formation of [Ru₂Cl(mopnapy)₂(O₂CMe)₂], [Ru₂(mopnapy)₂(O₂CMe)₂] and [Ru₂(mopnapy)₄].⁵ The reduction of the Ru₂⁵⁺ core in the first complex is accompanied by a change in the co-ordination mode of the bridging naphthyridine derivatives from N,O to N,N in [Ru₂(mopnapy)₂(O₂CMe)₂]. Steric interactions between adjacent phenyl substituents must then be regarded as being responsible for the adoption of the electronically less favourable N,O binding pattern⁵ by three of the bridging ligands in the fully substituted polar complex [Ru₂(mopnapy)₄].

The establishment of ambidentate properties for such asymmetrically substituted naphthyridine derivatives in dinuclear complexes with Ru–Ru bonds prompted us to investigate the ability of these ligands to adopt a trinucleating role in a suitable linear three-atom metal cluster. Only one complex, [Rh₃(μ-onap)₂(CO)₂(cod)₂]ClO₄ (Honap = 1,8-naphthyridin-2-one, cod = cycloocta-1,5-diene) is known,⁶ albeit without direct metal–metal bonding [Rh···Rh distances 2.907(3) and 2.912(2) Å], in which a naphthyridine derivative



co-ordinates three metal atoms. A tetranuclear complex [{Mo₂(O₂CBu^t)₃]₂(μ-donap)]·2thf (H₂donap = 1,8-naphthyridine-2,7-dione, thf = tetrahydrofuran), containing two discrete Mo–Mo quadruple bonds of length 2.10 Å, in close proximity at a central Mo–Mo distance of 3.17 Å, has recently been studied by Chisholm and co-workers⁷ as a molecular model for building blocks of stiff-chain polymers.

We now report the reaction of [Ru₃(CO)₁₂] with the naphthyridine derivatives Hbmnapy (2-benzylamino-7-methyl-1,8-naphthyridine), Hmonap and Hmopnapy in benzene at reflux, which leads to the formation of *cis*-[Ru₂(bmnapy)₂(CO)₄] **1**, *cis*-[Ru₃(monap)₂(CO)₆] **2** and *cis*-[Ru₃(mopnapy)₂(CO)₆] **3** respectively. The carbonyl ruthenium cluster [Ru₃(CO)₁₂] was chosen as the starting material as formation of a linear Ru–Ru–Ru chain might be expected to be possible with a minimum of reorganization of the original triangle of metal atoms.

† Supplementary data available: see Instructions for Authors, *J. Chem. Soc., Dalton Trans.*, 1995, Issue 1, pp. xxv–xxx.

Experimental

Solvents were dried and distilled under argon before use. The IR spectra were recorded as KBr discs on a Perkin-Elmer 1760 spectrometer, electronic spectra on a Perkin-Elmer Lambda 15 spectrometer, ^1H NMR spectra on a Bruker AM 400 spectrometer, and FAB mass spectra on a VG Autospec instrument with 3-nitrobenzyl alcohol as the matrix. Elemental analyses were performed on a Carlo Erba 1106 analyser. All reactions were carried out under argon by use of standard Schlenk techniques. The naphthyridine derivatives Hmonapy⁸ and Hmopnapy⁵ were synthesized according to literature procedures. The compound $[\text{Ru}_3(\text{CO})_{12}]$ was obtained from Heraeus and used as received.

Syntheses.—Hbmnapy. A solution of 2-chloro-7-methyl-1,8-naphthyridine (11.16 g, 62.5 mmol) in benzylamine (30 cm³) was refluxed for 8 h. After removal of the solvent in vacuum, the solid was recrystallized from toluene (20 cm³) to afford Hbmnapy in 86% yield (13.35 g) (Found: C, 77.2; H, 6.7; N, 16.6. Calc. for $\text{C}_{16}\text{H}_{15}\text{N}_3$: C, 77.1; H, 6.1; N, 16.8%). FAB mass spectrum: m/z 250 (100%, $[\text{M} - \text{H}]^+$). ^1H NMR (CD_2Cl_2): δ 2.63 (s, 3 H, CH_3), 4.79 (d, 2 H, CH_2), 5.66 (s, 1 H, NH), 6.67 (d, 1 H, H^3), 7.03 (d, 1 H, H^6), 7.36 (m, 5 H, C_6H_5), 7.76 (d, 1 H, H^4) and 7.82 (d, 1 H, H^5). IR (KBr disc): $\tilde{\nu}/\text{cm}^{-1}$ 3202w, 3030w, 1624s, 1522s, 1344s, 1148w, 802w and 734m. UV/VIS (CH_2Cl_2): $\lambda_{\text{max}}/\text{nm}$ ($\epsilon/\text{dm}^3 \text{ mol}^{-1} \text{ cm}^{-1}$) 344 (1347), 262 (1032) and 229 (1386).

***cis*- $[\text{Ru}_2(\text{bmnapy})_2(\text{CO})_4]$ 1.** The compound Hbmnapy (0.050 g, 0.2 mmol) and $[\text{Ru}_3(\text{CO})_{12}]$ (0.043 g, 0.067 mmol) were refluxed in benzene solution (20 cm³) for 1 h, during which time the colour changed from orange to red. After cooling, pentane (20 cm³) was added and the resulting precipitate dried under vacuum to afford complex 1 in 82% yield (0.066 g). Suitable crystals were grown by gas diffusion of pentane into a toluene solution (Found: C, 53.9; H, 4.4; N, 9.8. Calc. for $\text{C}_{36}\text{H}_{28}\text{N}_6\text{O}_4\text{Ru}_2$: C, 53.3; H, 3.5; N, 10.3%). FAB mass spectrum: m/z 811 (100, M^+), 783 (19, $[\text{M} - \text{CO}]^+$), 755 (10, $[\text{M} - 2\text{CO}]^+$) and 654 (49%, $[\text{M} - \text{Ru}(\text{CO})_2]^+$). ^1H NMR (CD_2Cl_2): δ 2.25 (s, 3 H, CH_3), 4.76 (s, 2 H, CH_2), 6.26 (d, 1 H, H^3), 6.37 (d, 1 H, H^6), 6.94 (m, 5 H, C_6H_5), 7.16 (d, 1 H, H^4) and 7.35 (d, 1 H, H^5). IR (KBr disc): $\tilde{\nu}/\text{cm}^{-1}$ 2018s, 1941s (CO), 1623m, 1555m, 1440w and 1026w. UV/VIS (CH_2Cl_2): $\lambda_{\text{max}}/\text{nm}$ ($\epsilon/\text{dm}^3 \text{ mol}^{-1} \text{ cm}^{-1}$) 402 (470), 348 (811) and 274 (1454).

***cis*- $[\text{Ru}_3(\text{monapy})_2(\text{CO})_6]$ 2.** A benzene solution (20 cm³) of Hmonapy (0.032 g, 0.2 mmol) and $[\text{Ru}_3(\text{CO})_{12}]$ (0.064 g, 0.1 mmol) was refluxed for 1 h changing from orange to red with the formation of a red precipitate, which was filtered off and dried under vacuum to afford complex 2 in 90% yield (0.071 g) (Found: C, 44.8; H, 3.0; N, 5.9. Calc. for $\text{C}_{24}\text{H}_{14}\text{N}_4\text{O}_8\text{Ru}_3$: C, 45.9; H, 2.4; N, 5.9%). FAB mass spectrum: m/z 791 (51, $[\text{M} + \text{H}]^+$), 763 (27, $[\text{M} + \text{H} - \text{CO}]^+$) and 633 (100%, $[\text{M} - \text{Ru}(\text{CO})_2]^+$). ^1H NMR (CD_2Cl_2): δ 2.62 (s, 3 H, CH_3), 6.98 (d, 1 H, H^3), 7.09 (d, 1 H, H^6), 7.66 (d, 1 H, H^4) and 7.84 (d, 1 H, H^5). IR (KBr disc): $\tilde{\nu}/\text{cm}^{-1}$ 2060s, 2017s, 1999s (CO), 1651m, 592m, 576m and 448w. UV/VIS (CH_2Cl_2): $\lambda_{\text{max}}/\text{nm}$ ($\epsilon/\text{dm}^3 \text{ mol}^{-1} \text{ cm}^{-1}$) 422 (2757), 342 (8757) and 276 (9233).

***cis*- $[\text{Ru}_3(\text{mopnapy})_2(\text{CO})_6]$ 3.** The compound Hmopnapy (0.047 g, 0.2 mmol) and $[\text{Ru}_3(\text{CO})_{12}]$ (0.064 g, 0.1 mmol) were refluxed in benzene solution (20 cm³) for 1 h, during which time the colour turned from orange to deep red. After cooling pentane (20 cm³) was added and the resulting precipitate dried under vacuum to afford complex 3 in 94% yield (0.089 g). Suitable crystals were grown by gas diffusion of pentane into a solution of 3 in 1,2,4-trichlorobenzene (Found: C, 37.4; H, 2.2; N, 6.4. Calc. for $\text{C}_{36}\text{H}_{22}\text{N}_4\text{O}_8\text{Ru}_3$: C, 36.5; H, 1.8; N, 7.1%). FAB mass spectrum: m/z 942 (100, M^+), 914 (15, $[\text{M} - \text{CO}]^+$), 886 (9, $[\text{M} - 2\text{CO}]^+$), 785 (44, $[\text{M} - \text{Ru}(\text{CO})_2]^+$) and 628 (23%, $[\text{M} - 2\text{Ru}(\text{CO})_2]^+$). ^1H NMR (CD_2Cl_2): δ 2.66 (s, 3 H, CH_3), 6.85 (d, 1 H, H^3), 7.27 (d, 1 H, H^6), 7.35, 7.82 (m, 5 H, C_6H_5) and 7.88 (d, 1 H, H^4). IR (KBr disc): $\tilde{\nu}/\text{cm}^{-1}$




Fig. 1 Molecular structure of *cis*- $[\text{Ru}_2(\text{bmnapy})_2(\text{CO})_4]$ 1

2042s, 1981s (CO), 1626m, 1575m, 1506m, 1442m, 1369m and 835w. UV/VIS (CH_2Cl_2): $\lambda_{\text{max}}/\text{nm}$ ($\epsilon/\text{dm}^3 \text{ mol}^{-1} \text{ cm}^{-1}$) 441 (11 665), 351 (23 239) and 282 (21 242).

X-Ray Crystallography.—Crystal data and details of data collection and structure refinement for complexes 1 and 3 are presented in Table 1. Cell constants were determined by least-squares refinement on diffractometer angles for 25 reflections automatically centred on a Siemens P4 diffractometer using graphite-monochromated Mo-K α radiation ($\lambda = 0.710 73 \text{ \AA}$). Data collection was performed in the ω mode with respective scan ranges of 1.0 and 1.1° and speed ranges of 2.0–14.6 and 3.0–14.6° min⁻¹. No significant alterations were observed in the control intensities registered every 100 reflections.

Structure analysis and refinement. Patterson synthesis (Ru atoms) followed by standard heavy-atom procedures. Full-matrix least squares on $|F|$ with hydrogen atoms for the naphthyridine ligands at calculated positions. The asymmetric unit of complex 1 contains 0.5 *cis*- $[\text{Ru}_2(\text{bmnapy})_2(\text{CO})_4]$ moieties and 0.5 toluene solvent molecules. That of 3 displays two independent *cis*- $[\text{Ru}_3(\text{mopnapy})_2(\text{CO})_6]$ molecules and three 1,2,4-trichlorobenzene solvent molecules. Anisotropic thermal parameters were introduced for the non-hydrogen atoms of *cis*- $[\text{Ru}_2(\text{bmnapy})_2(\text{CO})_4]$ 1 and the Ru and Cl atoms in the crystal lattice of 3. Weighting scheme: $w = [\sigma^2(F_o) + pF_o^2]^{-1}$, $R' = [\sum w(F_o - F_c)^2 / \sum wF_o^2]^{1/2}$, with $p = 0.0$ (1) and 0.0003 (3). Structure solution and refinement with SHELXS 86 and SELX 76.¹⁰ Fractional atomic coordinates for 1 and 3 are listed in Table 2, selected bond lengths and angles in Table 3.

Additional material available from the Cambridge Crystallographic Data Centre comprises H-atom coordinates, thermal parameters and remaining bond lengths and angles.

Results and Discussion

Reaction of Hbmnapy with $[\text{Ru}_3(\text{CO})_{12}]$ in refluxing benzene leads to the formation of $[\text{Ru}_2(\text{bmnapy})_2(\text{CO})_4]$ 1, independent of whether a 3:1 or 2:1 molar ratio is employed. As may be seen from Fig. 1, the bridging naphthyridine ligands in this diamagnetic diruthenium(II) complex display an N¹,N² coordination mode, in contrast to the N¹,N⁸ binding pattern preferred by dinuclear ruthenium(II) complexes, in the absence of pronounced steric interactions.⁵ Perusal of the bond lengths and angles in Table 3 indicates, however, that N(8) is involved in the co-ordination sphere of the ruthenium atom Ru(1), by

Table 1 X-Ray crystallographic data for complexes **1** and **3**

	1	3
Formula	C ₃₆ H ₂₈ N ₆ O ₄ Ru ₂ ·C ₆ H ₅ CH ₃	C ₃₆ H ₂₂ N ₄ O ₈ Ru ₃ ·1.5C ₆ H ₃ Cl ₃
<i>M</i>	902.9	1213.9
Crystal system	Monoclinic	Monoclinic
Space group	<i>C</i> 2/ <i>c</i>	<i>P</i> <i>n</i>
<i>T</i> /K	293	173
<i>a</i> /Å	18.640(4)	16.828(3)
<i>b</i> /Å	14.503(3)	12.533(3)
<i>c</i> /Å	15.829(3)	21.497(4)
β/°	113.67(3)	106.66(3)
<i>U</i> /Å ³	3919(1)	4343(1)
<i>Z</i>	4	4
<i>F</i> (000)	1824	2388
<i>D</i> _c /g cm ⁻³	1.530	1.856
Crystal size/mm	0.69 × 0.26 × 0.14	0.52 × 0.43 × 0.38
μ/cm ⁻¹	8.21	13.67
Absorption correction	ψ Scan	DIFABS ⁹
2θ Scan range/°	2θ ≤ 55	2θ ≤ 50
Reflections measured	5217	8344
Reflections independent	4863	7959
Observed reflections [<i>I</i> ≥ 2σ(<i>I</i>)]	2766	4248
<i>R</i> _{int}	0.018	0.006
<i>R</i>	0.053	0.050
<i>R</i> '	0.053	0.050
Residual electron density/e Å ⁻³	0.97, -0.72	0.57, -0.36

participation in a four-membered RuNCN chelate ring with Ru(1)–N(1) and Ru(1)–N(8) distances of respectively 2.066(6) and 2.377(7) Å. This N¹,N⁸ co-ordination mode has previously been observed in [Fe(napy)₄][ClO₄]₂¹¹ and [Ru(napy)₄]X₂ (X = PF₆ or Cl).¹² Without involvement of N(8), the dinuclear complex **1** would exhibit a total electron count of only 30. This suggests that the symmetry-related N(8) atoms may be regarded as occupying the axial sites of the dinuclear complex, leading thereby to the expected 34 electron count. The associated stabilization of the Ru₂²⁺ core will presumably outweigh the energetic disadvantage of the N¹,N² bridging mode.

Complex **1** represents the first dinuclear ruthenium(I) species without additional axial ligands to be characterized. Despite the extremely distorted octahedral geometry at Ru(1), the Ru(1)–N(8) distance of 2.377(7) Å is only 0.073 Å longer than the average Ru–N_{ax} distance to the axially co-ordinated pyridine-2-one (Hpyo) molecules in [Ru₂(μ-pyo)₂(CO)₄(Hpyo)₂].¹³ The Ru–Ru bond length of 2.707(1) Å in **1** is similar to distances observed in previously characterized diruthenium(I) complexes, for instance 2.670(1) Å in [Ru₂(μ-pyo)₂(CO)₄(Hpyo)₂]¹³ or 2.711(1) Å in [Ru₂(μ-pyo)₂(CO)₄(PPh₃)₂].¹⁴ As a result of the marked difference of 0.45 Å between the N(1)⋯N(2) and Ru(1)–Ru(1a) distances in the RuNCNRu five-membered ring, the equatorial co-ordination planes of Ru(1) and Ru(1a) are inclined away from one another at an angle of 17.6°. A similar value of 16.2° was observed in [Ru₂(μ-pyo)₂(CO)₄(PPh₃)₂].¹⁴

Evaluation of the molecular geometry of complex **1**, in which the asymmetrically 2,7-disubstituted naphthyridine derivative adopts the novel μ-1κN¹,N⁸:2κN² co-ordination mode, suggests that such ligands could be capable of adopting a trinucleating function for linear Ru₃ chains with similar Ru–Ru distances. However, FAB mass spectra provide no evidence for the formation of a trinuclear species when the reaction between Hbmnapy and [Ru₃(CO)₁₂] is carried out with a 3:1 molar ratio of the starting compounds. In contrast the diamagnetic triruthenium complexes *cis*-[Ru₃(monapy)₂(CO)₆] **2** and *cis*-[Ru₃(mopnapy)₂(CO)₆] **3** may be prepared under analogous conditions. The FAB mass spectra and elemental analyses are in both cases in accordance with the formulation as trinuclear species. Both complexes also display a characteristic loss of an Ru(CO)₂ fragment in their mass spectra, leading to the formation of the cation of the dinuclear species [Ru₂L₂(CO)₄]⁺ (L = monapy or mopnapy). Crystals of **3**




**1****2****3**

Table 2 Fractional atomic coordinates ($\times 10^4$) for complexes **1** and **3** with estimated standard deviations (e.s.d.s) in parentheses

Atom	x	y	z	Atom	x	y	z
Compound 1							
Ru(1)	−783(1)	1199(1)	2266(1)	C(27)	1501(4)	−1626(5)	3430(5)
C(1')	−1161(4)	2067(6)	1333(7)	C(3)	209(4)	−961(5)	4388(5)
O(1')	−1404(4)	2615(5)	766(5)	C(4)	−471(4)	−1280(6)	4367(5)
C(2')	−634(4)	2097(6)	3138(6)	C(5)	−1979(6)	−1100(7)	3580(6)
O(2')	−520(3)	2681(4)	3682(5)	C(6)	−2568(6)	−633(7)	2945(7)
N(1)	−440(3)	163(4)	3244(4)	C(7)	−2443(4)	58(7)	2459(7)
C(2)	250(4)	−207(5)	3818(5)	C(71)	−3109(5)	594(7)	1683(7)
N(2)	882(3)	141(4)	3747(4)	N(8)	−1719(3)	316(4)	2584(4)
C(21)	1657(4)	−181(5)	4421(5)	C(9)	−1121(4)	−150(5)	3214(5)
C(22)	1957(4)	−1031(5)	4128(5)	C(10)	−1204(4)	−872(5)	3730(5)
C(23)	2746(4)	−1244(6)	4614(5)	C(31)	0	6396(10)	2500
C(24)	3069(5)	−2024(6)	4384(6)	C(32)	−167(10)	4995(8)	1656(5)
C(25)	2596(5)	−2598(6)	3675(6)	C(33)	−111(10)	5956(8)	1678(5)
C(26)	1828(5)	−2395(5)	3222(6)	C(34)	0	4566(10)	2500
Compound 3							
Ru(11)	−1638(1)	1533(4)	−609(3)	O(232)	−1677(14)	5309(12)	3349(9)
Ru(13)	1626(1)	1587(4)	637(3)	N(31)	471(15)	2544(25)	4119(12)
Ru(12)	0	1348(4)	0	C(32)	1264(14)	2182(33)	4200(11)
C(111)	−1692(25)	573(22)	−1264(13)	O(32)	1795(13)	2274(13)	4777(8)
O(111)	−1618(17)	−132(16)	−1591(12)	C(33)	1442(20)	1509(28)	3712(13)
C(112)	−1974(18)	549(17)	−95(13)	C(34)	868(14)	1203(32)	3163(16)
O(112)	−2088(15)	−171(11)	203(9)	C(35)	−635(15)	1245(27)	2527(12)
C(121)	174(22)	371(19)	−582(13)	C(351)	−517(23)	615(24)	1996(14)
O(121)	366(15)	−217(17)	−927(10)	C(36)	−1425(18)	1671(27)	2542(13)
C(122)	−96(20)	138(21)	459(17)	C(37)	−1554(14)	2210(26)	3062(12)
O(122)	−24(13)	−568(11)	822(9)	C(371)	−2413(16)	2428(30)	3090(14)
C(131)	2041(22)	823(24)	62(14)	C(372)	−2665(17)	2092(27)	3623(14)
O(131)	2253(20)	309(20)	−306(12)	C(373)	−3484(16)	2363(23)	3583(12)
C(132)	1677(21)	450(19)	1185(14)	C(374)	−3935(17)	2941(21)	3042(10)
O(132)	1721(13)	−234(11)	1556(8)	C(375)	−3665(15)	3312(22)	2526(11)
N(11)	−199(14)	2561(25)	690(13)	C(376)	−2859(16)	3009(28)	2549(15)
C(12)	−988(13)	2957(33)	584(13)	N(38)	−867(13)	2510(30)	3553(12)
O(12)	−1620(13)	2666(15)	98(9)	C(39)	−99(11)	2178(20)	3590(10)
C(13)	−1231(20)	3562(28)	1070(13)	C(310)	28(13)	1519(21)	3072(11)
C(14)	−637(16)	3782(33)	1627(16)	N(41)	174(18)	2180(21)	5401(16)
C(15)	794(15)	3684(28)	2319(13)	C(42)	−590(14)	1693(22)	5277(13)
C(151)	671(26)	4245(27)	2872(16)	O(42)	−1219(14)	2101(11)	4834(9)
C(16)	1644(17)	3425(27)	2366(13)	C(43)	−752(21)	843(22)	5675(14)
C(17)	1790(14)	2863(25)	1861(12)	C(44)	−139(19)	465(34)	6179(18)
C(171)	2628(16)	2521(30)	1836(14)	C(45)	1281(20)	559(31)	6831(18)
C(172)	3247(12)	1931(21)	2272(13)	C(451)	1228(23)	−343(20)	7241(13)
C(173)	4035(13)	1771(19)	2197(11)	C(46)	2081(20)	1038(22)	6900(15)
C(174)	4291(16)	2193(20)	1685(11)	C(47)	2183(15)	1912(28)	6544(17)
C(175)	3694(19)	2780(32)	1231(18)	C(471)	3012(16)	2419(23)	6646(11)
C(176)	2910(23)	2891(35)	1325(17)	C(472)	3428(17)	2914(25)	7230(13)
N(18)	1203(13)	2586(30)	1294(12)	C(473)	4217(20)	3269(28)	7237(15)
C(19)	406(11)	2813(20)	1227(11)	C(474)	4466(19)	3391(25)	6676(11)
C(110)	191(17)	3401(30)	1739(13)	C(475)	4049(18)	2900(25)	6090(14)
N(21)	83(16)	2745(21)	−608(15)	C(476)	3320(18)	2387(26)	6108(12)
C(22)	853(14)	3221(22)	−438(12)	N(48)	1547(17)	2228(23)	6019(14)
O(22)	1431(14)	2983(10)	96(9)	C(49)	771(14)	1894(34)	5936(19)
C(23)	1011(19)	4199(22)	−733(13)	C(410)	656(21)	952(27)	6294(17)
C(24)	417(14)	4530(21)	−1265(13)	Cl(11)	−6312(10)	536(14)	−1230(9)
C(25)	−1047(19)	4402(31)	−2007(18)	Cl(12)	−6280(12)	2704(13)	−503(8)
C(251)	−965(24)	5224(21)	−2460(14)	Cl(14)	−3205(12)	1946(17)	928(8)
C(26)	−1843(19)	3887(22)	−2130(14)	C(11')	−5444(15)	963(22)	−624(15)
C(27)	−1917(16)	3158(28)	−1672(16)	C(12')	−5401(14)	1898(22)	−259(18)
C(271)	−2736(19)	2664(30)	−1712(12)	C(13')	−4667(15)	2195(22)	208(14)
C(272)	−3188(19)	2245(25)	−2310(14)	C(14')	−4047(13)	1420(19)	334(14)
C(273)	−3901(18)	1620(26)	−2417(15)	C(15')	−3984(16)	410(19)	78(14)
C(274)	−4305(19)	1646(26)	−1935(11)	C(16')	−4772(16)	288(25)	−361(15)
C(275)	−3850(17)	2089(26)	−1345(13)	Cl(21)	6594(11)	4495(14)	6102(7)
C(276)	−3065(17)	2559(26)	−1188(12)	Cl(22)	6552(10)	2214(13)	5424(9)
N(28)	−1275(16)	2739(23)	−1188(14)	Cl(24)	3474(10)	3088(12)	3899(5)
C(29)	−540(14)	3230(28)	−1054(16)	C(21')	5747(15)	4164(21)	5449(15)
C(210)	−383(16)	4047(22)	−1485(13)	C(22')	5737(15)	3123(18)	5210(14)
Ru(21)	1914(1)	3471(4)	5466(3)	C(23')	5107(14)	2757(22)	4675(15)
Ru(22)	274(1)	3654(4)	4863(1)	C(24')	4402(12)	3401(18)	4481(12)
Ru(23)	−1350(1)	3405(4)	4221(3)	C(25')	4420(16)	4381(18)	4796(12)

Table 2 (contd.)

C(211)	1963(17)	4582(15)	6030(11)	C(26')	5036(14)	4794(21)	5323(12)
O(211)	2090(14)	5234(13)	6429(8)	Cl(31)	1612(11)	1818(11)	8517(7)
C(212)	2109(19)	4517(17)	4919(13)	Cl(32)	1660(6)	3728(5)	7652(5)
O(212)	2320(14)	5001(14)	4535(8)	Cl(34)	-1542(16)	2965(24)	6370(10)
C(221)	119(22)	4567(19)	5486(12)	C(31')	694(10)	2096(15)	7916(9)
O(221)	83(15)	5221(14)	5856(10)	C(32')	759(9)	2971(16)	7527(10)
C(222)	321(20)	4813(19)	4347(17)	C(33')	38(12)	3193(19)	7026(10)
O(222)	458(14)	5627(12)	4139(10)	C(34')	-668(13)	2576(21)	6984(12)
C(231)	-1689(22)	4322(22)	4771(15)	C(35')	-765(13)	1690(21)	7350(12)
O(231)	-1971(15)	4863(17)	5089(10)	C(36')	-23(11)	1483(18)	7830(10)
C(232)	-1556(20)	4566(18)	3681(14)				

Table 3 Selected bond lengths (Å) and angles (°)

Compound 1			
Ru(1)–Ru(1a)	2.707(1)	Ru(1)–N(8)	2.377(7)
Ru(1)–N(1)	2.066(6)	Ru(1)–N(2a)	2.173(6)
Ru(1)–C(1')	1.851(9)	Ru(1)–C(2')	1.838(9)
Ru(1a)–Ru(1)–N(1)	79.6(2)	Ru(1a)–Ru(1)–N(8)	138.1(1)
Ru(1a)–Ru(1)–N(2a)	88.0(2)	Ru(1a)–Ru(1)–C(1')	104.0(3)
Ru(1a)–Ru(1)–C(2')	88.5(2)	N(1)–Ru(1)–N(8)	58.8(2)
N(1)–Ru(1)–N(2a)	87.1(2)	N(1)–Ru(1)–C(1')	175.2(3)
N(1)–Ru(1)–C(2')	92.5(3)	N(2a)–Ru(1)–N(8)	85.6(2)
N(2a)–Ru(1)–C(1')	90.0(3)	N(2a)–Ru(1)–C(2')	175.6(3)
C(1')–Ru(1)–C(2')	90.7(4)	C(1')–Ru(1)–N(8)	117.3(3)
C(2')–Ru(1)–N(8)	97.1(3)	Ru(1)–N(1)–C(2)	135.8(6)
Ru(1)–N(1)–C(9)	102.6(4)	Ru(1)–N(8)–C(9)	88.1(5)
Ru(1)–N(8)–C(7)	154.7(5)	Ru(1)–N(2a)–C(2a)	121.0(4)
N(1)–C(2)–N(2)	115.4(7)	N(1)–C(9)–N(8)	110.5(7)
Compound 3			
Ru(11)–Ru(12)	2.701(2)	Ru(12)–Ru(13)	2.702(3)
Ru(11)–N(28)	2.15(3)	Ru(11)–O(12)	2.08(2)
Ru(11)–C(111)	1.83(3)	Ru(11)–C(112)	1.83(3)
Ru(12)–N(11)	2.22(3)	Ru(12)–N(21)	2.21(3)
Ru(12)–C(121)	1.83(3)	Ru(12)–C(122)	1.84(3)
Ru(13)–N(18)	2.16(3)	Ru(13)–O(22)	2.07(2)
Ru(13)–C(131)	1.85(3)	Ru(13)–C(132)	1.84(3)
Ru(11)–Ru(12)–Ru(13)	168.6(3)	Ru(12)–Ru(11)–N(28)	85.5(6)
Ru(12)–Ru(11)–O(12)	84.5(6)	Ru(12)–Ru(11)–C(111)	97.8(12)
Ru(12)–Ru(11)–C(112)	95.6(8)	N(28)–Ru(11)–O(12)	89.7(10)
N(28)–Ru(11)–C(111)	88.7(13)	N(28)–Ru(11)–C(112)	177.2(12)
O(12)–Ru(11)–C(111)	177.1(12)	O(12)–Ru(11)–C(112)	87.8(10)
C(111)–Ru(11)–C(112)	93.7(13)	Ru(11)–Ru(12)–N(11)	86.4(6)
Ru(11)–Ru(12)–N(21)	82.8(7)	Ru(11)–Ru(12)–C(121)	93.8(10)
Ru(11)–Ru(12)–C(122)	95.5(10)	Ru(13)–Ru(12)–N(11)	84.9(6)
Ru(13)–Ru(12)–N(21)	89.2(7)	Ru(13)–Ru(12)–C(121)	94.7(11)
Ru(13)–Ru(12)–C(122)	92.9(10)	N(11)–Ru(12)–N(21)	84.3(11)
N(11)–Ru(12)–C(121)	94.3(12)	N(11)–Ru(12)–C(122)	98.8(13)
N(21)–Ru(12)–C(121)	94.3(12)	N(21)–Ru(12)–C(122)	176.4(13)
C(121)–Ru(12)–C(122)	82.6(14)	Ru(12)–Ru(13)–N(18)	85.4(6)
Ru(12)–Ru(13)–O(22)	80.5(6)	Ru(12)–Ru(13)–C(131)	97.2(10)
Ru(12)–Ru(13)–C(132)	95.2(11)	N(18)–Ru(13)–O(22)	81.3(11)
N(18)–Ru(13)–C(131)	175.3(13)	N(18)–Ru(13)–C(132)	89.6(14)
O(22)–Ru(13)–C(131)	95.2(11)	O(22)–Ru(13)–C(132)	170.3(13)
C(131)–Ru(13)–C(132)	94.0(14)		

suitable for an X-ray structural analysis were grown by gas diffusion of pentane into a solution of the complex in 1,2,4-trichlorobenzene. The asymmetric unit in the polar space group Pn contains three solvent molecules and two three-atom clusters, the structure of the first of which is depicted in Fig. 2. The positions of the Ru atoms of the two $[\text{Ru}_3(\text{mop-napy})_2(\text{CO})_6]$ molecules are related by a pseudo inversion centre. As the metrical data for the independent clusters are similar, only the first molecule will be discussed in detail.

Clusters **2** and **3** represent the first examples of three-atom chains bridged by naphthyridine derivatives. The ligands adopt a head-to-tail orientation and display a $\mu_3\text{-}\kappa\text{O}^2\text{:}2\kappa\text{N}^1\text{:}3\kappa\text{N}^8$

co-ordination mode. Inspection of the Ru–Ru bond lengths in **3** [2.701(2), 2.702(3) Å] indicates that these are effectively unchanged in comparison to the diruthenium(t) complex **1** [2.707(1) Å]. The Ru–Ru–Ru angle displays a value of 168.6(3)°. As a result of the blocking of the axial sites by the respective 7-methyl and 7-phenyl substituents in **2** and **3**, these clusters exhibit an unusual electron count of 46. Such linear three-atom chains can invariably be described by a localized bonding description and many examples are known of metal carbonyl clusters with 50 valence electrons,¹⁵ including the triruthenium compound $[\text{Ru}_3(\eta\text{-C}_5\text{H}_5)_2(\text{CO})_8]$.¹⁶ To our knowledge, with the exception of the unusual compound $[\text{Mn}_3(3\text{-MeC}_5\text{H}_6)_4]$




Fig. 2 Structure of the first independent molecule of *cis*-[Ru₃(mopnapy)₂(CO)₆] 3




Fig. 3 Projection of complex 3 parallel to the axis of the Ru₃ chain

(41 electrons),¹⁷ all examples of three-atom chains with less than 50 valence electrons contain at least one metal atom belonging to the platinum group.¹⁵ For metals which conform to the 16-electron rule, a linear three-atom sequence will be expected to be associated with 44 valence electrons. Only one cluster [Ni₃(C₁₂H₁₇)₂]²⁻ has previously been reported¹⁸ which exhibits an electron count of 46 similar to those of 2 and 3. The unusually low number of valence electrons in the trimetallic compounds presented in this work prompts the question as to whether agostic¹⁹ C-H...Ru interactions

between the 7-methyl or 7-phenyl substituents and the terminal ruthenium atoms might provide axial electron pairs. Perusal of the molecular structure of 3 in the direction of the Ru₃ chain (Fig. 3) suggests that this is not the case. Distances of 2.49 and 2.54 Å are observed for Ru(11)...H(276) and Ru(13)...H(176), with the hydrogen atoms markedly displaced from the respective axes of the Ru-Ru bonds. Corroboration for this interpretation is provided by the absence of pronounced ¹H NMR high-field shifts for the protons involved.¹⁹ The phenyl rings are twisted at respective angles of

56.9 and 52.7° relative to the planes of their naphthyridine ring systems.

A formal description as an $\text{Ru}^{\text{I}}\text{--Ru}^0\text{--Ru}^{\text{I}}$ chain is reasonable for cluster **3**. It is interesting that the Ru–Ru single bond lengths [2.701(2), 2.702(3) Å] in this 46-electron cluster are much shorter than in the 50-electron triruthenium cluster $[\text{Ru}_3(\eta\text{-C}_5\text{H}_5)_2(\text{CO})_8]^{16}$ [2.889(1) Å], presumably as a result of the absence of axially co-ordinated ligands in the former compound. The adoption of a trinucleating function by the bridging naphthyridine derivative in **3** leads to marked changes in the X(2)–Ru–Ru (X = N or O) and N(1)–Ru–Ru angles in comparison to those of the dinuclear complex **1**. For instance N(2a)–Ru(1)–Ru(1a) displays a value of 88.0(2)° in the latter compound, which is much larger than the average value of 82.5(6)° in **3**. At the same time the N(1)–Ru–Ru [X(2)] angle increases from 79.6(2) to 83.9(7)° on going from **1** to **3**. This means that the [N(1)] Ru–Ru [X(2)] bond shifts towards N(8) in **1** to allow for axial co-ordination of this nitrogen and away from N(8) in **3** to accommodate the third ruthenium atom in the Ru_3 chain. Significant increases are also observed for the angles N(1)–C(2)–X(2) [115.4(7) in **1**, 123(3)° in **3**], N(1)–C(9)–N(8) [110.5(7) in **1**, 119(3)° in **3**] and C(2)–X(2)–Ru [121.0(4) in **1**, 129(2)° in **3**]. The adoption of a very large C(2)–X(2)–Ru angle, which is necessary for a trinucleating co-ordination mode, will be associated with a relatively small energy disadvantage for an exocyclic oxygen O(2), as in mopnapy^- or monapy^- . In contrast such an angle would be energetically very unfavourable for the trigonal nitrogen atom N(2) in bmnapy^- , with the apparent consequence that this ligand is incapable of bridging a Ru_3 chain. The angle changes discussed above are also associated with pronounced increases in the distances X(2)···N(1) [2.27(1) in **1**, 2.37(3) Å in **3**] and N(1)···N(8) [2.20(1) in **1**, 2.35(3) Å in **3**] on going from the di- to the trinuclear complex. The analogous distances in $[\text{H}_2\text{mopnapy}]\text{Cl}^5$ are 2.257(5) and 2.297(5) Å. In this context, it is interesting that, despite the flexibility of the naphthyridine derivative in **3**, the N–Ru–Ru–N and N–Ru–Ru–O torsion angles in this cluster are relatively small (2.0–8.2°), as is also observed for **1** [N(1)–Ru(1)–Ru(1a)–N(2) 2.7°].

This work has demonstrated, for the first time, a trinucleating capability of substituted naphthyridine ligands for three-atom metal chains. The employment of asymmetrically 2,7-disubstituted derivatives with bulky groups such as methyl or phenyl appears to be essential to prevent polymerization. For instance, the analogous reaction of $[\text{Ru}_3(\text{CO})_{12}]$ with Honapy leads to

the formation of a precipitate, which cannot be dissolved in conventional organic solvents.

References

- 1 W. R. Tikkanen, E. Binamira-Soriaga, W. C. Kaska and P. C. Ford, *Inorg. Chem.*, 1983, **22**, 1147; 1984, **23**, 141; J. L. Bear, L. K. Chau, M. Y. Chavan, F. Lefoulon, R. P. Thummel and K. M. Kadish, *Inorg. Chem.*, 1986, **25**, 1516; R. P. Thummel, F. Lefoulon, D. Williamson and M. Y. Chavan, *Inorg. Chem.*, 1986, **25**, 1675.
- 2 E. Binamira-Soriaga, N. L. Keder and W. C. Kaska, *Inorg. Chem.*, 1990, **29**, 3167.
- 3 J. P. Collin, A. Jouaiti, J.-P. Sauvage, W. C. Kaska, M. A. McLoughlin, N. L. Keder, W. T. A. Harrison and G. D. Stucky, *Inorg. Chem.*, 1990, **29**, 2238.
- 4 W. S. Sheldrick and M. Mintert, *Inorg. Chim. Acta*, 1994, **219**, 23.
- 5 M. Mintert and W. S. Sheldrick, *Inorg. Chim. Acta*, 1995, **236**, 13.
- 6 A. Tiripicchio, F. J. Lahoz, L. A. Oro, M. A. Ciriano and B. E. Villaroya, *Inorg. Chim. Acta*, 1986, **111**, L1.
- 7 R. H. Clayton, M. H. Chisholm, J. C. Huffman and E. B. Lobkovsky, *Angew. Chem.*, 1990, **103**, 893; *J. Am. Chem. Soc.*, 1991, **113**, 8709.
- 8 E. V. Brown, *J. Org. Chem.*, 1965, **30**, 1607; M. Wozniak and M. Skiba, *Pol. J. Chem.*, 1981, **55**, 2429.
- 9 N. Walker and D. Stuart, *Acta Crystallogr., Sect. A*, 1983, **39**, 158.
- 10 G. M. Sheldrick, SHELXS 86, A Program for Structure Determination, University of Göttingen, 1986; SHELX 76, A Program for Structure Refinement, University of Cambridge, 1976.
- 11 A. Clearfield and P. Singh, *Chem. Commun.*, 1970, 389.
- 12 M. Mintert and W. S. Sheldrick, unpublished work.
- 13 P. L. Andreu, J. A. Cabeza, G. A. Carriedo, V. Riera, S. Garcia-Granda, J. Van der Maelen and G. Mori, *J. Organomet. Chem.*, 1991, **421**, 305.
- 14 S. J. Sherlock, M. Cowie, E. Singleton and M. M. de V. Steyn, *J. Organomet. Chem.*, 1989, **361**, 353.
- 15 D. M. P. Mingos and A. S. May, in *The Chemistry of Metal Cluster Complexes*, eds. D. F. Shriver, H. D. Kaesz and R. D. Adams, VCH, New York, 1990, ch. 2, pp. 29–31.
- 16 N. Cook, L. E. Smart, P. Woodward and J. D. Cotton, *J. Chem. Soc., Dalton Trans.*, 1979, 1032.
- 17 M. C. Böhm, R. D. Ernst, R. Gleiter and D. R. Wilson, *Inorg. Chem.*, 1983, **22**, 3815.
- 18 K. Jonas, C. Kruger and J. C. Sekutowski, *Angew. Chem., Int. Ed. Engl.*, 1979, **18**, 487.
- 19 M. Brookhart and M. L. H. Green, *J. Organomet. Chem.*, 1983, **250**, 395.

Received 14th March 1995; Paper 5/01598G