

Shunsaku Ohta*, Kentaro Sato, Ikuo Kawasaki, Yuko Yamaguchi, Satoko Nishio and
Masayuki Yamashita

Kyoto Pharmaceutical University, Misasagi Yamashinaku, Kyoto 607-8414, Japan

E-mail: sohta@mb.kyoto-phu.ac.jp

Received August 6, 2003

Reaction of 2-(1-chloro-2,2-dimethylpropyl)-1-methyl-1*H*-imidazoles with diethyl methylmalonate in the presence of NaH gave a normal S_N product, 2-[(1,1-diethoxycarbonyl)ethyl]-2,2-dimethylpropyl]-1-methyl-1*H*-imidazole (**2a**), and two *tele*-reaction products, 5-(1,1-diethoxycarbonyl)ethyl)-1-methyl-2-(2,2-dimethylpropyl)-1*H*-imidazole (**3a**) and *trans*-4,5-di-(1,1-diethoxycarbonyl)ethyl)-4,5-dihydro-1-methyl-2-(2,2-dimethylpropyl)-1*H*-imidazole (**4a**) in 5, 17 and 70 % yields, respectively. The scope and mechanism of this reaction are discussed.

J. Heterocyclic Chem., **41**, 335 (2004).

Introduction.

Electrophilic substitutions of an imidazole nucleus [1] and nucleophilic substitutions *via* lithioimidazoles [2] have been well known for the preparation of imidazole compounds, and we have applied the reactions of lithioimidazoles [3] and imidazolium salts [4] to the syntheses of several pharmaceutically interesting compounds and natural products [5]. On the other hand, nucleophilic reactions on the imidazole ring have been mainly performed through the activation by quaternization of the ring [6] or introducing an electron-withdrawing group such as halogen atom(s) into the nucleus [7], other examples have been quite rare [8,9]. In the preceding paper, we reported a nucleophilic addition of amines to the imidazole ring of 2-(1-chloro-2,2-dimethylpropyl)-1-methyl-1*H*-imidazole hydrochloride (**1a**) [10]. In this paper, we would like to present the first example and a possible mechanism of a C-C bond-forming nucleophilic *tele*-reaction [11] of carbon nucleophiles with **1a** and its derivatives without any activation steps of the imidazole nucleus.

Results and Discussion.

We first examined a reaction of 2-(1-chloro-2,2-dimethylpropyl)-1-methyl-1*H*-imidazole hydrochloride (**1a**) with diethyl methylmalonate in the presence of NaH (entry 1), and it was found the use of three equivalents of the malonate and NaH gave two *tele*-reaction products, 5-(1,1-diethoxycarbonyl)ethyl)-1-methyl-2-(2,2-dimethyl-


propyl)-1*H*-imidazole (**3a**) and *trans*-4,5-di(1,1-diethoxycarbonyl)ethyl)-4,5-dihydro-1-methyl-2-(2,2-dimethylpropyl)-1*H*-imidazole (**4a**) in 67 % and 11 % yields, respectively, together with a small amount of a normal S_N product [12], 2-[1-(1,1-diethoxycarbonyl)ethyl]-2,2-dimethylpropyl]-1-methyl-1*H*-imidazole (**2a**). The structure of **3a** was determined by HMBC correlations, and the *trans*-orientation of the 4- and 5- substituents of **4a** was confirmed by the similarity of the ¹H-NMR coupling constant (3.0 Hz) of H4 - H5 in the imidazoline ring as the reported value for 4,5-di(*N,N*-dimethylamino)-1-methyl-

Table 1
Nucleophilic Reaction of Malonates to Imidazole Derivatives

Entry	R ¹	R ²	1 / NaH / malonate	Yield (%) (Product)		
				2	3	4
1	<i>t</i> -Bu	Me	1 / 3 / 3	3 (2a)	67 (3a)	11 (4a)
2	<i>t</i> -Bu	Me	1 / 5 / 5	5 (2a)	26 (3a)	57 (4a)
3	<i>t</i> -Bu	Me	1 / 10 / 10	5 (2a)	17 (3a)	70 (4a)
4	<i>t</i> -Bu	Me	1 / 6.5 / 5	5 (2a)	34 (3a)	- [a]
5	<i>i</i> -Pr	Me	1 / 5 / 5	9 (2b)	10 (3b)	48 (4b)
6	Et	Me	1 / 3 / 3	35 (2c)	25 (3c)	30 (4c)
7	<i>t</i> -Bu	H	1 / 5 / 5	16 (2d)	61 (3d)	7 (4d)
8	<i>t</i> -Bu	Et	1 / 5 / 5	5 (2e)	32 (3e)	28 (4e)
9	<i>t</i> -Bu	CH ₂ CH=CH ₂	1 / 5 / 5	6 (2f)	89 (3f)	- [a]
10 [b]	<i>t</i> -Bu	Ph	1 / 2 / 2	3 (2g)	11 (3g)	- [a]

[a] Not detected in the reaction mixture; [b] Unexpected S_N product **5** was also obtained in 44 %.


Scheme 1



2-(2,2-dimethylpropyl)-2-imidazoline (3.8 Hz) [10]. To our knowledge, this is the first example of direct nucleophilic addition of a carbon nucleophile to the imidazole ring without any electron-withdrawing substituent on the nucleus.

We considered the reaction mechanism as shown in Scheme 2. Increasing of the amount of the nucleophile decreased the yield of **3a** and increased the yield of **4a** (entries 2 and 3). In the presence of some excess of NaH over that of diethyl methylmalonate, the di-adduct **4a** was not obtained at all (entry 4) because of the absence of the proton source. This result might suggest that protonation of the intermediate **6** by diethyl methylmalonate at the side chain of the 2-position is an important factor for the formation of **4a**, and in the case of entry 4 the reaction might stop at intermediate **6**, which is converted to **3a** during work-up.

Scheme 2



2-(1-Chloro-2-methylpropyl)-1-methyl-1*H*-imidazole hydrochloride (**1b**) was subjected to similar reaction conditions as that of entry 2 to decrease yield of the addition products **3b** and **4b** (10 and 48 %, respectively) (entry 5), and in the case of 2-(1-chloropropyl)-1-methyl-1*H*-imidazole hydrochloride (**1c**) increased yield of the S_N product **2c** (35 %) (entry 6). In entry 6, the nucleophile and the proton source can approach around the 2-position more easily than in entries 1 - 5 to increase yield of the S_N product **2c**. When the hydrochloride **1a** was treated with simple diethyl malonate instead of diethyl methylmalonate, the mono-adduct **3d** was obtained as the main product in 61 % yield together with some S_N product (**2d**, 16 %) and di-adduct (**4d**, 7 %) (entry 7). The yield of **3d** is comparable with that of **3a** in entry 1, and this indicates that the C5-H of the intermediate **6** ($R^2 = \text{Na}$) is abstracted by the adjacent carbanion and also the proton source $[\text{CH}_2(\text{COOEt})_2]$ can approach more easily around the 2-position of the intermediate. When the reaction of **1a** was carried out with more bulky malonate, diethyl ethylmalonate and diethyl allylmalonate than diethyl methylmalonate, yield of di-adducts was decreased very much whereas yield of mono-adducts was increased (entry 8 and 9). These results


show that selectivity between path-a and path-b seems to depend mainly on the steric hindrance around C5-H of the intermediate **6** and bulkiness of the nucleophile as well as basicity of the nucleophile.

Reaction of **1a** with diethyl sodiophenylmalonate gave surprisingly an unexpected other type of *tele*-product, diethyl 4-[1-(1-methyl-1*H*-imidazol-2-yl)-2,2-dimethylpropyl]phenylmalonate (**5**), which might be raised up from an S_N reaction on the 4-position of the phenyl group of the malonate, in 44 % yield accompanied with a small amount of **2g** and **3g** (entry 10). To our knowledge, this type of unexpected nucleophilic *tele*-reaction seems to be also a very rare example in diethyl phenylmalonate [13].

Furthermore, reactions of **1a** with organometallic reagents such as alkyl Grignard reagents and dialkylzinc were examined (Scheme 3). The *tele*-substitution reactions proceeded by treatment of **1a** with five equivalents of

methylmagnesium bromide or *n*-butylmagnesium chloride to give the corresponding tri-alkylated *tele*-reaction products **8a** (37 %) and **8b** (6 %) together with normal S_N products **7a** (21 %) and **7b** (13 %), respectively. This result might also show that the absence of acidic proton such as malonates in the reaction system was preferred as above-mentioned to provide the mono-adduct *via* path-a. On the other hand, the reaction of **1a** with four equivalents of diethylzinc gave only an S_N product **7c** in 27 % yield.

Scheme 3



Conclusion.

We have found a novel nucleophilic *tele*-reaction to the imidazole nucleus. Further investigations of the present reaction are currently underway.

EXPERIMENTAL

All melting points were measured with a Yanaco MP micro-melting-point apparatus and are uncorrected. The infrared spectra were taken with Shimadzu IR-435 spectrophotometer. ^1H nmr and ^{13}C nmr spectra were measured on Varian UNITY INOVA 400NB (^1H : 400 MHz, ^{13}C : 100 MHz), JEOL EX-300 (^1H : 300 MHz, ^{13}C : 75 MHz) or JEOL EX-270 (^1H : 270 MHz, ^{13}C : 68 MHz) spectrometer and the chemical shifts were expressed in parts per million downfield from tetramethylsilane as the internal standard. Mass spectra (MS) were measured on JEOL JMS BU-20 spectrometer using an electron impact mode (70 eV). Silica gel (Merck Art. 7734) for column chromatography was used.

2-(1-Chloro-2,2-dimethylpropyl)-1-methyl-1*H*-imidazole Hydrochloride (**1a**).

Thionyl chloride (SOCl_2 , 0.73 mL, 10 mmol) was added to a stirred solution of 2-(1-hydroxy-2,2-dimethylpropyl)-1-methyl-1*H*-imidazole [3b] (840 mg, 5 mmol) in CHCl_3 (5 mL) under N_2 at 0 °C. After stirring for 2 h at room temperature, solvents were evaporated to give a crystalline residue, which was recrystallized from EtOH–AcOEt. The compound was obtained as a colorless powder, 961 mg (86 %), mp 168–170 °C; ir (potassium bromide): 3396, 2913, 1591, 1514, 1463 cm^{-1} ; ^1H nmr (300 MHz, CD_3OD): δ 1.15 (9H, s), 4.00 (3H, s), 5.54 (1H, s), 7.62 (1H, d, $J = 2.2$ Hz), 7.66 (1H, d, $J = 1.8$ Hz); ^{13}C nmr (75 MHz, CD_3OD): δ 26.4 x 3, 36.3, 39.7, 59.7, 121.0, 125.4, 145.3; ms: m/z 188 (1.2), 186 (M^+ , 3.9), 171 (5), 151 (15), 132 (32), 130 (100), 121 (13), 95 (36), 81 (11), 57 (6); Hrms Calcd. for $\text{C}_9\text{H}_{15}\text{ClN}_2$: $[\text{M}]^+ = 186.0924$. Found: 186.0933.

Anal. Calcd. for $\text{C}_9\text{H}_{15}\text{ClN}_2$: C, 48.44; H, 7.23; N, 12.55. Found: C, 48.51; H, 7.34; N, 12.24.

2-(1-Chloro-2-methylpropyl)-1-methyl-1*H*-imidazole Hydrochloride (**1b**).

n-Butyl lithium (*n*-BuLi, 1.6 *M* in *n*-hexane; 32.8 mL, 52.5 mmol) was added to a stirred solution of 1-methylimidazole (4.4 mL, 55 mmol) in THF (50 mL) under N_2 at -78 °C. After stirring for 30 min at the same temperature, 2-methylpropanal (4.54 mL, 50 mmol) was added and the resulting mixture was stirred for 2 h at ambient temperature. The mixture was then acidified with HCl aq. (10 %, 50 mL) and washed with diethyl ether (30 mL x 2). The aqueous layer was basified with K_2CO_3 powder and extracted with AcOEt (30 mL x 3). The organic layer was dried over anhydrous sodium sulfate and evaporated to give an oily residue, which was purified by recrystallized from AcOEt–*n*-hexane to give 2-(1-hydroxy-2-methylpropyl)-1-methyl-1*H*-imidazole as colorless prisms, 3.58 g (47 %), mp 77–79 °C; ir (CHCl_3): 3099, 2942, 1489, 1464 cm^{-1} ; ^1H nmr (300 MHz, CDCl_3): δ 0.81 (3H, d, $J = 6.8$ Hz), 1.06 (3H, d, $J = 6.6$ Hz), 2.14 (1H, oct., $J = 6.7$ Hz), 3.67 (3H, s), 4.25 (1H, br s), 4.34 (1H, d, $J = 7.9$ Hz), 6.77 (1H, d, $J = 1.1$ Hz), 6.91 (1H, d, $J = 1.3$ Hz); ^{13}C nmr (75 MHz, CDCl_3): δ 18.6, 19.0, 32.9, 33.9, 72.3, 121.0, 126.6, 149.7; ms: m/z 154 (M^+ , 7.4), 137 (4), 111 (100), 83 (24); Hrms Calcd. for $\text{C}_8\text{H}_{14}\text{N}_2\text{O}$: $[\text{M}]^+ = 154.1106$. Found: 154.1103.

Anal. Calcd. for $\text{C}_8\text{H}_{14}\text{N}_2\text{O}$: C, 62.31; H, 9.15; N, 18.17. Found: C, 62.12; H, 9.26; N, 17.95.

The title compound was prepared in a similar manner as that used for preparation of **1a** except for use of 2-(1-hydroxy-2-methylpropyl)-1-methyl-1*H*-imidazole instead of 2-(1-hydroxy-2,2-dimethylpropyl)-1-methyl-1*H*-imidazole. After evaporation

of the solvents, the hydrochloride (**1b**) was obtained as a pale yellow crystalline residue, which was used in the next reaction without further purification; ir (CHCl_3): 3360, 3138, 2950, 1594, 1464 cm^{-1} ; ^1H nmr (270 MHz, CDCl_3): δ 0.96 (3H, d, $J = 6.6$ Hz), 1.29 (3H, d, $J = 6.4$ Hz), 2.63–2.77 (1H, m), 3.40 (1H, br), 4.10 (3H, s), 5.34 (1H, d, $J = 10.2$ Hz), 7.36 (1H, d, $J = 1.8$ Hz), 7.63 (1H, d, $J = 1.8$ Hz); ^{13}C nmr (68 MHz, CDCl_3): δ 19.8, 20.1, 34.1, 35.8, 55.4, 118.7, 124.2, 143.5; ms: m/z 174 (2.8), 172 (M^+ , 10), 137 (100), 132 (25), 130 (75), 121 (90), 96 (48), 81 (27); Hrms Calcd. for $\text{C}_8\text{H}_{13}\text{ClN}_2$: $[\text{M}]^+ = 172.0767$. Found: 172.0773.

2-(1-Chloropropyl)-1-methyl-1*H*-imidazole Hydrochloride (**1c**).

2-(1-Hydroxypropyl)-1-methyl-1*H*-imidazole was prepared in a similar manner as that used for preparation of 2-(1-hydroxy-2-methylpropyl)-1-methyl-1*H*-imidazole except for use of propanal instead of 2-methylpropanal. The crude product was purified by column chromatography (acetone) to give the alcohol as a yellow viscous oil, 4.41 g (63 %); ir (CHCl_3): 3133, 2941, 1489, 1455 cm^{-1} ; ^1H nmr (CDCl_3 , 400 MHz): δ 0.94 (3H, t, $J = 7.4$ Hz), 1.89 (2H, pent., $J = 7.3$ Hz), 3.69 (3H, s), 4.59 (1H, t, $J = 7.0$ Hz), 6.76 (1H, d, $J = 1.3$ Hz), 6.83 (1H, d, $J = 1.3$ Hz); ^{13}C nmr (CDCl_3 , 100 MHz): δ 10.1, 28.9, 32.8, 68.1, 121.3, 126.4, 149.7; ms: m/z 140 (M^+ , 29), 122 (36), 111 (100), 83 (51), 56 (19); Hrms Calcd. for $\text{C}_7\text{H}_{12}\text{N}_2\text{O}$: $[\text{M}]^+ = 140.0950$. Found: 140.0947.

The title compound was prepared in a similar manner as that used for preparation of **1a** except for use of 2-(1-hydroxypropyl)-1-methyl-1*H*-imidazole instead of 2-(1-hydroxy-2,2-dimethylpropyl)-1-methyl-1*H*-imidazole. After evaporation of the solvents, the crude hydrochloride **1c** was obtained as a pale yellow crystalline residue, which was purified by recrystallized from EtOH–AcOEt to give **1c** as a colorless powder, 865 mg (89 %), mp 138–140 °C; ir (potassium bromide): 3400, 2950, 2769, 1677, 1590, 1522, 1378 cm^{-1} ; ^1H nmr ($\text{DMSO}-d_6$, 400 MHz): δ 1.05 (3H, t, $J = 7.3$ Hz), 2.38 (2H, pent., $J = 7.3$ Hz), 3.92 (3H, s), 5.69 (1H, t, $J = 7.3$ Hz), 7.72 (1H, d, $J = 1.8$ Hz), 7.80 (1H, d, $J = 1.8$ Hz); ^{13}C nmr ($\text{DMSO}-d_6$, 100 MHz): δ 11.1, 28.2, 34.7, 50.9, 119.9, 124.6, 143.8; ms: m/z 160 (2.4), 158 (M^+ , 8.3), 132 (8), 130 (23), 123 (96), 107 (100), 96 (31), 81 (35), 56 (30); Hrms Calcd. for $\text{C}_7\text{H}_{11}\text{ClN}_2$: $[\text{M}]^+ = 158.0611$. Found: 158.0609.

Anal. Calcd. for $\text{C}_7\text{H}_{12}\text{Cl}_2\text{N}_2$: C, 43.10; H, 6.20; N, 14.36. Found: C, 43.38; H, 6.44; N, 14.09.

Synthesis of 2-[1-(1,1-Diethoxycarbonyl)ethyl]-2,2-dimethylpropyl-1-methyl-1*H*-imidazole (**2a**), 5-(1,1-Diethoxycarbonyl)ethyl-1-methyl-2-(2,2-dimethylpropyl)-1*H*-imidazole (**3a**) and (4*R**,5*R**)-4,5-Di(1,1-diethoxycarbonyl)ethyl-4,5-dihydro-1-methyl-2-(2,2-dimethylpropyl)-1*H*-imidazole (**4a**).

General Procedure (Entry 2 in Table 1 as an example).

Sodium hydride (NaH, 60 % in oil, 200 mg, 5 mmol) was added to a suspension of **1a** (223 mg, 1 mmol) in THF (1 mL) under N_2 and ice cooling, and the mixture was stirred for 5 min at 0 °C, then diethyl methylmalonate (0.86 mL, 5 mmol) was added to the mixture and stirring was continued for 4 h at ambient temperature. Water (1 mL) was added to the reaction mixture and the products were extracted with AcOEt (5 mL x 3) and the organic layer was dried over anhydrous sodium sulfate. The solvent was evaporated to give an oily residue, which was separated by column chromatography ($\text{CHCl}_3/\text{MeOH} = 30/1$) to give **2a** (first fraction, 16 mg, 5 %), **3a** (second fraction, 85 mg, 26 %) and **4a** (third fraction, 282 mg, 57 %).

Compound **2a** was obtained as a yellow viscous oil; ir (CHCl₃): 2939, 1720, 1478 cm⁻¹; ¹H nmr (400 MHz, CDCl₃): δ 1.00 (9H, s), 1.01 (3H, t, J = 7.1 Hz), 1.29 (3H, t, J = 7.1 Hz), 1.99 (3H, s), 3.70 (3H, s), 3.81-3.88 (2H, m), 3.95 (1H, s), 4.12-4.29 (2H, m), 6.75 (1H, d, J = 1.3 Hz), 6.94 (1H, d, J = 1.1 Hz); ¹³C nmr (100 MHz, CDCl₃): δ 13.80, 13.84, 17.0, 29.6 x 3, 33.4, 36.8, 47.0, 59.0, 61.4, 61.7, 119.8, 126.4, 147.5, 170.9, 172.1; ms: *m/z* 324 (M⁺, 5), 268 (12), 195 (100), 149 (25), 121 (22); Hrms Calcd. for C₁₇H₂₈N₂O₄: [M]⁺ = 324.2049. Found: 324.2048.

Compound **3a** was obtained as a yellow viscous oil; ir (CHCl₃): 2926, 1721, 1463 cm⁻¹; ¹H nmr (300 MHz, CDCl₃): δ 1.01 [9H, s, C(CH₃)₃], 1.26 [6H, t, J = 7.2 Hz, (OCH₂CH₃)₂], 1.90 [3H, s, CCH₃(CO₂Et)₂], 2.59 [2H, s, CH₂C(CH₃)₃], 3.49 (3H, s, NCH₃), 4.23 [4H, q, J = 7.3 Hz, (OCH₂CH₃)₂], 6.92 (1H, s, Im-H); ¹³C nmr (75 MHz, CDCl₃): δ 13.9 x 2 [(OCH₂CH₃)₂], 21.9 [CCH₃(CO₂Et)₂], 29.6 x 3 [C(CH₃)₃], 32.5 (NCH₃), 32.9 [C(CH₃)₃], 40.0 [CH₂C(CH₃)₃], 53.4 [CCH₃(CO₂Et)₂], 62.0 x 2 [(OCH₂CH₃)₂], 126.6 (C-4), 128.7 (C-5), 149.2 (C-2), 170.4 x 2 [(COOEt)₂]; ms: *m/z* 324 (M⁺, 8), 268 (100), 251 (29), 195 (96), 167 (10), 149 (35), 121 (19), 57 (10); Hrms Calcd. for C₁₇H₂₈N₂O₄: [M]⁺ = 324.2049. Found: 324.2047.

Compound **4a** was obtained as a yellow viscous oil; ir (CHCl₃): 2954, 1719 cm⁻¹; ¹H nmr (400 MHz, CDCl₃): δ 1.04 (9H, s), 1.22-1.29 (12H, m), 1.37 (3H, s), 1.39 (3H, s), 2.14 (1H, d, J = 14.3 Hz), 2.19 (1H, d, J = 14.5 Hz), 2.81 (3H, s), 4.08-4.27 (8H, m), 4.37 (1H, d, J = 3.0 Hz), 4.49 (1H, d, J = 3.0 Hz); ¹³C nmr (100 MHz, CDCl₃): δ 13.8, 13.87, 13.88, 13.91, 15.4, 16.8, 29.9 x 3, 31.4, 34.4, 40.2, 57.7, 57.9, 60.9, 61.1, 61.3, 61.4, 66.7, 69.9, 166.6, 170.0, 170.3, 170.4, 170.6; ms: *m/z* 498 (M⁺, 3), 325 (76), 268 (100), 251 (30), 195 (43), 153 (13), 97 (13); Hrms Calcd. for C₂₅H₄₂N₂O₈: [M]⁺ = 498.2941. Found: 498.2939.

2-[1-(1,1-Diethoxycarbonyl)ethyl]-2-methylpropyl]-1-methyl-1*H*-imidazole (**2b**), 5-(1,1-Diethoxycarbonyl)ethyl]-1-methyl-2-(2-methylpropyl)-1*H*-imidazole (**3b**) and (4*R**,5*R**)-4,5-Di(1,1-diethoxycarbonyl)ethyl]-4,5-dihydro-1-methyl-2-(2-methylpropyl)-1*H*-imidazole (**4b**).

These compounds were prepared in a similar manner as that used for preparation of **2a**, **3a** and **4a** except for use of **1b** instead of **1a**. Title compounds were separated by column chromatography (CHCl₃/MeOH = 20/1).

Compound **2b** was obtained in 9 % yield (28 mg) as a yellow viscous oil; ir (CHCl₃): 2940, 1720 cm⁻¹; ¹H nmr (400 MHz, CDCl₃): δ 0.65 (3H, d, J = 6.6 Hz), 0.94 (3H, d, J = 6.6 Hz), 1.03 (3H, t, J = 7.1 Hz), 1.29 (3H, t, J = 7.1 Hz), 1.80 (3H, s), 2.19-2.32 (1H, m), 3.66 (3H, s), 3.67 (1H, d, J = 8.3 Hz), 3.88-3.93 (2H, m), 4.17-4.30 (2H, m), 6.72 (1H, d, J = 1.3 Hz), 6.94 (1H, d, J = 1.1 Hz); ¹³C nmr (100 MHz, CDCl₃): δ 13.8, 13.9, 15.5, 21.5, 21.7, 31.6, 32.9, 45.3, 57.7, 61.4, 61.6, 119.5, 127.1, 148.4, 171.0, 171.5; ms: *m/z* 310 (M⁺, 11), 268 (12), 237 (9), 195 (55), 137 (100), 123 (10), 96 (46); Hrms Calcd. for C₁₆H₂₆N₂O₄: [M]⁺ = 310.1892. Found: 310.1897.

Compound **3b** was obtained in 10 % yield (30 mg) as a yellow viscous oil; ir (CHCl₃): 2936, 1722, 1460 cm⁻¹; ¹H nmr (400 MHz, CDCl₃): δ 0.97 (6H, d, J = 6.8 Hz), 1.27 (6H, t, J = 7.0 Hz), 1.90 (3H, s), 2.04-2.20 (1H, m), 2.55 (2H, d, J = 7.3 Hz), 3.48 (3H, s), 4.23 (4H, q, J = 7.0 Hz), 6.90 (1H, s); ¹³C nmr (100 MHz, CDCl₃): δ 13.9 x 2, 21.8, 22.5 x 2, 28.0, 31.9, 36.3, 53.3, 62.0 x 2, 126.6, 128.6, 150.4, 170.4 x 2; ms: *m/z* 310 (M⁺, 27), 268 (100), 237 (87), 195 (20), 163 (9); Hrms Calcd. for C₁₆H₂₆N₂O₄: [M]⁺ = 310.1892. Found: 310.1897.

Compound **4b** was obtained in 48 % yield (231 mg) as a yellow viscous oil; ir (CHCl₃): 2936, 1719 cm⁻¹; ¹H nmr (400 MHz, CDCl₃): δ 0.97 (3H, d, J = 6.4 Hz), 0.98 (3H, d, J = 6.6 Hz), 1.20-1.30 (12H, m), 1.35 (3H, s), 1.37 (3H, s), 1.93-2.03 (1H, m), 2.07 (2H, d, J = 6.6 Hz), 2.81 (3H, s), 4.08-4.28 (8H, m), 4.35 (1H, d, J = 2.9 Hz), 4.46 (1H, d, J = 2.9 Hz); ¹³C nmr (100 MHz, CDCl₃): δ 13.8 x 2, 13.9 x 2, 15.2, 16.7, 22.7, 23.1, 26.3, 33.7, 36.7, 57.8, 57.9, 61.0, 61.2, 61.39, 61.42, 66.7, 69.7, 167.8, 170.0, 170.3, 170.4, 170.6; ms: *m/z* 484 (M⁺, 3), 311(100), 268 (41), 237 (53), 129 (12), 74 (10); Hrms Calcd. for C₂₄H₄₀N₂O₈: [M]⁺ = 484.2784. Found: 484.2782.

2-[1-(1,1-Diethoxycarbonyl)ethyl]propyl]-1-methyl-1*H*-imidazole (**2c**), 5-(1,1-Diethoxycarbonyl)ethyl]-1-methyl-2-propyl-1*H*-imidazole (**3c**) and (4*R**,5*R**)-4,5-Di(1,1-diethoxycarbonyl)ethyl]-4,5-dihydro-1-methyl-2-propyl-1*H*-imidazole (**4c**).

These compounds were prepared in a similar manner as that used for preparation of **2a**, **3a** and **4a** except for use of **1c** instead of **1a**. Title compounds were separated by column chromatography (AcOEt).

Compound **2c** was obtained in 35% yield (104 mg) as a yellow viscous oil; ir (CHCl₃): 2940, 1721, 1456 cm⁻¹; ¹H nmr (400 MHz, CDCl₃): δ 0.69 (3H, t, J = 7.3 Hz), 1.08 (3H, t, J = 7.0 Hz), 1.28 (3H, t, J = 7.1 Hz), 1.63-1.71 (1H, m), 1.72 (3H, s), 1.88-2.00 (1H, m), 3.59 (1H, dd, J = 11.5, 2.9 Hz), 3.67 (3H, s), 3.94-4.02 (2H, m), 4.24 (2H, q, J = 7.1 Hz), 6.74 (1H, d, J = 1.1 Hz), 6.98 (1H, d, J = 1.1 Hz); ¹³C nmr (100 MHz, CDCl₃): δ 12.3, 13.8, 14.0, 15.8, 25.0, 32.8, 41.2, 57.9, 61.2, 61.5, 119.9, 127.3, 147.4, 171.3, 171.5; ms: *m/z* 296 (M⁺, 8), 251 (9), 223 (17), 195 (8), 123 (100), 96 (11); Hrms Calcd. for C₁₅H₂₄N₂O₄: [M]⁺ = 296.1736. Found: 296.1739.

Compound **3c** was obtained in 25 % yield (74 mg) as a yellow viscous oil; ir (CHCl₃): 2940, 1722, 1459 cm⁻¹; ¹H nmr (400 MHz, CDCl₃): δ 1.01 (3H, t, J = 7.3 Hz), 1.27 (6H, t, J = 7.1 Hz), 1.72-1.84 (2H, m), 1.89 (3H, s), 2.63 (2H, t, J = 7.7 Hz), 3.48 (3H, s), 4.23 (4H, q, J = 7.1 Hz), 6.88 (1H, s); ¹³C nmr (100 MHz, CDCl₃): δ 13.9 x 3, 20.7, 21.7, 29.4, 31.7, 53.2, 62.0 x 2, 126.5, 128.7, 150.8, 170.3 x 2; ms: *m/z* 296 (M⁺, 9), 268 (25), 223 (100), 195 (22), 149 (38); Hrms Calcd. for C₁₅H₂₄N₂O₄: [M]⁺ = 296.1736. Found: 296.1733.

Compound **4c** was obtained in 30 % yield (139 mg) as a yellow viscous oil; ir (CHCl₃): 2945, 1719, 1604, 1444 cm⁻¹; ¹H nmr (400 MHz, CDCl₃): δ 0.96 (3H, t, J = 7.4 Hz), 1.22-1.29 (12H, m), 1.33 (3H, s), 1.35 (3H, s), 1.53-1.66 (2H, m), 2.10-2.22 (2H, m), 2.81 (3H, s), 4.08-4.30 (8H, m), 4.32 (1H, d, J = 2.9 Hz), 4.47 (1H, d, J = 2.9 Hz); ¹³C nmr (100 MHz, CDCl₃): δ 13.8, 13.90, 13.92, 13.94, 14.2, 15.2, 16.4, 20.1, 30.0, 33.5, 57.7, 58.0, 61.0, 61.2, 61.39, 61.42, 66.6, 69.4, 168.4, 170.0, 170.4 x 2, 170.6; ms: *m/z* 470 (M⁺, 1), 297 (100), 268 (10), 223 (61), 149 (10), 74 (9); Hrms Calcd. for C₂₃H₃₈N₂O₈: [M]⁺ = 470.2628. Found: 470.2629.

2-[1-(1,1-Diethoxycarbonylmethyl)-2,2-dimethylpropyl]-1-methyl-1*H*-imidazole (**2d**), 5-(1,1-Diethoxycarbonylmethyl)-1-methyl-2-(2,2-dimethylpropyl)-1*H*-imidazole (**3d**) and (4*R**,5*R**)-4,5-Di(1,1-diethoxycarbonylmethyl)-4,5-dihydro-1-methyl-2-(2,2-dimethylpropyl)-1*H*-imidazole (**4d**).

These compounds were prepared in a similar manner as that used for preparation of **2a**, **3a** and **4a** except for use of diethyl malonate instead of diethyl methylmalonate. Title compounds were separated by column chromatography (AcOEt/*n*-hexane = 1/1).

Compound **2d** was obtained in 16 % yield (51 mg) as a yellow viscous oil; ir (CHCl₃): 2940, 1741, 1720, 1479 cm⁻¹; ¹H nmr (270 MHz, CDCl₃): δ 0.96 (9H, s), 1.03 (3H, t, J = 7.1 Hz), 1.30 (3H, t, J = 7.2 Hz), 3.54 (1H, d, J = 10.6 Hz), 3.70 (3H, s), 3.89 (2H, q, J = 7.1 Hz), 4.15 (1H, d, J = 10.6 Hz), 4.16-4.31 (2H, m), 6.72 (1H, d, J = 1.3 Hz), 6.89 (1H, d, J = 1.3 Hz); ¹³C nmr (75 MHz, CDCl₃): δ 13.8, 13.9, 27.9 x 3, 33.1, 35.4, 45.0, 54.2, 61.3, 61.7, 119.8, 126.7, 148.2, 168.3, 169.5; ms: *m/z* 310 (M⁺, 3), 254 (19), 181 (100), 135 (12), 96 (14); Hrms Calcd. for C₁₆H₂₆N₂O₄: [M]⁺ = 310.1892. Found: 310.1903.

Compound **3d** was obtained in 61 % yield (188 mg) as a yellow viscous oil; ir (CHCl₃): 2934, 1730, 1464 cm⁻¹; ¹H nmr (300 MHz, CDCl₃): δ 1.00 (9H, s), 1.28 (6H, t, J = 7.1 Hz), 2.59 (2H, s), 3.51 (3H, s), 4.15-4.35 (4H, m), 4.69 (1H, d, J = 0.3 Hz), 7.00 (1H, d, J = 0.5 Hz); ¹³C nmr (75 MHz, CDCl₃): δ 13.9 x 2, 29.5 x 3, 31.5, 32.8, 40.0, 49.4, 62.0 x 2, 122.5, 127.8, 148.3, 166.6 x 2; ms: *m/z* 310 (M⁺, 12), 254 (100), 182 (26), 110 (29); Hrms Calcd. for C₁₆H₂₆N₂O₄: [M]⁺ = 310.1892. Found: 310.1903.

Compound **4d** was obtained in 7 % yield (32 mg) as a yellow viscous oil; ir (CHCl₃): 2941, 1722, 1387 cm⁻¹; ¹H nmr (400 MHz, CDCl₃): δ 1.00 (9H, s), 1.24-1.29 (12H, m), 2.09 (1H, d, J = 13.9 Hz), 2.13 (1H, d, J = 13.9 Hz), 2.88 (3H, s), 3.67 (1H, d, J = 6.4 Hz), 3.70 (1H, d, J = 4.2 Hz), 4.12-4.25 (8H, m), 4.28-4.30 (1H, br s), 4.46 (1H, dd, J = 6.4, 5.1 Hz); ¹³C nmr (100 MHz, CDCl₃): δ 13.9 x 3, 14.0, 29.8 x 3, 31.5, 34.0, 39.8, 56.4, 56.8, 61.2, 61.4, 61.5, 61.6, 64.4, 68.0, 166.0, 167.2, 167.55, 167.64, 167.8; ms: *m/z* 470 (M⁺, 17), 414 (74), 342 (84), 311 (100), 270 (42), 254 (99), 133 (37), 115 (40); Hrms Calcd. for C₂₃H₃₈N₂O₈: [M]⁺ = 470.2628. Found: 470.2632.

2-[1-(1,1-Diethoxycarbonylpropyl)-2,2-dimethylpropyl]-1-methyl-1*H*-imidazole (**2e**), 5-(1,1-Diethoxycarbonylpropyl)-1-methyl-2-(2,2-dimethylpropyl)-1*H*-imidazole (**3e**) and (4*R*^{*}, 5*R*^{*})-4,5-Di(1,1-diethoxycarbonylpropyl)-4,5-dihydro-1-methyl-2-(2,2-dimethylpropyl)-1*H*-imidazole (**4e**).

These compounds were prepared in a similar manner as that used for preparation of **2a**, **3a** and **4a** except for use of diethyl ethylmalonate instead of diethyl methylmalonate. Title compounds were separated by column chromatography (CHCl₃/MeOH = 20/1).

Compound **2e** was obtained in 5 % yield (18 mg) as a yellow viscous oil; ir (CHCl₃): 2939, 1716, 1479 cm⁻¹; ¹H nmr (400 MHz, CDCl₃): δ 0.83 (3H, t, J = 7.3 Hz), 1.03 (9H, s), 1.21 (3H, t, J = 7.1 Hz), 1.33 (3H, t, J = 7.1 Hz), 1.81-1.95 (2H, m), 3.63 (3H, s), 3.72 (1H, s), 3.97-4.38 (4H, m), 6.76 (1H, d, J = 1.3 Hz), 7.02 (1H, d, J = 1.3 Hz); ¹³C nmr (100 MHz, CDCl₃): δ 10.4, 13.8, 13.9, 27.8, 29.6 x 3, 33.2, 37.2, 49.9, 60.9, 61.1, 61.7, 119.4, 126.9, 146.8, 170.5, 171.8; ms: *m/z* 338 (M⁺, 1), 209 (100), 163 (49), 151 (19), 135 (17); Hrms Calcd. for C₁₈H₃₀N₂O₄: [M]⁺ = 338.2205. Found: 338.2209.

Compound **3e** was obtained in 32 % yield (109 mg) as a yellow viscous oil; ir (CHCl₃): 2935, 1720, 1462 cm⁻¹; ¹H nmr (400 MHz, CDCl₃): δ 1.00 (9H, s), 1.01 (3H, t, J = 7.4 Hz), 1.26 (6H, t, J = 7.1 Hz), 2.39 (2H, q, J = 7.4 Hz), 2.59 (2H, s), 3.48 (3H, s), 4.23 (4H, q, J = 7.2 Hz), 6.96 (1H, s); ¹³C nmr (100 MHz, CDCl₃): δ 9.6, 13.9 x 2, 28.0, 29.6 x 3, 32.5, 32.8, 39.9, 57.7, 61.7 x 2, 127.3, 127.5, 148.9, 169.4 x 2; ms: *m/z* 338 (M⁺, 9), 282 (100), 265 (35), 209 (64), 163 (22), 135 (12), 111 (16); Hrms Calcd. for C₁₈H₃₀N₂O₄: [M]⁺ = 338.2205. Found: 338.2214.

Compound **4e** was obtained in 28 % yield (147 mg) as a yellow viscous oil; ir (CHCl₃): 2937, 1717 cm⁻¹; ¹H nmr (400 MHz,

CDCl₃): δ 0.90 (3H, t, J = 7.3 Hz), 0.97 (3H, t, J = 7.3 Hz), 1.03 (9H, s), 1.25-1.31 (12H, m), 1.70-1.80 (1H, m), 1.90-1.96 (1H, m), 2.08-2.18 (4H, m), 2.79 (3H, s), 4.05-4.27 (8H, m), 4.55 (1H, br s), 4.59 (1H, d, J = 2.4 Hz); ¹³C nmr (100 MHz, CDCl₃): δ 9.5, 9.6, 13.9, 13.96, 13.98, 14.04, 25.1, 25.5, 29.8 x 3, 31.3, 33.4, 40.3, 60.8 x 2, 61.15, 61.19, 61.23, 62.2, 68.0, 68.6, 166.1, 169.3, 170.0, 170.1, 170.3; ms: *m/z* 526 (M⁺, 1), 339 (100), 282 (74), 265 (38), 209 (13), 153 (19), 143 (11), 97(11); Hrms Calcd. for C₂₇H₄₆N₂O₈: [M]⁺ = 526.3254. Found: 526.3260.

2-[1-(1,1-Diethoxycarbonyl-3-butenyl)-2,2-dimethylpropyl]-1-methyl-1*H*-imidazole (**2f**) and 5-(1,1-diethoxycarbonyl-3-butenyl)-1-methyl-2-(2,2-dimethylpropyl)-1*H*-imidazole (**3f**).

These compounds were prepared in a similar manner as that used for the preparation of **2a**, **3a** and **4a** except for the use of diethyl allylmalonate instead of diethyl methylmalonate. Title compounds were separated by column chromatography (AcOEt/*n*-hexane = 1/1).

Compound **2f** was obtained in 6 % yield (20 mg) as a yellow viscous oil; ir (CHCl₃): 2943, 1722 cm⁻¹; ¹H nmr (270 MHz, CDCl₃): δ 1.04 (9H, s), 1.17 (3H, t, J = 7.1 Hz), 1.33 (3H, t, J = 7.1 Hz), 2.64 (2H, d, J = 7.1 Hz), 3.63 (3H, s), 3.71 (1H, s), 3.88-4.39 (4H, m), 4.89-4.95 (2H, m), 5.70-5.86 (1H, m), 6.75 (1H, d, J = 1.1 Hz), 7.01 (1H, d, J = 1.1 Hz); ¹³C nmr (68 MHz, CDCl₃): δ 13.9, 14.0, 29.8 x 3, 33.4, 37.4, 39.3, 49.4, 61.10, 61.14, 61.5, 117.2, 119.4, 126.9, 134.7, 146.6, 170.2, 171.0; ms: *m/z* 350 (M⁺, 1), 253 (18), 221 (100), 207 (33), 175 (20), 147 (17), 137 (33); Hrms Calcd. for C₁₉H₃₀N₂O₄: [M]⁺ = 350.2205. Found: 350.2210.

Compound **3f** was obtained in 89 % yield (310 mg) as a yellow viscous oil; ir (CHCl₃): 2936, 1724, 1467 cm⁻¹; ¹H nmr (400 MHz, CDCl₃): δ 1.00 (9H, s), 1.25 (6H, t, J = 7.1 Hz), 2.59 (2H, s), 3.10 (2H, d, J = 7.3 Hz), 3.49 (3H, s), 4.21 (4H, q, J = 7.1 Hz), 5.07-5.16 (2H, m), 5.87-5.97 (1H, m), 6.99 (1H, s); ¹³C nmr (100 MHz, CDCl₃): δ 14.0 x 2, 29.6 x 3, 32.6, 32.9, 39.4, 40.0, 57.5, 61.9 x 2, 118.9, 127.4 x 2, 133.0, 149.1, 169.1 x 2; ms: *m/z* 350 (M⁺, 22), 309 (84), 294 (100), 277 (67), 253 (67), 221 (27), 207 (77), 122 (23), 57 (27); Hrms Calcd. for C₁₉H₃₀N₂O₄: [M]⁺ = 350.2205. Found: 350.2210.

2-[1-(1,1-Diethoxycarbonyl-1-phenylmethyl)-2,2-dimethylpropyl]-1-methyl-1*H*-imidazole (**2g**), 5-(1,1-Diethoxycarbonyl-1-phenylmethyl)-1-methyl-2-(2,2-dimethylpropyl)-1*H*-imidazole (**3g**) and 2-{1-[4-(1,1-Diethoxycarbonylmethyl)phenyl]-2,2-dimethylpropyl}-1-methyl-1*H*-imidazole (**5**).

These compounds were prepared in a similar manner as that used for preparation of **2a**, **3a** and **4a** except for the use of diethyl phenylmalonate instead of diethyl methylmalonate. Title compounds were separated by column chromatography (AcOEt/*n*-hexane = 1/1).

Compound **2g** was obtained in 3 % yield (10 mg) as colorless needles (*n*-hexane), mp 113-115 °C; ir (CHCl₃): 2942, 1730, 1479 cm⁻¹; ¹H nmr (400 MHz, CDCl₃): δ 0.72 (9H, s), 1.04 (3H, t, J = 7.1 Hz), 1.22 (3H, t, J = 7.1 Hz), 3.59 (3H, s), 3.74-3.82 (1H, m), 3.92-4.00 (1H, m), 4.20 (1H, s), 4.21-4.35 (2H, m), 6.70 (1H, d, J = 1.1 Hz), 6.98 (1H, d, J = 1.1 Hz), 7.25-7.34 (3H, m), 7.88 (2H, d, J = 7.0 Hz); ¹³C nmr (100 MHz, CDCl₃): δ 13.7 x 2, 29.2 x 3, 33.3, 37.2, 48.1, 61.4, 61.6, 67.1, 119.6, 126.6, 126.7 x 2, 127.3 x 2, 131.9, 135.1, 147.7, 169.2, 171.7; ms: *m/z* 386 (M⁺, 16), 329 (100), 257 (28), 151 (71); Hrms Calcd. for C₂₂H₃₀N₂O₄: [M]⁺ = 386.2205. Found: 386.2204.

Anal. Calcd. for $C_{22}H_{30}N_2O_4$: C, 68.37; H, 7.82; N, 7.25. Found: C, 68.12; H, 7.82; N, 7.38.

Compound **3g** was obtained in 11 % yield (42 mg) as a yellow viscous oil; ir (CHCl₃): 2937, 1724, 1464 cm⁻¹; ¹H nmr (400 MHz, CDCl₃): δ 0.99 (9H, s), 1.28 (6H, t, J = 7.1 Hz), 2.56 (2H, s), 3.01 (3H, s), 4.31 (4H, q, J = 7.1 Hz), 7.11 (1H, s), 7.26-7.36 (5H, m); ¹³C nmr (100 MHz, CDCl₃): δ 13.9 x 2, 29.6 x 3, 32.5, 32.9, 39.9, 62.4 x 2, 63.3, 127.4, 128.1, 128.3 x 2, 128.6 x 2, 128.9, 136.1, 149.0, 168.4 x 2; ms: *m/z* 386 (M⁺, 14), 330 (100), 313 (50), 257 (51), 211 (30), 183 (12), 115 (13), 57 (16); Hrms Calcd. for $C_{22}H_{30}N_2O_4$: [M]⁺ = 386.2205. Found: 386.2199.

Compound **5** was obtained in 44 % yield (171 mg) as colorless plates (*n*-hexane), mp 118-120 °C; ir (CHCl₃): 2936, 1722 cm⁻¹; ¹H nmr (400 MHz, CDCl₃): δ 1.06 (9H, s), 1.24 (3H, t, J = 7.1 Hz), 1.25 (3H, t, J = 7.1 Hz), 3.46 (3H, s), 3.77 (1H, s), 4.13-4.26 (4H, m), 4.58 (1H, s), 6.70 (1H, d, J = 1.3 Hz), 7.00 (1H, d, J = 1.1 Hz), 7.30 (2H, d, J = 8.2 Hz), 7.46 (2H, d, J = 8.2 Hz); ¹³C nmr (100 MHz, CDCl₃): δ 13.9 x 2, 28.1 x 3, 32.8, 35.6, 52.5, 57.4, 61.6 x 2, 119.7, 126.5, 128.4 x 2, 130.5 x 2, 130.8, 138.7, 148.3, 168.1 x 2; ms: *m/z* 386 (M⁺, 2), 330 (100), 257 (52), 227 (9), 183 (13), 170 (9); Hrms Calcd. for $C_{22}H_{30}N_2O_4$: [M]⁺ = 386.2205. Found: 386.2197.

Anal. Calcd. for $C_{22}H_{30}N_2O_4$: C, 68.37; H, 7.82; N, 7.25. Found: C, 68.17; H, 7.76; N, 7.33.

1-Methyl-2-(1,2,2-trimethylpropyl)-1*H*-imidazole (**7a**) and 1,5-Dimethyl-2-(2,2-dimethylpropyl)-1*H*-imidazole (**8a**).

These compounds were prepared in a similar manner as that used for preparation of **2a**, **3a** and **4a** except for use of a methylmagnesium bromide (0.9 M solution in THF) instead of NaH and diethyl methylmalonate. Title compounds were separated by column chromatography (CHCl₃/MeOH = 10/1).

Compound **7a** was obtained in 21 % yield (35 mg) as a yellow viscous oil; ir (CHCl₃): 2932, 1473, 1359 cm⁻¹; ¹H nmr (400 MHz, CDCl₃): δ 0.95 (9H, s), 1.29 (3H, d, J = 7.1 Hz), 2.69 (1H, q, J = 7.1 Hz), 3.59 (3H, s), 6.73 (1H, d, J = 1.1 Hz), 6.97 (1H, d, J = 1.1 Hz); ¹³C nmr (100 MHz, CDCl₃): δ 15.2, 27.5 x 3, 33.0, 35.0, 40.0, 119.4, 126.7, 151.3; ms: *m/z* 166 (M⁺, 8), 151 (8), 110 (100), 96 (14), 71 (16); Hrms Calcd. for $C_{10}H_{18}N_2$: [M]⁺ = 166.1470. Found: 166.1465.

Compound **8a** was obtained in 37 % yield (61 mg) as a yellow viscous oil; ir (CHCl₃): 2927, 1464, 1435, 1400, 1361 cm⁻¹; ¹H nmr (400 MHz, CDCl₃): δ 1.00 (9H, s), 2.17 (3H, d, J = 0.9 Hz), 2.57 (2H, s), 3.43 (3H, s), 6.71 (1H, d, J = 1.1 Hz); ¹³C nmr (100 MHz, CDCl₃): δ 10.0, 29.6 x 3, 30.6, 32.8, 40.2, 124.4, 127.0, 146.3; ms: *m/z* 166 (M⁺, 16), 151 (7), 110 (100), 95 (10), 71 (7); Hrms Calcd. for $C_{10}H_{18}N_2$: [M]⁺ = 166.1470. Found: 166.1474.

1-Methyl-2-[1-(1,1-dimethylethyl)pentyl]-1*H*-imidazole (**7b**) and 5-Butyl-1-methyl-2-(2,2-dimethylpropyl)-1*H*-imidazole (**8b**).

These compounds were prepared in a similar manner as that used for preparation of **2a**, **3a** and **4a** except for use of a *n*-butylmagnesium chloride (0.9 M solution in THF) instead of NaH and diethyl methylmalonate. Title compounds were separated by column chromatography (CHCl₃/MeOH = 20/1).

Compound **7b** was obtained in 13 % yield (26 mg) as a yellow viscous oil; ir (CHCl₃): 2927, 1474, 1361 cm⁻¹; ¹H nmr (400 MHz, CDCl₃): δ 0.82 (3H, t, J = 7.3 Hz), 0.89-0.97 (2H, m), 0.94 (9H, s), 1.20-1.30 (2H, m), 1.65-1.73 (1H, m), 1.87-1.97 (1H, m), 2.45 (1H, dd, J = 11.8, 2.8 Hz), 3.57 (3H, s), 6.74 (1H, d, J = 1.1

Hz), 7.00 (1H, d, J = 1.3 Hz); ¹³C nmr (100 MHz, CDCl₃): δ 14.0, 22.7, 27.9 x 3, 29.7, 30.8, 32.8, 35.1, 46.8, 119.3, 126.8, 150.6; ms: *m/z* 208 (M⁺, 18), 189 (46), 151 (100), 137 (22), 123 (17), 109 (91), 96 (34); Hrms Calcd. for $C_{13}H_{24}N_2$: [M]⁺ = 208.1939. Found: 208.1938.

Compound **8b** was obtained in 6 % yield (13 mg) as a yellow viscous oil; ir (CHCl₃): 2929, 2920, 1463, 1351 cm⁻¹; ¹H nmr (400 MHz, CDCl₃): δ 0.95 (3H, t, J = 7.4 Hz), 1.00 (9H, s), 1.37-1.46 (2H, m), 1.56-1.64 (2H, m), 2.49 (2H, dt, J = 0.7, 7.7 Hz), 2.56 (2H, s), 3.43 (3H, s), 6.70 (1H, s); ¹³C nmr (100 MHz, CDCl₃): δ 13.9, 22.4, 24.5, 29.7 x 3, 30.2, 30.6, 32.8, 40.1, 123.8, 131.7, 146.3; ms: *m/z* 208 (M⁺, 11), 193 (14), 151 (100), 137 (19), 123 (12), 110 (32), 58 (14); Hrms Calcd. for $C_{13}H_{24}N_2$: [M]⁺ = 208.1939. Found: 208.1934.

2-(1-Ethyl-2,2-dimethylpropyl)-1-methyl-1*H*-imidazole (**7c**).

This was prepared in a similar manner as that used for the preparation of **2a**, **3a** and **4a** except for use of a diethyl zinc (1.0 M solution in *n*-hexane) instead of NaH and diethyl methylmalonate. The title compound was purified by column chromatography (Et₂O) and obtained as a pale yellow viscous oil, 49 mg (27 %); ir (CHCl₃): 2933, 1475, 1361 cm⁻¹; ¹H nmr (400 MHz, CDCl₃): δ 0.65 (3H, t, J = 7.2 Hz), 0.94 (9H, s), 1.73-1.96 (2H, m), 2.39 (1H, dd, J = 11.7, 3.1 Hz), 3.58 (3H, s), 6.74 (1H, d, J = 1.1 Hz), 7.00 (1H, d, J = 1.3 Hz); ¹³C nmr (100 MHz, CDCl₃): δ 13.0, 22.9, 27.9 x 3, 32.8, 35.0, 48.8, 119.3, 126.8, 150.3; ms: *m/z* 180 (M⁺, 8), 165 (9), 151 (8), 123 (97), 109 (100), 96 (20), 58 (13); Hrms Calcd. for $C_{11}H_{20}N_2$: [M]⁺ = 180.1626. Found: 180.1634.

Acknowledgements.

We are grateful for financial supports in part by the Frontier Research Program from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan, and a Grant-In-Aid for the promotion of the advancement of education and research in graduate schools in subsidies for ordinary expenses of private schools from the Promotion and Mutual Aid Corporation for Private Schools.

REFERENCES AND NOTES

- [1] M. R. Grimmett, in *Advances in Heterocyclic Chemistry*, Vol 27, A. R. Katritzky and A. Boulton, eds, Academic Press, New York, 1980, pp 241.
- [2] B. Iddon and R. I. Ngochindo, *Heterocycles*, **38**, 2487 (1994).
- [3a] S. Ohta, T. Yamamoto, I. Kawasaki, M. Yamashita, H. Katsuma, R. Nasako, K. Kobayashi and K. Ogawa, *Chem. Pharm. Bull.*, **40**, 2681 (1992); [b] S. Ohta, T. Yamamoto, I. Kawasaki, M. Yamashita, Y. Nagashima and T. Yoshikawa, *Chem. Pharm. Bull.*, **42**, 821 (1994).
- [4a] S. Nakamura, N. Tsuno, M. Yamashita, I. Kawasaki, S. Ohta and Y. Ohishi, *J. Chem. Soc., Perkin Trans 1*, 429 (2001); [b] I. Kawasaki, S. Nakamura, S. Yanagitani, A. Kakuno, M. Yamashita and S. Ohta, *J. Chem. Soc., Perkin Trans 1*, 3095 (2001).
- [5a] I. Kawasaki, N. Taguchi, T. Yamamoto, M. Yamashita and S. Ohta, *Tetrahedron Lett.*, **36**, 8251 (1995); [b] I. Kawasaki, M. Yamashita and S. Ohta, *Chem. Pharm. Bull.*, **44**, 1831 (1996); [c] I. Kawasaki, N. Taguchi, M. Yamashita and S. Ohta, *Chem. Pharm. Bull.*, **45**, 1393 (1997); [d] I. Kawasaki, H. Katsuma, Y. Nakayama, M. Yamashita and S. Ohta, *Heterocycles*, **48**, 1887 (1998); [e] S. Ohta, N. Tsuno, K. Maeda, S. Nakamura, N. Taguchi, M. Yamashita and I. Kawasaki, *Tetrahedron Lett.*, **41**, 4623 (2000); [f] I. Kawasaki, N. Sakaguchi, N. Fukushima, N. Fujioka, F. Nikaido, M. Yamashita and S. Ohta, *Tetrahedron Lett.*, **43**,

4377 (2002); [g] S. Nakamura, I. Kawasaki, M. Kunimura, M. Matsui, Y. Noma, M. Yamashita and S. Ohta, *J. Chem. Soc., Perkin Trans 1*, 1061 (2002); [h] S. Nakamura, I. Kawasaki, M. Yamashita and S. Ohta, *Heterocycles*, **60**, 583 (2003).

[6a] M. Begtrup, *Angew. Chem., Int. Ed.*, **13**, 347 (1974); [b] M. Grimmett in *Comprehensive Organic Chemistry*, Vol **4**, P. G. Sammers, ed, Pergamon, Oxford, 1979, pp 357; [c] T. Itoh, H. Hasegawa, K. Nagata and A. Ohsawa, *J. Org. Chem.*, **59**, 1319 (1994); [d] T. Itoh, M. Miyazaki, H. Hasegawa, K. Nagata and A. Ohsawa, *Chem. Commun.*, 1217 (1996).

[7a] E. S. Hand and W. W. Paudler, *J. Org. Chem.*, **43**, 2900 (1978); [b] R. Achour, E. M. Essassi and R. Zniber, *Tetrahedron Lett.*, **29**, 195 (1988); [c] H. Yamanaka, S. Ohba and T. Sakamoto, *Heterocycles*, **31**, 1115 (1990); [d] K. Burger, E. Höß and K. Geith, *Synthesis*, 360 (1990); [e] R. P. Subrayan and P. G. Rasmussen, *Tetrahedron*, **51**, 6167 (1995); [f] G. Tennant, C. J. Wallis and G. W. Weaver, *J. Chem. Soc., Perkin Trans 1*, 629 (1999).

[8] A. Olofson, K. Yakushijin and D. A. Horne, *J. Org. Chem.*, **63**, 1248 (1998).

[9] T. Nakano, W. Rodríguez, S. Z. De Roche, J. M. Larrauri, C. Rivas and C. Pérez, *J. Heterocyclic Chem.*, **17**, 1777 (1980).

[10] S. Ohta, T. Osaki, S. Nishio, A. Furusawa, M. Yamashita and I. Kawasaki, *Tetrahedron Lett.*, **41**, 7503 (2000).

[11a] J. Suwinski and K. Swierczek, *Tetrahedron*, **57**, 1639 (2001); [b] J. F. Arens, *Bull. Soc. Chim. Fr.*, 3037 (1968).

[12a] R. I. Fryer, R. F. Lauer, E. J. Trybulski, S. Vitone, A. Walser and G. Zenchoff, *J. Heterocyclic Chem.*, **20**, 1605 (1983); [b] T. S. Manoharan and R. S. Brown, *J. Org. Chem.*, **54**, 1439 (1989); [c] Y. Njoya, N. Boufatah, A. Gellis, P. Rathelot, M. P. Crozet and P. Vanelle, *Heterocycles*, **57**, 1423 (2002).

[13] Representative examples: [a] O. A. Attanasi, L. De Crescentini, P. Filippone and F. Mantellini, *Synlett*, 554 (2001); [b] R. E. Banks, M. K. Besheesh, S. N. Mohialdin-Khaffaf and I. Sharif, *J. Chem. Soc., Perkin Trans 1*, 2069 (1996); [c] W. E. Barnette, *J. Am. Chem. Soc.*, **106**, 452 (1984); [d] T. S. Everett, S. T. Purrington and C. L. Bumgardner, *J. Org. Chem.*, **49**, 3702 (1984); [e] C. G. Kruse, A. Wijsman and A. Van der Gen, *J. Org. Chem.*, **44**, 1847 (1979).