ORIGINAL ARTICLE



3, 3,5 and 2,6 Expanded Aza-BODIPYs Via Palladium-Catalyzed Suzuki-Miyaura Cross-Coupling Reactions: Synthesis and Photophysical Properties

Halil Yılmaz¹ · Gökhan Sevinç² · Mustafa Hayvalı¹

Received: 9 September 2020 / Accepted: 30 October 2020 © Springer Science+Business Media, LLC, part of Springer Nature 2020

Abstract

Novel symmetrical aza-borondipyrromethene (aza-BODIPY) compounds bearing 4-methoxyphenyl, 4-methoxybiphenyl, 2,4dimethoxybipheny, 4-bromophenyl and N,N-diphenyl-4-biphenylamine groups on the 3, 3,5 and 2,6 positions of aza-BODIPY core were synthesized via Suzuki-Miyaura coupling reactions while unsymmetrical analogues were obtained from the starting mono Br-substituted aza-BODIPY material which was obtained from nitrosolated pyrrole derivative. The characterizations were performed by means of ¹H-NMR, ¹³C-NMR, FTIR and HRMS-TOF-ESI techniques. The spectral properties of the aza-BODIPY derivatives were investigated using absorption and fluorescence spectroscopy. The novel compounds with extended conjugation have broadband absorption in near infrared region and show significant shifts on their absorption and fluorescence spectra compared to unsubstituted analogues. The highest bathochromic shifts were observed π -extended and strong electron donating groups at 3,5 positions of the aza-BODIPY scaffold. Depend on substitution positions of attached groups to the indacene core, the fluorescence quantum yields of chromophores were determined to be drastic changes. The singlet oxygen generation capability of the compounds were evaluated and 2,6-bromine substituted compounds AA1 and CC1 showed high singlet oxygen quantum yields (71% and 74%, respectively). Enhanced photophysical properties such as intense absorption, extended conjugation and singlet oxygen production make the investigated aza-BODIPY's promising candidates for photodynamic therapy applications and organic photovoltaic cells in NIR region.

Keywords Aza-BODIPY · Unsymmetrical aza-BODIPY · Azadipyrromethenes · Suzuki-Miyaura coupling · NIR region dyes

Introduction

Although, the first reaction of tetraarylazadipyrromethenes with boron (Aza-BODIPY) occurred in the early 1990s [1] these dyes have been recognized by the studies of the O'Shea group since 2002 [2].Aza-BODIPY dyes are azaanalogous of difluorobora-diaza-s-indacenes (BODIPY) and known as attractive fluorophores because of their

Supplementary Information The online version contains supplementary material available at https://doi.org/10.1007/s10895-020-02646-4.

advantageous features. Sharp absorption and fluorescence bands, large molar extinction coefficients, high fluorescence quantum yields and high photostability [3-5] all contribute to the appeal of these interesting compounds. Aza-BODIPY compounds are well-known NIR dyes because of their large NIR region absorption (700–1100 nm) [6]. This property depends on a large scale 1,3,5,7 aryl substituents. Particularly electron donating groups on 3,5-aryl substituents increase extinction coefficients and absorption maximum, however on 1,7-aryl substituted electron donating groups show minor effects. Furthermore, 2,6-substituted (β-pyrrole) groups may cause hypsochromic shifts on the absorption wavelength and quench fluorescence intensity of aza-BODIPY dyes [7]. Currently, aza-BODIPY dyes are used in diverse applications, such as photodynamic therapy (PDT) [8–10], NIR-emitting chemosensors for heavy metal determination [11], in telecommunication applications with strong two-photon absorption (TPA) properties [12–14], as a chromophore group in pH sensors [15, 16], photosensitized singlet oxygen generation

Mustafa Hayvalı Mustafa.Hayvali@science.ankara.edu.tr

¹ Department of Chemistry, Faculty of Science, Ankara University, Anadolu, 06100 Ankara, Turkey

² Department of Chemistry, Faculty of Science and Literature, Bilecik Şeyh Edebali University, 11230 Bilecik, Turkey

Photodynamic therapy (PDT) is an effective and applicable treatment for various types of cancers and also it has been researched for other diseases. Photosensitizer, light and oxygen are essential components for PDT treatments. Photosensitizers are specific chromophores that produces biological cytotoxic singlet oxygen $({}^{1}O_{2})$ from the ${}^{3}O_{2}$ using the light. This makes the chromophores prominent investigating area for the improvement of PDT applications. So that, there have been many research papers for dye molecules such as porphyrin derivatives [23, 24]. Porphyrins were the first photosensitizers for using clinic treatment [25], which has several disadvantageous. First, absorption and emission wavelength aren't tunable with bonding side groups. Second, photons do not penetrate tissue beyond a few millimeters due to short absorption wavelength and low absorption leading to skin photosensitivity and low excretion rate [26-28]. Because of these disadvantages it has been designed new generation dyes that have intense absorption in near IR region like BODIPY [29-31] and aza-BODIPY [17, 18] derivatives.

Herein, we present new red/NIR absorbing aza-BODIPY series which are decorated with 4-methoxyphenyl, 2,4dimethoxy phenyl and triphenylamine (TFA) precursors via Palladium-catalyzed Suzuki-Miyaura cross-coupling reactions. Due to mild reaction conditions and commercial availability of starting boronic acid reagents, we preferred the Suzuki-Miyaura coupling. In addition diverse boronic acids in commercial catalogs are safer than the other organometallic reagents [32]. The Suzuki-Miyaura coupling has been successfully employed on 3, 3,5 and 2,6 positions of aza-BODIPY core. 3 and 3,5 substituted aza-BODIPY dyes showed absorption and emission in the NIR region. On the other hand 2,6 substituted aza-BODIPY derivatives showed quenching fluorescence intensity and hypsochromic shifts compared to unsubstituted form. Electron donating groups and extended conjugations dramatically have changed photophysical properties of aza-BODIPY core depending on substitution position. As a consequence, these compounds seem to display enhanced photophysical properties, which will be helpful for developing novel versatile apparatus for organic solar cells, cellular imaging, PDT and two photon absorption applications in longer wavelengths in chemical and biological fields.

Experimental

Materials and Characterization

Compounds 1a, 2a, 3a, 1b, 2b, A, B and B1 were prepared according to a literature procedures [14, 33–35]. All solvents were purchased from common commercial sources. The melting points of the synthesized compounds were determined in

open capillares using Barnstead Electrothermal IA9100. Fluorimetric measurements were applied on Perkin Elmer LS55 Spectrometer and electronic excitation spectra were recorded on Shimadzu UV1800 spectrophotometer. Mass spectral analyses were performed on an Agilent 6224 HRMS spectrometer. ¹H-NMR spectra were recorded on VARIAN Mercury 400 MHz spectrometer. ¹H NMR chemical shifts (δ) are given in ppm down fieldfrom Me₄Si, determined by chloroform (δ = 7.26 ppm. ¹³C NMR spectra were recorded on VARIAN Mercury 100 MHz spectrometer. ¹³C NMR chemical shifts (δ) are reported in ppm with the internal CDCl₃ δ 77.0 ppm as standard. Perkin–Elmer 100 spectrometer (equippedwith ATR unit) was used for FT-IR spectra of the compounds in the range 650–4000 cm⁻¹.

Determination of Fluorescence Quantum Yields (Φ_F)

In order to determine the fluorescence quantum yields of the dyes, comparative method (Eq. 1) [36] was applied.

$$\Phi_F = \Phi_F(\text{Std}) \frac{\text{F.A}_{\text{Std}}.n^2}{F_{\text{Std}}.A.n_{\text{Std}}^2}$$
(1)

Where Φ_F (Std) is the fluorescence quantum yield of BF₂ Chelate of (3,5-Diphenyl-1H-pyrrol-2-yl)(3,5diphenylpyrrol-2-ylidene) amine (Φ_F = 0.34 in chloroform) [7]. F and F_{Std} denote the areas under the fluorescence emission curves of samples and the standard, respectively. A and A_{Std} are the respective absorbances of the samples and standard compound at the excitation wavelengths, respectively. The refractive indices, n and n_{Std} of the solvents were applied. The concentration of the dilute solutions at the excitation wavelengths fixed at 5×10^{-6} M in CHCl₃. All spectra were recorded at 25 °C on non-deaerated samples.

The Parameters for Singlet Oxygen Quantum Yields (Φ_{Δ})

To study the singlet oxygen production properties of compounds, experiments were carried out in CH₂Cl₂ with 1,3diphenyl isobenzofuran (DPBF) as a chemical singlet oxygen trap [37]. The absorbance of DPBF was tuned around 1.0 at 414 nm and the absorbance of the sensitizers was tuned between from 0.2 to 0.3 in CH₂Cl₂. A solution of the aza-BODIPY derivatives and DPBF was irradiated with a 630 nm Perkin Elmer L5S55 fluorescent light and the decrease in the absorption band of DPBF was monitored at the intervals of 2 s. Singlet oxygen quantum yields (Φ_{Δ}) were calculated using the standard, methylene blue (MB) (Φ_{Δ} = 0.57 in CH₂Cl₂), by plotting the Δ OD of DPBF against the irradiation time. The quantum yield Φ_{Δ} was calculated by the following equation [37]:

$$\Phi_{\Delta sam} = \Phi_{\Delta std} \left(\frac{m_{sam}}{m_{std}} \right) \left(\frac{F_{std}}{F_{sam}} \right)$$
(2)

Where sam and std. designate the aza-BODIPY photosensitizer and methylene blue (MB), respectively. m is the slope of difference in change in absorbance of DPBF (414 nm) with the irradiation time, F is the absorption correction factor, which is given by $F = 1-10^{-OD}$. The photosensitizers were irradiated with monochromatic light at the peak absorption wavelength for 2 s. Absorbance was measured for several times after each irradiation. The slope of the graph of absorbance maxima of DPBF at 414 nm versus the photoirradiation time for each photosensitizer was calculated.

Synthesis

Synthesis of 2-(4-Bromophenyl)-4-(4-Methoxyphenyl) -1H-Pyrrole (3b)

1-(4- bromophenyl)-4-nitro-3-(4- methoxyphenyl)butan-1one (2b) (7.02 g, 18.6 mmol) was dissolved in 180 mL MeOH/THF (1:1) mixture at room temperature. A stirred solution was treated with KOH (5.24 g, 93.45 mmol). After 1 h the mixture was added drop wise to a solution of 200 mL %33 H₂SO₄/MeOH (v/v) at 0 °C and the solution was stirred for further 1 h at room temperature. Water (250 mL) and ice (250 mL) were added and the mixture was neutralized with aqueous 5 M NaOH and extracted with $CHCl_3$ (100 mL \times 3). The solvent was evaporated in vacuo and obtained oil was held in vacuum drying over the night. The oil was treated with acetic acid (174 mL) and NH₄OAc (14.41 g, 0.19 mol) and the solution was heated at 100 °C for 1 h. The reaction mixture was cooled to room temperature, ice (340 mL) was added, and the mixture carefully neutralized with aqueous 5 M NaOH. The resulting solid was collected by filtration, washed with distilled water, dried and crystallized from CH₃Cl, resulting the product as a light blue solid (3.45 g, %56), mp228–231 °C (from chloroform). ¹HNMR (400 MHz, DMSO) δ 11.42 (s, 1H), 7.61 (d, J= 8.0 Hz, 2H), 7.50 (d, 2H), 7.23 (s, 1H), 6.89 (s, 1H), 6.88 (d, *J* = 8.0 Hz, 2H), 3.73 (s, 3H). ¹³CNMR (100 MHz, DMSO) δ 157.11, 131.93, 131.39, 130.76, 128.18, 125.49, 125.19, 124.74, 118.02, 116.16, 113.93, 103.62, 54.93. IR (ATR, cm⁻¹) 3437.1, 3014.7, 2835.4, 1242.1. HRMS (TOF-ESI): Calcd ($C_{17}H_{14}BrNO$) [M + H]⁺: m/z = 328.03371; found, m/z = 328.03328.

Synthesis of 3,5-Bis(4-Methoxyphenyl)-2-Nitroso-1H-Pyrrole (4a)

2,4-bis(4-methoxyphenyl)-1H-pyrrole (3a) (1.95 g, 7.00 mmol) was dissolved in EtOH (70 mL). To a stirred

solution was added concentrated HCl(1.40 mL), and followed by a drop wise addition of aqueous NaNO₂ (560 mg, 8.05 mmol, in 14 mL of H₂O). The reaction mixture was stirred for 30 min and cooled to 0 °C, and another portion of concentrated HCl (7.00 mL) was added. The solution was allowed to stir for 1 h, and the resulting red solid was collected by filtration and washed with Et₂O. The solid was dissolved in minimal EtOH, an excess of aqueous NaOAc and ice was added, and the solution was stirred for 1 h. The resulting solid was collected by filtration and purified by chromatography on silica (1:1, EtOAc/hex), resulting the product as a green solid (1.68 g, %78), mp. 186–187 °C.¹HNMR (400 MHz, CDCl₃) δ 8.11 (d, J = 8.8 Hz, 2H), 7.83 (d, J = 8.8 Hz, 2H), 7.07 (s, 1H), 7.01 (d, J = 8.8 Hz, 2H), 6.99 (d, J = 8.4 Hz, 2H), 3.89 (s, 3H), 3.87 (s, 3H). ¹³CNMR (100 MHz, CDCl₃) δ 162.61, 161.08, 130.83, 129.08, 124.39, 122.17, 114.85, 114.32, 113.79, 55.53, 55.38. IR (ATR, cm⁻¹) 3228.8, 3012.8, 2841.1, 1600.9, 1246.0. HRMS (TOF-ESI): Calcd (C18H16N2O3) $[M + H]^+$: m/z = 309.12394; found, m/z = 309.12540.

Synthesis of BF₂ Chelate of [3,5-Bis(4-Methoxyphenyl) -1H-Pyrrol-2-Yl]-3,5-Bis(4- Methoxyphenyl)Pyrrol-2-Ylidene] Amine (A1)

A solution of [3,5-bis(4-methoxyphenyl)-1H-pyrrol-2-yl]-3,5-bis(4-methoxyphenyl)pyrrol-2-ylidene]amine (A) (2.00 g, 3.90 mmol) in dry CH₂Cl₂ (300 mL) was treated with N,N-diisopropylethylamine (6.08 mL, 39.0 mmol) and the mixture was stirred for 10 min at room temperature. Then, boron trifluoride diethyl etherate (7.22 mL, 58.5 mmol) was added and the reaction mixture was stirred at room temperature for 24 h. The mixture was washed with water, and the organic layer dried over sodium sulfate. Removal of solvent gave a residue, which was purified by column chromatography on silica eluting with hexane/ EtOAc (2:1, V/V) resulting the product as a dark blue solid (1.84 g, 76%), mp 265–266 °C.¹HNMR (400 MHz, CDCl₃) δ 8.05 (t, J = 8.2 Hz, 4H), 6.99 (d, J = 8.0 Hz, 2H), 6.98 (d, J = 8.0 Hz, 2H), 6.91 (s, 1H), 3.88 (s, 3H), 3.87 (s, 3H) ¹³CNMR (100 MHz, CDCl₃) δ 161.68, 160.87, 157.78, 145.14, 144.65, 142.74, 131.49, 131.45, 131.40, 130.73, 125.49, 124.38, 117.09, 117.08, 114.13, 114.08, 55.40, 55.38. IR (ATR,m⁻¹) 3005.1, 2837.3, 1597.1, 1384.9, 1251.8. HRMS (TOF-ESI): Calcd $(C_{36}H_{30}BF_2N_3O_4)$ $[M]^+$: m/z = 617.22976; found, m/z = 617.23289.

Synthesis of BF₂ Chelate of [4-Bromo-3-(4-Methoxyphenyl) -5-(4- Methoxyphenyl)-1H-Pyrrol-2-Yl] -[4-Bromo-3-(4-Methoxyphenyl)-5-(4-Methoxyphenyl) Pyrrol-2-Ylidene]Amine (AA1)

A solution of BF_2 chelate of [3,5-bis(4-methoxyphenyl)-1H-pyrrol-2-yl]-3,5-bis(4-methoxyphenyl)pyrrol-2-

vlidene]amine (A1)(1.23 g, 2.00 mmol) in dry CH₂Cl₂ (100 mL) was added to N-bromosuccinimide (NBS) (0.78 g, 4.40 mmol), and the mixture was stirred at room temperature for 24 h. After the reaction was completed, the solvent was evaporated in vacuo. Afterwards, the residue was washed with water, and dried over sodium sulfate. The residual was purified by column chromatography on silica eluting with hexane/EtOAc (2:1) which gave the product as a dark brown solid (1.31 g, 84%), mp >350 °C.¹HNMR (400 MHz, CDCl₃) δ 7.90 (d, J = 8.0 Hz, 2H), 7.76 (d, J=8.8 Hz, 2H), 6.98 (d, J=9.2 Hz, 2H), 6.98 (d, J = 9.2 Hz, 2H), 3.88 (s, 3H), 3.85 (s, 3H). ¹³CNMR (100 MHz, CDCl₃) δ 161.55, 160.80, 157.14, 144.01, 141.99, 141.98, 132.40, 132.36, 131.38, 123.51, 121.98, 113.59, 113.52, 55.38, 55.27. IR (ATR, cm⁻¹) 3003.2, 2835.4, 1599.0, 1379.1, 1249.9. HRMS (TOF-ESI): Calcd $(C_{36}H_{28}BBr_2F_2N_3O_4)$ $[M + H]^+$: m/z = 776.05657; found, m/z = 776.06088.

Synthesis of BF₂ Chelate of [4-Methoxyphenyl-3-(4-Methoxyphenyl) -5-(4-Methoxyphenyl)-1H-Pyrrol-2-Yl] -[4-Methoxyphenyl-3-(4-Methoxyphenyl) -5-(4-Methoxyphenyl)Pyrrol-2-Ylidene]Amine (A2)

A mixture of BF₂ chelate of [4-bromo-3-(4methoxyphenyl)-5-(4-methoxyphenyl)-1H-pyrrol-2-yl]-[4bromo-3-(4-methoxyphenyl)-5-(4-methoxyphenyl)pyrrol-2-ylidene]amine (AA1) (0.23 g, 0.30 mmol), 4methoxyphenylboronic acid (105 mg, 0.69 mmol), Pd(PPh₃)₄ (15.6 mg, 0.0014 mmol) and K₂CO₃ (0.28 g, 2.00 mmol) were added to a round bottomed flask, and then degassed argon for 10 min. The mixture of toluene (4 mL), ethanol (2 mL) and water (2 mL) were injected under argon atmosphere. The solution was stirred at 85 °C for 24 h. After cooling to room temperature, the mixture was washed with water and extracted with CH_2Cl_2 (75 mL \times 2). The solvent was evaporated under reduced pressure. Afterwards, the residual was purified by column chromatography on silica eluting with toluene, which gave the product as a purple solid (82 mg, 33%) mp > 350 °C.¹HNMR (400 MHz, CDCl₃) δ 7.46 (d, J = 8.8 Hz, 2H), 7.41 (d, J = 8.4 Hz, 2H), 6.90 (d, J = 8.8 Hz, 2H), 6.79 (d, J = 8.8 Hz, 2H), 6.77 (d, J =9.2 Hz, 2H1), 6.74 (d, J=8.4 Hz, 2H), 3.81 (s, 3H), 3.78 (s, 6H); ¹³CNMR (100 MHz, CDCl₃) δ 160.57, 159.73, 158.61, 157.86, 155.85, 155.81, 145.13, 139.93, 138.46, 132.44, 132.34, 131.78, 125.76, 124.72, 123.38, 113.25, 113.22, 55.20, 55.11. IR (ATR, cm⁻¹) 3034.0, 2835.4, 1599.0, 1373.3, 1247.9. HRMS (TOF-ESI): Calcd $(C_{50}H_{42}BF_2N_3O_6)$ $[M + H]^+$: m/z = 830.32133; found, m/z = 830.31900.

Synthesis of BF₂ Chelate of [4-N,N-Diphenylaminophenyl-3-(4-Methoxyphenyl) -5-(4-Methoxyphenyl)-1H-Pyrrol-2-Yl] -[4-N,N-Diphenylaminophenyl-3-(4-Methoxyphenyl) -5-(4-Methoxyphenyl)Pyrrol-2-Ylidene]Amine (A4)

Compound A4 was prepared from BF₂ Chelate of [4-Bromo-3-(4-methoxyphenyl)-5-(4-methoxyphenyl)-1H-pyrrol-2 $y_{1} = [4 - bromo - 3 - (4 - methoxypheny_{1}) - 5 - (4 - methoxypheny_$ methoxyphenyl)pyrrol-2-ylidene]amine (AA1) (0.19 g, 0.25 mmol), 4-(diphenylamino)phenylboronic acid (173.5 mg, 0.60 mmol), Pd(PPh₃)₄ (13 mg, 0.0012 mmol) and K₂CO₃ (0.23 g, 1.66 mmol) according to the method for A2. The residual was purified by column chromatography on silica eluting with toluene which gave the product as a purple solid (53 mg, 19%) mp > 350 °C. ¹HNMR (400 MHz, CDCl₃) δ 7.53 (d, J = 8.8 Hz, 2H), 7.46 (d, J = 8.8 Hz, 2H), 7.25 (t, J = 8.0 Hz, 4H), 7.07 (d, J = 7.2 Hz, 2H), 7.01 (t, J = 7.4 Hz, 2H), 6.89 (d, J = 8.8 Hz, 2H), 6.84 (d, J = 8.4 Hz, 2H), 6.83 (d, J =8.4 Hz, 2H), 6.80 (d, J = 8.4 Hz, 2H), 3.84 (s, 3H), 3.81 (s, 3H); ¹³CNMR (100 MHz, CDCl₃) δ 160.68, 159.78, 157.89, 147.48, 146.66, 145.16, 139.78, 132.44, 129.21, 127.36, 124.76, 124.58, 124.44, 123.26, 123.04, 122.97, 113.23, 113.14, 55.24, 55.14. IR (ATR, cm⁻¹) 3032.1, 2835.4, 1599.0, 1375.2, 1251.8. HRMS (TOF-ESI): Calcd $(C_{72}H_{56}BF_2N_5O_4)$ [M] ⁺: m/z = 1103.43936; found, m/z = 1103.44170.

Synthesis of BF₂ Chelate of [5-(4-Methoxyphenyl) -3-(4-Methoxybiphenyl)-1H-Pyrrol-2-Yl] -[5-(4-Methoxyphenyl)-3-(4-Methoxybipheny) Pyrrol-2-Ylidine]Amine (B2)

A mixture of [3-(4-methoxyphenyl)-5-(4-bromophenyl)-1Hpyrrol-2-yl][3-(4-methoxyphenyl)-5-(4-bromophenyl)pyrrol-2-yliden]amine (B) (0.23 g, 0.30 mmol), 4methoxyphenylboronic acid (105 mg, 0.69 mmol), Pd(PPh₃)₄ (15.6 mg, 0.0014 mmol) and K₂CO₃ (0.28 g, 2.00 mmol) were added to a round bottomed flask, and then degassed argon for 10 min. The mixture of toluene (4 mL), ethanol (2 mL) and water (2 mL) were injected under argon atmosphere. The solution was stirred at 85 °C for 24 h. After cooling to room temperature, the mixture was washed with water and extracted with CH_2Cl_2 (75 mL \times 2). The solvent was evaporated under reduced pressure. The residual was carried into the next stage without further purification. The resulting solid was dissolved in dry CH₂Cl₂ (100 mL). Then, the reaction mixture was treated with N, Ndiisopropylethylamine (0.54 mL, 3.00 mmol) and the mixture was stirred for 10 min at room temperature. Afterwards, boron trifluoride diethyl etherate (0.45 mL, 4.50 mmol) was added and the reaction mixture was stirred at room temperature for 24 h. The mixture was washed with water, and the organic layer dried over sodium sulfate.

The solvent is removed in vacuo to obtain a residue, which was purified by column chromatography on silica eluting with toluene, resulting the product as a dark brown solid (61 mg, 26%), mp 279–280 °C.¹HNMR (400 MHz, CDCl₃) δ 8.14 (d, J = 8.8 Hz, 2H), 8.07 (d, J = 8.8 Hz, 2H), 7.67 (d, J = 8.4 Hz, 2H), 7.60 (d, J = 8.8 Hz, 2H), 7.00 (s, 1H), 7.00 (d, J = 8.8 Hz, 2H), 6.99 (d, J = 8.8 Hz, 2H), 3.90 (s, 3H), 3.86 (s, 6H). ¹³CNMR (100 MHz, CDCl₃) δ 160.84, 159.67, 158.22, 145.57, 143.28, 143.25, 142.85, 137.25, 132.62, 130.85, 130.14, 130.10, 130.05, 129.99, 128.23, 126.64, 125.41, 125.27, 117.50, 114.32, 114.16, 109.99, 55.43, 55.35. IR (ATR, cm⁻¹) 3111.2, 2833.4, 1597.1, 1383.0, 1244.1. HRMS (TOF-ESI): Calcd (C₄₈H₃₈BF₂N₃O₄) [M]⁺: m/z = 769.29236; found, m/z = 769.29487.

Synthesis of BF₂ Chelate of [5-(4-Methoxyphenyl) -3-(2,4-Dimethoxybiphenyl)-1H-Pyrrol-2-Yl] -[5-(4-Methoxyphenyl)-3-(2,4-Dimethoxybipheny) Pyrrol-2-Ylidine]Amine (B3)

B3 was prepared from [3-(4-methoxyphenyl)-5-(4bromophenyl)-1H-pyrrol-2-yl][3-(4-methoxyphenyl)-5-(4bromophenyl)pyrrol-2-yliden]amine (B) (0.23 g, 0.30 mmol), 2,4-dimethoxyphenylboronic acid (125.6 mg, 0.69 mmol), Pd(PPh₃)₄ (15.6 mg, 0.0014 mmol) and K₂CO₃ (0.28 g, 2.00 mmol) according to the method for B2.The obtained residue was purified by column chromatography on silica eluting with toluene which gave the product as a purple solid (85 mg, 34%), mp 273–274 °C.¹HNMR (400 MHz, CDCl₃) δ 8.11 (d, J = 8.4 Hz, 2H), 8.07 (d, J = 8.8 Hz, 2H), 7.65 (d, J = 8.4 Hz, 2H), 7.33 (d, J = 8.0 Hz, 1H), 7.01 (s, 1H), 7.01 (d, J = 8.8 Hz, 2H), 6.60-6.58 (m, 2H), 3.91 (s, 3H), 3.86 (s, 6H), 3.83 (s, 6H). ¹³CNMR (100 MHz, CDCl₃) δ 160.78, 160.73, 158.50, 157.71, 145.54, 143.11, 142.36, 140.88, 131.34, 130.82, 129.73, 129.50, 129.31, 129.27, 125.49, 122.65, 117.69, 114.14, 109.99, 104.81, 99.01, 55.53, 55.43. IR (ATR, cm⁻¹) 3117.0, 2831.5, 1600.9, 1379.1, 1259.5. HRMS (TOF-ESI): Calcd $(C_{50}H_{42}BF_2N_3O_6)$ [M + H]⁺: m/z = 830.32133; found, m/z = 830.32234.

Synthesis of BF₂ Chelate of [5-(4-Methoxyphenyl) -3-(4-N,N-Diphenylaminobiphenyl)-1H-Pyrrol-2-Yl] -[5-(4-Methoxyphenyl)-3-(4-N,N-Diphenylaminobiphenyl) Pyrrol-2-Ylidine]Amine (B4)

B4 was prepared from $[3-(4-methoxyphenyl)-5-(4-bromophenyl)-1H-pyrrol-2-yl][3-(4-methoxyphenyl)-5-(4-bromophenyl)pyrrol-2-yliden]amine (B) (0.23 g, 0.30 mmol), 4-(diphenylamino)phenylboronic acid (199.5 mg, 0.69 mmol), Pd(PPh_3)_4 (15.6 mg, 0.0014 mmol) and K_2CO_3 (0.28 g, 2.00 mmol) according to the method for B2.The residual$

was purified by column chromatography on silica eluting with toluene which gave the product as a cobalt blue solid (178 mg, 57%), mp > 350 °C.¹HNMR (400 MHz, CDCl₃) δ 8.15 (d, J = 8.4 Hz, 2H), 8.07 (d, J = 8.8 Hz, 2H), 7.69 (d, J = 8.4 Hz, 2H), 7.54 (d, J = 8.8 Hz, 2H), 7.29 (t, J = 7.8 Hz, 4H), 7.15 (d, J = 7.6 Hz, 4H), 7.13 (d, J = 8.8 Hz, 2H), 7.06 (t, J = 7.4 Hz, 2H), 7.01 (s, 2H), 6.99 (s, 1H),3.90 (s, 3H). ¹³CNMR (100 MHz, CDCl₃) δ 160.83, 158.06, 147.83, 147.47, 145.62, 143.13, 142.68, 133.58, 130.85, 130.16, 130.12, 130.08, 130.04, 129.32, 127.78, 126.53, 125.42, 124.72, 123.33, 123.21, 117.53, 114.16, 55.43. IR (ATR, cm⁻¹) 3034.0, 2835.4, 1587.4, 1386.8, 1251.8. HRMS (TOF-ESI): Calcd (C₇₀H₅₂BF₂N₅O₂) [M + H]⁺: m/z 1044.42605; found, m/z = 1044.42421.

Synthesis of [5-(4-Methoxyphenyl)-3-(4-Methoxyphenyl) -1H-Pyrrol-2-Yl]-[5-(4-Methoxyphenyl)-3-(4-Bromophenyl) Pyrrol-2-Ylidene]Amine (C)

3,5-bis(4-methoxyphenyl)-2-nitroso-1H-pyrrole (4a) (1.54 g, 5.0 mmol) and 2-(4-bromophenyl)-4-(4-methoxyphenyl)-1Hpyrrole (3b) (1.64 g, 5.00 mmol) were dissolved in AcOH (25 mL). The reaction mixture was added acetic anhydride (5 mL) and heated to 100 °C for 1 h. Ice (100 mL) and 5 M NaOH (100 mL) was added and the mixture was stirred for 30 min, extracted with CH_2Cl_2 (100 mL \times 3). The combined organics were dried over Na₂SO₄ and the solvent was removed under reduced pressure. The crude product was purified on a silica gel column chromatography (1:2, EtOAc/hex), resulting the product as a dark green solid (2.98 g, %96), mp > 350 °C.¹HNMR (400 MHz, CDCl₃) δ 8.02 (d, J = 8.8 Hz, 2H), 7.99 (d, J = 9.2 Hz, 2H), 7.95 (d, J = 8.0 Hz, 2H), 7.68 (d, J = 8.4 Hz, 2H), 7.62 (d, J = 9.2 Hz, 2H), 7.14 (s, 1H), 7.05 (d, J = 8.0 Hz, 2H), 6.96 (s, 1H), 6.95 (d, J = 8.0 Hz, 2H), 3.92 (s, 3H), 3.88 (s, 3H), 3.85 (s, 3H). IR (ATR, cm-1) 3074.5, 2831.8, 1589.0, 1246.0. HRMS (TOF-ESI): Calcd $(C_{35}H_{28}BrN_{3}O_{3})$ [M + H]⁺: m/z = 620.13720; found, m/z = 620.13950.

Synthesis of BF₂ Chelate of [5-(4- Methoxyphenyl)-3-(4-Methoxyphenyl)-1H-Pyrrol-2-Yl]-[5-(4-Methoxyphenyl) -3-(4-Bromophenyl)Pyrrol-2-Ylidine]Amine (C1)

C1 was prepared from [5-(4-methoxyphenyl)-3-(4-methoxyphenyl)-1H-pyrrol-2-yl]-[5-(4-methoxyphenyl)-3-(4-bromophenyl)pyrrol-2-ylidene]amine (C) (1.08 g, 1.75 mmol),*N*,*N*-diisopropylethylamine (3.05 mL, 17.5 mmol),boron trifluoride diethyl etherate (3.24 mL, 26.25 mmol) and dry CH₂Cl₂ (300 mL) according to the method for B1.The residual was purified by column chromatography on silica eluting with hexane/EtOAc (2:1,*v* $/v) which gave the product as a dark brown solid (1.02 g, 87%), mp 245–246 °C.¹HNMR (400 MHz, CDCl₃) <math>\delta$ 8.09 (d, J=9.2 Hz,

2H), 8.06 (d, J = 9.2 Hz, 2H), 8.02 (d, J = 9.2 Hz, 2H), 7.88 (d, J = 8.4 Hz, 2H), 7.59 (d, J = 8.4 Hz, 2H), 7.01 (d, J = 8.8 Hz, 2H), 6.99 (d, J = 8.8 Hz, 2H), 6.99 (t, 1H), 6.98 (d, J = 8.8 Hz, 2H), 6.84 (s, 1H), 3.90 (s, 3H), 3.89 (s, 3H), 3.88 (s, 3H).¹³CNMR (100 MHz, CDCl₃) δ 162.35, 161.10, 160.65, 160.38, 154.90, 131.89, 131.84, 131.79, 131.67, 131.19, 130.96, 130.85, 130.80, 130.76, 130.68, 125.58, 125.02, 124.71, 123.72, 117.96, 117.93, 116.43, 114.32, 114.19, 114.14, 55.45, 55.43, 55.41. IR (ATR, cm⁻¹) 3003.2, 2835.4, 1597.1, 1379.1, 1253.7. HRMS (TOF-ESI): Calcd (C₃₅H₂₇BBrF₂N₃O₃) [M]⁺: m/z = 666.13753; found, m/z = 666.13917.

Synthesis of BF₂ Chelate of [4-Bromo-3-(4-Methoxyphenyl) -5-(4-Methoxyphenyl)-1H-Pyrrol-2-Yl] -[4-Bromo-3-(4-Methoxyphenyl)-5-(4-Bromophenyl) Pyrrol-2-Ylidene]Amine (CC1)

CC1 was prepared from BF₂ chelate of [5-(4methoxyphenyl)-3-(4- methoxyphenyl)-1H-pyrrol-2yl]-[5-(4-methoxyphenyl)-3-(4-bromophenyl)pyrrol-2ylidine]amine (C1) (0.40 g, 0.60 mmol), N-bromosuccinimide (NBS) (0.22 g, 1.26 mmol) and CH₂Cl₂ (150 mL) according to the method for AA1. The residual was purified by column chromatography on silica eluting with hexane/EtOAc (2:1, V/ V) which gave the product as a dark brown solid (0.41 g, 83%), mp 292–293 °C. ¹HNMR (400 MHz, CDCl₃) δ 7.90 (d, J = 8.8 Hz, 4H), 7.88 (d, J = 9.2 Hz, 4H), 7.77 (d, J =9.2 Hz, 2H), 7.59 (s, 4H), 6.99 (d, J = 8.8 Hz, 4H), 6.98 (d, J = 9.2 Hz, 2H), 3.89 (s, 3H), 3.88 (s, 3H), 3.86 (s, 3H). ¹³CNMR (100 MHz, CDCl₃) δ 162.01, 161.11, 160.80, 159.54, 154.22, 144.92, 144.91, 143.37, 143.36, 141.17, 141.16, 132.53, 132.48, 132.33, 131.96, 131.26, 128.81, 125.11, 123.52, 123.19, 121.52, 113.68, 113.64, 55.40, 55.38, 55.33. IR (ATR, cm⁻¹) 3001.2, 2835.4, 1599.0, 1377.2, 1246.0. HRMS (TOF-ESI): Calcd $(C_{35}H_{25}BBr_{3}F_{2}N_{3}O_{3})$ [M + H]⁺: m/z = 823.95650; found, m/z = 823.95874.

Synthesis of BF₂ Chelate of [5-(4-Methoxypheny) -3-(4-Methoxypheny)-1H-Pyrrol-2-Yl]-[5-(4-Methoxypheny) -3-(4-Methoxybiphenyl)Pyrrol-2-Ylidine]Amine (C2)

C2 was prepared from[5-(4-methoxyphenyl)-3-(4-methoxyphenyl)-1H-pyrrol-2-yl]-[5-(4-methoxyphenyl)-3-(4-bromophenyl)pyrrol-2-ylidene]amine (C) (0.18 g, 0.30 mmol), 4-methoxyphenylboronic acid (51.47 mg, 0.33 mmol), Pd(PPh_3)_4 (7.80 mg, 0.0007 mmol) and K₂CO₃ (0.28 g, 2.00 mmol) according to the method for B2.The obtained residue was purified by column chromatography on silica eluting with toluene, which gave the product as a purple solid (106 mg, 50%), mp 261 °C.¹HNMR (400 MHz, CDCl₃) δ 8.10 (t, J = 8.4 Hz, 4H), 8.06 (d, J = 9.2 Hz, 2H), 8.05 (d, J = 8.8 Hz, 2H), 7.66 (d, J = 8.4 Hz, 2H), 7.60 (d, J = 8.8 Hz, 2H), 7.00 (d, J = 8.0 Hz, 4H), 6.99 (d, J = 8.4 Hz, 4H), 6.96 (s, 1H), 6.96 (s, 1H), 3.89 (s, 6H), 3.87 (s, 3H), 3.86 (s, 3H); ¹³CNMR (100 MHz, CDCl₃) δ 161.98, 160.85, 160.63, 159.61, 158.99, 156.85, 145.74, 144.97, 143.54, 143.52, 142.49, 142.35, 142.33, 137.86, 132.71, 131.71, 131.66, 131.61, 130.85, 130.73, 130.26, 130.01, 129.96, 129.92, 129.02, 128.20, 126.59, 125.59, 125.28, 124.10, 117.50, 117.15, 117.10, 117.09, 114.31, 114.21, 114.12, 114.10, 55.40, 55.34. IR (ATR, cm⁻¹) 3032.1, 2833.4, 1599.0, 1384.9, 1251.8. HRMS (TOF-ESI): Calcd (C₄₂H₃₄BF₂N₃O₄) [M + H]⁺: m/z =: 694.26888; found, m/z = 694.27082.

Synthesis of BF₂ Chelate of [5-(4-Methoxypheny) -3-(4-Methoxypheny)-1H-Pyrrol-2-YI]-[5-(4-Methoxypheny) -3-(2,4-Dimethoxybiphenyl)Pyrrol-2-Ylidine]Amine (C3)

C3 was prepared from [5-(4-methoxyphenyl)-3-(4methoxyphenyl)-1H-pyrrol-2-yl]-[5-(4-methoxyphenyl)-3-(4-bromophenyl)pyrrol-2-ylidene]amine (C) (0.18 g, 0.30 mmol), 2,4-dimethoxyphenylboronic acid (60.05 mg, 0.33 mmol), Pd(PPh₃)₄ (7.80 mg, 0.0007 mmol) and K₂CO₃ (0.28 g, 2.00 mmol) according to the method for B2. The obtained residue was purified by column chromatography on silica eluting with toluene which gave the product as a dark brown solid (89 mg, 41%), mp 228–229 °C.¹HNMR (400 MHz, CDCl₃) δ 8.09 (d, J=9.2 Hz, 2H), 8.07 (t, J= 8.6 Hz, 4H), 8.06 (d, J=9.2 Hz, 2H), 7.64 (d, J=8.8 Hz, 2H), 7.33 (d, J = 8.4 Hz, 1H), 7.00 (d, J = 8.8 Hz, 2H), 6.99 (d, J = 8.8 Hz, 4H), 6.61–6.57 (m, 2H), 6.98 (s, 1H), 6.96 (s, 1H), 3.90 (s, 6H), 3.88 (s, 3H), 3.87 (s, 3H), 3.83 (s, 3H). ¹³CNMR (100 MHz, CDCl₃) δ 161.90, 160.81, 160.70, 160.84, 157.69, 157.03, 154.03, 132.51, 131.62, 131.61, 131.55, 131.33, 130.82, 130.74, 129.91, 129.46, 129.23, 129.20, 129.11, 125.36, 124.19, 120.37, 117.41, 117.36, 114.19, 114.13, 114.11, 109.99, 104.79, 99.01, 55.53, 55.48, 55.43, 55.40. IR (ATR, cm^{-1}) 3066.8, 2835.3, 1597.0, 1383.0, 1271.1. HRMS (TOF-ESI): Calcd $(C_{43}H_{36}BF_2N_3O_5)$ [M + Na] ⁺: m/z = 746.26140; found, m/z = 746.26448.

Synthesis of BF₂ Chelate of [5-(4-Methoxypheny) -3-(4-Methoxypheny)-1H-Pyrrol-2-Yl]-[5-(4-Methoxypheny) -3-(4-N,N-Diphenylaminobiphenyl)Pyrrol-2-Ylidine]Amine (C4)

C4 was prepared from[5-(4-methoxyphenyl)-3-(4-methoxyphenyl)-1H-pyrrol-2-yl]-[5-(4-methoxyphenyl)-3-(4-bromophenyl)pyrrol-2-ylidene]amine (C) (0.18 g, 0.30 mmol), 4-(diphenylamino)phenylboronic acid (95.41 mg, 0.33 mmol), Pd(PPh_3)_4 (7.80 mg, 0.0007 mmol) and K₂CO₃ (0.28 g, 2.00 mmol) according to the method for B2. The obtained residue was purified by column chromatography on silica eluting with toluene which gave the product as a dark brown solid (113 mg. 45%),mp > 350 °C.¹HNMR (400 MHz, CDCl₃) δ 8.13 (d, J = 8.4 Hz, 2H), 8.10 (d, J = 9.6 Hz, 2H), 8.07 (d, J = 8.8 Hz, 2H), 8.06 (d, J=9.2 Hz, 2H), 7.68 (d, J= 8.8 Hz, 2H), 7.54 (d, J = 8.4 Hz, 2H), 7.29 (t, J =8.0 Hz, 4H), 7.17-7.14 (m, 4H), 7.13 (s, 1H), 7.06 (t, J = 7.4 Hz, 2H), 7.01 (d, J = 9.2 Hz, 4H), 7.00 (d, J =8.8 Hz, 4H), 6.97 (s, 1H), 3.90 (s, 6H), 3.88 (s, 3H). ¹³CNMR (100 MHz, CDCl₃) δ 162.76, 161.97, 160.87, 160.85, 147.50, 142.37, 137.61, 136.59, 135.78, 135.69, 131.64, 131.60, 131.34, 130.29, 130.05, 129.99, 129.94, 129.31, 127.76, 126.51, 125.60, 125.32, 124.70, 123.38, 123.18, 114.22, 114.14, 114.12, 55.43, 55.41. IR (ATR, cm⁻¹) 3032.1, 2833.4, 1589.4, 1386.8, 1251.8. HRMS (TOF-ESI): Calcd $(C_{53}H_{41}BF_2N_4O_3)$ [M]⁺: m/z = 830.32400; found, m/z = 830.32609.

Synthesis of BF₂ Chelate of [4-Methoxyphenyl-3-(4-Methoxyphenyl) -5-(4-Methoxyphenyl)-1H-Pyrrol-2-Yl] -[4-Methoxyphenyl-3-(4-Methoxyphenyl)-5-(4-Bromophenyl) Pyrrol-2-Ylidene]Amine (CC2)

CC2 was prepared from BF₂chelate of [4-Bromo-3-(4methoxyphenyl)-5-(4-methoxyphenyl)-1H-pyrrol-2-yl]-[4bromo-3-(4-methoxyphenyl)-5-(4-bromophenyl)pyrrol-2ylidene]amine (CC1) (0.21 g, 0.25 mmol), 4methoxyphenylboronic acid (136.8 mg, 0.95 mmol), Pd(PPh₃)₄ (19.5 mg, 0.0018 mmol) and K₂CO₃ (0.35 g, 2.50 mmol) according to the method for A2. The obtained residue was purified by column chromatography on silica eluting with toluene which gave the product as a dark brown solid (48 mg, 21%) mp > 350 °C.¹HNMR (400 MHz, CDCl₃) δ 7.48 (d, J = 9.2 Hz, 2H), 7.46 (d, J = 8.4 Hz, 2H), 7.42 (d, J = 8.8 Hz, 2H), 7.38, J = 8.4 Hz, 2H), 7.31 (d, J = 8.8 Hz, 2H), 6.90 (d, J = 8.8 Hz, 4H), 6.88 (d, J = 8.8 Hz, 2H), 6.79 (d, J = 8.8 Hz, 2H), 6.78 (d, J = 8.8 Hz, 2H), 6.75 (d, J = 8.8 Hz, 2H), 6.74 (d, J =8.8 Hz, 2H), 3.81 (s, 6H), 3.79 (s, 3H), 3.78 (s, 3H), 3.78 (s, 3H). ¹³CNMR (100 MHz, CDCl₃) δ 161.01, 160.39, 160.00, 159.79, 158.79, 158.65, 154.7, 154.65, 146.05, 144.43, 141.14, 139.43, 139.40, 139.39, 132.51, 132.42, 132.11, 131.74, 131.68, 131.12, 131.07, 131.04, 130.92, 130.15, 129.02, 128.20, 125.36, 125.34, 125.28, 124.72, 124.36, 122.75, 113.92, 113.88, 113.32, 55.22, 55.16, 55.12. IR (ATR, cm⁻¹) 3072.6, 2837.3, 1599.0, 1365.6, 1246.0. HRMS (TOF-ESI): Calcd $(C_{49}H_{39}BBrF_2N_3O_5)$ [M + H]⁺: m/z = 880.21922; found, m/z = 880.22060.

Results and Discussion

Synthesis and Coupling Reactions

Symmetrical aza-BODIPY compounds were initially synthesized successfully by condensation of acetophenone and aromatic aldehyde derivatives. Then, unsymmetrical analogues were obtained by pyrrole-nitrosolated pyrrole coupling reactions to compare the photophysical properties of the different substitution patterns as shown in Schemes 1 and 2. Diverse electron donor groups are directly attached at the -2, -3, -5and - 6 position of aza-BODIPYs. O'Shea's procedures [2, 7] were used in the synthesis of symmetrical aza-BODIPY compounds A1 and B1. Unsymmetrical azadipyrromethene compound C was synthesized by condensation of 2-(4bromophenyl)-4-(4-methoxyphenyl)-1H-pyrrole(3b) and 3,5-(4methoxyphenyl)-2-nitroso-1H-pyrrole (4a) and afforded in high (96%) yield. The compounds A1 and C1 compounds were treated with 2.1 equiv. of N-bromosuccinimide (NBS) in CH₂Cl₂ at room temperature. AA1 and CC1 were obtained with 83% and 84% yields, respectively. -3, -3,5, -2,6 aryl bromine substituted azadipyrromethene ligands and aza-BODIPY complexes were designed as starting materials for Suzuki-Miyaura coupling reactions. Reactions were performed with 4-methoxyphenylboronic acid, 2,4-Dimethoxyphenylboronic acid, 4-(diphenylamino)phenylboronic acid as electron donating groups in the presence of tetrakis(triphenylphosphine)palladium(0) [Pd(PPh₃)₄] in H₂O/EtOH/Toluene (1:1:2, by volume) mixture at 85 °C for 24 h. Compounds B2-B4 and C2-C4 were obtained by one pot synthesis using aryl bromine substituted azadipyrromethene ligands. Compounds A2, A4 and CC2 were synthesized 2,6-dibromine, 2,6-dibromine and 3,4-bromophenyl substituted aza-BODIPY derivatives (Schemes 1 and 2). All reactions have been monitored by TLC analysis and reaction mixture chromatographed by using silica gel.

The yields of the coupling reactions were found to change depending on the reactivity of 3, 3,5 and 2,6 positions of the aza-BODIPY core (Scheme 3). 3 positions of the aza-BODIPY compounds C2, C3 and C4 have been obtained with 50, 41 and 45% yields, respectively. 3,5 positions of compound **B4** has been obtained higher yield in 57%. On the other hand 3,5 position analogue compounds B2 and B3 have been obtained a relatively smaller yield of 26 and 34%. It was observed that the lower yields arise from the poor solubility of the compounds in reaction solvents. Similarly, coupling reactions on the 2,6 substituted aza-BODIPY compounds A2 and A4 gave the relatively poor yields (33% and 19%, respectively). The poor coupling yields may also be attributed to steric hindrance of the 2,6 positions of the indacene core caused by the existence of 1,7 phenyl groups on the indacene core. Apart from that, when the Suzuki-Miyaura coupling reaction has been performed with 2,3,6 tri-bromine substituted compound CC1 with excess boronic acid derivative, no all



Scheme 1 Synthetic pathway of the symmetrical aza-BODIPY Dyes (i) Pd(PPh_3)₄, K₂CO₃, ArB(OH)₂, 85 °C, 24 h. (ii) BF₃OEt₂, DIEA, CH₂Cl₂, rt., 24 h

target products have been obtained. Only 2,6 4methoxyphenyl substituted aza-BODIPY compound has been isolated from this reaction and afforded in 21% yield (Scheme 3). This may indicates the region selectivity in coupling reactions on 2,6 positions compared to 3 or 3,5 positions on the aza-BODIPY scaffold [38, 39].

Spectroscopy and Photophysical Properties

Identification of the compounds was performed through ¹H-NMR, ¹³C-NMR, FTIR and HRMS. In ¹HNMR of symmetrical aza-BODIPYs, the 2,6 pyrrole protons appear as a singlet between 6.91 to 7.00 ppm. However, unsymmetrical aza-BODIPY compounds, the pyrrole protons appear as two singlet signals in different ppm and these protons were disappeared in 2,6 phenyl or bromine substituted aza-BODIPY compounds. 1,3,5,7 aryl protons of compounds A1, B1, B2, B3, B4, C2, C3, C4 were observed between 8.05 to 8.15 ppm. On the other hand, 2,6 phenyl substituted A2, A4 and CC2 compounds were monitored to upfield shift between 7.53 to

7.46, compared to 2,6 unsubstituted aza-BODIPY compounds. These results show that the electronic properties of aza-BODIPY core at 2,6 positions could be changed markedly [40]. Interestingly, ¹HNMR of 3 bromine substituted aza-BODIPY compound **CC1** was observed only one sharp singlet peak in 7.59 ppm corresponding to four aryl protons (Fig. 1). Similar singlet peak was observed our previous study at 2,3,5,6 bromine substituted aza-BODIPY compounds [41]. This effect can be observed in para-disubstituted benzene having two identical substituents, because all of the hydrogens are magnetically equivalent stemming from the symmetry in the molecule.

High resolutions mass spectra (HRMS-ESI-TOF) of the compounds A1, A4, B2, C1 and A, A2, B3, B4,C, C2-C4 showed molecular and protonated molecular ion peaks, respectively and compound C3 gave [M-Na]⁺ molecular ion peaks. Isotope peaks were observed depending on the number of bromine atoms to bromine substituted aza-BODIPY compounds consisted with the assigned formulations. For instance, the spectrum for compound CC1 consisting of three



Scheme 2 Synthetic pathway of the unsymmetrical aza-BODIPY dyes (i) $Pd(PPh_3)_4$, K_2CO_3 , $ArB(OH)_2$, 85 °C, 24 h. (ii) BF_3OEt_2 , DIEA, CH_2Cl_2 , rt., 24 h

bromine atoms was revealed M, M + 2, M + 4 and M + 6 peak groups as shown in Fig. 2.

The photophysical parameters, measured in CHCl₃ solution $(5 \times 10^{-6} \text{ M})$, are reported in Table 1. The absorption spectra of all aza-BODIPY derivatives exhibited two intense peaks, In the UV region (between 311 to 351 nm, corresponding to the $\pi \rightarrow \pi^*$ transitions and the major characteristic absorption bands in the NIR region (between 672 to 718 nm, S₀ \rightarrow S₁ transitions) (Fig. 3). The extinction coefficients for newly synthesized series are high, in the range of 20,400–51,600 M⁻¹ cm⁻¹ for high energy region and in the range of 46,600–147,600 M⁻¹ cm⁻¹ for the low energy region.

The Stokes shifts for these new series are generally narrow $(600-1045 \text{ cm}^{-1})$, except for the compound **B4** (1443cm⁻¹). Furthermore, substitution of bromine atoms in the 2,6

positions (AA1 and CC1) involves 12 and 11 nm hypsochromic shift compared to A1 and C1. Owing to the introduction of methoxyphenyl and triphenylamine groups onto the 2 and 6 positions in the aza-BODIPY moiety, compounds A2 and A4 shows lower hypsochromic shift compared to A1. However, 10 nm bathochromic shift was observed when compared to AA1 compound. In emission spectra, all 2,6-substituted compounds gave reduced fluorescence quantum yields. Besides, quenched fluorescence was observed for the compound A4 due to the strong electron donating behavior of directly linked triphenylamine group to the aza-BODIPY core.

The 3 and $3,5 \pi$ -extended aza-BODIPYs (**B2-B4** and **C2-C4**) containing electron donating groups showed significant bathochromic shifts on both absorption and emission spectra



Scheme 3. Synthesis of the compound CC2



Fig. 1 ¹H NMR spectra of the compound CC1

depicted in Figs.3 and 4, respectively. 3,5 substituted series **B2/B3** (24 and 22 nm) and 3 substituted series compound **C2/C3** (12 and 11 nm) showed systematic bathochromic shifts compared to reference compounds **B1** and **C1**, respectively. The largest red-shift in the series was observed for the compound **B4** ($\lambda_{abs} = 718$ nm), which has two strong electron-donating and π -extended triphenylamine groups attached to the 3,5positions of the aza-BODIPY moiety. Additionally, the absorption bands of the triphenylamine substituted compounds relatively broader and the Stokes shifts are larger compared to those of methoxy substituted counterparts.

Compounds C2, C3 and C4 have electron donating groups on 3 position of the aza-BODIPY core that exhibited red shifted absorption maxima compared to unsubstituted compound C1. Note that, the highest molar absorption coefficient was observed 3 π -extended and 2,4-dimethoxyphenyl substituted unsymmetrical aza-BODIPY compound C3 (ϵ =147,000 M⁻¹ cm⁻¹). And also unsymmetrical compound **C2** showed the highest fluorescence quantum yield ($\Phi_F = 0.44$).

It's clear from the graphs that the triphenylamine group as an electron donor forms a charge transfer state [14]. The triphenylamine substituted compounds have lower molar absorption coefficients and quantum yields. Conversely, the aza-BODIPY derivatives with weaker electron donating groups or unsubstituted aza-BODIPYs have higher values (see Table 1). Note that, the fluorescence quantum yields of the compounds **A1**, **B1** and **C1** which have only difference on the 3 position of the core structure as bromine and methoxy group don't differ greatly. This may be explained by the equally capable of donating electrons of relevant ones to the aza-BODIPY core when the appended to 3,5 positions of the core. As a general result, compared to unsubstituted analogues, the bathochromic shifts between 40 and 50 nm both in absorption



 Table 1
 Summary of the

 photophysical properties of the
 aza-BODIPY dyes. All spectra

 were measured in chloroform
 in chloroform

Compound	$\lambda_{abs(nm)}$	$\lambda_{em(nm)}$	$\epsilon[M^{-1} \ cm^{-1}]$	FWHM (nm)	Stokes Shifts (cm ⁻¹)	$\Phi_{\rm F}$
A	324/628	_	46,400/60200	60/90	_	_
A1	336/692	722	20,400/89400	51	600	0.360
AA1	680	718	61,600	57	77	0.015
A2	690	738	46,600	83	943	0.004
A4	311/690	-	42,000/52600	130	-	-
В	315/615	-	54,200/54400	46/90	-	_
B1	330/674	708	20,200/62400	63	712	0.310
B2	344/698	744	22,800/77400	66	886	0.430
B3	315/696	738	27,400/75200	66	819	0.410
B4	351/718	801	51,600/61000	91	1443	0.010
С	318/619	-	33,800/39600	63/97	-	_
C1	331/683	714	24,800/72600	54	636	0.360
CC1	672	711	85,800	58	816	0.004
C2	339/695	729	21,400/68800	59	671	0.440
C3	310/694	733	43,400/147600	58	767	0.230
C4	344/701	735	41,600/83000	73	660	0.036
CC2	680	732	66,200	89	1045	0.001

and emission spectra were provided by coupling reactions for the new dyes.

In order to determine the singlet oxygen production features of the novel compounds, singlet oxygen generation study with 1,3-diphenylisobenzofuran (DPBF) scavenger was carried out in solution (Figs. 5 and 6).

The formation of triplet state by intersystem crossing (ISC) in emissive molecules causes the fluorescence quenching that makes them potential candidates in singlet oxygen production. In this regard, singlet oxygen generation was expected to obtain only non-emissive compounds [42]. Therefore, compounds A1, B1-B3 and C1-C3 were not expected any singlet oxygen efficiency due to the high fluorescence quantum yields (Table 1). Thus, we have tested non-emissive symmetrical and asymmetrical compounds which substituted 3, 3,5 and 2,6 π -extended and electron donating groups (A2, A4, B4, C4 and CC2). Our singlet oxygen determination studies shows that compounds A2, A4, B4, C4 and CC2 have not

changed singlet oxygen efficient of aza-BODIPY dyes, although these compounds have showed weak emission features. This may due to radiationless decay to the ground state instead of energy or electron transfer to the triplet state of designed aza-BODIPY dyes [43]. However, 2,6 dibromine substituted aza-BODIPY compounds AA1 and CC1 were showed high singlet oxygen quantum yields because of the heavy atom effect of bromine atoms [7, 41]. So that compounds AA1 and CC1 solution with DPBF were observed decrease in absorption spectra at 414 nm (Figs. 5 and 6, respectively). The Singlet oxygen quantum yields of the compounds AA1 and CC1 were calculated 71% and 74% in DCM against methylene blue which was used as standard, respectively. Based on the results from literature, halogenated aza-BODIPY compounds generally show higher singlet oxygen production values about %70 ($\Phi\Delta = 0.70$ in DCM) [8, 10, 44]. From this point, it can be concluded that the compounds AA1 and CC1 also have to some degree elevated singlet oxygen



Fig. 3 Normalized absorption spectra of aza-BODIPY dyes $(5 \times 10^{-6} \text{ M}, \text{ in CHCl}_3)$



Fig. 4 Emission spectra of the aza-BODIPY dyes $(5 \times 10^{-6} \text{ M}, \text{ in CHCl}_3)$

production capabilities. Note that, even the compounds have close features in the production of singlet oxygen, novel compounds could be excited with lower energy. In this respect, extending the π -electron system and asymmetric structure of the aza-BODIPYs via Suzuki-Miyaura coupling do not make them efficient sensitizer in photodynamic therapy applications.

Conclusions

In summary, we prepared 3, 3,5 and 2,6 substituted symmetrical and unsymmetrical aza-BODIPY dyes via Pd-catalyzed Suzuki-Miyaura cross coupling reactions. The Suzuki-Miyaura coupling reactions on aza-BODIPY derivatives in 3,5 [44–46] and 2,6 [40] positions are very limited and also 3 positions reactions of the unsymmetrical aza-BODIPY compounds were the first example in the literature. In this paper, we reported that at 3, 3,5 and 2,6 positions Suzuki-Miyaura

reactions effect of photophysical and spectroscopic properties. 2,6 substituted aza-BODIPY compounds showed drastically change in the electronic properties as well as fluorescence. absorption intensities and stokes shifts were altered dramatically. And also these series were synthesized with poor yields compared to 3 and 3,5 positions because of the steric hindrance. The highest bathochromic shifts were observed π extended and strong electron donating groups at 3,5 positions. Our previous work indicated that extended π -conjugation and strong electron donating groups cause increased two-photon absorption properties of aza-BODIPY dyes at telecommunication wavelengths. Furthermore, 2,6-dibromine substituted aza-BODIPY compounds AA1 and CC1 were exhibited significantly high singlet oxygen quantum yields were found to be 71 and 74% respectively. These data may open an effective perspective and these compounds could be good candidates for diverse applications such as photodynamic therapy (PDT) and two photon absorption (2PA). In particular, synthesized compounds may be applicable for n-type semiconductors on



Fig. 5 5Absorption spectra of DPBF upon irradiation in the presence of the compound AA1 for 14 s (recorded at 2 s interval) and plot of change in absorbance of DPBF at 414 nm vs irradiation time ($\lambda_{ex} = 630$ nm) against methylene blue (MB) as the standard in CH₂Cl₂



Fig. 6 Absorption spectra of DPBF upon irradiation in the presence of the compound CC1 for 14 s (recorded at 2 s interval) and plot of change in absorbance of DPBF at 414 nm vs irradiation time ($\lambda_{ex} = 630$ nm) against methylene blue (MB) as the standard in CH₂Cl₂

organic photovoltaic cells (OPVs) due to the several desirable characteristics, for instance, long wavelength absorption, high FWHM and high molar absorption coefficients, as a subject of future study.

Funding This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Compliance with Ethical Standards

Conflicts of Interest/Competing Interests The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- Sathyamoorthi G, Soong ML, Ross TW, Boyer JH (1993) Fluorescent Tricyclic Beta-Azavinamidine - BF₂ Complexes. Heteroat Chem 4:603–608
- 2. Killoran J, Allen L, Gallagher JF, Gallagher WM, O'Shea DF (2002) Synthesis of BF_2 chelates of tetraarylazadipyrromethenes and evidence for their photodynamic therapeutic behaviour. Chem Commun 8:1862–1863
- Treibs A, Kreuzer FH (1968) Di- and tri-Pyrrylmethene complexes with Di-Fluoro boron. Annalen Der Chemie-Justus Liebig 718: 208–223
- Loudet A, Burgess K (2007) BODIPY dyes and their derivatives: syntheses and spectroscopic properties. Chem Rev 107:4891–4932
- Wood TE, Thompson A (2007) Advances in the chemistry of dipyrrins and their complexes. Chem Rev 107:1831–1861
- McDonnell SO, O'Shea DF (2006) Near-infrared sensing properties of dimethlyamino-substituted BF₂-azadipyrromethenes. Org Lett 8:3493–3496
- Gorman A, Killoran J, O'Shea C, Kenna T, Gallagher WM, O'Shea DF (2004) In vitro demonstration of the heavy-atom effect for photodynamic therapy. J Am Chem Soc 126:10619–10631

- Gallagher WM, Allen LT, O'Shea C, Kenna T, Hall M, Gorman A, Killoran J, O'Shea DF (2005) A potent nonporphyrin class of photodynamic therapeutic agent: cellular localisation, cytotoxic potential and influence of hypoxia. Br J Cancer 92:1702–1710
- McDonnell SO, Hall MJ, Allen LT, Byrne A, Gallagher WM, O'Shea DF (2005) Supramolecular photonic therapeutic agents. J Am Chem Soc 127:16360–16361
- O'Connor AE, Mc Gee MM, Likar Y, Ponomarev V, Callanan JJ, O'Shea DF, Byrne AT, Gallagher WM (2012) Mechanism of cell death mediated by a BF₂-chelated tetraaryl-azadipyrromethene photodynamic therapeutic: dissection of the apoptotic pathway in vitro and in vivo. Int J Cancer 130:705–715
- Coskun A, Yilmaz MD, Akkaya EU (2007) Bis(2-pyridyl)substituted boratriazaindacene as an NIR-emitting chemosensor for hg(II). Org Lett 9:607–609
- Bouit P-A, Kamada K, Feneyrou P, Berginc G, Toupet L, Maury O, Androud C (2009) Two-photon absorption-related properties of functionalized BODIPY dyes in the infrared range up to telecommunication wavelengths. Adv Mater 21:1151–1154
- Kucukoz B, Hayvali M, Yilmaz H, Uguz B, Kurum U, Yaglioglu HG, Elmali A (2012) Synthesis, optical properties and ultrafast dynamics of aza-boron-dipyrromethene compounds containing methoxy and hydroxy groups and two-photon absorption crosssection. J Photoch Photobio A 247:24–29
- Tekin S, Kucukoz B, Yilmaz H, Sevinc G, Hayvali M, Yaglioglu HG, Elmali A (2013) Enhancement of two photon absorption properties by charge transfer in newly synthesized aza-borondipyrromethene compounds containing triphenylamine, 4-ethynyl-N,N-dimethylaniline and methoxy moieties. J Photoch Photobio A 256:23–28
- Killoran J, O'Shea DF (2006) Impact of a conformationally restricted receptor on the BF₂ chelated azadipyrromethene fluorosensing platform. Chem Commun:1503–1505
- Jokic T, Borisov SM, Saf R, Nielsen DA, Kuehl M, Klimant I (2012) Highly Photostable near-infrared fluorescent pH indicators and sensors based on BF₂-chelated Tetraarylazadipyrromethene dyes. Anal Chem 84:6723–6730
- Adarsh N, Avirah RR, Ramaiah D (2010) Tuning photosensitized singlet oxygen generation efficiency of novel Aza-BODIPY dyes. Org Lett 12:5720–5723

- Batat P, Cantuel M, Jonusauskas G, Scarpantonio L, Palma A, O'Shea DF, McClenaghan ND (2011) BF₂-Azadipyrromethenes: probing the excited-state dynamics of a NIR Fluorophore and photodynamic therapy agent. J Phys Chem A 115:14034–14039
- Mueller T, Gresser R, Leo K, Riede M (2012) Organic solar cells based on a novel infrared absorbing aza-bodipy dye. Sol Energ Mat Sol C 99:176–181
- Min J, Ameri T, Gresser R, Lorenz-Rothe M, Baran D, Troeger A, Sgobba V, Leo K, Riede M, Guldi DM, Brabec CJ (2013) Two similar near-infrared (IR) absorbing Benzannulated Aza-BODIPY dyes as near-IR sensitizers for ternary solar cells. ACS Appl Mater Interfaces 5:5609–5616
- 21. Murtagh J, Frimannsson DO, O'Shea DF (2009) Azide Conjugatable and pH responsive near-infrared fluorescent imaging probes. Org Lett 11:5386–5389
- Collado D, Vida Y, Najera F, Perez-Inestrosa E (2014) PEGylated aza-BODIPY derivatives as NIR probes for cellular imaging. RSC Adv 4:2306–2309
- Sharman WM, Allen CM, van Lier JE (1999) Photodynamic therapeutics: basic principles and clinical applications. Drug Discov Today 4:507–517
- O'Connor AE, Gallagher WM, Byrne AT (2009) Porphyrin and Nonporphyrin photosensitizers in oncology: preclinical and clinical advances in photodynamic therapy. Photochem Photobiol 85: 1053–1074
- 25. Dolmans D, Fukumura D, Jain RK (2003) Photodynamic therapy for cancer. Nat Rev Cancer 3:380–387
- Ochsner M (1997) Photodynamic therapy: the clinical perspective review on applications for control of diverse tumorous and nontumorous diseases. Arzneim-Forsch Drug Res 47:1185–1194
- MacDonald IJ, Dougherty TJ (2001) Basic principles of photodynamic therapy. J Porphyrins Phthalocyanines 5:105–129
- Hajri A, Wack S, Meyer C, Smith MK, Leberquier C, Kedinger M, Aprahamian M (2002) In vitro and in vivo efficacy of Photofrin (R) and pheophorbide a, a bacteriochlorin, in photodynamic therapy of colonic cancer cells. Photochem Photobiol 75:140–148
- Awuah SG, Polreis J, Biradar V, You Y (2011) Singlet oxygen generation by novel NIR BODIPY dyes. Org Lett 13:3884–3887
- Duman S, Cakmak Y, Kolemen S, Akkaya EU, Dede Y (2012) Heavy atom free singlet oxygen generation: doubly substituted configurations dominate S-1 states of Bis-BODIPYs. J Organomet Chem 77:4516–4527
- Huang L, Yu X, Wu W, Zhao J (2012) Styryl Bodipy-C-60 dyads as efficient heavy-atom-free organic triplet photosensitizers. Org Lett 14:2594–2597
- Kotha S, Lahiri K, Kashinath D (2002) Recent applications of the Suzuki-Miyaura cross-coupling reaction in organic synthesis. Tetrahedron 58:9633–9695
- Hall MJ, McDonnell SO, Killoran J, O'Shea DF (2005) A modular synthesis of unsymmetrical tetraarylazadipyrromethenes. J Organomet Chem 70:5571–5578
- Qian Y, Ma G-Y, Yang Y, Cheng K, Zheng Q-Z, Mao W-J, Shi L, Zhao J, Zhu H-L (2010) Synthesis, molecular modeling and

biological evaluation of dithiocarbamates as novel antitubulin agents. Bioorg Med Chem 18:4310–4316

- 35. Sakamoto R, Kusaka S, Kitagawa Y, Kishida M-a, Hayashi M, Takara Y, Tsuchiya M, Kakinuma J, Takeda T, Hirata K, Ogino T, Kawahara K, Yagi T, Ikehira S, Nakamura T, Isomura M, Toyama M, Ichikawa S, Okumura M, Nishihara H (2012) Fluorescent azadipyrrinato zinc(II) complex: hybridisation with a dipyrrinato ligand. Dalton T 41:14035–14037
- Fery-Forgues S, Lavabre D (1999) Are fluorescence quantum yields so tricky to measure? A demonstration using familiar stationery products. J Chem Educ 76:1260–1264
- 37. Ma J, Yuan X, Kucukoz B, Li S, Zhang C, Majumdar P, Karatay A, Li X, Yaglioglu HG, Elmali A, Zhao J, Hayvali M (2014) Resonance energy transfer-enhanced rhodamine-styryl Bodipy dyad triplet photosensitizers. J Mater Chem C 2:3900–3913
- Feng Z, Jiao L, Feng Y, Yu C, Chen N, Wei Y, Mu X, Hao E (2016) Regioselective and stepwise syntheses of functionalized BODIPY dyes through palladium-catalyzed cross-coupling reactions and direct C-H Arylations. J Organomet Chem 81:6281–6291
- Zhao Q, Li C, Senanayake CH, Tang W (2013) An efficient method for sterically demanding Suzuki-Miyaura coupling reactions. Chemistry 19:2261–2265
- 40. Kumar S, Khan TK, Ravikanth M (2015) Synthesis and properties of hexaarylated AzaBODIPYs. Tetrahedron 71:7608–7613
- 41. Karatay A, Miser MC, Cui X, Kucukoz B, Yilmaz H, Sevinc G, Akhuseyin E, Wu X, Hayvali M, Yaglioglu HG, Zhao J, Elmali A (2015) The effect of heavy atom to two photon absorption properties and intersystem crossing mechanism in aza-borondipyrromethene compounds. Dyes Pigments 122:286–294
- 42. Xu Y, Feng T, Yang T, Wei H, Yang H, Li G, Zhao M, Liu S, Huang W, Zhao Q (2018) Utilizing Intramolecular Photoinduced Electron transfer to enhance Photothermal tumor treatment of Aza-BODIPY-based near-infrared nanoparticles. ACS Appl Mater Interfaces 10:16299–16307
- 43. Xu Y, Zhao M, Zou L, Wu L, Xie M, Yang T, Liu S, Huang W, Zhao Q (2018) Highly stable and multifunctional Aza-BODIPYbased phototherapeutic agent for anticancer treatment. ACS Appl Mater Interfaces 10:44324–44335
- 44. Naresh Balsukuri NM, Lone MY, Mori S, Das A, Sen P, Gupta I (2020) Donor-acceptor architectures of tetraphenylethene linked aza-BODIPYs: Synthesis, crystal structure, energy transfer and computational studies. Dyes Pigments 176:17
- Khan TK, Sheokand P, Agarwal N (2014) Synthesis and studies of Aza-BODIPY-based pi-conjugates for organic electronic applications. Eur J Org Chem 2014:1416–1422
- 46. Liu S, Shi Z, Xu W, Yang H, Xia N, Liu X, Zhao Q, Huang W (2014) A class of wavelength-tunable near-infrared aza-BODIPY dyes and their application for sensing mercury ion. Dyes Pigments 103:145–153

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.