

Amine induced carbonylation of alkynes to cyclobutenediones using $\text{Fe}_3(\text{CO})_{12}$

Chellappan Rameshkumar and Mariappan Periasamy *

School of Chemistry, University of Hyderabad, Central University PO, Hyderabad-500 046, India


Received 1 December 1999; accepted 9 February 2000

Abstract

Iron carbonyl species, prepared in situ in THF using $\text{Fe}_3(\text{CO})_{12}$, react with alkynes at 25°C, in the presence of certain amines, to give the corresponding cyclobutenediones in moderate to good yields (25–61%) after $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$ oxidation. © 2000 Elsevier Science Ltd. All rights reserved.

Keywords: alkynes; amines; carbonylation; cyclobutenediones.

Iron carbonyls have been increasingly used in organic synthesis in recent years.^{1–5} The readily accessible, inexpensive $\text{Fe}(\text{CO})_5$ is not reactive towards organic substrates under ambient conditions. The $\text{Fe}(\text{CO})_5$ reacts with amine oxides to give reactive coordinatively unsaturated ' $\text{Fe}(\text{CO})_4$ ' species.^{6–8} Also, it was reported that amines such as *n*-butylamine, pyrrolidine and piperidine form addition complexes with $\text{Fe}(\text{CO})_5$.^{9,10} In continuation of studies on the synthetic applications of iron carbonyl species,^{11–13} we became interested in the application of the $\text{Fe}_3(\text{CO})_{12}$ for the preparation of reactive iron carbonyl




Scheme 1.

* Corresponding author.

species. The $\text{Fe}_3(\text{CO})_{12}$ species is a precursor to a variety of organoiron compounds.^{14–17} Under photolysis, it dissociates into ' $\text{Fe}(\text{CO})_4$ '.^{18,19} Recently, it was reported that the Pauson–Khand reaction of alkyne– $\text{Co}_2(\text{CO})_6$ species is facilitated by certain amines.^{20,21} Accordingly, we have examined the effect of amines on the reactivity of alkynes with $\text{Fe}_3(\text{CO})_{12}$. We report here that amines facilitate the reaction of alkynes with $\text{Fe}_3(\text{CO})_{12}$ to give the corresponding cyclobutenediones after $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$ oxidation (Scheme 1).

The $\text{Fe}_3(\text{CO})_{12}$ reacts with alkynes in the presence of *n*-butylamine at 25°C to give the corresponding cyclobutenediones in moderate to good yields (25–61%) after $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$ oxidation (Table 1). Several alkynes were converted to the corresponding cyclobutenediones. It was observed that other amines such as *n*-octylamine, benzylamine, pyrrolidine, piperidine, triethylamine and DMF also give the same cyclobutenediones (28–45%). However, it was observed that the use of *n*-butylamine gives better yields (25–61%) (Scheme 1).

Evidently, this reagent system can tolerate unmasked functional groups such as hydroxyl and trimethylsilyl (entries 5, 6, 7 and 8 Table 1). Also, in the case of *n*-butylamine (entries 3 and 7 Table 1) traces of cyclic imides (**3c** and **7c**) are formed (2–5%) under these conditions.²²




The following is the typical procedure: The $\text{Fe}_3(\text{CO})_{12}$ was prepared following a reported procedure.²³ The $\text{Fe}_3(\text{CO})_{12}$ (3.02 g, 6 mmol) and *n*-butylamine (0.201 g, 2.75 mmol) in THF (50 mL) under N_2 atmosphere was stirred for 30 min at 25°C. 1-Heptyne (0.120 g, 1.25 mmol) was added and the stirring was continued for a further 12 h. The metal carbonyl complex was decomposed using $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$ (1.7 g mmol) in acetone (10 mL). Saturated aq. NaCl (25 mL) was added, and the resulting solution was extracted with ether (100 mL), washed with brine (20 mL), dried over anhydrous MgSO_4 and concentrated. The residue was subjected to column chromatography (silica gel, hexane/ethyl acetate). Ethyl acetate (2%) in hexane eluted cyclobutenedione **1a** (0.116 g, 61%).²⁴

A tentative mechanistic pathway for the formation of cyclobutenediones can be visualised as shown in Scheme 2. Presumably, the amine may help in the splitting of the $\text{Fe}_3(\text{CO})_{12}$ to give (amine) $\text{Fe}(\text{CO})_4$ and ' $\text{Fe}_2(\text{CO})_8$ ' species, which on further reaction with alkynes could give the intermediate complex of the type **I**. Oxidation of **I** with $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$ would produce the corresponding cyclobutenediones.¹³


Yields are of products isolated by column chromatography using hexane:ethyl acetate (98:2) as eluent and based on the amount of alkynes used.





To examine the possibility whether coordinatively unsaturated ' $\text{Fe}(\text{CO})_4$ ' species can be prepared



Scheme 2.

Table 1
Reaction of $\text{Fe}_3(\text{CO})_{12}$ with alkyne in the presence of *n*-butylamine in THF



S.No.	Alkyne	Product ^a	Yield (%) ^b
1.	R= C ₅ H ₁₁ , R'= H	1a	61
2.	R= C ₆ H ₁₃ , R'= H	2a	54
3.	R= C ₆ H ₅ , R'= H	3a	41
4.	R, R'= Ph	4a	45
5.	R= C ₅ H ₁₁ , R'= SiMe ₃	5a	28
6.	R= C ₈ H ₁₇ , R'= SiMe ₃	6a	25
7.		 7a	37
8.		 8a	33

a) Products were identified from spectral data (IR, ¹H NMR, ¹³C NMR and Mass).²² The spectral data of product **1a** and **4a** were identical to the reported data.²⁴ The spectral data of products **2a** and **3a** are comparable to the spectral data reported for compound **1a**. For the cyclobutenediones, **5a** and **7a** HRMS data were also obtained.

b) Yields are of products isolated by column chromatography using hexane/ethyl acetate (98:2) as eluent and based on the amount of alkynes used.

using $\text{Fe}(\text{CO})_5$, we carried out the reaction with $\text{Fe}(\text{CO})_5$ (6 mmol), *n*-butylamine (2.75 mmol) and phenylacetylene (1.25 mmol).^{9,10} However, only the starting alkyne was recovered.

In conclusion, the simple, convenient procedure for the double carbonylation of alkynes to obtain the corresponding cyclobutenediones using a readily available iron carbonyl reagent system described here should be attractive for synthetic applications.

Acknowledgements

We are grateful to the UGC, CSIR and DST (New Delhi) for financial support. We are also thankful to the UGC for support under Special Assistance Program.

References

- Schore, N. E. *Chem. Rev.* **1988**, *88*, 1081.
- Brunet, J. J. *Chem. Rev.* **1990**, *90*, 1041.
- Liebeskind, L. S.; Bausdon, S. L.; South, M. S.; Iyer, S.; Leeds, J. F. *Tetrahedron* **1985**, *41*, 5839.
- Khumtaveepron, K.; Alper, H. *Acc. Chem. Res.* **1995**, *28*, 414.
- Noyori, R.; Hayakawa, Y. *Tetrahedron* **1985**, *41*, 5879.
- Alper, H.; Edward, J. T. *Can. J. Chem.* **1970**, *48*, 1543.
- Shvo, Y.; Hazum, E. *J. Chem. Soc., Chem. Commun.* **1975**, 829.
- Elzinga, J.; Hogeveen, H. *J. Chem. Soc., Chem. Commun.* **1977**, 705.
- Edgell, W. F.; Yang, M. T.; Bulkin, B. J.; Bayer, R.; Koizumi, N. *J. Am. Chem. Soc.* **1965**, *87*, 3080.
- Edgell, W. F.; Bulkin, B. J. *J. Am. Chem. Soc.* **1966**, *88*, 4839.
- Periasamy, M.; Radhakrishnan, U.; Brunet, J.-J.; Chauvin R, Elzaizi, A. *Chem. Commun.* **1996**, 1499.
- Periasamy, M.; Rameshkumar, C.; Radhakrishnan, U. *Tetrahedron Lett.* **1997**, *38*, 7229.
- Periasamy, M.; Rameshkumar, C.; Radhakrishnan, U.; Brunet J.-J. *J. Org. Chem.* **1998**, *63*, 4930.
- Hubel, W.; Braye E. H.; Clauss, A.; Weiss, E.; Kruerke, U.; Brown, D. A.; King, G. S. D.; Hoogzand, C. *Inorg. & Nuclear Chem.* **1959**, *9*, 204.
- Hubel, W.; Braye, E. H. *J. Inorg. & Nuclear Chem.* **1959**, *10*, 250.
- King, R. B.; Manuel, T. A.; Stone, F. G. A. *Inorg. & Nuclear Chem.* **1961**, *16*, 233.
- King, R. B.; Treichel, P. M.; Stone, F. G. A. *J. Am. Chem. Soc.* **1961**, *83*, 3600.
- Carr, M. D.; Kane, V. V.; Whiting, M. C. *Proc. Chem. Soc.* **1964**, 408.
- Kane, V. V.; Light, J. R. C.; Whiting, M. C. *Polyhedron* **1985**, *4*, 533.
- Rajesh, T.; Periasamy, M. *Tetrahedron Lett.* **1998**, *39*, 117.
- Sugihara, T.; Yamada, M.; Ban, H.; Yamaguchi, M.; Kaneko, C. *Angew. Chem., Int. Ed. Engl.* **1997**, *36*, 2801.
- Spectral data: ^{13}C NMR (CDCl_3) data: (δ ppm): (**1a**) 13.7, 22.1, 25.6, 27.1, 31.2, 184.8, 196.6, 199.9, 208.3; MS (EI): m/z 152 [M^+ , 13%], 81 [$(\text{M}^+ - \text{C}_5\text{H}_{11})$, 20%]. (**2a**) 13.9, 25.9, 26.8, 28.9, 29.6, 31.8, 184.9, 196.7, 199.9, 208.3. (**3a**) 14.0, 22.6, 26.0, 27.5, 29.3, 29.7, 31.7, 31.9, 184.8, 196.6, 199.7, 208.4. (**4a**) 128.7, 129.7, 131.2, 134.6, 187.4, 196.1. (**3c**) 10.1, 18.6, 21.5, 27.7, 30.6, 36.2, 38.5, 124.0, 132.4, 172.5, 175.9. (**5a**) -2.1, 13.7, 22.2, 26.5, 29.0, 31.8, 200.1, 201.4, 207.7, 211.3. MS (EI): m/z : 224 (M), 154 (B); HRMS; m/z calcd: 224.123258, found: 224.123138. (**6a**) -2.1, 14.0, 22.5, 26.8, 29.0, 29.6, 31.5, 31.7, 200.0, 207.6, 211.2. (**7a**) 13.7, 22.1, 26.9, 28.4, 29.3, 31.8, 74.9, 124.9, 125.3, 128.3, 128.8, 196.4, 198.5, 201.2, 202.9. MS (EI); m/z : 272 (M^+ , 4%). HRMS; m/z calcd: 272.141243 found: 272.141843. (**7c**) 13.4, 13.7, 19.9, 22.1, 24.6, 25.1, 29.5, 31.2, 31.3, 38.3, 42.7, 57.0, 74.8, 125.6, 127.7, 128.2, 143.6, 178.5, 179.1. MS (EI); m/z : 347. Anal. calcd for $\text{C}_{21}\text{H}_{31}\text{O}_3\text{N}$: C, 73.04; H, 8.98; N, 4.05. Found: C, 73.10; H, 9.15; N, 4.11. (**8a**) 13.8, 13.9, 22.2, 22.4, 24.9, 26.0, 26.9, 28.9, 31.5, 31.8, 35.0, 68.6, 197.3, 198.4, 201.8, 202.7. MS (EI); m/z : 266 ($\text{M}^+ + 1$, 5%), 195 ($\text{M}^+ - \text{C}_5\text{H}_{11}$, 16%).
- King, R. B.; Stone, F. G. A. *Inorg. Synth.* **1960**, *7*, 193.
- Parker, M. S. A.; Rizzo, C. J. *Synth. Commun.* **1995**, *25*, 2781.