

Investigative Studies on the Formation of the Imidazo [4',5':3,4]pyrido[2,3-b]indole Ring: Formal Synthesis of the Alkaloids Grossularines-1 and 2. X-Ray Crystal Structures of 5-Indol-3-yl-imidazole and Bisimidazo-carbazole Derivatives.

Pedro Molina^a, Pilar M. Fresneda^a, Miguel A. Sanz^a, Concepción Foces-Foces^b, M. Carmen Ramirez de Arellano^c

 a) Departamento de Química Orgánica, Facultad de Química, Universidad de Murcia, Campus de Espinardo, E-30071 Murcia, Spain

 b) Departamento de Cristalografía, Instituto de Química-Física 'Rocasolano', CSIC, Serrano 119, E-28006 Madrid, Spain

c) Departamento de Química Inorgánica, Facultad de Química, Universidad de Murcia, Campus de Espinardo, E-30071 Murcia, Spain

Received 2 February 1998; revised 16 March 1998; accepted 29 May 1998

Abstract

Reactions of several 2,3-disubstituted indoles, bearing a chlorine or amino function at position 2 and an 1,2-dicarbonyl group or bromoacetyl substituent at position 3, with N,N-dimethylguanidine is reported. In general, 5-indol-3-yl imidazole derivatives are found to be the reaction products, however when the 3-bromoacetyl-2-tertbutoxycarbonylaminoindole is used the 2-dimethylamino-4-hydroxy-6-methoxymethyl-3H-imidazo[4',5':3,4]pyrido[2,3-b]indole is obtained through a step-wise formation of the pyridine and the imidazole rings. The crystal molecular structure of $5.H_2O$ and $7.HBr.H_2O.3/2$ EtOH have been determined by X-Ray analysis. The water molecules in 5. H_2O act as both donor (O-H...N) and acceptor (N-H...O) and are responsible for the formation of strands along the b axis. © 1998 Elsevier Science Ltd. All rights reserved.

Keyword: Indoles, imidazoles, imidazolines, sponges, X-Ray crystal structures

Marine organisms are among the most promising sources of new and biologically active molecules. Certain secondary metabolites are non-traditional guanidine-based alkaloids¹ that possess a broad spectrum of powerful biological activities. The guanidine moiety is most frequently found in the guise of a 2-aminoimidazole ring that is generally substituted with alkyl groups on carbon or nitrogen.²⁻⁸

Marine alkaloids grossularines, possessing a cyclic guanidine moiety in the form of 2dimethylaminoimidazole subunit, were isolated in 1984 from the marine tunicate *Dendrodoa grossularia* (Stylidae) collected in the coast of Brittany.⁹⁻¹⁰ These compounds, which are

Dedicated to Prof. Alan R. Katritzky for his important contribution to Organic Chemistry

the first examples of naturally ocurring pyrido[2,3-b]indoles (α -carbolines), exhibit marked cytotoxicity toward murine and human tumor cells.¹¹



Recently, two approaches have been developed for the construction of the tetracyclic imidazo[4',5':3,4]pyrido[2,3-b]indole framework of the titled compounds. The first is based on the palladium-catalyzed cross coupling between the appropriately substituted imidazo[4, 5-c]pyridine and a stannane aniline followed by base-promoted intramolecular cyclization of the resulting biaryl compound to construct the α -carboline ring system.¹²

In the second approach, the formation of the α -carboline ring is based on the thermal electrocyclic ring-closure of a 2-azahexatriene system, including the indole 2,3-bond and the imidazole 4,5-bond. The starting material 3-imidazolylindole derivative was prepared by cross-coupling reaction between 3-iodoindole-2-carboxylate and the directed metallation-derived imidazole followed by hydrolysis of the ester group and further Curtius rearrangement with diphenyl phosphorazidate (DPPA).¹³⁻¹⁴ In the first total synthesis of grossularines, the tetracyclic ring system was elaborated into the target compounds through a sequence involving triflate formation and subsequent arylation by way of the Stille or Suzuki cross-coupling methodologies with or without carbon monoxide insertion.¹⁵

Following our programme directed toward the synthesis of imidazole-containing alkaloids from marine origin,¹⁶⁻¹⁸ we wish to report an efficient synthesis of the 2-dimethylamino-4-hydroxy-6-methoxymethyl-3H-imidazo[4',5':3,4]pyrido[2,3-b]indole, successfully used in the above mentioned synthesis of the alkaloids grossularines-1 and 2. Our approach is based on the sequential formation of the imidazole and pyridine rings starting from a 2,3-disubstituted indole possessing suitable functionalities. In this sense, the substituent at position 3 of the starting indole must be suitable to build-up the desired 2-dimethylaminoimidazole ring, whereas the pyridine ring could be formed by cyclization of the appropriate nitrogen functionality at position 2 onto the preformed imidazole ring and thus completing the assemblage of the gross ring system of the target molecule.

A survey of the literature reveals that classical methods for the preparation of 2aminoimidazole derivatives involve condensation of α -aminocarbonyl compounds with cyanamide,¹⁹⁻²² reaction of α -dicarbonyl compounds with guanidine followed by reduction,²³⁻²⁴ and reaction of α -haloketones with N-acetylguanidine.⁷



Thus, we initially required a N-protected 2,3-disubstituted indole and the starting material of choice was the N-methoxymethyl-3-acetyl-2-chloroindole 1, which was prepared in 85% yields from 3-acetyl-2-chloroindole²⁵ and methoxymethyl chloride in the presence of sodium hydride. Oxidation of compound 1 with selenium dioxide afforded the 3-indolylglyoxal hydrate 2 in 96% yield which was converted into the corresponding azide 3 in 80% yield by treatment with polymeric quaternary ammonium azide.²⁶ The reaction of 2 and 3 with N,N-dimethylguanidine in ethanol at -30°C gave 4 (91%) and 5 (95%) respectively. The unexpected formation of the aminocompound 5 possessing an unusual 4H-imidazol-4-one ring could be understood by initial formation of an intermediate 3-indolyl dihydroimidazolone like 4 which under the reactions conditions underwent dehydrogenation of the dihydroimidazole ring with concomitant reduction of the azido group to give 5. It is noteworthy that the inmuno-modulating 2-(dimethylamino)-5-(1H-indol-3-yl)-4H-imidazol-4-one has been isolated from *Dendrodoa grossularia*²⁷ and represents the first naturally ocurring member of the 4H-imidazol-4-one ring.

Suitable crystals of 5 allowed an X-Ray structure analysis. The monohydrate of 5, as it exists in the solid state, is depicted in Fig.1a together with the numbering scheme. As far as we know [Cambridge Structural Database²⁸ (October 1997 release; CSD hereinafter)], it is noteworthy that it is the first time that the crystal structure of an imidazol-4-one ring is reported (Fig. 1, Table 1). The N4-C11 bond shows double-bond character while a certain degree of electron delocalization is observed in the other N-C bonds. The N atoms of the amino groups (N2 and N5) are sp² hybridized (Table 1). A strong intramolecular hydrogen bond (Table 1) is found between one N-H group and O=C one of the five-membered ring (Fig. 1a) that cause remarkable variation, mainly, in the bond distances and angles of the



Fig. 1.- (a) Displacement ellipsoid plot (30% probability level) of 5.H₂O showing the atom-numbering.
(b) A view of one hydrogen-bonded strand of dimers along the b axis. Dotted lines represent hydrogen bond interactions.

NI-CI	1.363(3)	N1-C8	1.406(3)	N1-C9	1.461(4)
N2-C1	1.312(3)	C1-C2	1.442(3)	C2-C3	1.457(2)
C3-C8	1.408(3)	C2-C11	1.396(3)	C11-C12	1.527(2)
N3-C12	1.354(4)	N3-C13	1.353(3)	N4-C13	1.382(3)
N4-C11	1.338(3)	C12-O2	1.232(3)	C13-N15	1.324(4)
N5-C14	1.458(3)	N5-C15	1.454(3)		
C8-N1-C1	110.2(2)	C8-N1-C9	125.3(3)	C1-N1-C9	124.2(2)
NI-C1-C2	108.6(2)	N1-C1-N2	122.2(2)	C2-C1-N2	129.2(2)
C1-C2-C3	105.7(2)	C1-C2-C11	128.9(2)	C3-C2-C11	125.3(2)
C2-C3-C8	107.5(2)	C3-C8-N1	107.9(2)	N4-C11-C12	108.0(2)
C11-C12-N3	107.2(2)	C11-C12-O2	127.0(2)	N3-C12-O2	125.9(2)
C12-N3-C13	103.8(2)	N3-C13-N4	117.7(2)	C13-N4-C11	103.3(2)
N3-C13-N5	122.3(2)	N4-C13-N5	120.0(2)	C13-N5-C14	121.4(2)
C13-N5-C15	122.1(2)	C14-N5-C15	116.6(2)		
C8-N1-C9-O1	68.7(3)	C14-N5-C13-N3	2.1(4)	C1-C2-C11-C12	-3.3(4)
N1-C9-O1-C10	68.6(3)	C15-N5-C13-N4	3.8(3)		
Hydrogen interactions					
Х-НҮ		Х-Н	XY	HY	Х-НҮ
N2-H21O1		0.82(4)	3.211(3)	2.83(3)	111(3)
N2-H21O3(x,y-1,z)		0.82(4)	3.042(3)	2.23(4)	174(3)
N2-H22O2		0.88(3)	2.712(3)	1.91(4)	151(3)
N2-H22O2(-x,-y,1-z)		0.88(3)	3.034(2)	2.46(3)	124(3)
O3-H31N3(-x,1-y,1-z)	0.92(3)	2.854(2)	1.94(3)	170(5)
O3-H32N4		0.93(5)	3.141(3)	2.28(4)	153(3)
C4-H4N4	······	0.99(3)	3.050(3)	2.45(2)	119(2)

Table 1. Selected geometrical parameters $(Å, \circ)$ for compound 5.H₂O.

pseudo-seven-membered ring formed. The C2-C11, C1-C2 and C2-C3 bonds are significanly elongated as compared to the Csp^2-Csp^2 value²⁹ [C=C-C(=O)-C=C: 1.478(31Å, the sample standard deviation being in parenthesis] and to the average values corresponding to 47 N-indole derivatives retrieved from CSD [1.371(39) and 1.437(11)Å for C1-C2 and C2-C3 respectively]. The external angles at C1, C2 and C11, that correspond to the internal angles in the pseudo-seven-membered ring are widened and the differences between the two external angles at each atom are highly significant (Table1). The indole and the imidazole systems are almost coplanar aided by the intramolecular interactions.

As illustrated in Fig. 1b, the water molecules link molecules of 5 to form dimers related by inversion centres that are further joined through N-H...OH₂ and N-H...O=C bonds to give strands along the **b** axis.

All attemps to reduce the dihydroimidazolone ring in compound 4 with a wide variety of reagents (NaBH₄ / TiCl₄; NaBH₄ / 'BuOOH; BH₃·SM₂; H₂ / Pd/C; LiAlH₄ / Et₂O; NaBH₄ / CH₃SO₃K / DMSO; DIBAL-H) were unsuccessful and complex mixtures were obtained. Unfortunately, the treatment of compound 5 with the same reagents led to the unchanged starting material.



Reagents and Conditions: a) Br_2 , Et_2O , $AlCl_3$, $0^{\circ}C \rightarrow r.t.$; b) DMG, Et_3N , DMF, $0^{\circ}C \rightarrow r.t.$

Scheme 2

These frustrating results led us to select a less oxidizable substituent at position 3. To this end, bromination of indole derivative 1 in the presence of $AlCl_3$ at 0°C provide the 2-bromoacetyl-3-chloroindole 6 in 75% yield. When an ethanolic solution of 6 was treated with N,N-dimethylguanidine at room temperature the bisimidazocarbazole derivative 7 was obtained in 60% yield in almost pure form, instead of the expected 5-(3-indolyl)imidazole (Scheme 2). An X-ray structure determination confirmed the proposed structure. (Fig 2, Table 2).

It was tempting to speculate about the mechanism for the conversion $6\rightarrow7$. A tentative mechanism could involve initial formation of the enamine 8 which undergoes alkylation by action of the second equivalent of 6 to give 9 which, by loss of HBr, led to the 1-azadiene 10. An intramolecular Michael addition takes place to give 11 which undergoes cyclization across the tautomeric form 12 to afford the bisimidazolyl 13. Dehydrogenation followed by intramolecular nucleophilic displacement of the chlorine atom at position 2 of the indole



Fig. 2.- Structure of 7.H₂O. 3/2 CH₃CH₂OH showing the atom numbering scheme. Ethanol molecules have been omitted for clarity.

Table 2. Selected b	ond lengths [Å] a	nd angles [⁰] for 7	7-H2O-3/2CH3CH2OH

CI-C16	1.737(4)	N1-C1	1.373(5)	N1-C9	1.398(5)
N2-C7	1.380(5)	N2-C2	1.477(5)	N3-C3	1.347(5)
N3-C7	1.361(5)	N4-C8	1.355(5)	N4-C4	1.390(5)
N5-C5	1.352(5)	N5-C8	1.370(5)	N7-C7	1.300(5)
N7-C23	1.447(6)	N7-C24	1.450(6)	N8-C8	1.330(5)
N8-C25	1.452(6)	N8-C26	1.458(5)	C1-C6	1.378(5)
C1-C2	1.514(6)	C2-C3	1.539(5)	C2-C15	1.546(5)
C3-C4	1.360(5)	C4-C5	1.405(6)	C5-C6	1.443(6)
C6-C10	1.421(6)	C15-C16	1.385(6)		
C1-N1-C27	128.9(3)	C8-N8-C25	118.4(3)	N7-C7-N2	123.9(4)
C9-N1-C27	122.4(3)	C7-N2-C2	107.8(4)	N3-C7-N2	113.6(4)
C3-N3-C7	105.6(3)	C8-N4-C4	106.4(3)	N8-C8-N4	122.3(4)
C5-N5-C8	103.9(3)	C16-N6-C17	107.2(3)	N8-C8-N5	124.9(4)
C16-N6-C29	128.9(4)	C17-N6-C29	123.6(3)	N4-C8-N5	112.7(3)
C7-N7-C23	121.9(4)	C7-N7-C24	120.9(4)	C14-C9-N1	129.4(4)
C8-N8-C26	119.8(4)	C29-O3-C30	112.7(3)	N1-C9-C10	108.0(4)
N1-C1-C6	109.0(3)	N1-C1-C2	127.6(3)	C11-C10-C6	134.6(4)
C6-C1-C2	122.6(4)	N2-C2-C1	117.2(4)	C14-C13-C12	121.2(4)
N2-C2-C3	98.0(3)	C1-C2-C3	109.0(3)	N6-C17-C18	109.1(3)
N2-C2-C15	112.4(3)	C1-C2-C15	110.3(3)	C16-C15-C2	126.6(3)
C3-C2-C15	109.0(3)	N3-C3-C4	129.4(4)	C18-C15-C2	128.0(3)
N3-C3-C2	111.9(3)	C4-C3-C2	118.6(3)	C15-C16-Cl	130.5(3)
C3-C4-N4	132.1(4)	C3-C4-C5	121.5(4)	N6-C17-C22	127.8(4)
N4-C4-C5	105.1(3)	N5-C5-C4	111.7(4)	C19-C18-C15	136.5(4)
N5-C5-C6	128.2(4)	C4-C5-C6	119.8(4)	C17-C18-C15	106.3(4)
C1-C6-C5	118.9(4)	C12-C11-C10	118.0(4)	O2-C27-N1	113.2(3)
C10-C6-C5	133.0(4)	N7-C7-N3	122.4(4)		

Hydrogen bonds.

X-HY	XY	HY	XHY	
O1W-H01Br	3.274(3)	2.42(4)	162.2(4.9)	
O1W-H02N3	2.905(5)	2.14(5)	140.2(4.9)	
N4-H4O1W	2.805(5)	2.02(1)	147.6(0.1)	
N2-H03Br (x-1, y, z)	3.447(4)	2.66(5)	172.57(4.9)	

by the enamine-vinylogous system provided the final product. Another plausible pathway would involve formation of the expected 5-(3-indolyl)imidazole which undergoes cyclodimerization through a Diels-Alder reaction, whereby the C=C double bond of the indole ring and the C=C double bond of the imidazole ring at position 3 functions as a diene, and the C=C double bond of the imidazole ring of the other molecule has taken the role of dienophile. Loss of HCl from the cycloadduct, followed by an oxidative aromatization, furnishes 7.(Scheme 3).



Scheme 3

The behaviour of indole 6 towards N,N-dimethylguanidine led us to reexamine the reaction sequence. Conversion of 1 into the azidoindole 15 was performed in 65% yield by reaction with sodium azide in DMSO at 45°C. Staundinger reaction of 15 with triphenylphosphine in diethyl ether at room temperature, and further treatment with THF/H₂O in the presence of catalytic amounts of hydrochloric acid provided 16 in 85% yield. Treatment of 16 with (Boc)₂O/DMAP at room temperature afforded 17 in 70% yield. Bromination of compound 17 in methanol gave 18 in 84% yield, which by treatment with trifluoroacetic acid in dichloromethane at room temperature provided 19 in 81% yield. However, when compound 17 was treated with Br₂ in chloroform at 50°C underwent bromination with concomitant N-Boc deprotection to give directly 19 in 70% yield. The reaction of 19 with N,N-dimethylguanidine in ethanol at room temperature directly provided 20 in 80% yield.



Scheme 4

Taking into account that a) the formation of isocyanates from carbamates usually requires either strong acid conditions or high temperatures³⁰ and b) in the previously reported synthesis¹⁵ the electrocyclization of the isocyanate group at position 2 onto the preformed imidazole at position 3 to give the pyridine ring is carried out at high temperature, we reasonably thought that the conversion $19\rightarrow 20$ does not involve the formation of the intermediate isocyanate. A tentative mechanism for this conversion could involve initial formation of a dihydropyridone ring by nucleophilic attack of the enolate ion derived from the bromoacetyl substituent at position 3 on the carbamate group at position 2 and further formation of the imidazole ring accross the remaining α -bromocarbonyl moiety.

Formation of compound 20 constitutes a formal total synthesis of grossularines-1 and -2, since 20 may be easily converted into the target molecules in a straightforward manner.

EXPERIMENTAL SECTION

General Methods.

All melting points were determined on a Kofler hot-plate melting point aparatus and are uncorrected. IR spectra were obtained as Nujol emulsion or films on a Nicolet Impact 400 spectrophotometer. NMR spectra were recorded on a Bruker AC200 (200MHz) or a varian Unity 300 (300MHz). Mass spectra were recorded on a Hewlett-Packard 5993C spectrometer or a Fisson AUTOSPEC5000 VG. Microanalyses were performed on a Perkin Elmer 240C instrument.

X-ray Analysis.- A summary of data collection and refinement process is given in Table 3. The structure of $5.H_2O$ was solved by direct methods $(SIR92)^{31}$ and refined by least-squares procedures. All hydrogens were obtained from difference Fourier synthesis and refined isotropically in the last cycles. Scattering factors for $5.H_2O$, were taken from the International Tables for X-Ray Crystallography³² and the calculations were carried out with the XTAL,³³ PESOS³⁴ and PARST³⁵ set of programs running on a DEC3000-300X workstation.

Cyssial data Formula $C_{13}H_{17}S_0^2$, H_0^0 $C_{30}H_{12}BrClN_8O_2$, H_2O , $3/2CH_1CH_2OH$ Crystal bize (mm) 0 43 to 30 to 1/7 0.62 to 20 to 0.13 Symmetry Triclinic, P-1 Triclinic, P-1 Unit cell dimensions ($\hat{\Lambda}$, $\hat{\gamma}$) assignments from 58 Leas-squares from 74 reflections (2.0-60<45.0°) reflections (112-60-25.1°) as 10.7719(14) bassignments bassignments as 0.7719(14) bassignments Unit cell dimensions ($\hat{\Lambda}$, $\hat{\gamma}$) as 2.372(7) as 10.7719(14) bassignments unit cell dimensions ($\hat{\Lambda}$, $\hat{\gamma}$) as 2.372(7) cs 14.680(2) cs 10.680(10) genometry or 10.797(7) cs 10.680(10) genometry genometry for 7.11, 2 1933.2(5), 2 10.707 gelgen?), Arthofo 3.73, 317.34, 336 1.270, 739.11, 770 pelcon apetitie oriented genometry Graphite oriented Graphite monochromator. os cans Detector apetities 1 x 1°, 1 min/reflex. MoKa Assistant 1.7° genometry Graphite monochromator. os cans		5.H ₂ O	7.H ₂ O.3/2CH ₃ CH ₂ OH
Formula $C_{13}H_{17}N_5O_2$. H_5O $C_{30}H_{32}BClN_5O_2$. H_5O . $3/2CH_1CH_2OH$ Crystal bakinOrange, prismRed BlockCrystal size (ma) $6/3$ to 30 to 1/7Red BlockSymmetryTinclinic, P.1Triclinic, P.1Unit cell determination:Least-squares fit form 58Least-squares fit form 74reflections (2.0-00 $2/20-00FileUnit cell dimensions (Å?)a=0.2719(14)$	Crystal data		
Crystal habit Crystal habit Crystal habit (rystal habit SymmetryOrage, prism 0.43 × 0.20 × 0.17 0.43 × 0.20 × 0.13 0.45 × 0.20 × 0.14Red Block 0.45 × 0.20 × 0.13 0.45 × 0.20 × 0.13 0.45 × 0.20 × 0.14Unt cell diterminationLeast-squares fit from 58 reflections (2.0064 0.45 × 0.20 × 0.21 × 0.13 a=0.0719(14) b=8.9473(7) b=14.178(2) c=11.930(11) c=14.386(2) a=10.6500(10) b=100.440(10) p=06.500(13) b=100.440(10) p=06.500(13) b=100.440(10) p=06.500(13) b=100.440(10) p=06.500(13) p=100.440(10) p=06.500(13) p=100.440(10) p=06.500(13) p=100.440(10) p=06.500(13) p=100.440(10) p=06.500(13) p=100.440(10) p=06.500(13) p=100.440(10) p=06.500(13) p=100.440(10) p=06.500(13) p=100.440(10) p=06.500(13) p=100.440(10) p=10	Formula	$C_{15}H_{17}N_5O_2 \cdot H_2O$	C ₃₀ H ₃₂ BrClN ₈ O ₂ .H ₂ O. 3/2CH ₃ CH ₂ OH
Crystal size (mm) 0.43 \times 0.30 \times 0.17 0.63 \times 0.20 \times 0.13 Symmetry Triclinic, P-1 Triclinic, P-1 Unit cell determination Least-squares fit from 58 reflections (1.0.5664.50°) Least-squares fit from 74 reflections (1.1.2-66.2.5.1°) Unit cell dimensions (Å, °) Least-squares fit from 58 reflections (2.0.5664.50°) reflections (1.1.2-66.2.5.1°) Unit cell dimensions (Å, °) B=39.372(7) B=10.719(14) B=39.372(7) G=10.590(10) c=14.386(2) q=10.591(10) q=10.530(8) g=10.591(10) q=0.6509(13) y=106.530(8) g=10.719(14) Detegore N, M, F0000 1.373.31.7.4, 336 1.270.739.11, 770 gains Cut/Ca Mox Ca Graphite monochromator. Detector apertures 1 x 1°.1 min/reflex. Stantaria Stantaria Radiation Cuk a Mox Ca Stantaria Scan width: 1.7° 1.3° Stantaria Stantaria 246(0.03) 6571 25° Termerature (K) 244(0.03) 6571 25° Stantaria 246(0.03) 6571 246 <td>Crystal habit</td> <td>Orange, prism</td> <td>Red Block</td>	Crystal habit	Orange, prism	Red Block
Symmetry Unit cell determination Least-quares fit from 78 reflections (2.0 eq. 20.50 eq. 20.50 eq. 20.50 eq. 20.50 eq. 20.50 eq. 20.50 	Crystal size (mm)	0.43 x 0.30 x 0.17	0.62 x 0.20 x 0.13
Unit cell determination: Least-squares fit from 74 reflections (2.0-6645.0°) Least-squares fit from 74 reflections (2.0-6645.0°) Unit cell dimensions (Å,*) a=8.3712(7) a=10.7719(14) b=8.9473(7) c=14.386(2) c=11.5910(1) c=14.386(2) c=10.90,979(7) G=106.680(10) b=103.591(10) b=104.40(10) y=06.590(13) y=106.530(8) pecking: V(Å), Z 767.7(1), 2 De(gem*), M, F(000) 1.373.317.34, 336 1.270, 739.11, 770 geme* Jaro Toricle diffractometer. Scifert XRD3000-S Siemens P4 four circle diffractometer Graphite monochromator. $\omega/20$ scans. ω scans ω scans Detector apertures 1 x 1°.1 min/reflex. scans ω scans Scan width: 1.7° 1.3° 1.78 25° 173 Temperature (K) 295 173 Number of reflexions: 276 25° Collected 2648 7787 Independent (Rint.) 24640.033) 6571 (0.023) Observed (201 criterion) 126 6571 Sca	Symmetry	Triclinic, P-1	Triclinic, P-1
$ \begin{array}{llllllllllllllllllllllllllllllllllll$	Unit cell determination:	Least-squares fit from 58	Least-squares fit from 74
		reflections $(2.0 < \theta < 45.0^{\circ})$	reflections ($11.2 < \theta < 25.1^{\circ}$)
	Unit cell dimensions (Å,°)	a=8.2372(7)	a=10.7719(14)
		b=8.9473(7)	b=14.178(2)
		c=11.5910(11)	c=14.386(2)
		$\alpha = 109.079(7)$	$\alpha = 106.680(10)$
packing: V(Å), Z 767.7(1), 2 93.2(5), 2 De(g/cm ³), M, F(000) 1.373, 317.34, 336 1.270, 739.11, 770 µ(cm ³) 8.175 11.78 Exprimental data Four circle diffractometer: Seifert XRD3000-S Siemens P4 four circle diffractometer Bisecting geometry. Graphite oriented monochromator. W20 scans. ω scans ω scans Detector apertures 1 x 1°. 1 min/reflex. MoK α Scans Radiation CuK α NoK α Scans Scan width: 1.7° 1.3° ϕ_{max} 65° 5° Scans ω scans θ_{max} 65° 25° Trans Independent (Rint.) 246(0.03) 6578 (0.023) 6571 Observed (2σ(1) criterion) 1986 6571 3 every 250 reflect. No variation Solution and refinement Solution severy 100 reflect. 3 every 250 reflect. 3 every 250 reflect. Solution Direct methods Direct methods - - Radiator of the difference synthesis - 412 - Solution and		$\beta = 103.591(10)$	$\beta = 100.440(10)$
Packing: V(A), Z767.7 (1), 21933.2(5), 2Detg(grm), M, F(000)1.373.317.34.3361.270.739.11.770Utrom 18.1751.178Experimental dataTechniqueFour circle diffractometer: Seifert XRD3000-SSiemens P4 four circle diffractometermonochromator. $U20$ scans.Detector apertures 1 x 1°.1 min./reflex.Graphite monochromator.RadiationCuK α MoK α Scan width:1.7°1.3° ϕ_{max} 65°25°Temperature (K)295173Number of reflexions:CuCleted2648Collected26486571Independent (Rint.)2446(0.033)6578 (0.023)Observed (2011) criterion)19866571Standard reflexions:2reflexions every 100 reflect.3 every 250 reflect.Solution and refinementDirect methodsF², full matrixSolution and refinement:Fo, full matrixF², full matrixSolution and refinement:-412Solution and refinement:960.003Hatoms-96Number of variables285396Number of variables285396Number of variables9.90 (0.003)H atomsFrom difference synthesisrigid methyls, others ridingWurber of variables29.5396Number of variables29.5396Number of variables29.5396Number of variables29.5396Number of variables9.90 (0.00		γ = 96.509(13)	$\gamma = 106.530(8)$
$ \begin{array}{c} Detg(cm^n), M, F(000) & 1.373, 317.34, 336 & 1.270, 739.11, 770 \\ \mu(cm^{-1}) & 8.175 & 11.78 \\ \hline \\ Experimental data \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ $	Packing: V(A ³), Z	767.7(1), 2	1933.2(5), 2
μ (cm°,')8.17511.78Experimental dataTechniqueSolution circle diffractometer: Seifert XRD3000-S Bisecting geometry. Graphite oriented monochromator. $\omega/2\theta$ scans. Detector apertures 1 x 1°. 1 min./reflex.Siemens P4 four circle diffractometer Graphite monochromator. ω scans Detector apertures 1 x 1°. 1 min./reflex.RadiationCuKoaMoK α Scan width:1.7°1.3° θ_{max} 65°25°Temperature (K)295173Number of reflexions:Culced2648Collected26487787Independent (Rint.)2446(0.033)6578 (0.023)Observed (2g(1) criterion)19866571Standard reflexions:2 reflexions every 100 reflect. 	$Dc(g/cm^3), M, F(000)$	1.373, 317.34, 336	1.270, 739.11, 770
Experimental dataTechniqueFour circle diffractometer: Seifert XRD3000-5 Bisecting geometry. Graphite oriented monochromator. $M20$ scans. Detector apertures 1 × 1°. 1 min./reflex.Siemens P4 four circle diffractometer Graphite monochromator. $M20$ scans. M RadiationCuKoMoKoScan width:1.7° θ_{max} 65°Temperature (K)295Vumber of reflexions: Collected25°Collected2648Independen (Rint.)2466.033)Number of reflexions: Novariation6571Vumber of reflexions: Collected21°Variation and refinement: Secondary extinction (X)2660SolutionDirect methods F ² , full matrix AvariationSolutionDirect methods F ² , full matrix 206(6)Number of variables25°Number of variables25° <td>μ(cm⁻¹)</td> <td>8.175</td> <td>11.78</td>	μ(cm ⁻¹)	8.175	11.78
TechniqueFour circle diffractometer: Seifert XRD300-S Bisecting geometry. Graphite oriented monochromator. $\omega/20$ scans. ω scansSiemens P4 four circle diffractometer Graphite monochromator. ω scans ω scansRadiationCuK α McK α Scan width:1.7° 6 5°1.3° 25° θ_{max} 65° 25°25° 7Temperature (K)295173Number of reflexions: Collected2648 2648 2648 0bserved (2σ(1) criterion)2446(0.03) 1986 25°116578 25°11Standard reflexions: No variation22787 20023) 20024 20024 200247877 20023) 20023)Solution and refinement: Solution and refinement: Secondary extinction (x10 ⁴)Direct methods 2006(6) 2005(6)Direct methods 27°, full matrix 412 2003(2003)Parameters: Number of variables Aumber of restraints Parameters:Parameters: 412 412 412Madwer difference synthesis, rigid methyls, others ridingWeighting-scheme Final <afr </afr paeks and holes(A^{-3}) paeks and holes(A^{-3}) 0039,0.040Parameters: 40039,0.040U33(C10)=0.128(3)	Experimental data		
Bisecting geometry. Graphite oriented monochromator. $\omega/2\theta$ scans. Detector apertures 1 x 1°. 1 min./reflex.Graphite monochromator. ω scansRadiationCuK α MoK α Scan width:1.7°1.3° θ_{max} 65°25°Temperature (K)295173Number of reflexions: Collected26487787Independent (Rint.)2446(0.033)6578 (0.023)Observed (20(1) criterion)19866571Standard reflexions: 2 reflexions every 100 reflect. No variation3 every 250 reflect. 276 decaySolutionDirect methods Fo, full matrix 0.206(6)Direct methods F ² , full matrix PParameters: Number of variables285396Number of variables285396Number of variables285396Number of restraints e no-412Final <shift error="">0.0050.003H atomsFrom difference synthesis rigid methyls, others ridingWeighting-schemeEmpirical as to give no trends in exad/F> vs. w¹ = $\sigma^2(F^2) + (aP)^2 + bP. 3P = (2Fc^2 + Fo^2)$ a and bare constantsMax. thermal value (Å) Final AF peaks and holes(eÅ⁻³) Equation0.20, -0.27 0.0400.646, -0.614 0.054, 0.145</shift>	Technique	Four circle diffractometer: Seifert XRD3000-S	Siemens P4 four circle diffractometer
monochromator. $\omega/2\theta$ scans. Detector apertures 1 x 1°, 1 min./reflex. ω scansRadiation Scan width:LXK α MoK α Scan width:1.7°1.3° θ_{max} 65°25°Temperature (K)295173Number of reflexions: Collected2446(0.033)6578 (0.023)Observed (20(1) criterion)19866571Standard reflexions: No variation2446(0.033)6578 (0.023)Observed (20(1) criterion)19866571Standard reflexions: No variation2% decaySolution and refinement22% decaySolutionDirect methodsDirect methodsRefinement: Secondary extinction (x10°)0.206(6)-Parameters: Number of variables285396Number of restraints Final <shift error="">-412Final <shift error="">0.0050.003H atomsFrom difference synthesis rigid methyls, others ridingWeighting-schemeEmpirical as to give no trends in <math>\omega \Delta F > vs. < Großis) and $<\sin \theta A >$Max. thermal value (Å³) Final AF203(C(10)=0.128(3)U33(C24)=0.087(5)Final AR Number of Name No.0030.0050.005Solution103(S(24)=0.087(5)and bare constantsParameters: Number of restraints Final AR0.0040.054, 0.145</math></shift></shift>		Bisecting geometry. Graphite oriented	Graphite monochromator.
Detector apertures 1 x 1°. 1 min/reflex.MoK α Reading Scan width:CuK α MoK α Scan width: 1.3° MoK α θ_{max} 65°25°Temperature (K)295173Number of reflexions:TotalStandard (0.03)Collected26487787Independent (Rint.)2446(0.033)6571Observed (2011) criterion19865771Standard reflexions:2 reflexions every 100 reflect.3 every 250 reflect.Solution and refinementDirect methodsDirect methodsSolutionDirect methodsDirect methodsRefinement:60, 60(6)-SolutionStandard (x10°)0.206(6)Parameters:Number of variables285Number of variables285396Number of referations: $-$ A12Wamber of variables $-$ SolutionNumber of restrains $-$ Anameters: $-$ A12Number of variables285396Number of variables $-$ Solution $-$ A12Final <From difference synthesisrigid methyls, others ridingHardsCum difference synthesisrigid methyls, others ridingWeighting-schemeEmpirical as to give no trends in $cud F> vs. < Fobsl> and < sin\theta/A>- A12Mark difference synthesisrigid methyls, others ridingPark and holes(d_1^{A^{-3}})Q32(C10)=0.128(3)Q33(C24)=0.087(5)Final A and RwQ309, Q,040Q346, Q,04Q346, Q,041$		monochromator. ω/2θ scans.	ωscans
Radiation Scan width: LT MoKα Scan width: 1.7° 1.3° θ_{max} 65° 25° Temperature (K) 295 173 Number of reflexions: 7787 Collected 2648 7787 Independent (Rint.) 2446(0.033) 6578 (0.023) Observed (2σ(f) criterion) 1986 6571 Standard reflexions: 2 reflexions every 100 reflect. 3 every 250 reflect. No variation 2% decay 200 Solution and refinement 57, full matrix F², full matrix Solution Direct methods prevent methods Refinement: Fo, full matrix F², full matrix Secondary extinction (x10 ⁴) 0.206(6) - Parameters: Number of variables 285 Number of restraints - 412 Final <shift error=""> 0.005 0.003 H atoms From difference synthesis rigid methyls, others riding weighting-scheme Empirical as to give no trends in couAF> and b are constants</shift>		Detector apertures 1 x 1°. 1 min./reflex.	
Scan width: 1.7° 1.3° θ_{max} 65° 25° Temperature (K) 295 173 Number of reflexions: 787 Collected 2648 7787 Independent (Rint.) 2446(0.033) 6578 (0.023) Observed (20(1) criterion) 1986 6571 Standard reflexions: 2 reflexions every 100 reflect. 3 every 250 reflect. No variation 2% decay 2% decay Solution and refinement: Fo, full matrix F ² , full matrix Secondary extinction (x10 ⁴) 0.206(6) - Parameters: 96 9003 Number of variables 285 396 Number of variables 280 couo stimthesis rigd methyls, others r	Radiation	CuKa	ΜοΚα
$θ_{max}$ 65° 25° Temperature (K) 295 173 Number of reflexions:	Scan width:	1.7°	1.3°
Temperature (K)295173Number of reflexions: Collected26487787Independent (Rint.)2446(0.033)6578 (0.023)Observed (2c(1) criterion)19866571Standard reflexions:2 reflexions every 100 reflect. No variation3 every 250 reflect. 2% decaySolution and refinement2% decaySolution and refinement:Fo, full matrix F^2 , full matrixSolution (x10 ⁴)0.206(6)-Parameters:9090Number of variables285396Number of restraints-412Final <shift error="">0.0050.003H atomsFrom difference synthesisN-H and water difference synthesis, rigid methyls, others ridingWeighting-schemeEmpirical as to give no trends in $w^{-1} = \sigma^2(F^2) + (aP)^2 + bP, 3P = (2Fc^2 + Fo^2)$ a and b are constantsMax. thermal value (Å³)U33(C(10))=0.128(3)U33(C24)=0.087(5)Final AF peaks and holes(cÅ⁻³)0.20, -0.270.646, -0.614 0.054, 0.145</br></shift>	θ _{max}	65°	25°
Number of reflexions:Z6487787Collected26487787Independent (Rint.)2446(0.033)6578 (0.023)Observed (20(1) criterion)19866571Standard reflexions:2 reflexions every 100 reflect.3 every 250 reflect.No variation2 reflexions every 100 reflect.3 every 250 reflect.No variation0 zore2 reflexionsSolution and refinement:Eo, full matrixDirect methodsSecondary extinction (x104)0.206(6)-Parameters:-412Number of variables285396Number of restraints-412Final <shif error="">0.0050.003H atomsFrom difference synthesisN-H and water difference synthesis, rigid methyls, others ridingWeighting-schemeEmpirical as to give no trends in <cod>w⁻¹ = $\sigma^2(F^2) + (aP)^2 + bP.$ 3P = $(2Fc^2 + Fo^2)$ a and b are constantsMax. thermal value (Å2)U33[C(10)]=0.128(3)U33(C24)=0.087(5)Final $A A Rw$0.209, 0.0400.054, 0.145</cod></shif>	Temperature (K)	295	173
Collected Independent (Rint.)26487787Independent (Rint.)2446(0.033)6578 (0.023)Observed (2 σ (I) criterion)19866571Standard reflexions:2 reflexions every 100 reflect.3 every 250 reflect.No variation2% decaySolution and refinement:Fo, full matrix2% full matrixSolution (x10 ⁴)0.206(6)-Parameters:-412Number of variables285396Number of restraints-412Final <shift error="">0.0050.003H atomsFrom difference synthesisN-H and water difference synthesis, rigid methyls, others ridingWeighting-schemeEmpirical as to give no trends in avital-counciesMax. thermal value (Å²)U33(C(10)]=0.128(3)U33(C24)=0.087(5)Final AF peaks and holes(eÅ⁻³)0.20, -0.27 0.039, 0.0400.646, -0.614 0.054, 0.145</shift>	Number of reflexions:		
Independent (Rint.)2446(0.033)6578 (0.023)Observed (2 $\sigma(I)$ criterion)19866571Standard reflexions:2 reflexions every 100 reflect.3 every 250 reflect.No variation2% decaySolution and refinementDirect methodsDirect methodsRefinement:Fo, full matrixF ² , full matrixSecondary extinction (x10 ⁴)0.206(6)-Parameters:Number of variables285Number of variables285396Number of restraints-Final <shift error="">0.0050.003H atomsFrom difference synthesisrigid methyls, others ridingWeighting-schemeEmpirical as to give no trends in < $\alpha Q^2F>$ vs. <ifobsi> and <sin<math>\theta/A>wa⁻¹ = $\sigma^2(F^2) + (aP)^2 + bP$. 3P = $(2Fc^2 + Fo^2)$ a and b are constantsMax. thermal value (Å³)U33[C(10)]=0.128(3)U33(C24)=0.087(5)Final A AF peaks and holes(eÅ⁻³)0.20, -0.27 0.039, 0.0400.646, -0.614 0.054, 0.145</sin<math></ifobsi></shift>	Collected	2648	7787
Observed $(2\sigma(I) \text{ criterion})$ 19866571Standard reflexions:2 reflexions every 100 reflect. No variation3 every 250 reflect. 2% decaySolution and refinement	Independent (Rint.)	2446(0.033)	6578 (0.023)
Standard reflexions:2 reflexions every 100 reflect. No variation3 every 250 reflect. 2% decaySolution and refinement 2% decaySolutionDirect methodsRefinement:Fo, full matrixSecondary extinction (x10 ⁴)0.206(6)Parameters:0.206(6)Number of variables285Solution96Number of restraints-412Final <shift error="">0.0050.0050.003H atomsFrom difference synthesisWeighting-schemeEmpirical as to give no trends in <cod>$w^{-1} = \sigma^2(F^2) + (aP)^2 + bP. 3P = (2Fc^2 + Fo^2)$ a and b are constantsMax. thermal value (Å2)U33[C(10)]=0.128(3)U33(C24)=0.087(5)Final ΔF peaks and holes(eÅ^{-3})0.20, -0.27 0.039, 0.0400.646, -0.614 0.054, 0.145</cod></shift>	Observed $(2\sigma(I) \text{ criterion})$	1986	6571
No variation 2% decaySolution and refinementDirect methodsDirect methodsRefinement:Fo, full matrixF², full matrixSecondary extinction (x10 ⁴)0.206(6)-Parameters:Number of variables285396Number of variables285396Number of restraints-Final <shift error="">0.0050.003H atomsFrom difference synthesisN-H and water difference synthesis, rigid methyls, others ridingWeighting-schemeEmpirical as to give no trends in cw⁻¹ = $\sigma^2(F^2) + (aP)^2 + bP, 3P = (2Fc^2 + Fo^2)$Max. thermal value (Å²)U33[C(10)]=0.128(3)U33(C24)=0.087(5)Final ΔF0.20, -0.270.646, -0.614peaks and holes(eÅ⁻³)0.20, -0.270.646, -0.614Final R and Rw0.039, 0.0400.054, 0.145</shift>	Standard reflexions:	2 reflexions every 100 reflect.	3 every 250 reflect.
Solution and refinementDirect methodsDirect methodsSolutionDirect methodsDirect methodsRefinement:Fo, full matrix F^2 , full matrixSecondary extinction (x10 ⁴)0.206(6)-Parameters:-Number of variables285396Number of restraints-412Final <shift error="">0.0050.003H atomsFrom difference synthesisN-H and water difference synthesis, rigid methyls, others ridingWeighting-schemeEmpirical as to give no trends in <cod²f> vs. <ifobsl> and <sinθ λ="">w⁻¹ = $\sigma^2(F^2) + (aP)^2 + bP. 3P = (2Fc^2 + Fo^2)$ a and b are constantsMax. thermal value (Ų) Final ΔF peaks and holes(eÅ⁻³)0.20, -0.27 0.20, -0.270.646, -0.614 0.054, 0.145</sinθ></ifobsl></cod²f></shift>		No variation	2% decay
Solution Refinement: Secondary extinction (x104)Direct methods Fo, full matrix 0.206(6)Direct methods F2, full matrix - -Parameters: Number of variables285 285 - -396 412 0.003Number of restraints Final <shift error="">-412 0.005H atomsFrom difference synthesis <math>< \infty \Delta^2 F> vs. <lfobsl> and <sin0 <math="">\lambda</sin0></lfobsl></math>>N-H and water difference synthesis, rigid methyls, others ridingWeighting-schemeEmpirical as to give no trends in <<math>< \infty \Delta^2 F> vs. <lfobsl> and <sin0 <math="">\lambda</sin0></lfobsl></math>>water constantsMax. thermal value (Å2) Final ΔF peaks and holes(eÅ-3) Final R and Rw0.20, -0.27 0.039, 0.0400.646, -0.614 0.054, 0.145</shift>	Solution and refinement		
Refinement: Secondary extinction (x104)Fo, full matrix 0.206(6) F^2 , full matrix -Parameters: Number of variables285396Number of variables285396Number of restraints Final <shift error="">-412O.0050.0030.003H atomsFrom difference synthesis e^{2}, Find as to give no trends in $e^{-2}(F^2) + (aP)^2 + bP$, $3P = (2Fc^2 + Fo^2)$ a and b are constantsMax. thermal value (Å2) Final ΔF peaks and holes(eÅ^{-3}) Final R and Rw0.20, -0.27 0.039, 0.0400.646, -0.614 0.054, 0.145</shift>	Solution	Direct methods	Direct methods
Secondary extinction $(x10^4)$ 0.206(6)-Parameters:-Number of variables285Number of restraints-Final <shift error="">0.005H atomsFrom difference synthesisWeighting-schemeEmpirical as to give no trends in <model <math="" <sin0="" and="" for="" synthesis="" the="">\lambda>Max. thermal value (Ų)U33[C(10)]=0.128(3)Max. thermal value (Ų)0.20, -0.27peaks and holes(eÅ^{-3})0.20, -0.27Final R and Rw0.039, 0.040</model></shift>	Refinement:	Fo, full matrix	F^2 , full matrix
Parameters:396Number of variables285396Number of restraints-412Final <shift error="">0.0050.003H atomsFrom difference synthesisN-H and water difference synthesis, rigid methyls, others ridingWeighting-schemeEmpirical as to give no trends in <model p=""> vs. <ifobst> and <sin0 <math="">\lambda>w⁻¹ = $\sigma^2(F^2) + (aP)^2 + bP.$ $3P = (2Fc^2 + Fo^2)$ a and b are constantsMax. thermal value (Ų)U33[C(10)]=0.128(3)U33(C24)=0.087(5)Final ΔF peaks and holes(eÅ⁻³)0.20, -0.270.646, -0.614Final R and Rw0.039, 0.0400.054, 0.145</sin0></ifobst></model></shift>	Secondary extinction $(x10^4)$	0.206(6)	-
Number of variables285396Number of restraints-412Final <shift error="">0.0050.003H atomsFrom difference synthesisN-H and water difference synthesis, rigid methyls, others ridingWeighting-schemeEmpirical as to give no trends in <md>$w^{-1} = \sigma^2(F^2) + (aP)^2 + bP.$ $3P = (2Fc^2 + Fo^2)$ a and b are constantsMax. thermal value (Ų)U33[C(10)]=0.128(3)U33(C24)=0.087(5)Final ΔF peaks and holes(eÅ^{-3})0.20, -0.270.646, -0.614Final R and Rw0.039, 0.0400.054, 0.145</md></shift>	Parameters:		
Number of restraints-412Final <shift error="">0.0050.003H atomsFrom difference synthesisN-H and water difference synthesis, rigid methyls, others ridingWeighting-schemeEmpirical as to give no trends in <mod2f> vs. <ifobst> and <sin0 <math="">\lambda>w⁻¹ = $\sigma^2(F^2) + (aP)^2 + bP.$ $3P = (2Fc^2 + Fo^2)$ a and b are constantsMax. thermal value (Ų)U33[C(10)]=0.128(3)U33(C24)=0.087(5)Final ΔF peaks and holes(eÅ⁻³)0.20, -0.270.646, -0.614Final R and Rw0.039, 0.0400.054, 0.145</sin0></ifobst></mod2f></shift>	Number of variables	285	396
Final <shift error="">0.0050.003H atomsFrom difference synthesisN-H and water difference synthesis, rigid methyls, others ridingWeighting-schemeEmpirical as to give no trends in <mod¥f> vs. <lfobsl> and <sinθ <math="">\lambda>w⁻¹ = $\sigma^2(F^2) + (aP)^2 + bP$, $3P = (2Fc^2 + Fo^2)$ a and b are constantsMax. thermal value (Ų)U33[C(10)]=0.128(3)U33(C24)=0.087(5)Final ΔF peaks and holes(eÅ⁻³)0.20, -0.270.646, -0.614Final R and Rw0.039, 0.0400.054, 0.145</sinθ></lfobsl></mod¥f></shift>	Number of restraints	-	412
H atomsFrom difference synthesisN-H and water difference synthesis, rigid methyls, others ridingWeighting-schemeEmpirical as to give no trends in $< \omega \Delta^2 F > vs. < Fobsl> and < \sin\theta/\lambda >w^{-1} = \sigma^2 (F^2) + (aP)^2 + bP. 3P = (2Fc^2 + Fo^2)a and b are constantsMax. thermal value (Ų)Final \Delta Fpeaks and holes(eÅ^{-3})U33[C(10)]=0.128(3)U33(C24)=0.087(5)Final R and Rw0.039, 0.0400.054, 0.145$	Final <shift error=""></shift>	0.005	0.003
Weighting-schemeEmpirical as to give no trends in $< \omega \Delta^2 F>$ vs. < Fobst> and <sin<math>\theta/\lambda>$w^{-1} = \sigma^2 (F^2) + (aP)^2 + bP.$ 3P = $(2Fc^2 + Fo^2)$ a and b are constantsMax. thermal value (Ų)U33[C(10)]=0.128(3)U33(C24)=0.087(5)Final ΔF peaks and holes(eÅ⁻³)0.20, -0.270.646, -0.614Final R and Rw0.039, 0.0400.054, 0.145</sin<math>	H atoms	From difference synthesis	N-H and water difference synthesis
Weighting-schemeEmpirical as to give no trends in $<\omega\Delta^2 F>$ vs. $< Fobs >$ and $<\sin\theta/\lambda>$ $w^{-1} = \sigma^2(F^2) + (aP)^2 + bP.$ $3P = (2Fc^2 + Fo^2)$ a and b are constantsMax. thermal value (Ų)U33[C(10)]=0.128(3)U33(C24)=0.087(5)Final ΔF peaks and holes(eÅ^{-3})0.20, -0.270.646, -0.614Final R and Rw0.039, 0.0400.054, 0.145		······································	rigid methyls, others riding
$\langle \omega \Delta^2 F \rangle$ vs. $\langle Fobs \rangle$ and $\langle \sin \theta / \lambda \rangle$ a and b are constants Max. thermal value (Å ²) U33[C(10)]=0.128(3) U33(C24)=0.087(5) Final ΔF peaks and holes(eÅ ⁻³) 0.20, -0.27 0.646, -0.614 Final R and Rw 0.039, 0.040 0.054, 0.145	Weighting-scheme	Empirical as to give no trends in	$w^{-1} = \sigma^2(F^2) + (aP)^2 + bP \ 3P = (2Fc^2 + Fc^2)$
Max. thermal value (Ų)U33[C(10)]=0.128(3)U33(C24)=0.087(5)Final ΔF peaks and holes(eÅ ⁻³)0.20, -0.270.646, -0.614Final R and Rw0.039, 0.0400.054, 0.145		$<\omega\Delta^2 F>$ vs. $< Fobs >$ and $<\sin\theta/\lambda>$	a and b are constants
Final ΔF 0.20, -0.27 0.646, -0.614 Final R and Rw 0.039, 0.040 0.054, 0.145	Max. thermal value (Å ²)	U33[C(10)]=0.128(3)	U33(C24)=0.087(5)
peaks and holes(eÅ ⁻³) 0.20, -0.27 0.646, -0.614 Final R and Rw 0.039, 0.040 0.054, 0.145	Final ΔF		
Final R and Rw 0.039, 0.040 0.054, 0.145	peaks and holes(eÅ ⁻³)	0.20, -0.27	0.646, -0.614
	Final R and Rw	0.039, 0.040	0.054, 0.145

The structure of $7.H2O.3/2CH_3CH_2OH$ was solved by direct methods and refined anisotropically on F² (SHELXTL).³⁶ Hydrogen atoms for the water molecule and the NH group were located in a difference Fourier synthesis and refined with restrained O-H and N-H bond lengths. Other hydrogen atoms were included as rigid methyl groups or using a riding model. The ethanol molecules were not well resolved, one of them being disordered over two sites and ethanol hydrogen atoms were not considered. The final R(F) was 0.0541, for 439 parameters, 4181 observed reflections and 412 restraints (Table 3) to local symmetry and U components of neighbouring light atoms. The scattering factors were taken from International Tables for Crystallography.³⁷

N-Methoxymethyl-3-acetyl-2-chloroindole **1**. To a suspension of sodium hydride (1.06 g, 2.65 mmol) in dry dimethylformamide (30 mL), was added dropwise a solution of 3-acetyl-2-chloroindole (4.32 g, 2.23 mmol) in the same solvent (60 mL). The mixture was stirred at room temperature for 1 h. After this time, the solution was cooled to 0°C and chloromethylmethyl ether (2.13 g, 2.65 mmol) was slowly added. The solution was allowed to warm at room temperature and stirred for 1 h. Then it was poured into ice/water and the precipitated solid was collected by filtration, air-dried and recrystallized from ethyl acetate/n-hexane to give **1** (85%) m.p. 88-90°C (colourless prisms). ¹H n.m.r. (200 MHz CDCl₃) δ 2.69 (s, 3H, CH₃O), 3.32 (s, 3H, CH₃O), 5.57 (s, 2H, CH₂O), 7.25-7.35 (m, 2H, H-5 + H-6), 7.35-7.5 (m, 1H, H-7), 8.30-8.4 (m, 1H, H-4). ¹³C n.m.r. (50 MHz, CDCl₃) δ 30.9 (CH₃), 56.4 (CH₃O), 74.15 (CH₂O), 109.73 (C-7), 114.3 (C-3), 122.1 and 124.0 (C-5 or C-6), 123.4 (C-4). IR (nujol) v 1642 (C=O) cm⁻¹. MS: m/z (%) (EI positive) 239 (M+2, 31), 237 (M, 100). Anal. Calcd. for C₁₂H₁₂CINO₂ : C, 60.64; H, 5.09; N, 5.89. Found; C, 60.43; H, 4.95; N, 5.97.

[3-(*N*-Methoxymethyl)-2-chloroindolyl]glyoxal hydrate **2**. To a solution of SeO₂ (1.90 g, 16.7 mmol) in H₂O (5 mL) a solution of **1** (2.0 g, 8.4 mmol) in dioxane (40 mL) was added. The resultant solution was heated at 100°C for 24 h and then filtered. The solvent was removed from the filtrate and the crude product was recrystallized from diethyl ether/n-hexane to give **2** (96%) m.p. 96-97°C (oranges prisms) ¹H n.m.r. (300 MHz, DMSO-d₆) δ 3.28 (s, 3H, CH₃O), 5.70 (s, 2H, CH₂O), 5.86 (t, 1H, J=8.1 Hz, CH(OH)₂), 6.53 (d, 2H, J=8.1 Hz, OH), 7.2-7.4 (m, 2H, H-5 + H-6), 7.72 (d, 1H, 1H, J=8.1 Hz, H-7), 8.24 (dd, 1H, J=7.2, 1.2 Hz, H-4). ¹³C n.m.r. (75 MHz, DMSO-d₆) δ 56.0 (CH₃O), 74.0 (CH₂O), 87.8 (CH(OH)₂), 110.4 (C-3), 110.9 (C-7), 121.4 (C-4), 123.0 and 123.9 (C-5 or C-6), 125.3 (C-3a), 132.0 (C-2), 135.5 (C-7a), 189.2 (C=O). IR (nujol) v : 3346 (m), 1641(s) cm⁻¹. MS: m/z (%) (FAB positive, NBA) 252 (M-H₂O+H, 67), 222 (100). Anal. Calcd. for C₁₂H₁₂ClNO₄ : C, 53.44; H, 4.49; N, 5.19. Found: C, 53.57; H, 4.29; N, 5.31. [3-(*N*-Methoxymethyl)-2-azidoindolyl]glyoxal hydrate **3**. A mixture of compound **2** (0.53

[3-(N-Methoxymethyl)-2-azidoindolyl]glyoxal hydrate 3. A mixture of compound 2 (0.53 g, 1.96 mmol), polymeric azide (9.22 g) and dry acetonitrile (50 mL) was stirred at room temperature for 48 h. The mixture was filtered and the solvent was removed under reduced pressure from the filtrate. The crude product was chromatographed on a silica gel column, with diethyl ether as eluent and then recrystallized from diethyl ether/n-hexane to give 3

(80%) m.p. 84°C (orange prisms) ¹H n.m.r. (200 MHz, DMSO-d₆) δ 3.25 (s, 3H, CH₃O), 5.57 (s, 2H, CH₂O), 5.71 (t, 1H, J= 8Hz, CH(OH)₂), 6.62 (d, 2H, J=8 Hz, OH), 7.25-7.31 (m, 2H, H-5, H-6), 7.58-7.65 (m, 1H, H-7), 7.9-8.1 (m, 1H, H-4). ¹³C n.m.r. (50 MHz, DMSO-d₆) δ 56.1 (CH₃O), 73.0 (CH₂O), 88.9 (CH(OH)₂), 104.0 (C-3), 110.8 (C-7), 121.5 (C-4), 122.7 and 123.1 (C-5 or C-6), 123.8 (C-3a), 133.5 (C-7a), 142.1 (C-2), 191.7 (C=O). IR (nujol) v 3334 (m), 2164 (s), 1640 (s) cm⁻¹. MS: m/z (%) (EI positive) 258 (M-H₂O, 20), 230 (26), 143 (100). Anal. Calcd. for $C_{12}H_{12}N_4O_4$: C, 52.17; H, 4.38; N, 20.28. Found: C, 52.29; H, 4.63; N, 20.13.

2-Dimethylamino-5-[3-(N-methoxymethyl)-2-chloroindolyl]-1,5-dihydro-4H-imidazol-4-

one 4. To a cooled -30°C solution of N,N-dimethylguanidine (0.138 g, 1.58 mmol) in anhydrous ethanol (10 mL) were added glacial acetic acid (1 mL) and a solution of compound 2 (0.40 g, 1.58 mmol) in anhydrous ethanol (10 mL). The resultant mixture was stirred at that temperature for 12 h. Then, it was poured in ice-water and extracted with chloroform (5x15 mL). The combined organic layers were concentrated to dryness and the crude product was chromatographed on a silica gel column using first ethyl acetate/nhexane (3:2) and then methanol. Concentration to dryness of the methanolic fractions provided 4. (91%) m.p. 183°C (yellow prisms) ¹H n.m.r. (300 MHz, DMSO-d_c) δ 2.96 (s, 3H, CH₃N), 3.15 (s, 3H, CH₃N), 3.23 (d, 3H, J=0.9 Hz, CH₃O), 5.11 (d, 1H, J=0.9 Hz, H-5), 5.55 (d, 1H, J=11.5 Hz, CH₂O), 5.58 (d, 1H, J=11.5 Hz, CH₂O), 7.09 (dt, 1H, J=7.2, 0.9 Hz, H-6'), 7.20 (d, 1H, J=7.5 Hz, H-7'), 7.24 (dt, 1H, J=7.2, 1,2 Hz, H-5'), 7.63 (d, 1H, J= 8.1 Hz, H-4'). ¹³C n.m.r. (75 MHz, DMSO-d₆) δ 36.0 (CH₃N), 38.0 (CH₃N), 55.7 (CH₃O), 57.5 (C-5), 73.4 (CH₂O), 108.6 (C-3'), 110.4 (C-7'), 118.1 (C-4'), 120.9 and 122.9 (C-5' or C-6'), 124.75 and 124.78 (C-3a' or C-2'), 135.8 (C-7a'), 171.25 (C-2), 185.6 (C-4). IR (nujol) v : 3177(m), 1688 (s), 1602 (s) cm⁻¹. MS: m/z (%) (FAB positive, NBA) 323 (M+2+H, 10), 322 (M+2, 22), 321 (M+H, 54), 286 (M-Cl+H, 4), 285 (M-Cl, 4), 320 (M, 13), 67 (100); (EI positive) 322 (M+2, 1), 320 (M, 3). Anal. Calcd. for $C_{15}H_{17}ClN_4O_2$: C. 56.17; H, 5.34; N, 17.47. Found: C, 56.08; H, 5.43; N, 12.61.

2-Dimethylamino-5-[3-(N-methoxymethyl)-2-aminoindolyl]-4H-imidazol-4-one 5. To a cooled -30°C solution of N,N-dimethylguanidine (0.268 g, 3.08 mmol) in anhydrous ethanol (20 mL) was added dropwise a solution of compound **3** (0.85 g, 3.08 mmol) in anhydrous ethanol/dry tetrahydrofuran (1:1) (20 mL). The mixture was stirred at that temperature for 4 h. The solvent was removed under reduced pressure and the residue was recrystallized from tetrahydrofuran/n-hexane (1:1) to give **5** (95%) m.p. 256°C (red prisms) ¹H n.m.r. (300 MHz, DMSO-d₆) δ 3.29 (s, 3H,CH₃O), 3.31 (s, 3H, CH₃N), 3.44 (s, 3H, CH₃N), 5.47 (s, 2H, CH₂O), 7.10-7.22 (m, 2H, H-5' + H-6'), 7.28-7.38 (m, 1H, H-7'), 8.28-8.38 (m, 1H, H-4'). ¹³C n.m.r. (75 MHz, DMSO-d₆) δ 37.0 (CH₃N), 38.1 (CH₃N), 55.7 (CH₃O), 72.1 (CH₂O), 95.6 (C-3'), 109.1 (C-7'), 122.7 (C-5'), 123.1 (C-4'), 123.8 (C-6'), 126.1 (C-3a'), 138.2 (C-7a'), 157.6 (C-2'), 166.1 (C-2), 181.4 (C-4), 183.3 (C-5). IR (nujol) v 3180 (m), 1663 (m), 1605 (m), 1555 (s) cm⁻¹. MS: m/z (%) (EI positive) 300 (M+1, 6), 299 (M, 34), 99 (100).Anal. Calcd. for C₁₅H₁₇N₅O₂ : C, 60.19; H, 5.72; N, 23.40. Found: C, 59.93; H, 5.87; N, 23.22.

N-Methoxymethyl-3-bromoacetyl-2-chloroindole **6**. To a mixture of compound **1** (0.35 g, 1.47 mmol), AlCl₃ (0.004 g) and dry diethyl ether (20 mL), Br₂ (0.235 g, 1.47 mmol) was slowly added at 0°C under nitrogen. The mixture was allowed to warm to room temperature and stirred for 1 h, and concentrated to *ca*. 10 mL. After cooling, the precipitated solid was collected by filtration and chromatographed on a silica gel column with ethyl acetate/n-hexane (3:2) as eluent to give **6** (75%) m.p. 109-110°C (colourless prisms from ethyl ether/n-hexane). ¹H n.m.r. (300 MHz, CDCl₃) δ 3.34 (s, 3H, CH₃O), 4.55 (s, 2H, CH₂Br), 5.59 (s, 2H, CH₂O), 7.29-7.38 (m, 2H, H-5 + H-6), 7.50-7.41 (m, 1H, H-7), 8.38-8.26 (m, 1H, H-4). ¹³C n.m.r. (75 MHz, CDCl₃) δ 56.5 (CH₂Br), 56.5 (CH₃O), 74.4 (CH₂O), 110.0 (C-7), 111.7 (C-3), 122.2 and 124.5 (C-5 or C-6), 123.8 (C-4), 126.0 (C-3a), 131.1 (C-2), 135.6 (C-7a), 185.7 (C=O). IR (nujol) v 1641 (C=O) cm⁻¹. MS: m/z (%) (EI positive) 319 (M+4, 19), 318 (M+3, 10), 317 (M+2, 56), 316 (M+1, 9), 315 (M, 48). Anal. Calcd. for C₁₂H₁₁BrClNO₂ : C, 45.53; H, 3.50; N, 4.42. Found: C, 45.42; H, 3.39; N, 4.61.

Preparation of the Salt 7. To a mixture of N,N, dimethylguanidine (0.063 g, 0.727 mmol), triethylamine (0.073 g, 0.727 mmol) and dry dimethylformamide (15 mL), a solution of compound 6 (0.23 g, 0.727 mmol) in the same solvent (15 mL) was added at 0°C under nitrogen. The resultant mixture was allowed to warm to room temperature and stirred for 24 h. The solvent was removed under reduced pressure and the solid residue was washed with water (3x15 mL) and extracted with chloroform (3x20 mL). The combined organic layers were dried over anhydrous $MgSO_4$. Filtration and elimination of the solvent gave a crude product which was chromatographed on a silica gel column using methanol/dichloro methane (1:4) as eluent to give 7 (60%) m.p. 235-240°C (d) (red prisms from acetone) ¹H n.m.r. (300 MHz, DMSO-d₆) δ 3.17 (s, 6H, (CH₃)₂N), 3.22 (s, 3H, CH₃O), 3.25 (s, 3H, CH₃O), 3.38 (s, 3H, CH₃N), 3.52 (s, 3H, CH₃N), 5.56 (d, 1H, J=12 Hz, CH₂O), 5.60 (d, 1H, J=12 Hz, CH₂O), 5.75 (s, 3H, CH₂O + H-4'), 6.76 (dd, 1H, J=8.2, 7.6 Hz, H-5'), 7.15 (dd, 1H, J=8.2, 7.4 Hz, H-6'), 7.39 (dd, 1H, J=6.7, 6.7 Hz, H-10), 7.43 (dd, 1H, J=7.6, 7.0 Hz, H-9), 7.66 (d, 1H, J=8.4 Hz, H-7'), 7.78 (d, 1H, J=7.5 Hz, H-8), 8.17 (dd, 1H, J=7.3, 1.5, H-11), 9.57 (s, 1H, NH), 11.9 (s, 1H, NH). ¹³C n.m.r. (75 MHz, DMSO-d₆), 38.0 (N(CH₃)₂), 38.6 (1C, N(CH₃)₂), 39.3 (1C, N(CH₂)₃), 55.5 (CH₃O), 55.9 (CH₃O), 71.9 (C-6a), 73.3 (CH₂O), 74.3 (CH₂O), 105.9 (C-11b), 107.9 (C-3'), 110.9 (C-7'), 112.3 (C-8), 115.4 (g), 117.9 (C-4'), 117.9 (C-4'), 120.9 (C-11), 121.5 (C-5'), 122.7 (C-6'), 122.9 (C-9 + 1C(q)), 123.3 (q), 124.4 (C-10), 125.4 (q), 135.4 (q), 137.3 (s), 143.4 (q), 158.7 (q), 161.6 (q), 171.0 (q), 174.4 (q). IR (nujol) v 3478 (m), 3395 (m) 1591 (m), 1463 (s) cm⁻¹.MS: m/z (%) (FAB positive, NBA) 573 (M+2+H,16), 572 (M+1+H,40), 571 (M+H, 100), 570 (M-1+H, 20); HREIMS C₃₀H₃₁ClN₈O₂ calcd. 570.2258, found 570.2203. Anal. Calcd. for C₁₀H₃₁ClN₈O₂:C, 63.10; H, 5.47; N, 19.62. Found: C, 62.95; H, 5.33; N, 19.51. Suitable crystals for X-ray analysis were obtained by recrystallization from tetrahydrofuran/ ethanol/n-hexane.

N-Methoxymethyl-3-acetyl-2-azidoindole 15. To a solution of 1 (1.50 g, 6.3 mmol) in dimethylsulfoxide (30 mL), NaN₃ (1.0 g, 16.6 mmol) was added. The mixture was stirred

at 45°C for 24 h then poured into ice-water and extracted with diethyl ether (4 x 50 mL). The combined organic layers were dried over anhydrous MgSO₄. Filtration and concentration to dryness afforded the crude product, which was chromatographed on a silica gel column using diethyl ether/n-hexane as eluent to give **15** as oil (65%) ¹H n.m.r. (200 MHz, CDCl₃) δ 2.79 (s, 3H, CH₃CO), 3.18 (s, 3H, CH₃O), 5.50 (s, 2H, CH₂O), 7.37-7.25 (m, 2H, H-5 + H-6), 7.5-7.4 (m, 1H, H-7), 7.9-7.8 (m, 1H, H-4). ¹³C n.m.r. (50 MHz, CDCl₃) δ 31.1 (CH₃CO), 56.3 (CH₃O), 73.0 (CH₂O), 107.0 (C-2), 110.4 (C-7), 119.9 (C-4), 122.8 and 122.9 (C-5 or C-6), 125.0 (C-3a), 133.4 (C-7a), 140.6 (C-2), 193.3 (C=O). IR (CHCl₃) v : 2140 (s), 1722 (s) 1649 (s) cm⁻¹. MS: m/z (%) (FAB positive, NBA) 245 (M+H, 65). Anal. Calcd. for C₁₂H₁₂N₄O₂ : C, 59.01; H, 4.95; N, 22.94. Found: C, 58.77; H, 4.72; N, 23.20.

N-Methoxymethyl-3-acetyl-2-aminoindole 16. To a cooled 0°C solution of triphenyl phosphine (0.53 g, 2.04 mmol) in dry diethyl ether (10 mL), a solution of compound 15 (0.5 g, 2.04 mmol) in the same solvent (10 mL) was added dropwise under nitrogen. The mixture was allowed to warm to room temperature and stirred for 12 h. After cooling, the precipitated solid was collected by filtration and recrystallized from benzene/n-hexane to give the N-methoxymethyl-2-acetyl-3-(triphenylphosphoranylidene) aminoindole : 0.88 g (90%) m.p. 153-152°C (colourless prisms) ¹H n.m.r. (200 MHz, CDCl₃) δ 2.15 (s, 3H, CH₃CO), 3.24 (s, 3H, CH₃O), 5.62 (s, 2H, CH₂O), 7.0-7.15 (m, 2H, H-5 + H-6) 7.20-7.33 $(m, 1H, H-7), 7.34-7.48 (m, 9H, H_m + H_p), 7.49-7.57 (m, H-4), 7.58-7.72 (m, 6H, H_p).$ ¹³C n.m.r. (50 MHz, CDCl₃) δ 30.0 (CH₃CO), 55.8 (CH₃O), 71.7 (CH₂O), 100.9 (C-3), 108.7 (C-7), 118.4 (C-4), 120.0 (C-6), 121.2 (C-5), 127.2 (C-3a), 128.1 $({}^{3}J_{C,P} = 12.8 \text{ Hz}, C_{m})$, 131.0 (${}^{4}J_{C,P}$ = 2.9 Hz, C_n), 132 (${}^{2}J_{C,P}$ = 9.9 Hz, C_o), 133.3 (${}^{1}J_{C,P}$ = 108.4 Hz, C_i), 133.9 ($J_{C,P}$ = 1.24 Hz, C-7a), 154.0 (J_{C-P} = 10.9 Hz, C-2), 189.3 (C=O). ³¹P n.m.r. (CDCl₃) δ 11.74. IR (nujol) v :1619 (C=O)cm⁻¹. MS: m/z (%) (EI positive) 480 (M+2, 6), 479 (M+1, 32), 478 (M,100).Anal. Calcd. for C₃₀H₂₇N₂O₂P : C, 75.30; H, 5.69; N, 5.85. Found: C, 75.18; H, 5.78; N, 5.69.

To a solution of the above mentioned iminophosphorane (1.0 g, 2.1 mmol) in tetrahydrofuran (20 mL), 5% HCl (20 mL) was added. The resultant mixture was stirred at room temperature for 24 h and then concentrated to dryness. A mixture of the remaining solid and 5% HCl (20 mL) was stirred at room temperature for 30 min and then extracted with diethyl ether (2x20 mL). To the aqueous phase, was added 5% NaOH (25 mL) and then extracted with ethyl acetate (3x25 mL). The combined organic layers were dried over anhydrous MgSO₄. Filtration, concentration to dryness, and recrystallization from ethyl acetate gave **16** (98%) m. p. 186-188°C (colourless prisms) ¹H n.m.r. (200 MHz, DMSO-d₆) δ 2.44 (s, 3H, CH₃CO), 3.23 (s, 3H, CH₃O), 5.47 (s, 2H, CH₂O), 7.01 (ddd, 1H, J=7.8, 7.5, 1.5 Hz, H-6), 7.09 (ddd, 1H, J=7.5, 7.5, 0.9 Hz, H-5), 7.32 (d, 1H, J= 7.5 Hz, H-7), 8.04 (s, 2H, NH₂).¹³C n.m.r. (50 MHz, DMSO-d₆) δ 29.7 (CH₃CO), 55.3 (CH₃O), 71.5 (CH₂O), 95.7 (C-3), 109.1 (C-7), 117.7 (C-4), 119.9 (C-6), 121.7 (C-5), 125.9 (C-3a), 133.8 (C-7a), 154.1 (C-2), 190.4 (C=O). IR (nujol) ν : 3304 (m), 3277 (m), 1636 (s) cm⁻¹.

MS: m/z (%) (EI positive) 219 (M+1, 6), 218 (M, 51), 143 (100). Anal. Calcd. for $C_{12}H_{14}N_2O_2$: C, 66.04; H, 6.47; N, 12.84. Found: C, 65.89; H, 6.35; N, 12.71.

N-Methoxymethyl-3-acetyl-2-di(tertbutoxycarbonyl)aminoindole 17. To a cooled at 0°C solution of Boc₂O (2.2 g, 9.2 mmol) in dry tetrahydrofuran (25 mL) a solution of compound 16 (1.0 g, 4.6 mmol) and 4-dimethylaminopyridine (1.12 g, 9.2 mmol) in the same solvent (100 mL) was added dropwise. The resultant solution was stirred at room temperature for 18 h. The solvent was removed under reduced pressure. The solid residue was taken up in diethyl ether (50 mL), washed with 0.1N HCl (20 mL), H₂O (20 mL) and dried over anhydrous $MgSO_4$. After filtration, the solvent was removed under reduced pressure to give a solid, which recrystallized from ethyl acetate/n-hexane gave 17 (85%) m.p. 122-125 (colourless prisms) 1 H n.m.r. (300 MHz, CDCl₃) δ 1.42 (s, 18H, ((CH₃)₃C), 2.54 (s, 3H, CH₃CO), 3.30 (s, 3H, CH₃O), 5.38 (s, 2H, CH₂O), 7.26-7.38 (m, 2H, H-5 + H-6), 7.48-7.56 (m, 1H, H-7), 8.28-8.34 (m, 1H, H-4). ¹³C n.m.r . (75 MHz, CDCl₃) δ 27.8 ((CH₃)₃C), 24.6 (CH₃CO), 56.4 (CH₃O), 73.9 (CH₂O), 84.3 ((CH₃)₃C), 110.1 (C-7), 112.4 (C-3), 122.6 (C-4), 122.9 and 124.0 (C-5 or C-6), 125.0 (C-3a), 134.3 (C-7a), 136.8 (C-2), 149.9 (COO'Bu), 192.8 (C=O). IR (nujol) v : 1805 (s), 1728 (s), 1640 (s) cm⁻¹. MS : m/z (%) (EI positive) 420 (M+2, 3), 419 (M+1, 21), 418 (M, 100), 218 (91). Anal. Calcd. for C₂₂H₃₀N₂O₆ : C, 63.14; H, 7.23; N, 6.69. Found: C, 62.98; H, 7.38; N, 6.55.

N-Methoxymethyl-3-bromoacetyl-2-di(tertbutoxycarbonyl)aminoindole 18. To a heated at 50°C solution of compound 17 (0.1 g, 0.24 mmol) in dry dimethylformamide (10 mL) a solution of Br₂ (0.077 g, 0.48 mmol) in anhydrous methanol (10 mL) was added dropwise. The mixture was stirred at that temperature for 1h and then at room temperature for 12h. After addition of dichloromethane (20 mL), the solution was washed with saturated aqueous NaHCO₃ (2x10 mL). The organic layer was dried over anhydrous MgSO₄ and filtered. Concentration to dryness yielded a crude material which was chromatographed on a silica gel column with diethyl ether/n-hexane as eluent to give 18 (84%) m.p. 118-120°C (colourless prisms from diethyl ether/n-hexane). ¹H n.m.r. (300 MHz, CDCl₃) δ 1.40 (s, 18H, (CH₃)₃C), 3.27 (s, 3H, CH₃O), 4.41 (s, 2H, CH₂Br), 5.38 (s, 2H, CH₂O), 7.28-7.40 (m, 2H, H-5 + H-6), 7.52-7.60 (m, 1H, H-7), 8.09-7.17 (m, 1H, H-4). ¹³C n.m.r. (75 MHz, $CDCl_{1}$ δ 27.7 ((CH₃)₃C), 34.2 (CH₂Br), 56.4 (CH₃O), 74.0 (CH₂O), 84.5 ((CH₃)₃C) , 109.5 (C-3), 110.7 (C-7), 122.0 (C-4), 123.3 and 124.2 (C-5 or C-6), 124.3 (C-3a), 134.2 (C-7a), 138.0 (C-2), 149.6 (COO'Bu), 185.5 (C=O). IR (nujol) v: 3452 (m), 1798 (s), 1766 (m), 1722 (m), 1675 (s) cm⁻¹. MS : m/z (%) (IE positive) 498 (M+2, 2), 496 (M, 2), 398 (18), 396 (19), 296 (55), 143 (57), 57 (100). Anal. Calcd. for C₂₂H₂₀BrN₂O₆: C, 53.13; H, 5.88; N, 5.63. Found: C, 53.18; H, 5.91; N, 5.70.

N-Methoxymethyl-3-bromoacetyl-2-tertbutoxycarbonylaminoindole 19.

Method A. To a heated at 50°C solution of compound 17 (0.1 g, 0.24 mmol) in dry chloroform (15 mL) was added a solution of Br_2 (0.038 g, 0.24 mmol) in the same solvent (5 mL). After addition, the solution was stirred at room temperature for 12 h, then washed with saturated aqueous NaHCO₃ (10 mL) and water (10 mL). The organic layer was dried over MgSO₄. After filtration, the solvent was removed under reduced pressure and the

resulting solid was chromatographed on a silica gel column using diethyl ether/n-hexane (7:3) as eluent to give 19 in 70% yield.

Method B. To a cooled at 0°C solution of compound 18 (0.4 g, 0.80 mmol) in dry dichloromethane (10 mL), trifluoroacetic acid (0.27 g, 2.4 mmol) was added dropwise under nitrogen. After addition, the solution was allowed to warm to room temperature and stirring continued for 2 h. Then, dichloromethane (10 mL) was added and the resultant solution was washed with saturated aqueous NaHCO₃ (20 mL) and brine (20 mL). The organic layer was dried over anhydrous MgSO4. After filtration, the solvent was removed under reduced pressure and the crude product was chromatographed on a silica gel column using diethyl ether/n-hexane (7:3) as eluent to give 19 as oil in 81% yield: ¹H n.m.r. (200 MHz, $CDCl_3$) δ 1.44 (s, 9H, (CH₃)₃C), 3.03 (s, 3H, CH₃O, 4.39 (s, 2H, CH₂Br), 5.49 (s, 1H, CH₂O), 7.1-7.3 (m, 2H, H-5 + H-6), 7.37-7.46 (m, 1H, H-7), 7.51-7.65 (m, 1H, H-4). ¹³C n.m.r. (50 MHz, CDCl₃) & 27.9 ((CH₃)₃C), 35.0 (CH₂Br), 56.2 (CH₃O), 76.5 (CH₂O), 82.6 ((CH₃)₃C), 102.1 (C-3), 118 (C-7), 120.0 (C-4), 122.9 and 123.2 (C-5 or C-6), 123.7 (C-3a), 133.8 (C-7a), 143.9 (C-2), 152.6 (COO'Bu), 187.4 (C=O). IR (nujol) v : 3268 (m), 1733 (s), 1620 (s) cm⁻¹. MS : m/z (%) (EI positive) 398 (M+2, 33), 396 (M, 34) 296 (75). Anal. Calcd. for C₁₇H₂₁BrN₂O₄: C, 51.40; H, 5.33; N, 7.05. Found: C, 51.27; H, 5.21; N, 7.22.

2-Dimethylamino-6-methoxymethylimidazo[4',5':3,4]pyrido[2,3-b]indol-4-one **20**. To a solution of the indole derivative **19** (0.397 g. 1mmol) in anhydrous ethanol (50 mL) was added dropwise a solution of N,N-dimethylguanidine (0.174 g, 2 mmol) in the same solvent (100 mL) at 0°C under nitrogen. The reaction mixture was allowed to warm at room temperature and stirred for 3 h. The solvent was removed under reduced pressure and the residual material was chromatographed on a silica gel column eluting with dichloromethane /ethanol (9:1) to give **12** (0.25 g, 80%). ¹H n.m.r.(300 MHz, DMSO-d₆) δ 3.10 (s, 6H, (CH₃)₂N) 3.24 (s, 3H, CH₃O), 5.75 (s, 2H, CH₂O), 7.19 (dt, J=7.5, 7.5, 0.9 Hz, H-9), 7.30 (dt, J=7.5, 7.5, 0.9 Hz, H-8), 7.54 (d, J=8.1 Hz, H-7), 8.12 (d, J=7.5 Hz, H-10). ¹³C n.m.r. (75 MHz, DMSO-d₆) δ 38.0 (CH₃)₂N, 55.6 (CH₃O), 72.0 (CH₂O), 99.4 (q), 109.9 (C-7), 116.0 (q), 119.8 (C-9), 120.7 (C-10), 121.0 (q), 121.9 (q), 122.9 (C-8), 136.9 (q), 148.2 (q), 151.5 (q), 154.6 (q). IR (nujol) v : 3344(m), 1684(m), 1597(s) cm⁻¹. MS: m/z(%) (FAB positive, NBA) 312 (M+H, 100); (EI positive) 312 (M+1, 23), 311 (M, 100), 280 (49), 266 (22). Anal. Calcd. for C₁₆H₁₇N₅O₂ : C, 61.72; H, 5.50; N, 22.49. Found: C, 61.58; H, 5.63; N, 22.60.

Acknowledgements:

We gratefully acknowledge the financial support of the Dirección General de Investigación Científica y Técnica (PB95-1019 and PB96-0001-C03-02). One of us (M.A.S.) also thanks the Ministerio de Educación y Cultura for a scholarship.

References:

- Albizati KF, Martin VA, Agharahimi MR, Stolze DA Synthesis of Marine Natural Products in Biorganic Marine Chemistry. Scheuer PJ, editors. Berlin: Springer, 1992;6:158-248.
- (2) Kobayashi J, Tsuda M, Murayama T, Nakamura H, Ohizumi Y, Ishibashi M, Iwamura M, Otha T, Nozoe, S Tetrahedron 1990;46:5579-5586.
- (3) Commerço A, Gueremy C Tetrahedron Lett. 1991;32:1419-1422.
- (4) He H, Faulkner DJ, Lee A, Clardy J J. Org. Chem. 1992;57:2176-2178.
- (5) Faulkner DJ Nat. Prod. Rep. 1992;9:323-364.
- (6) Alvi KA, Peters BM, Hunter LM, Crews P Tetrahedron 1993;49:329-336.
- (7) Little TL, Webber SE J. Org. Chem. 1994;59:7299-7305.
- (8) Chan GW, Mong S, Hemling ME, Freyer AJ, Offen PH, DeBrosse CW, Sarau HM, Westley JW J. Nat. Prod. 1993;56:116-121.
- (9) Moquin C, Guyot M Tetrahedron Lett. 1984;25:5047-5048.
- (10) Moquin-Pattey C, Guyot, M. Tetrahedron 1989;45:3445-3450.
- (11) Helbecque N, Berbier JL, Henichard JP, Moquin-Pattey C, Guyot M Cancer Biochem. Biophys. 1987; 9: 271-279.
- (12) Achab S, Guyot M, Potier P Tetrahedron Lett. 1993;34:2127-2130.
- (13) Achab S, Guyot M, Potier P Tetrahedron Lett. 1995;36:2615-2618.
- (14) Choshi T, Yamada S, Sugino E, Kuwada T, Hibino S Synlett. 1995;147-148.
- (15) Choshi T, Yamada S, Sugino E, Kuwada T, Hibino S J Org. Chem. 1995;60:5899-5904.
- (16) Molina P, Fresneda PM, Almendros P Tetrahedron Lett. 1992;33:4491-4494.
- (17) Molina P, Almendros P, Fresneda PM Tetrahedron 1994;50:2241-2254.
- (18) Molina P, Almendros P, Fresneda PM Tetrahedron Lett. 1994;35:2235-2236.
- (19) Lawson A J. Chem. Soc. 1956:307-310
- (20) Storey B, Sullivan WW, Moyer CL J. Org. Chem. 1964;29:3118-3120
- (21) Laucini GC, Lazzari E J. Heterocyclic. Chem. 1966;3:152-154.
- (22) Cavallieri B, Ballotta R, Lancini GC J. Heterocyclic. Chem. 1972;9:979-984
- (23) Nishimura T, Kitajima K J. Org. Chem. 1979;44:818-824
- (24) Dalkafouki A, Ardisson J, Kunesch N, Lacombe L, Poisson JE Tetrahedron Lett. 1991;32:5325-5328
- (25) Monge A, Palop J, Ramirez C, Font M, Fernandez-Alvarez E Eur. J. Med. Chem. 1991;26,179-188.
- (26) Hassner A, Stern M Angew. Chem. Int. Ed. Eng. 1986;25:478-479.
- (27) Guyot M, Meyer M Tetrahedron Lett. 1986;27:2621-2622.
- (28) Allen FH, Davies JE, Galloy JJ, Johnson O, Kennard O, Macrae CF, Mitchell EM, Mitchell JF, Smith JM, Watson DG J. Chem. Info. Compt. Sci. 1991;31:187-204.
- (29) Allen FH, Kennard O, Watson DG, Brammer L, Orpen AG, Taylor R J. Chem. Soc. Perkin Trans 2 1987;
- (30) Molina P, Tarraga A. Comprehensive Organic Functional Group Transformations. Katritzky AR, Meth-Cohn O, Rees CW editors. Cambridge, Pergamon 1995;5:961-1020.
- (31) Altomare A, Burla MC, Carmalli M, Cascarano G, Giacovazzo, Guagliardi A, Polidori G SIR92, J. Appl. Cryst. 1994;27:435.
- (32) International Tables for X-Ray Crystallography.Birmingham, Kynoch Press.1974.
- (33) Hall SR, Flack HD, Stewart JM. 'Xtal3.2'. University of Western Australia editors.Lamb: Perth, 1994.
- (34) Martínez-Ripoll JM, Cano FH. "PESOS" Unpublished Program.
- (35) Nardelli M Comput. Chem. 1983;7:95-98.
- (36) Sheldrick GM. University of Göttingen. 1993.
- (37) International Tables for Crystallography. Vol C, table 6.1.1.4. 1992.