

## PAPER

View Article Online  
View Journal | View IssueCite this: *RSC Adv.*, 2017, 7, 18690Singlet oxygen generation properties of an inclusion complex of cyclic free-base porphyrin dimer and fullerene C<sub>60</sub><sup>†</sup>Yousuke Ooyama,<sup>a</sup> Toshiaki Enoki,<sup>a</sup> Joji Ohshita,<sup>a</sup> Takuya Kamimura,<sup>b</sup> Shuwa Ozako,<sup>b</sup> Taro Koide<sup>b</sup> and Fumito Tani<sup>\*b</sup>

To gain insight into the singlet oxygen (<sup>1</sup>O<sub>2</sub>) generation properties of supramolecular complexes of cyclic free-base porphyrin dimer with fullerene C<sub>60</sub>, we evaluated the <sup>1</sup>O<sub>2</sub> quantum yield ( $\Phi_{\Delta}$ ) and rate constant ( $K_{\text{obs}}$ ) of <sup>1</sup>O<sub>2</sub> generation for a cyclic free-base porphyrin dimer (CPD) linked by butadiyne bearing four 4-pyridyl groups and its inclusion complex (C<sub>60</sub>⊂CPD) with C<sub>60</sub>. We demonstrate that CPD and C<sub>60</sub>⊂CPD possess the ability to generate <sup>1</sup>O<sub>2</sub> under visible light irradiation. Moreover, it was found that the  $\Phi_{\Delta}$  value of C<sub>60</sub>⊂CPD is lower than that of CPD. Based on the kinetic and thermodynamic consideration concerning the electron transfer processes between the porphyrin dimer and C<sub>60</sub>, this work revealed that the lower  $\Phi_{\Delta}$  value of the C<sub>60</sub> inclusion complex would be attributed to the formation of the charge-separated state  $\text{C}_{60}^{\cdot-}\text{-CPD}^{\cdot+}$ , leading to a low intersystem crossing (ISC) efficiency for the formation of the triplet excited state <sup>3</sup>(CPD)\*.

Received 6th March 2017  
Accepted 22nd March 2017

DOI: 10.1039/c7ra02699d

rsc.li/rsc-advances

## Introduction

Photosensitizers possessing the ability to generate singlet oxygen (<sup>1</sup>O<sub>2</sub>) have created considerable interest in recent years from the viewpoint of fundamental studies in photochemistry and their potential applications in photodynamic therapy (PDT).<sup>1–6</sup> These photosensitizers generally produce <sup>1</sup>O<sub>2</sub> through the following processes: initially the photosensitizer absorbs light ( $h\nu$ ) to generate the singlet excited state of the photosensitizer (<sup>1</sup>S\*), then the photoexcited sensitizer (<sup>1</sup>S\*) undergoes intersystem crossing (ISC) to generate the triplet excited state (<sup>3</sup>S\*). Subsequent energy transfer from the photoexcited sensitizer (<sup>3</sup>S\*) to triplet oxygen (<sup>3</sup>O<sub>2</sub>) produces <sup>1</sup>O<sub>2</sub>. Thus, to generate <sup>3</sup>S\* efficiency is one of the most effective strategies to give high <sup>1</sup>O<sub>2</sub> quantum yield ( $\Phi_{\Delta}$ ). In particular, porphyrin dyes have been regarded as promising candidates for photosensitizers as a result of their strong Soret (400–500 nm) and moderate Q band (500–700 nm) absorption properties, as well as their electrochemical, photochemical and thermal stabilities. The typical photosensitizer, free-base tetraphenylporphyrin (H<sub>2</sub>TPP) exhibits  $\Phi_{\Delta}$  value of 0.6–0.7 (in benzene).<sup>7</sup> Much effort in

molecular design and development of porphyrin photosensitizers have been made to further improve the  $\Phi_{\Delta}$  value so far.<sup>7–11</sup> Some researchers reported that porphyrin–fullerene C<sub>60</sub> dyads as well as boron dipyrromethene (BODIPY)-C<sub>60</sub> dyads exhibits a higher  $\Phi_{\Delta}$  value than each porphyrin or C<sub>60</sub>, which is attributed to the formation of long-lived triplet excited state (<sup>3</sup>S\*) by the effective ISC due to intramolecular energy transfer or




Fig. 1 Chemical structures of cyclic free-base porphyrin dimer (CPD) linked by butadiyne bearing four 4-pyridyl groups, its inclusion complex (C<sub>60</sub>⊂CPD) with C<sub>60</sub> and ABAB porphyrin monomer H<sub>2</sub>PyP as a reference.

<sup>a</sup>Department of Applied Chemistry, Graduate School of Engineering, Hiroshima University, Higashi-Hiroshima 739-8527, Japan. E-mail: yooyama@hiroshima-u.ac.jp; Fax: +81-82-424-5494

<sup>b</sup>Institute for Materials Chemistry and Engineering, Kyushu University, 744 Motooka, Nishi-ku, Fukuoka 819-0395, Japan. E-mail: tanif@ms.ifoc.kyushu-u.ac.jp; Fax: +81-92-802-6224

<sup>†</sup> Electronic supplementary information (ESI) available. See DOI: 10.1039/c7ra02699d

electron transfer (charge separation) between photosensitizer and  $C_{60}$ .<sup>11,12</sup>

Recently, we have designed and developed cyclic free-base porphyrin dimer **CPD** linked by butadiyne bearing four 4-pyridyl groups and its inclusion complex  $C_{60} \subset \text{CPD}$  with  $C_{60}$  (Fig. 1).<sup>13</sup> It was expected that  $C_{60} \subset \text{CPD}$  has favorable photochemical and electrochemical properties for PDT through the electrochemical measurements and the transient absorption spectroscopy, based on the fact that the singlet excited state  $C_{60}^1(\text{CPD})^*$  undergoes intrasupramolecular electron transfer to give a completely charge-separated state  $C_{60}^{\cdot-} \text{CPD}^{\cdot+}$ . Thus, in this work, to gain insight into the  $^1\text{O}_2$  generation properties of supramolecular complex of cyclic free-base porphyrin dimer with  $C_{60}$ , we evaluated the  $\Phi_{\Delta}$  and rate constant ( $K_{\text{obs}}$ ) of  $^1\text{O}_2$  generation for **CPD** and  $C_{60} \subset \text{CPD}$ . Here we reveal that cyclic free-base porphyrin dimer and its inclusion complex with fullerene  $C_{60}$  possess the ability to generate  $^1\text{O}_2$  under visible light irradiation, based on the kinetic and thermodynamic consideration concerning the electron transfer processes between the porphyrin dimer and  $C_{60}$ .

## Results and discussion

The cyclic free-base porphyrin dimer **CPD** in  $\text{CH}_2\text{Cl}_2/\text{MeOH}$  exhibits strong Soret band at around 420 nm and relatively weak Q band in the range 500–650 nm (Fig. 2,  $\lambda_{\text{max}}^{\text{abs}}/\text{nm}$  ( $\epsilon/\text{M}^{-1} \text{cm}^{-1}$ ) = 416 (708 000), 514 (31 200), 548 (7400), 587 (9000), 642 (2900)). The molar extinction coefficients ( $\epsilon$ ) of Soret and Q bands for **CPD** are higher than those of **H2PyP** ( $\lambda_{\text{max}}^{\text{abs}}/\text{nm}$  ( $\epsilon/\text{M}^{-1} \text{cm}^{-1}$ ) = 418 (419 000), 513 (19 200), 548 (6200), 588 (6000), 643 (3000))<sup>13c</sup> as an ABAB porphyrin monomer with two pyridyl groups and two phenyl groups. The fact is attributed to the porphyrin dimer structure of **CPD** with two porphyrin units. For the  $C_{60}$  inclusion complex  $C_{60} \subset \text{CPD}$ , it is difficult to obtain its exact absorption spectra because the 1 : 1 complex of **CPD** with  $C_{60}$  is in dissociation equilibrium in solution of  $10^{-5}$  to  $10^{-6}$  M concentration which is suitable for the measurement of photoabsorption spectra of porphyrins. In our previous work, however, we have demonstrated that upon addition of  $C_{60}$  to the solution of the cyclic porphyrin dimer, its Soret band was redshifted with a decrease in intensity, whereas its Q band was slightly redshifted but increased in intensity.<sup>13</sup>  $^1\text{O}_2$  generation by **CPD**,  $C_{60} \subset \text{CPD}$  or **H2PyP** in  $\text{CH}_2\text{Cl}_2/\text{MeOH}$  (=1/1, v/v) was evaluated by monitoring




Fig. 2 Photoabsorption spectra of **CPD** and **H2PyP** in  $\text{CH}_2\text{Cl}_2/\text{MeOH}$ .

the photoabsorption spectral change of the known  $^1\text{O}_2$  scavenger 1,3-diphenylisobenzofuran (DPBF) accompanied by the reaction of DPBF with the generated  $^1\text{O}_2$ , that is, DPBF can trap  $^1\text{O}_2$  through its photooxidation.<sup>14</sup>  $\text{CH}_2\text{Cl}_2/\text{MeOH}$  was bubbled with air for 15 min. The air-saturated solution containing **CPD**,  $C_{60} \subset \text{CPD}$  or **H2PyP** and DPBF was irradiated with 509 nm ( $300 \mu\text{W cm}^{-2}$ ,  $\epsilon = 27\,300 \text{ M}^{-1} \text{cm}^{-1}$  @  $\lambda^{\text{abs}} = 509 \text{ nm}$  for **CPD** and  $\epsilon = 17\,000 \text{ M}^{-1} \text{cm}^{-1}$  @  $\lambda^{\text{abs}} = 509 \text{ nm}$  for **H2PyP**, respectively) obtained by passage of xenon light through monochromator. For both **CPD** and  $C_{60} \subset \text{CPD}$  as well as **H2PyP** the absorption band of DPBF at around 410 nm decreased with the increase in the photoirradiation time (Fig. 3), which indicate the reaction of DPBF with  $^1\text{O}_2$  generated upon the excitation of the porphyrin dimer. To gain insight into the effect of the cyclic porphyrin dimers on the efficiency of DPBF photooxidation, the changes in optical density ( $\Delta\text{OD}$ ) of DPBF are plotted against the photoirradiation time (Fig. 4a), and the slope ( $m_{\text{sl}}$ ) is used to estimate the  $\Phi_{\Delta}$  value for **CPD**,  $C_{60} \subset \text{CPD}$  and **H2PyP**. The  $m_{\text{sl}}$  value ( $-1.5 \times 10^{-2}$ ) of **H2PyP** is larger than those of **CPD** ( $-1.2 \times 10^{-2}$ ) and  $C_{60} \subset \text{CPD}$  ( $-9.8 \times 10^{-3}$ ). Moreover, it was revealed that the  $m_{\text{sl}}$  value of **CPD** is larger than that of  $C_{60} \subset \text{CPD}$ . Thus, the  $\Phi_{\Delta}$  values of **CPD**,  $C_{60} \subset \text{CPD}$  and **H2PyP** were estimated by the relative method using Rose Bengal (RB) ( $\Phi_{\Delta} = 0.80$ ,  $m_{\text{sl}} = -1.5 \times 10^{-2}$ , see Fig. S1†) in methanol<sup>15</sup> as the standard (Table 1). The  $\Phi_{\Delta}$  value of **CPD**,  $C_{60} \subset \text{CPD}$  and **H2PyP** is 0.62, 0.52 and 0.91 respectively, which is in good agreement with the  $m_{\text{sl}}$  value. This result suggests that as for the ABAB porphyrin monomer **H2PyP** the ISC efficiency from  $^1\text{S}^*$  to the  $^3\text{S}^*$  may be higher than in the cyclic free-base porphyrin dimer **CPD**. Moreover, it is worth noting that the  $\Phi_{\Delta}$  value of  $C_{60} \subset \text{CPD}$  is lower than that of **CPD**. Our previous work demonstrates that the decay of photoexcited





Fig. 3 Photoabsorption spectral changes for the photooxidation of DPBF (Abs. = ca. 1.0) using (a) **CPD** ( $1.3 \times 10^{-6} \text{ M}$ ), (b)  $C_{60} \subset \text{CPD}$  ( $1.0 \times 10^{-6} \text{ M}$ ) and (c) **H2PyP** ( $1.4 \times 10^{-6} \text{ M}$ ) as photosensitizer under photoirradiation with 509 nm ( $300 \mu\text{W cm}^{-2}$ ) in  $\text{CH}_2\text{Cl}_2/\text{MeOH}$  (=1/1, v/v).






**Fig. 4** (a) Plots of  $\Delta OD$  for DPBF against the photoirradiation time for the photooxidation of DPBF using CPD,  $C_{60}$ -CPD, H2PyP or Rose Bengal as photosensitizers under photoirradiation with 509 nm ( $300 \mu W cm^{-2}$ ) in  $CH_2Cl_2/MeOH$  (=1/1, v/v) and MeOH, respectively. (b) Plots of  $\ln(C_t/C_0)$  for DHN against the photoirradiation time for the photooxidation of DHN using CPD,  $C_{60}$ -CPD, H2PyP or Rose Bengal as photosensitizers under photoirradiation with visible light ( $>385$  nm,  $30 mW cm^{-2}$ ) in  $CH_2Cl_2/MeOH$  (=1/1, v/v).

**Table 1**  $^1O_2$  quantum yield ( $\Phi_\Delta$ ) and first-order rate constant ( $K_{obs}$ ) for the photooxidation of DPBF and DHN in  $CH_2Cl_2/MeOH$  (=1/1, v/v), respectively, using CPD,  $C_{60}$ -CPD or H2PyP as photosensitizer

Photosensitizer	$\Phi_\Delta^a$	$K_{obs}^b/min^{-1}$
CPD	0.62	$6.6 \times 10^{-3}$
$C_{60}$ -CPD	0.52	$5.8 \times 10^{-3}$
H2PyP	0.91	$9.5 \times 10^{-3}$

<sup>a</sup>  $^1O_2$  quantum yield (relative decomposition rate of DPBF), with Rose Bengal (RB) as standard ( $\Phi_\Delta = 0.80$  in methanol,<sup>15</sup> see Fig. S1) and 3-diphenylisobenzofuran (DPBF) as  $^1O_2$  scavenger. <sup>b</sup> First-order rate constant for the reaction of DHN with  $^1O_2$  generated upon photoexcitation of the photosensitizer. The  $K_{obs}$  for RB is  $5.3 \times 10^{-3} min^{-1}$  (see Fig. S2).


state of  $C_{60}$ -CPD has two steps (Fig. 5): the first step has a lifetime of 18 ps, which corresponds to the disappearance of the singlet excited state of  $C_{60}^1(CPD)^*$  (ca.  $-3.6$  eV), that is,  $C_{60}^1(CPD)^*$  undergoes intrasupramolecular electron transfer to give a completely charge-separated state  $C_{60}^{1-}-CPD^{2+}$  (ca.  $-3.7$  eV).<sup>13</sup>  $C_{60}^{1-}-CPD^{2+}$  decays with a lifetime of 470 ps to the ground state. The singlet excited state  $C_{60}^1(CPD)^*$  has a slower ISC to its triplet excited state  $C_{60}^3(CPD)^*$  (ca.  $-4.0$  eV), in addition, the  $C_{60}^3(CPD)^*$  would undergo energy transfer to  $C_{60}$ , leading to the



**Fig. 5** The photodynamics of  $C_{60}$ -CPD and  $^1O_2$  generation.

formation of  $^3C_{60}^*-(CPD)$ . Thus, on the basis of the photodynamics of the cyclic free-base porphyrin dimer and its inclusion complex with  $C_{60}$ , the lower  $\Phi_\Delta$  value of the  $C_{60}$  inclusion complex would be attributed to the formation of charge-separated state, leading to low ISC efficiency because the ISC is in kinetically competition with the intrasupramolecular electron transfer, that is, the formation of triplet excited state  $^3(CPD)^*$  is in kinetically unfavorable compared to that of the charge-separated state  $C_{60}^{1-}-CPD^{2+}$ .

In order to evaluate the photosensitizing ability of the cyclic free-base porphyrin dimer and its inclusion complex with  $C_{60}$ , the  $\ln(C_t/C_0)$  is plotted against the photoirradiation time, where  $C_t$  is a concentration of 1,5-dihydroxynaphthalene (DHN) at the reaction time ( $t$ ) and  $C_0$  is the initial concentration of DHN before photoirradiation.  $CH_2Cl_2/MeOH$  (=1/1, v/v) were bubbled with air for 15 min. The air-saturated solution containing CPD or  $C_{60}$ -CPD and DHN was irradiated with visible light ( $>385$  nm,  $30 mW cm^{-2}$ ) obtained by passage of xenon light through a 385 nm long path filter. The photoabsorption spectral changes for the photooxidation of DHN using CPD,  $C_{60}$ -CPD or H2PyP under photoirradiation with the visible light in  $CH_2Cl_2/MeOH$  (=1/1, v/v) are shown in Fig. 6. Evidently, the absorption band of DHN at around 300 nm decreased with the increase in the photoirradiation time. The plots of  $\ln(C_t/C_0)$  against the photoirradiation time indicate that for CPD,  $C_{60}$ -CPD and H2PyP the  $\ln(C_t/C_0)$  decreased almost linearly with the increase in the photoirradiation time (Fig. 4b). Thus, this result indicates the  $\ln(C_t/C_0)$  bears a linear relationship with the photoirradiation time to provide the first-order rate constants ( $K_{obs}$ ) for the photooxidation of DHN using the cyclic free-base porphyrin dimer or its inclusion complex with  $C_{60}$  as the



**Fig. 6** Photoabsorption spectral changes for the photooxidation of DHN ( $1.0 \times 10^{-4}$  M) using (a) CPD ( $2.5 \times 10^{-6}$  M), (b)  $C_{60}$ -CPD ( $2.5 \times 10^{-6}$  M) and (c) H2PyP ( $5.0 \times 10^{-6}$  M) as photosensitizer under photoirradiation with visible light ( $>385$  nm,  $30 mW cm^{-2}$ ) in  $CH_2Cl_2/MeOH$  (=1/1, v/v).



photosensitizer (Table 1). Obviously, the higher  $K_{\text{obs}}$  values of the cyclic free-base porphyrin dimer and its inclusion complex with  $\text{C}_{60}$  relative to RB (see Fig. S2†) are due to the contribution of the strong Soret band of the cyclic free-base porphyrin skeleton, although the  $K_{\text{obs}}$  values of **CPD** and  $\text{C}_{60}\text{CPD}$  are lower than that of **H2PyP** ( $9.5 \times 10^{-3} \text{ min}^{-1}$ ). It is worth noting here that the  $K_{\text{obs}}$  value ( $6.6 \times 10^{-3} \text{ min}^{-1}$ ) of **CPD** is greater than that ( $5.8 \times 10^{-3} \text{ min}^{-1}$ ) of  $\text{C}_{60}\text{CPD}$ . Therefore, this result demonstrates that **CPD** exhibits more efficient photosensitizing ability due to the effective ISC compared to  $\text{C}_{60}\text{CPD}$ .

In addition, we performed an electron paramagnetic resonance (EPR) method with 2,2,6,6-tetramethyl-4-piperidone (4-oxo-TEMP) as the spin-trapping agent, which can react with  $^1\text{O}_2$  to produce 4-oxo-TEMPO as a stable nitroxide radical.<sup>16</sup> When the air-saturated solution containing **CPD**,  $\text{C}_{60}\text{CPD}$  or **H2PyP** and 4-oxo-TEMP was irradiated with visible light ( $>385 \text{ nm}$ ,  $30 \text{ mW cm}^{-2}$ ) obtained by passage of xenon light through a 385 nm long path filter, for both the free-base porphyrin dimer and its inclusion complex with  $\text{C}_{60}$  as well as **H2PyP** the ESR spectra of 4-oxo-TEMPO were clearly observed as a characteristic 1 : 1 : 1 triplet (Fig. 7). Consequently, this work demonstrated that the cyclic free-base porphyrin dimer and its inclusion complex with  $\text{C}_{60}$  possess the ability to generate  $^1\text{O}_2$  under visible light irradiation.




Fig. 7 The ESR spectra of 4-oxo-TEMPO which is formed by the reaction of 4-oxo-TEMP with  $^1\text{O}_2$  which was generated by (a) **CPD**, (b)  $\text{C}_{60}\text{CPD}$  and (c) **H2PyP** under irradiation with visible light (temperature 298 K, microwave power 1 mW, microwave frequency 9.439 GHz, and field modulation 0.2 mT at 100 kHz). The air-saturated  $\text{CH}_2\text{Cl}_2$  solution containing **CPD** ( $2.5 \times 10^{-6} \text{ M}$ ),  $\text{C}_{60}\text{CPD}$  ( $2.5 \times 10^{-6} \text{ M}$ ) or **H2PyP** ( $5.0 \times 10^{-6} \text{ M}$ ) as the photosensitizer and 4-oxo-TEMP (50 mM) as the spin-trapping agent was irradiated with visible light ( $>385 \text{ nm}$ ,  $30 \text{ mW cm}^{-2}$  for 1 h) obtained by passage of xenon light through a 385 nm long path filter, where  $\text{CH}_2\text{Cl}_2$  as low polar solvent was used because it was difficult to obtain a clear ESR signal in polar solvent such as  $\text{CH}_2\text{Cl}_2/\text{MeOH}$  (=1/1, v/v).

## Conclusions

To investigate singlet oxygen ( $^1\text{O}_2$ ) generation properties of cyclic free-base porphyrin dimer and its inclusion complex with fullerene  $\text{C}_{60}$ , we evaluated the  $^1\text{O}_2$  quantum yield ( $\Phi_{\Delta}$ ) and rate constant ( $K_{\text{obs}}$ ) of  $^1\text{O}_2$  generation for cyclic free-base porphyrin dimer **CPD** and its inclusion complex  $\text{C}_{60}\text{CPD}$  with  $\text{C}_{60}$ . It was found that the  $\Phi_{\Delta}$  value of  $\text{C}_{60}\text{CPD}$  is lower than that of **CPD**. The lower  $\Phi_{\Delta}$  value of the  $\text{C}_{60}$  inclusion complex would be attributed to the formation of charge-separated state  $\text{C}_{60}^{\cdot-}\text{CPD}^{\cdot+}$ , leading to low intersystem crossing (ISC) efficiency for the formation of triplet excited state  $^3(\text{CPD})^*$ , although it was expected that the formation of  $\text{C}_{60}^{\cdot-}\text{CPD}^{\cdot+}$  is favorable for  $^1\text{O}_2$  generation. Consequently, this work demonstrates that the cyclic free-base porphyrin dimer and its supramolecular complex with  $\text{C}_{60}$  possess the ability to generate  $^1\text{O}_2$  under visible light irradiation.

## Experimental

### Evaluation of $^1\text{O}_2$ quantum yield

Quantum yield ( $\Phi_{\Delta}$ ) for singlet oxygen ( $^1\text{O}_2$ ) generation by cyclic free-base porphyrin dimer **CPD**, its inclusion complex  $\text{C}_{60}\text{CPD}$  with fullerene  $\text{C}_{60}$  and **H2PyP** in  $\text{CH}_2\text{Cl}_2/\text{MeOH}$  (=1/1, v/v) was evaluated by monitoring the photoabsorption spectral change of the known  $^1\text{O}_2$  scavenger 1,3-diphenylisobenzofuran (DPBF) accompanied by the reaction of DPBF with the generated  $^1\text{O}_2$ , that is, DPBF can trap  $^1\text{O}_2$  through its photooxidation.  $\text{CH}_2\text{Cl}_2/\text{MeOH}$  was bubbled with air for 15 min. The absorbance of DPBF was adjusted to around 1.0 in air-saturated solvent. Concentration of **CPD**,  $\text{C}_{60}\text{CPD}$  or **H2PyP** was adjusted with an absorbance of ca. 0.03 at the irradiation wavelength (509 nm). The air-saturated solution containing the photosensitizer (**CPD**,  $\text{C}_{60}\text{CPD}$  or **H2PyP**) and DPBF was irradiated with 509 nm ( $300 \mu\text{W cm}^{-2}$ ) obtained by passage of xenon light through monochromator. The photoabsorption spectral change of DPBF with the photoirradiation was monitored with an interval of 1 min up to 10 min. The absorption band of DPBF at around 410 nm decreased with the increase in the photoirradiation time. The changes in optical density ( $\Delta\text{OD}$ ) of DPBF are plotted against the photoirradiation time, and the slope is used to estimate the  $\Phi_{\Delta}$  values of **CPD**,  $\text{C}_{60}\text{CPD}$  and **H2PyP**. The  $\Phi_{\Delta}$  values of **CPD**,  $\text{C}_{60}\text{CPD}$  and **H2PyP** were estimated by the relative method using Rose Bengal (RB) ( $\Phi_{\Delta} = 0.80$ ) in methanol as the standard. Therefore, the  $^1\Phi_{\Delta}$  values were calculated according to the following eqn (1):

$$\Phi_{\Delta\text{sam}} = \Phi_{\Delta\text{ref}} \times [(m_{\text{sam}}/m_{\text{ref}}) \times (L_{\text{ref}}/L_{\text{sam}})] \quad (1)$$

where  $\Phi_{\Delta\text{sam}}$  and  $\Phi_{\Delta\text{ref}}$  are the  $^1\text{O}_2$  quantum yield of photosensitizer (**CPD**,  $\text{C}_{60}\text{CPD}$  or **H2PyP**) and RB, respectively,  $m_{\text{sam}}$  and  $m_{\text{ref}}$  are the slope of the difference ( $\Delta\text{OD}$ ) in the change in the absorption maximum wavelength of DPBF (around 410 nm) which are plotted against the photoirradiation time,  $L_{\text{sam}}$  and  $L_{\text{ref}}$  are the light harvesting





efficiency, which is given by  $L = 1 - 10^{-A}$  ("A" is the absorbance at the photoirradiation wavelength).

### Photosensitizing ability

Photosensitizing ability of CPD, C<sub>60</sub>CPD and H2PyP in CH<sub>2</sub>Cl<sub>2</sub>/MeOH was evaluated by plotting the  $\ln(C_t/C_0)$  against the photoirradiation time, where  $C_t$  is a concentration of 1,5-dihydroxynaphthalene (DHN) at the reaction time ( $t$ ) and  $C_0$  is the initial concentration of DHN before photoirradiation. CH<sub>2</sub>Cl<sub>2</sub>/MeOH was bubbled with air for 15 min. The air-saturated solution containing the photosensitizer ( $2.5 \times 10^{-6}$  M for CPD and C<sub>60</sub>CPD,  $5.0 \times 10^{-6}$  M for H2PyP and  $2.5 \times 10^{-6}$  M for RB) and DHN ( $1.0 \times 10^{-4}$  M) was irradiated with visible light ( $>385$  nm,  $30 \text{ mW cm}^{-2}$ ) obtained by passage of xenon light through a 385 nm long path filter. The photooxidation of DHN with the photoirradiation was monitored by following the decrease in the photoabsorption at around 300 nm with an interval of 1 min up to 10 min and then an interval of 5 min up to 30 min. The concentration ( $C_t$ ) of DHN at the reaction time ( $t$ ) was calculated based on Lambert-Beer law ( $A_{\text{DPBF}} = \epsilon cl$ ). The  $\ln(C_t/C_0)$  decreased almost linearly with the increase in the photoirradiation time due to the photooxidation of DHN, that is, the slope was used to estimate the rate constants ( $K_{\text{obs}}$ ).

### <sup>1</sup>O<sub>2</sub> detection by EPR spin-trapping method with 4-oxo-TEMP

The EPR spectra were recorded on a JEOL JES-RE1X spectrometer under the following experimental conditions: temperature 298 K, microwave power 1 mW, microwave frequency 9.439 GHz, and field modulation 0.2 mT at 100 kHz. The air-saturated CH<sub>2</sub>Cl<sub>2</sub> solution containing CPD ( $2.5 \times 10^{-6}$  M), C<sub>60</sub>CPD ( $2.5 \times 10^{-6}$  M) or H2PyP ( $5.0 \times 10^{-6}$  M) as the photosensitizer and 4-oxo-TEMP (50 mM) as the spin-trapping agent was irradiated with visible light ( $>385$  nm,  $30 \text{ mW cm}^{-2}$  for 1 h) obtained by passage of xenon light through a 385 nm long path filter. The ESR spectrum of 4-oxo-TEMPO which is formed by the reaction of 4-oxo-TEMP with <sup>1</sup>O<sub>2</sub>, was clearly observed as a characteristic 1 : 1 : 1 triplet (Fig. 7).

## Notes and references

- 1 J. F. Lovell, T. W. B. Liu, J. Chen and G. Zheng, *Chem. Rev.*, 2010, **110**, 2839.
- 2 M. C. DeRosa and R. J. Crutchley, *Coord. Chem. Rev.*, 2002, **233–234**, 351.
- 3 M. Pawlicki, H. A. Collins, R. G. Denning and H. L. Anderson, *Angew. Chem., Int. Ed.*, 2009, **48**, 3244.
- 4 K. A. Leonard, M. I. Nelen, L. T. Anderson, S. L. Gibson, R. Hilf and M. R. Detty, *J. Med. Chem.*, 1999, **42**, 3942.
- 5 J. M. Dąbrowski and L. G. Arnaut, *Photochem. Photobiol. Sci.*, 2015, **14**, 1765.
- 6 T. Patrice, *Photodynamic Therapy*, Royal Society of Chemistry, 2003.
- 7 (a) R. Bonnett, *Chem. Soc. Rev.*, 1995, **24**, 19; (b) M. Ethirajan, Y. Chen, P. Joshi and R. K. Pandey, *Chem. Soc. Rev.*, 2011, **40**, 340.
- 8 (a) J. P. Belair, C. J. Ziegler, C. S. Rajesh and D. A. Modarelli, *J. Phys. Chem. A*, 2002, **106**, 6445; (b) P. C. Lo, J. D. Huang, D. Y. Y. Cheng, E. Y. M. Chan, W. P. Fong, W. H. Ko and D. K. P. Ng, *Chem.-Eur. J.*, 2004, **10**, 4831; (c) A. Karotki, M. Khurana, J. R. Lepock and B. C. Wilson, *Photochem. Photobiol.*, 2006, **82**, 443; (d) L. Delanaye, M. A. Bahri, F. Tfibel, M.-P. Fontaine-Aupart, A. Mouithys-Mickalad, B. Heine, J. Piette and M. Hoebeke, *Photochem. Photobiol. Sci.*, 2006, **5**, 317; (e) M. Morone, L. Beverina, A. Abboto, F. Silvestri, E. Collini, C. Ferrante, R. Bozio and G. A. Pagani, *Org. Lett.*, 2006, **8**, 2719; (f) M. Khurana, H. A. Collins, A. Karotki, H. L. Anderson, D. T. Cramb and B. C. Wilson, *Photochem. Photobiol.*, 2007, **84**, 1441.
- 9 (a) A. P. Thomas, P. S. S. Babu, S. A. Nair, S. Ramakrishnan, D. Ramaiah, T. K. Chandrashekar, A. Srinivasan and M. R. Pillai, *J. Med. Chem.*, 2012, **55**, 5110; (b) K. Hirakawa, Y. Nishimura, T. Arai and S. Okazaki, *J. Phys. Chem. B*, 2013, **117**, 13490; (c) H. Horiuchi, M. Hosaka, H. Mashio, M. Terata, S. Ishida, S. Kyushin, T. Okutsu, T. Takeuchi and H. Hiratsuka, *Chem.-Eur. J.*, 2014, **20**, 6054; (d) Q. Yu, E. M. Rodriguez, R. Naccache, P. Forgione, G. Lamoureux, F. Sanz-Rodriguez, D. Scheglman and J. A. Capobianco, *Chem. Commun.*, 2014, **50**, 12150; (e) D. Yao, V. Hugues, M. Blanchard-Desce, O. Mongin, C. O. Paul-Roth and F. Paul, *New J. Chem.*, 2015, **39**, 7730.
- 10 (a) M. Prein and W. Adam, *Angew. Chem., Int. Ed.*, 2014, **53**, 6938; (b) H. Shinmori, F. Kodaira, S. Matsugo, S. Kawabata and A. Osuka, *Chem. Lett.*, 2005, **34**, 322; (c) L. G. Arnaut, M. M. Pereira, J. M. Dąbrowski, E. F. F. Silva, F. A. Schaberle, A. R. Abreu, L. B. Rocha, M. M. Barsan, K. Urbńska, G. Stoché and C. M. A. Brett, *Chem.-Eur. J.*, 2014, **20**, 5346; (d) F. Hammerer, G. Garcia, S. Chen, F. Royer, S. Achelle, C. Fiorini-Debuisschert, M.-P. Telulade-Fichou and P. Maillard, *J. Org. Chem.*, 2014, **79**, 1406; (e) J. Schmitt, V. Heitz, A. Sour, F. Bolze, H. Ftouni, J.-F. Nicoud, L. Flamigni and B. Ventura, *Angew. Chem., Int. Ed.*, 2015, **54**, 169.
- 11 M. E. Milanesio, M. G. Alvarez, V. Rivarola, J. J. Silber and E. N. Durantini, *Photochem. Photobiol.*, 2005, **81**, 891.
- 12 (a) L. Huang, X. Yu, W. Wu and J. Zhao, *Org. Lett.*, 2012, **14**, 2594; (b) A. Kamkaew, S. H. Lim, H. B. Lee, L. V. Kiew, L. Y. Chung and K. Burgess, *Chem. Soc. Rev.*, 2013, **42**, 77; (c) L. Huang, X. Cui, B. Therrien and J. Zhao, *Chem.-Eur. J.*, 2013, **19**, 17472.
- 13 (a) H. Nobukuni, Y. Shimazaki, H. Uno, Y. Naruta, K. Okubo, T. Kojima, S. Fukuzumi, S. Seki, H. Sakai, T. Hasobe and F. Tani, *Chem.-Eur. J.*, 2010, **16**, 11611; (b) H. Nobukuni, T. Kamimura, H. Uno, Y. Shimazaki, Y. Naruta and F. Tani, *Bull. Chem. Soc. Jpn.*, 2012, **85**, 862; (c) K. Sakaguchi, T. Kamimura, H. Uno, S. Mori, S. Ozako, H. Nobukuni, M. Ishida and F. Tani, *J. Org. Chem.*, 2014, **79**, 2980; (d) T. Kamimura, K. Ohkubo, Y. Kawashima, H. Nobukuni, Y. Naruta, F. Tani and S. Fukuzumi, *Chem. Sci.*, 2013, **4**, 1451; (e) T. Kamimura, K. Ohkubo, Y. Kawashima, S. Ozako, K. Sakaguchi, S. Fukuzumi and F. Tani, *J. Phys. Chem. C*, 2015, **119**, 25634; (f) Y. Ooyama,



- K. Uenaka, T. Kamimura, S. Ozako, M. Kanda, T. Koide and F. Tani, *RSC Adv.*, 2016, **6**, 16150.
- 14 K. Golinick and A. Griesbeck, *Tetrahedron*, 1985, **41**, 2057.
- 15 W. Wu, J. Sun, X. Cui and J. Zhao, *J. Mater. Chem. C*, 2013, **1**, 4577.
- 16 (a) Y. Yamakoshi, N. Umezawa, A. Ryu, K. Arakane, N. Miyata, Y. Goda, T. Masumizu and T. Nagano, *J. Am. Chem. Soc.*, 2003, **125**, 12803; (b) S. Oriana, S. Aroua, J. O. B. Söllner, X.-J. Ma, Y. Iwamoto and Y. Yamakoshi, *Chem. Commun.*, 2013, **49**, 9302.

