

Regioselective Acylation of (5-Alkylidene-1,3-cyclohexadiene)tricarbonyliron Complexes by the Reaction with Organometallic Reagents


Mitsuhiko Iwakoshi, Soo Ho Ban, Yujiro Hayashi, and Koichi Narasaka*

Department of Chemistry, School of Science, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033


(Received January 22, 1998; CL-980055)

Reaction of [(1,2,3,4- η)-5-alkylidene-1,3-cyclohexadiene]tricarbonyliron and organometallic reagents generates acyl-[(1,2,3,4,5- η)-1-alkylcyclohexadienyl]dicarbonyliron. The acyl group in these complexes migrates to the cyclohexadienyl moiety under carbon monoxide atmosphere, providing [(1,2,3,4- η)-6-acyl-2-alkyl-1,3-cyclohexadiene]tricarbonyliron, which are further transformed to *m*-alkylphenyl ketones by the oxidation with trimethylamine *N*-oxide.

Recently, we have reported the generation of acyl(η^5 -cyclohexadienyl)iron complexes **2a,b** by the reaction of tricarbonyl[(2,3,4,5- η)-2,4-cyclohexadien-1-one]iron **1a** and its *O*-benzyl oxime **1b** with higher order cuprates.^{1,2} In these reactions, **2a,b** are generated by nucleophilic attack of cuprates on the ligand carbon monoxide and the successive treatment with acetic anhydride. These complexes **2a,b** are synthetically important intermediates, and, in fact, they can be converted to *m*-acylated phenols and anilines, which are difficult to be prepared by the Friedel-Crafts acylation.




Generally acyl(η^5 -cyclohexa, and cycloheptadienyl)iron complexes are synthesized by several steps from the readily available tricarbonyl(η^4 -diene)iron derivatives.^{3,4} As the acyl-(η^5 -cyclohexadienyl)iron complexes **2** are generated from tricarbonyl(η^4 -cyclohexadiene)iron derivatives **1** directly, similar acyl(η^5 -cyclohexadienyl)iron complex is expected to be formed from [(1,2,3,4- η)-5-alkylidene-1,3-cyclohexadiene]tricarbonyliron. Based on this consideration, we have examined the conversion of [(1,2,3,4- η)-5-alkylidene-1,3-cyclohexadiene]tricarbonyliron derivatives to acyl(η^5 -cyclohexadienyl)iron complexes, and this method was applied to the regioselective formation of *m*-substituted phenyl ketone derivatives.



[(1,2,3,4- η)-5-Butylidene-1,3-cyclohexadiene]tricarbonyliron (**5**) was easily prepared from tricarbonyl[(2,3,4,5- η)-2,4-


cyclohexadiene-1-one]iron (**1a**).⁵ The reaction of **1a** with butyllithium at -78°C for 1 h afforded an alcohol **3** in good yield. The alcohol **3** was treated with aq. HBF_4 in propionic anhydride, giving tetrafluoroborate **4**, which was converted to the triene complex **5** as an inseparable *E* and *Z*-mixture (*E* : *Z* = 3 : 1) after heating in H_2O at 80°C for 3 h.

Reaction of [(1,2,3,4- η)-5-butylidene-1,3-cyclohexadiene]tricarbonyliron (**5**) and methyl lithium in THF at -35°C for 1 h, followed by the addition of aq. NH_4Cl , afforded a thermally unstable iron complex **6a** after extraction with Et_2O and evaporation of the solvent at 0°C . The low field carbonyl peak (260.8 ppm) of ^{13}C -NMR of **6a**⁶ indicates the existence of an acyliron moiety,⁷ which suggests the structure of **6a** as acetyl[(1,2,3,4,5- η)-1-butylcyclohexadienyl]dicarbonyliron. The acetyl group of the dicarbonyliron complex **6a** migrated selectively to the C_5 position of the cyclohexadienyl ligand under carbon monoxide atmosphere in Et_2O at room temperature, and [(1,2,3,4- η)-6-acetyl-2-butyl-1,3-cyclohexadiene]tricarbonyliron (**7a**)⁸ and its isomer **8a** were obtained in 83% total yield (**7a**:**8a** =>10:1) from **5** after Florisil column chromatography.



The dieneiron complexes **7a** and **8a** were transformed to *m*-butylphenyl methyl ketone by oxidation.⁹ That is, when the mixture of **7a** and **8a** was treated with trimethylamine *N*-oxide in *N,N*-dimethylacetamide (DMA) at room temperature, *m*-butylphenyl methyl ketone (**9a**; R = Me) was obtained in 68% yield. On the whole, *m*-butylphenyl methyl ketone (**9a**) was synthesized in 56% total yield from **5** (method A). Alternatively, **9a** was also synthesized regioselectively from **5** almost in the same yield without the conversion of the acyl(dicarbonyl)iron complex **6a** to the tricarbonyliron complex **7a** and **8a**. That is, the direct oxidation of the crude **6a** with trimethylamine *N*-oxide afforded **9a** in 57% yield from **5** (method B).


Various *m*-butylphenyl ketones **9** were prepared by these two methods as listed in Table 1.¹⁰ In addition to methyl and butyllithiums, bulky *s*-butyl and *t*-butyllithiums reacted with **5** to afford **9** in moderate yield. Phenyllithium could be also employed to give a benzophenone derivative **9c**. Except for the reaction of phenyllithium, methods A and B gave the similar results.

**Table 1.** Preparation of *m*-butylphenyl ketones **9** from **5**


R	Yield / % by method A		Yield / % by method B	
	7+8	9	total yield	9
Me ^b	83	68	56	57
<i>n</i> -Bu	91	77	70	72
<i>s</i> -Bu	77	60	46	52
<i>t</i> -Bu	94	64	60	53
Ph	92	81	75	45

^aIn all cases reaction temperature is -78 °C unless otherwise noted.^bReaction temperature is -35 °C.

The reaction of tricarbonyl[(1,2,3,4-η)-5-(1,3-dithian-2-ylidene)-1,3-cyclohexadiene]iron (**10**) and butyllithium was carried out by method A, giving butyl *m*-(1,3-dithian-2-yl)phenyl ketone **11** in 50% yield.



For the reaction of tricarbonyl[5-ethoxycarbonylmethylidene-1,3-cyclohexadiene]iron (**12**), higher order cuprates were found to be the choice of the nucleophile.¹¹ Higher order cuprates prepared from primary, secondary, and tertiary alkyl-lithiums and copper(I) iodide reacted with **12** to provide ethyl (*m*-acylphenyl)acetate derivatives **13** in moderate yield.

**Table 2.** Preparation of ethyl (*m*-acylphenyl)acetate **13** from **12**

R	Yield / %
Me	58
<i>n</i> -Bu	75
<i>s</i> -Bu	60
<i>t</i> -Bu	58

It has been reported that stabilized anions such as 2-lithio-2-methylpropionitrile or 2-lithio-1,3-dithiane react with tricarbonyl-(η⁴-1,3-cyclohexadiene)iron to afford alkylated cyclohexenes by the attack of these anions to the diene moiety. Non-stabilized organolithiums such as butyllithium react with the same iron complex but give several products such as 4-pentenoylcyclo-

hexene, 3-butylcyclohexene, and pentanal in low yield.¹² In the present reaction of [(1,2,3,4-η)-5-alkylidene-1,3-cyclohexadiene]tricarbonyliron, non-stabilized organolithiums or higher order cuprates attack not onto the triene moiety but onto the carbon monoxide of the tricarbonyliron complexes **5**, **10**, and **12** selectively. This acyl group migrates regioselectively to the C₅ position of the cyclohexadienyl ligand, affording *m*-alkylphenyl ketone or ethyl *m*-acylphenylacetate derivatives after the oxidation with trimethylamine *N*-oxide. It should be also mentioned that the obtained *m*-alkylphenyl ketone and ethyl *m*-acylphenylacetate derivatives are both difficult to be synthesized by the Friedel-Crafts acylation from alkylbenzenes and ethyl phenylacetate, respectively.¹³

References and Notes

- S. H. Ban, H. Sakurai, Y. Hayashi, and K. Narasaka, *Chem. Lett.*, **1997**, 699.
- S. H. Ban, Y. Hayashi, and K. Narasaka, preceding paper in this issue.
- a) G. M. Williams and D. E. Rudisill, *J. Am. Chem. Soc.*, **107**, 3357 (1985); b) G. M. Williams, R. A. Fisher, and R. H. Heyn, *Organometallics*, **5**, 818 (1986).
- A. V. Malkov and G. R. Stephenson, *J. Organomet. Chem.*, **489**, C44 (1995); A. J. Pearson and K. Srinivasan, *J. Org. Chem.*, **57**, 3965 (1992).
- A. Salzer, H. Schmalle, R. Stauber and S. Streiff, *J. Organomet. Chem.*, **408**, 403 (1991); R. J. H. Cowles, B. F. G. Johnson, J. Lewis, and A. W. Parkins, *J. Chem. Soc., Dalton Trans.*, **1972**, 1768.
- 6a**, ¹H NMR (500 MHz, CDCl₃, -10 °C) δ = 0.84 (3H, t, *J* = 6.9 Hz), 1.13-1.35 (3H, m), 1.51-1.70 (2H, m), 1.93-2.05 (1H, m), 1.99 (1H, d, *J* = 14.3 Hz), 2.32 (3H, s), 2.50 (1H, dd, *J* = 14.3, 5.7 Hz), 3.41-3.48 (1H, m), 4.26-4.35 (1H, m), 4.59-4.68 (1H, m), 6.12 (1H, t, *J* = 5.1 Hz); ¹³C NMR (125 MHz, CDCl₃, -10 °C) δ = 260.8 (RC=O), 220.1, 209.4 (CO); IR (KBr) 2000, 1940, 1645 cm⁻¹.
- The ¹³C-resonance of the acetyl carbon of acetyldicarbonylcyclopentadienyliron is reported to be 254.4 ppm. L. F. Farnell, E. W. Randall, and E. Rosenberg, *J. Chem. Soc., Chem. Commun.*, **1971**, 1078.
- 7a**, ¹H NMR (500 MHz, CDCl₃) δ = 0.95 (3H, t, *J* = 7.34 Hz), 1.39-1.46 (2H, m), 1.52-1.60 (1H, m), 1.62-1.70 (1H, m), 1.86 (1H, ddd, *J* = 15.3, 8.8, 2.3 Hz), 2.05 (1H, ddd, *J* = 15.2, 5.3, 3.3 Hz), 2.12 (3H, s), 2.27 (2H, t, *J* = 7.7 Hz), 2.44 (1H, ddd, *J* = 8.9, 5.4, 1.5 Hz), 3.08-3.10 (1H, m), 3.23-3.24 (1H, m), 5.13 (1H, d, *J* = 6.3 Hz); ¹³C NMR (125 MHz, CDCl₃) δ = 211.1, 206.8, 105.2, 85.4, 61.9, 58.9, 47.9, 36.2, 33.4, 28.3, 27.5, 22.5, 13.9; IR (KBr) 2038, 1969, 1709 cm⁻¹.
- Other oxidation methods, such as cerium(IV) ammonium nitrate (CAN) in MeOH and iodine in THF gave **9a** in lower yield.
- General experimental procedure is as follows: To a THF solution (4.5 ml) of **5** (103 mg, 0.38 mmol) was added *n*-BuLi (1.56 mol dm⁻³; 0.61 ml, 0.95 mmol) at -78 °C. After 1 h, the reaction was quenched with aq. NH₄Cl (25 ml). The combined ether extracts (10 ml x 3) were washed with brine and dried over anhydrous Na₂SO₄ at 0 °C to afford an ether solution of the crude **6**.
Method A: The ether solution of the crude **6** was stirred under carbon monoxide atmosphere for 18 h at room temperature. After removal of the solvent, the mixture of **7b** and **8b** (122.9 mg, 0.34 mmol, 91%) was obtained by the purification by Florisil column chromatography. To a DMA solution (10 ml) of **7b** and **8b** was added Me₃NO (0.3 g, 4 mmol) at 0 °C and the reaction mixture was stirred for 12 h at room temperature. The crude products were purified by TLC (hexane:ethyl acetate=10:1) to afford butyl *m*-butylphenyl ketone **9b** (57.2 mg, 0.26 mmol, 77% (70% for 2 steps)).
Method B: After removal of the solvent from the ether solution of the crude **6** at 0 °C, DMA (10 ml) and Me₃NO was added. The reaction mixture was stirred for 12 h. **9b** was obtained after the purification by TLC (48.1 mg, 0.22 mmol, 72% from **5** (83.3 mg, 0.30 mmol)).
- Organolithium reagent reacts with ethoxycarbonyl group of **12**.
- M. F. Semmelhack and J. W. Herndon, *Organometallics*, **2**, 363 (1983).
- H. Heaney, "The Bimolecular Aromatic Friedel-Crafts Reaction," in "Comprehensive Organic Synthesis," ed by B. M. Trost, Pergamon Press, Oxford (1991), Vol. 2, p 733.