

# Four-component reaction between aryl aldehydes, ethyl cyanoacetate and dimethyl acetylenedicarboxylate in the presence of cyclohexyl isocyanide

Mohammad Anary-Abbasinejad<sup>a</sup>, Alireza Hassanabadi<sup>b\*</sup> and Farzaneh Aiinparast<sup>c</sup>

<sup>a</sup>Young Researchers Club, Islamic Azad University, Anar Branch, Anar, Iran

<sup>b</sup>Department of Chemistry, Islamic Azad University, Zahedan Branch, PO Box 98135-978, Zahedan, Iran

<sup>c</sup>Department of Chemistry, Islamic Azad University, Yazd Branch, PO Box 89195-155, Yazd, Iran

A four-component reaction between aryl aldehydes, ethyl cyanoacetate and dimethyl acetylenedicarboxylate in the presence of cyclohexyl isocyanide has been developed. The reaction furnishes highly functionalised cyclopentadienes in moderate to good yields in a one-pot process.

**Keywords:** cyclohexyl isocyanide, four-component reaction, dimethyl acetylenedicarboxylate, cyclopentadienes


Cyclopentadienes (Cps) or fulvenes are highly useful synthetic intermediates in the field of organic and organometallic chemistry as ligands in coordination chemistry and homogeneous catalysis for olefin polymerisation.<sup>1</sup> They are useful not only as a reactive diene component in the Diels–Alder reaction<sup>2</sup> but also as precursors for the preparation of transition-metal complexes with Cp-type ligands.<sup>3</sup> However, the preparation of well-defined, highly substituted cyclopentadienes is not necessarily easy owing to the absence of general methods<sup>4</sup> and also to the facile migration of the endocyclic double bonds.<sup>5</sup> Among the MCRs, isocyanide based multi-component reactions (IMCRs) have gained the most attention by organic chemists. Ugi four component reactions (U-4CR)<sup>6–8</sup> and Passerini three component reactions (P-3CR)<sup>9</sup> are among the most important IMCRs. U-4CR and P-3CR describe the reactions of isocyanides with carboxylic acids in the presence of imines or aldehydes, respectively.

Recently, another kind of IMCR has been developed and extensively investigated. Isocyanides react easily with electron-deficient acetylenic diesters such as dimethyl acetylenedicarboxylate (DMAD) to produce a reactive zwitterionic intermediate which can be trapped by an electrophile. Recently,

a wide variety of electrophiles have been applied to trap isocyanide-DMAD intermediates, among them are carbon electrophiles such as aldehydes, imines, quinonoids,<sup>10</sup> 1,2-diketones,<sup>11</sup> 1,2,3-tricarbonyl compounds,<sup>12</sup> isocyanates,<sup>13</sup> and hydrogen electrophiles such as pyrrole,<sup>14</sup> amides,<sup>15</sup> hydroxycoumarin,<sup>16</sup> phenols,<sup>17</sup> phthalic anhydride,<sup>18</sup> and isatoic anhydride.<sup>19</sup> Treatment of the isocyanide-DMAD zwitterion with aromatic carboxylic acids has been reported to produce unsaturated amides.<sup>20</sup> Reaction of the isocyanide-DMAD adduct with aromatic-substituted acetic acids has been reported to afford 2,5-diaminofuran derivatives in the presence of two equivalents of an isocyanide.<sup>21</sup> In the context of our previous works on IMCRs,<sup>14–16,22</sup> we now report the results of our investigations on the reaction of cyclohexyl isocyanide **1** with dimethyl acetylenedicarboxylate **2**, aryl aldehydes **3**, and ethyl cyanoacetate **4** to afford aminocyclopentadienes **5**.

## Results and discussion

Treatment of the three reactants and cyclohexyl isocyanide in dichloromethane for 24 h at room temperature, after silica gel column chromatography afforded highly functionalised cyclopentadienes **5** in excellent yields (Scheme 1).




5	Ar	%Yield <sup>*</sup>
a	phenyl	90
b	p-CH <sub>3</sub> O-phenyl	92
c	p-Cl-phenyl	90
d	p-NO <sub>2</sub> -phenyl	89
e	o-NO <sub>2</sub> -phenyl	88
f	m-NO <sub>2</sub> -phenyl	85
g	o-Br-phenyl	90

\* Isolated yields

**Scheme 1** Reaction between aryl aldehydes, ethyl cyanoacetate and dimethyl acetylenedicarboxylate in the presence of cyclohexyl isocyanide.

\* Correspondent. E-mail: ar\_hasanabadi@yahoo.com



Scheme 2 Suggested mechanism for formation compound 5.

The structures of compounds **5a–g** were deduced from their elemental analyses and IR,  $^1\text{H}$  NMR, and  $^{13}\text{C}$  NMR spectra. The  $^1\text{H}$  NMR spectrum of compound **5a** was very simple and exhibited two sharp signals at 3.79 and 3.81 ppm for methoxycarbonyl groups, and one NH group ( $\delta = 8.22$  ppm, disappeared with addition of  $\text{D}_2\text{O}$ ). Protons of the ethyl group are observed as a triplet at  $\delta = 1.18$  ppm ( $^3J_{\text{HH}} = 7$  Hz) and a quartet at  $\delta = 4.22$  ( $^3J_{\text{HH}} = 7$  Hz). Cyclohexyl fragment protons resonated as multiplets at  $\delta = 1.23$ – $1.67$  and a multiplet at  $\delta = 3.64$  ppm and aromatic protons resonated at  $\delta = 7.03$ – $7.68$  ppm. The  $^{13}\text{C}$  NMR spectrum of compound **5a** showed 23 distinct resonances in agreement with the proposed structure. The IR spectrum showed an absorption band at  $3355\text{ cm}^{-1}$  for the NH group. The carbonyl stretching vibrations observed as strong absorption bands at  $1744$  and  $1675\text{ cm}^{-1}$ . The nitrile stretching vibrations were observed in an absorption band at  $2236\text{ cm}^{-1}$ .

On the basis of the well established chemistry of isocyanides<sup>6–8,18</sup> it is reasonable to assume that intermediate **7** is produced by initial Knoevenagel condensation of aldehyde **3** with ethyl cyanoacetate **4**. The compound **5** is produced by initial attack of isocyanide-DMAD zwitterion intermediate **6**. The zwitterions can add to the intermediate of **7** resulting in the formation of **8**. The latter then isomerises to the final product **5** via a [1,5]H shift. (Scheme 2)

In conclusion, we report here the four-component reaction between aryl aldehydes, ethyl cyanoacetate and dimethyl acetylenedicarboxylate with cyclohexyl isocyanide as a simple and efficient route for the synthesis of functionalised cyclopentadiene derivatives. The advantages of the reported method are that it is inexpensive with easily available starting materials, it has simple and neutral reaction conditions, gives high yields, is a single-product reaction and is a simple work-up process.

## Experimental

Melting points were determined with an Electrothermal 9100 apparatus. Elemental analyses were performed using a Costech ECS 4010 CHNS-O analyzer at analytical laboratory of Islamic Azad University Yazd branch. IR spectra were recorded on a Shimadzu IR-470 spectrometer.  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra were recorded on Bruker DRX-500 Avance spectrometer at solution in  $\text{CDCl}_3$  using TMS as internal standard. The chemicals used in this work were purchased from Fluka (Buchs, Switzerland) and were used without further purification.

### General procedure

To a magnetically stirred solution of ethyl cyanoacetate (1 mmol) and aryl aldehydes (1 mmol) in dichloromethane (10 mL) was added a mixture of dimethyl acetylenedicarboxylate (1 mmol) in dichloromethane (2 mL) at room temperature. The reaction mixture was then stirred for 1 min. Cyclohexyl isocyanide (1 mmol) was added and the reaction mixture was stirred for more than 24 h. The solvent was removed and the residue was purified by silica gel column

chromatography using hexane-ethyl acetate (3:1) as eluent. The solvent was removed under reduced pressure to give the product.

**1,2-Dimethyl 3-ethyl 3-cyano-5-cyclohexylamino-4-(phenyl)-cyclopenta-1,4-diene-1,2,3-tricarboxylate (5a):** Yellow oil, yield: 90%; IR (KBr) ( $\nu_{\text{max}}$ ,  $\text{cm}^{-1}$ ): 3355 (NH), 1744, 1675 (carbonyl groups). Anal. Calcd for  $\text{C}_{25}\text{H}_{28}\text{N}_2\text{O}_6$ : C, 66.36; H, 6.24; N, 6.19. Found: C, 66.45; H, 6.16; N, 6.25%.  $^1\text{H}$  NMR (500.1 MHz,  $\text{CDCl}_3$ ):  $\delta = 1.18$  (3H, t,  $^3J_{\text{HH}} = 7$  Hz,  $\text{CH}_3$ ), 1.23–1.67 (10 H, m, 5  $\text{CH}_2$  of cyclohexyl), 3.64 (1 H, m, CH of cyclohexyl), 3.79 and 3.81 (6 H, 2s, 2  $\text{OCH}_3$ ), 4.22 (2 H, q,  $^3J_{\text{HH}} = 7$  Hz,  $\text{OCH}_2$ ), 7.03–7.68 (4H, m, aromatic), 8.22 (1 H, d,  $^3J_{\text{HH}} = 11$  Hz, NH).  $^{13}\text{C}$  NMR (125.7 MHz,  $\text{CDCl}_3$ ):  $\delta = 14.32$  ( $\text{CH}_3$ ), 23.12, 24.75, 25.68, 29.68 and 30.21 (5  $\text{CH}_2$  of cyclohexyl), 39.05 (CH of cyclohexyl), 51.86 and 52.94 (2  $\text{OCH}_3$ ), 57.87 (C), 64.40 ( $\text{OCH}_2$ ), 99.68 (CN), 113.92, 120.42, 144.38 and 160.22 (4C olefinic), 127.71, 128.80, 130.13 and 134.48 (aromatic), 163.23, 165.42 and 166.29 (3CO).

**1,2-Dimethyl 3-ethyl 3-cyano-5-cyclohexylamino-4-(4-methoxyphenyl)-cyclopenta-1,4-diene-1,2,3-tricarboxylate (5b):** Yellow oil, yield: 92%; IR (KBr) ( $\nu_{\text{max}}$ ,  $\text{cm}^{-1}$ ): 1743, 1657 (carbonyl groups). Anal. Calcd for  $\text{C}_{26}\text{H}_{30}\text{N}_2\text{O}_7$ : C, 64.72; H, 6.27; N, 5.81. Found: C, 64.85; H, 6.38; N, 5.74%.  $^1\text{H}$  NMR (500.1 MHz,  $\text{CDCl}_3$ ):  $\delta = 1.24$ – $1.80$  (10 H, m, 5  $\text{CH}_2$  of cyclohexyl), 1.33 (3 H, t,  $^3J_{\text{HH}} = 7$  Hz,  $\text{CH}_3$ ), 3.64 and 3.76 (6 H, 2s, 2  $\text{OCH}_3$ ), 3.71 (1 H, m, CH of cyclohexyl), 3.89 (3 H, s,  $\text{OCH}_3$ ), 4.33 (2 H, q,  $^3J_{\text{HH}} = 7$  Hz,  $\text{OCH}_2$ ), 6.88 (2H, d,  $^3J_{\text{HH}} = 7$  Hz, aromatic), 7.09 (2H, d,  $^3J_{\text{HH}} = 7$  Hz, aromatic), 8.24 (1 H, d,  $^3J_{\text{HH}} = 11$  Hz, NH).  $^{13}\text{C}$  NMR (125.7 MHz,  $\text{CDCl}_3$ ):  $\delta = 14.34$  ( $\text{CH}_3$ ), 23.01, 24.23, 25.84, 29.75 and 30.07 (5  $\text{CH}_2$  of cyclohexyl), 39.16 (CH of cyclohexyl), 53.10 and 53.19 (2  $\text{OCH}_3$ ), 56.92 ( $\text{OCH}_3$ ), 63.45 (C), 64.60 ( $\text{OCH}_2$ ), 114.50 (CN), 115.00, 127.81, 129.65 and 144.31 (aromatic), 114.27, 129.79, 144.58 and 160.49 (4C olefinic), 162.75, 163.82 and 165.49 (3CO).

**1,2-Dimethyl 3-ethyl 3-cyano-5-cyclohexylamino-4-(4-chlorophenyl)-cyclopenta-1,4-diene-1,2,3-tricarboxylate (5c):** Yellow oil, yield: 90%; IR (KBr) ( $\nu_{\text{max}}$ ,  $\text{cm}^{-1}$ ): 1746, 1666 (carbonyl groups). Anal. Calcd for  $\text{C}_{25}\text{H}_{27}\text{ClN}_2\text{O}_6$ : C, 61.66; H, 5.59; N, 5.75. Found: C, 61.79; H, 5.43; N, 5.68%.  $^1\text{H}$  NMR (500.1 MHz,  $\text{CDCl}_3$ ):  $\delta = 1.25$ – $2.02$  (10 H, m, 5  $\text{CH}_2$  of cyclohexyl), 1.37 (3 H, t,  $^3J_{\text{HH}} = 7$  Hz,  $\text{CH}_3$ ), 3.66 and 3.91 (6 H, 2s, 2  $\text{OCH}_3$ ), 3.76 (1 H, m, CH of cyclohexyl), 4.35 (2 H, q,  $^3J_{\text{HH}} = 7$  Hz,  $\text{OCH}_2$ ), 7.28 (2H, d,  $^3J_{\text{HH}} = 7$  Hz, aromatic), 7.38 (2H, d,  $^3J_{\text{HH}} = 7$  Hz, aromatic), 8.28 (1 H, d,  $^3J_{\text{HH}} = 11$  Hz, NH).  $^{13}\text{C}$  NMR (125.7 MHz,  $\text{CDCl}_3$ ):  $\delta = 14.44$  ( $\text{CH}_3$ ), 23.09, 24.82, 25.84, 29.76 and 30.09 (5  $\text{CH}_2$  of cyclohexyl), 39.15 (CH of cyclohexyl), 51.73 and 52.83 (2  $\text{OCH}_3$ ), 58.37 (C), 64.66 ( $\text{OCH}_2$ ), 99.87 (CN), 113.92, 120.42, 129.33 and 160.92 (4C olefinic), 129.44, 130.07, 134.69 and 139.37 (aromatic), 163.35, 165.50 and 166.38 (3CO).

**1,2-Dimethyl 3-ethyl 3-cyano-5-cyclohexylamino-4-(4-nitrophenyl)-cyclopenta-1,4-diene-1,2,3-tricarboxylate (5d):** Yellow oil, yield: 89%; IR (KBr) ( $\nu_{\text{max}}$ ,  $\text{cm}^{-1}$ ): 3320 (NH), 1741, 1643 (carbonyl groups). Anal. Calcd for  $\text{C}_{25}\text{H}_{27}\text{N}_3\text{O}_8$ : C, 60.36; H, 5.47; N, 8.45. Found: C, 60.19; H, 5.63; N, 8.33%.  $^1\text{H}$  NMR (500.1 MHz,  $\text{CDCl}_3$ ):  $\delta = 1.20$  (3H, t,  $^3J_{\text{HH}} = 7$  Hz,  $\text{CH}_3$ ), 1.25–2.16 (10 H, m, 5  $\text{CH}_2$  of cyclohexyl), 3.74 and 3.76 (6 H, 2s, 2  $\text{OCH}_3$ ), 3.58 (1 H, m, CH of cyclohexyl), 4.18 (2 H, q,  $^3J_{\text{HH}} = 7$  Hz,  $\text{OCH}_2$ ), 7.31 (2H, d,  $^3J_{\text{HH}} = 7$  Hz, aromatic), 7.44 (2H, d,  $^3J_{\text{HH}} = 7$  Hz, aromatic), 8.29 (1 H, d,  $^3J_{\text{HH}} = 11$  Hz, NH).  $^{13}\text{C}$  NMR (125.7 MHz,  $\text{CDCl}_3$ ):  $\delta = 14.02$  ( $\text{CH}_3$ ), 23.17,

24.87, 25.79, 29.64 and 30.13 (5CH<sub>2</sub> of cyclohexyl), 39.19 (CH of cyclohexyl), 51.90 and 52.75 (2 OCH<sub>3</sub>), 58.46 (C), 64.39 (OCH<sub>2</sub>), 99.77 (CN), 114.08, 120.63, 129.42 and 160.87 (4C olefinic), 129.21, 130.19, 133.98 and 139.12 (aromatic), 163.17, 165.43 and 166.52 (3CO).

**1,2-Dimethyl 3-ethyl 3-cyano-5-cyclohexylamino-4-(2-nitrophenyl)-cyclopenta-1,4-diene-1,2,3-tricarboxylate (5e):** Yellow oil, yield: 88%; IR (KBr) ( $\nu_{\max}$ , cm<sup>-1</sup>): 3380 (NH), 1734, 1671 (carbonyl groups). Anal.Calcd for C<sub>25</sub>H<sub>27</sub>N<sub>3</sub>O<sub>8</sub>: C, 60.36; H, 5.47; N, 8.45. Found: C, 60.19; H, 5.63; N, 8.33%. <sup>1</sup>H NMR (500.1 MHz, CDCl<sub>3</sub>):  $\delta$  = 1.21 (3H, t, <sup>3</sup>J<sub>HH</sub> = 7 Hz, CH<sub>3</sub>), 1.25–2.19 (10 H, m, 5 CH<sub>2</sub> of cyclohexyl), 3.32 (1 H, m, CH of cyclohexyl), 3.57 and 3.71 (6 H, 2s, 2 OCH<sub>3</sub>), 4.24 (2 H, q, <sup>3</sup>J<sub>HH</sub> = 7 Hz, OCH<sub>2</sub>), 7.53–7.99 (4H, m, aromatic), 8.44 (1 H, d, <sup>3</sup>J<sub>HH</sub> = 11 Hz, NH). <sup>13</sup>C NMR (125.7 MHz, CDCl<sub>3</sub>):  $\delta$  = 14.14 (CH<sub>3</sub>), 24.10, 24.77, 25.44, 32.76 and 34.23 (5CH<sub>2</sub> of cyclohexyl), 39.15 (CH of cyclohexyl), 51.60 and 53.36 (2 OCH<sub>3</sub>), 55.52 (C), 65.11 (OCH<sub>2</sub>), 98.57 (CN), 114.05, 125.18, 140.87 and 160.95 (4C olefinic), 126.29, 129.69, 130.52, 133.34, 134.67 and 149.08 (aromatic), 163.49, 164.85 and 165.62 (3CO).

**1,2-Dimethyl 3-ethyl 3-cyano-5-cyclohexylamino-4-(3-nitrophenyl)-cyclopenta-1,4-diene-1,2,3-tricarboxylate (5f):** Yellow oil, yield: 85%; IR (KBr) ( $\nu_{\max}$ , cm<sup>-1</sup>): 3330 (NH), 1744, 1670 (carbonyl groups). Anal.Calcd for C<sub>25</sub>H<sub>27</sub>N<sub>3</sub>O<sub>8</sub>: C, 60.36; H, 5.47; N, 8.45. Found: C, 60.19; H, 5.63; N, 8.33%. <sup>1</sup>H NMR (500.1 MHz, CDCl<sub>3</sub>):  $\delta$  = 1.18 (3H, t, <sup>3</sup>J<sub>HH</sub> = 7 Hz, CH<sub>3</sub>), 1.23–1.78 (10 H, m, 5 CH<sub>2</sub> of cyclohexyl), 3.63 (1 H, m, CH of cyclohexyl), 3.76 and 3.83 (6 H, 2s, 2 OCH<sub>3</sub>), 4.25 (2 H, q, <sup>3</sup>J<sub>HH</sub> = 7 Hz, OCH<sub>2</sub>), 7.50–8.26 (4H, m, aromatic), 8.39 (1 H, d, <sup>3</sup>J<sub>HH</sub> = 11 Hz, NH). <sup>13</sup>C NMR (125.7 MHz, CDCl<sub>3</sub>):  $\delta$  = 14.51 (CH<sub>3</sub>), 23.09, 25.35, 29.76, 33.05 and 34.61 (5CH<sub>2</sub> of cyclohexyl), 39.22 (CH of cyclohexyl), 51.90 and 53.15 (2 OCH<sub>3</sub>), 55.89 (C), 65.10 (OCH<sub>2</sub>), 99.96 (CN), 113.63, 125.61, 135.68 and 161.35 (4C olefinic), 118.37, 122.57, 130.28, 133.32, 141.44 and 148.86 (aromatic), 163.20, 165.41 and 165.13 (3CO).

**1,2-Dimethyl 3-ethyl 3-cyano-5-cyclohexylamino-4-(2-bromophenyl)-cyclopenta-1,4-diene-1,2,3-tricarboxylate (5g):** Yellow oil, yield: 90%; IR (KBr) ( $\nu_{\max}$ , cm<sup>-1</sup>): 3320 (NH), 1743, 1667 (carbonyl groups). Anal.Calcd for C<sub>25</sub>H<sub>27</sub>BrN<sub>3</sub>O<sub>6</sub>: C, 56.51; H, 5.12; N, 5.27. Found: C, 56.65; H, 5.03; N, 5.35%. <sup>1</sup>H NMR (500.1 MHz, CDCl<sub>3</sub>):  $\delta$  = 1.12 (3H, t, <sup>3</sup>J<sub>HH</sub> = 7 Hz, CH<sub>3</sub>), 1.18–1.77 (10 H, m, 5 CH<sub>2</sub> of cyclohexyl), 3.60 (1 H, m, CH of cyclohexyl), 3.74 and 3.76 (6 H, 2s, 2 OCH<sub>3</sub>), 4.18 (2 H, q, <sup>3</sup>J<sub>HH</sub> = 7 Hz, OCH<sub>2</sub>), 7.32–7.70 (4H, m, aromatic), 8.33 (1 H, d, <sup>3</sup>J<sub>HH</sub> = 11 Hz, NH). <sup>13</sup>C NMR (125.7 MHz, CDCl<sub>3</sub>):  $\delta$  = 14.42 (CH<sub>3</sub>), 23.15, 25.40, 29.85, 34.83 and 35.25 (5CH<sub>2</sub>

of cyclohexyl), 39.19 (CH of cyclohexyl), 51.65 and 52.73 (2 OCH<sub>3</sub>), 55.62 (C), 64.59 (OCH<sub>2</sub>), 99.92 (CN), 113.88, 122.90, 139.41 and 160.98 (4C olefinic), 120.43, 128.50, 129.18, 129.65, 132.26 and 149.45 (aromatic), 163.34, 165.48 and 166.34 (3CO).

We gratefully acknowledge financial support from the Research Council of Islamic Azad University of Yazd and The Islamic Azad University of Zahedan of Iran.

Received 27 July 2010; accepted 6 September 2010

Paper 1000266 doi: 10.3184/030823410X12875740077344

Published online: 24 November 2010

## References

- 1 R. Halterman, *Chem. Rev.*, 1992, **92**, 965.
- 2 H. Yoon and W. Chae, *Tetrahedron Lett.*, 1997, **38**, 5169.
- 3 U. Siemeling, *Chem. Rev.*, 2000, **100**, 1495.
- 4 B.J. Rausch, R. Gleiter and F. J Rominger, *Organomet. Chem.*, 2002, **658**, 242.
- 5 C.G. Venier and E.W. Casserly, *J. Am. Chem. Soc.*, 1990, **112**, 2808.
- 6 I. Ugi, B. Werner and A. Domling, *Molecules*, 2003, **8**, 53.
- 7 A. Domling, *Curr. Opin. Chem. Biol.*, 2000, **4**, 318.
- 8 A. Domling, *Chem. Rev.*, 2006, **106**, 17.
- 9 M. Passerini, *Gazz. Chim. Ital.*, 1921, **51** II, 126.
- 10 V. Nair, R.S. Menon and V. Sreekumar, *Pure Appl. Chem.*, 2005, **77**, 1191.
- 11 V. Nair, R.S. Menon, A. Deepthi, B.R. Devi and A.T. Biju, *Tetrahedron Lett.*, 2005, **46**, 1337.
- 12 V. Nair and A. Deepthi, *Tetrahedron Lett.*, 2006, **47**, 2037.
- 13 A. Alizadeh, S. Rostamnia, N. Zohreh and H.R. Bijanzadeh, *Month.Chem.*, 2008, **139**, 49.
- 14 M. Anary-Abbasinejad, M.H. Mosslemin, H. Anaraki-Ardakani and S. Tahan, *J. Chem. Res.*, 2006, 306.
- 15 M. Anary-Abbasinejad, M.H. Mosslemin, S. Tahan and H. Anaraki-Ardakani, *J. Chem. Res.*, 2006, 170.
- 16 M. Anary-Abbasinejad, H. Anaraki-Ardakani, F. Rastegari and A. Hassanabadi, *J. Chem. Res.*, 2007, 602.
- 17 I. Yavari, H. Djahaniani and F. Nasiri, *Tetrahedron*, 2003, **59**, 9409.
- 18 A. Shaabani, M.B. Teimouri and H.R. Bijanzadeh, *J. Chem. Res.*, 2002, 381.
- 19 A. Shaabani, M.B. Teimouri, P. Mirzaei and H.R. Bijanzadeh, *J. Chem. Res.*, 2003, 82.
- 20 A. Alizadeh, S. Rostamnia and L.G. Zho, *Tetrahedron*, 2006, **62**, 5641.
- 21 A. Alizadeh, S. Rostamnia and M.L. Hu, *Synlett*, 2006, 1592.
- 22 M. Anary-Abbasinejad and M. Kamali-Gharamaleki, *J. Chem. Res.*, 2008, 383.

Copyright of Journal of Chemical Research is the property of Science Reviews 2000 Ltd. and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.