N-Metallation of MS_2N_2 rings. X-ray crystal structures of $[(\eta^5-C_5Me_5)Ir(S_2N_2)Au(PPh_3)][ClO_4], [\{(\eta^5-C_5Me_5)Ir(S_2N_2)Au\}_2 (\mu_2-dppm)][ClO_4]_2 and [Au(dppeS-$ *P* $)Cl]_2$

Stephen M. Aucott, Pravat Bhattacharyya, Heather L. Milton, Alexandra M. Z. Slawin and J. Derek Woollins*

School of Chemistry, University of St Andrews, Fife, Scotland KY16 9ST. E-mail: jdw3@st-and.ac.uk; Fax: +44-1334-463384; Tel: +44-1334-463861

Received (in London, UK) 16th June 2003, Accepted 1st August 2003 First published as an Advance Article on the web 29th August 2003

Bimetallic complexes $[(\eta^5-C_5R_5)M(S_2N_2)Au(PPh_3)][ClO_4]$ and tetrametallic species $[\{(\eta^5-C_5R_5)M(S_2N_2)Au\}_2(\mu_2-P^AP)][ClO_4]_2$ (R = H, M = Co; R = Me, M = Ir; P^P = dppm or dppe) can be prepared by treatment of $[(\eta^5-C_5R_5)M(S_2N_2)]$ with gold(1) electrophiles generated by chloride abstraction from $[AuCl(PPh_3)]$ or $[(AuCl)_2(\mu_2-P^AP)]$. X-Ray crystallography of $[(\eta^5-C_5Me_5)Ir(S_2N_2)Au\}_2(\mu_2-dppm)][ClO_4]_2$ confirms auration of the metal-bound nitrogen atom of the MS_2N_2 ring. π -Stacking of the MS_2N_2 rings occurs within both structures.

Introduction

The chemistry of sulfur–nitrogen heterocycle and cage systems has elicited considerable attention from both theoretical and synthetic standpoints. The general development of this fascinating area has been hampered by the paucity of stable and easily handled S–N reagents. Thus, despite its renowned explosive tendencies, tetrasulfur tetranitride (S₄N₄) has been the most commonly used material for the synthesis of binary S–N systems,¹ organic heterocycles² and metalla-sulfur–nitrogen complexes.^{3,4}

We recently reported the preparation of the tin(IV) disulfurdinitrido complex [ⁿBu₂Sn(S₂N₂)]₂, which shows great promise as a metathesis reagent for the synthesis of ES₂N₂ (E = metal or non-metal) rings.^{5,6} For example [(η^5 -C₅Me₅)Ir(S₂N₂)], only the second crystallographically characterised metal cyclopentadienyl S₂N₂ complex, is accessible from [(η^5 -C₅Me₅)IrCl₂(PPh₃)] or [(η^5 -C₅Me₅)IrCl(μ -Cl)]₂.⁶ Moreover [(η^5 -C₅Me₅)Ir(S₂N₂)] undergoes metallation at the *non metalbound nitrogen atom* by [(η^5 -C₅Me₅)IrCl(PPh₃)]⁺ to give [(η^5 -C₅Me₅)₂Ir₂(S₂N₂)Cl(PPh₃)]⁺. Conversely, the metal-bound nitrogen atom is the site of protonation in metal–[S₂N₂H]⁻ complexes^{7,8} and of metallation in [M(S₂N₂)(PPh₃)]₂ (M = Pt or Pd),⁹ [R₂Sn(S₂N₂)]₂ (R = ⁿBu, Me or^tBu)^{6,10,11} and [Ph₄P]₂[Ni₃(S₂N₂)₂],¹² which contain M₂(μ ₂-N)₂ cores.

Auration of MS_2N_2 rings by gold(1) electrophiles has not been previously reported. In this paper we describe the *N*metallation of $[(\eta^5-C_5H_5)Co(S_2N_2)]$ and $[(\eta^5-C_5Me_5)Ir(S_2N_2)]$ by cations generated by halide abstraction from gold(1)-chloride precursors, including representative X-ray crystal structures.

Results and discussion

Chloride abstraction from [AuCl(PPh₃)] and [(AuCl)₂(μ_2 -P^AP)] (P^AP = dppm or dppe) with silver(1) perchlorate in dichloromethane, followed by addition of stoichiometric quantities of [(η^5 -C₅H₅)Co(S₂N₂)] or [(η^5 -C₅Me₅)Ir(S₂N₂)] generated the heterometallic complexes **1**–**6** (Fig. 1) in 24–68% yields, the dppe complexes **5**, **6** being isolated in lowest yields.



Fig. 1 Structures of complexes 1–6.

The air- and moisture- stable complexes are soluble in dichloromethane, thf and acetone. The X-ray crystal structures of $[(\eta^{5}-C_{5}Me_{5})Ir(S_{2}N_{2})Au(PPh_{3})][ClO_{4}]$ and $[\{(\eta^{5}-C_{5}Me_{5})Ir(S_{2} N_2$ Au $_2(\mu_2$ -dppm)][ClO₄]₂ (see below) confirm that metallation occurs, as expected, at the more electron-rich metal-bound nitrogen atom. An attempt to metallate both nitrogen centres of $[(\eta^5-C_5H_5)Co(S_2N_2)]$ with two equivalents of $[Au(PPh_3)]^+$ in dichloromethane led to an intractable mixture of products. FAB⁺ MS for 1 and 2 contained peaks for $[(arene)M(S_2 - C_2)]$ N_2 Au(PPh₃)]⁺ at m/z 879 and 675 respectively, but the dicationic complexes 3-6 were less straightforward. Peaks at m/z813 for 3, 4 and m/z 827 for 5, 6 corresponded to [Au₂(P^P)Cl]⁺, with higher mass peaks of weaker intensity for 3-5 assigned to $[(P^P)Au_2Cl(S_2N_2)M(arene)]^+$ (m/z 1233,1030 and 1246 respectively). Since elemental analyses and NMR spectroscopy have indisputably established the purity of 1-6, we cannot explain the peculiar mass spectrometric data for 3-6. The IR spectra of the complexes are dominated by bands arising from the phosphine co-ligands and the perchlorate anions (ca. 1100, 625 cm⁻¹), bands from the S₂N₂ groups could not be assigned with certainty.

The crystal structure of 1 (Fig. 2) confirms metallation of the metal-bound nitrogen atom N(1). The IrS_2N_2 ring retains its planarity,⁶ the N(1)-Au(1)-P(1) axis is approximately linear



Fig. 2 X-Ray crystal structure of 1 (C–H bonds and perchlorate counterion omitted for clarity). Selected bond lengths (Å) and angles (°) (esd's in parentheses): Au(1)–N(1) 2.059(4), Au(1)–P(1) 2.2400(15), N(1)–S(1) 1.584(4), N(1)–Ir(1) 2.984(4), S(1)–N(2) 1.568(5), N(2)–S(2) 1.664(5), S(2)–Ir(1) 2.2060(15), N(1)–Au(1)–P(1) 173.71(12), S(1)–N(1)–Ir(1) 120.0(3), S(1)–N(1)–Au(1) 115.4(2), Ir(1)–N(1)–Au(1) 1124.6(2), N(2)–S(1)–N(1) 110.5(2), S(1)–N(2)–S(2) 116.2(3), N(2)–S(2)–Ir(1) 107.42(18), N(1)–Ir(1)–S(2) 85.81(13).

 $[173.71(12)^{\circ}]$ with Au(1)–N(1) and Au(1)–P(1) distances of 2.059(4) and 2.2400(15) Å respectively. The IrS₂N₂ rings are stacked [interplanar separation *ca.* 3.4 Å] in the manner of other MS₂N₂ structures,^{7,13} but intermolecular Au···Au interactions or close cation–anion contacts are absent.

A comparison of metallacycle bond lengths for $[(\eta^5-C_5Me_5)Ir(S_2N_2)]$, **1** and $[(\eta^5-C_5Me_5)_2Ir_2(S_2N_2)Cl(PPh_3)][PF_6]$ (Table 1), indicates that metallation appears to change the IrS₂N₂ bond lengths and angles in a similar fashion to protonation.⁸ The Ir–S(2) distances in **1** is 0.02 Å longer than in $[(\eta^5-C_5Me_5)Ir(S_2N_2)]$, with N(1)–S(1) *ca*. 0.05–0.07 Å longer in the bimetallic complexes compared to $[(\eta^5-C_5Me_5)Ir(S_2N_2)]$. N(2)–S(2) is also lengthened in the binuclear complexes.

The crystal structure of $3 \cdot CH_2Cl_2$ shows two $[(\eta^5 C_5Me_5$ $Ir(S_2N_2)$ units linked by $[Au_2(\mu_2-dppm)]^{2+}$ (Fig. 3), selected bond lengths and angles appearing in Table 2. The two N-Au-P axes [N(1)-P(1)-Au(1) 172.1(3)°, N(3)-Au(3)-P(3) 174.9(2)°] are parallel, the IrS₂N₂ rings adopting a headto-tail stacking arrangement with respect to one another. The Au(1)-Au(3) distance [3.1931(9) Å] is slightly longer than generally found for digold(1) complexes containing a bridging diphosphine ligand [2.576-3.173 Å].^{14–22} The molecular conformation enforced by the Au---Au interaction results in a rare intramolecular example of MS_2N_2 ring π -stacking, the $N(1) \cdots N(3)$ and $S(2) \cdots S(4)$ distances being 3.56 and 3.90 Å respectively, additionally the face-to-face phenyl rings have an interplanar separation of 3.77 Å. There is one short $Au \cdots O$ interaction with a perchlorate anion [Au(1) \cdots O(2) 3.31 Å], but no significant intercation contacts. Compound 3 represents an interesting oligomer or building subunit for polymeric M-S-N organometallic chains.

We have demonstrated that $[{}^{n}Bu_{2}Sn(S_{2}N_{2})]_{2}$ can form $MS_{2}N_{2}$ chelate rings by metathetical exchange of *cis*-chloride

 $\begin{array}{l} \textbf{Table 1} A \ comparison \ of \ metallacycle \ bond \ lengths \ in \ [(\eta^5-C_5Me_5)Ir(S_2N_2)], \ 1 \ and \ [(\eta^5-C_5Me_5)_2Ir_2(S_2N_2)Cl(PPh_3)][PF_6].^{''} \end{array}$

	$[(\eta^{5}-C_{5}Me_{5})Ir(S_{2}N_{2})]$	1	$\begin{array}{l} [(\eta^{5}\text{-}C_{5}Me_{5})_{2}Ir_{2}\\ (S_{2}N_{2})Cl(PPh_{3})]^{+} \end{array}$
Ir-N(1)	1.962(6)	1.983(4)	1.961(1)
Ir-S(2)	2.175(2)	2.206(1)	2.179(2)
N(1)-S(1)	1.509(7)	1.584(4)	1.558(10)
S(1) - N(2)	1.563(9)	1.568(4)	1.531(9)
N(2)–S(2)	1.632(8)	1.664(4)	1.685(9)
^{a} N(1) and	S(2) are the metal-bound	atoms of the	$[S_2N_2]^{2-}$ chain



Fig. 3 X-Ray crystal structure of $3 \cdot CH_2Cl_2$ (C–H bonds, solvent molecule and perchlorate counterions omitted for clarity).

ligands from a metal centre;^{5,6} we were intrigued as to whether the gold(1) centres of $[(AuCl)_2(\mu_2-P^{A}P)]$ (P^AP = dppm or dppe) could be linked with an $[S_2N_2]^{2-}$ unit to give nine- or ten-membered rings. To this end, $[{}^{n}Bu_{2}Sn(S_{2}N_{2})]_{2}$ and $[(AuCl)_{2}(\mu_{2}-\mu_{2})]_{2}$ P^P)] (1:1 molar ratio) were refluxed together in thf for four hours. ³¹P{¹H} NMR spectra of crude reaction mixtures contained one major component, characterised by two doublets $(J_{\rm PP} 16 \, \text{Hz} \text{ for dppm}, 63 \, \text{Hz} \text{ for dppe})$ and several weaker singlets. Gel permeation chromatography (Biobeads SX-8, dichloromethane eluant) enabled purification of the mixtures, monitored by ${}^{31}P{}^{1}H$ NMR spectroscopy of 1 cm³ fractions collected prior to, and including, the yellow eluate. Fractions whose ${}^{31}\bar{P}{}^{1}H$ NMR spectra did not contain doublets were not studied further. For the dppe complex, colourless crystals were grown from dichloromethane-hexane by solvent diffusion. X-Ray crystallography revealed, unexpectedly, a dimeric structure [Au(dppeS-P)Cl]2 (Fig. 4) i.e. one AuCl unit had been abstracted from [(AuCl)₂(µ₂-dppe)] and the uncoordinated phosphorus(III) centre converted into the phosphorus(v) sulfide by reaction with $[S_2N_2]^{2-}$, two such [Au(dppeS-P)Cl] units being linked by an Au-Au interaction. The structure of [Au(dppeS-P)Cl]₂ contains approximately linear P-Au-Cl units [P(1)-Au(1)-Cl(1) 174.93(8)°, P(3)-Au(2)-Cl(2) 172.69(8)°] whose axes are gauche to one another [P(1)-Au(1)-Au(2)- $Cl(2) - 58^{\circ}$, P(3)-Au(2)-Au(1)-Cl(1) - 62^{\circ}]. The Au(1)-Au(2) separation [3.2626(5) Å] is at the upper limit of the range observed for unsupported (i.e. without bridging ligands) Au(1)-Au(1) contacts [3.031(1)-3.264(2) Å].²⁴⁻³¹ FAB⁺ MS for the dppm and dppe complexes contained strong peaks at m/z 613 and 627 respectively for $[Au(P^PS)]^+$

In conclusion, metallation of $[(\eta^5-C_5H_5)Co(S_2N_2)]$ and $[(\eta^5-C_5Me_5)Ir(S_2N_2)]$ by gold(1) electrophiles takes place at the more electron-rich metal-bound nitrogen atom; we have obtained bimetallic and tetrametallic complexes. X-Ray crystallography reveals that the metallacycle experiences some distortion upon *N*-metallation but planarity of the MS₂N₂ ring is retained. Furthermore, intramolecular π -stacking of the MS₂N₂ rings occurs in the tetrametallic systems.

Experimental

Reactions were conducted under dinitrogen, subsequent procedures were performed in air. Dichloromethane was dried and distilled from calcium hydride prior to use, all other solvents and reagents were used as received. $[(\eta^5-C_5Me_5)Ir(S_2N_2)]$ and $[(\eta^5-C_5H_5)Co(S_2N_2)]$ were prepared by literature methods,^{6,23} chlorogold(1)–phosphine complexes were prepared by treating

Table 2 Selected bond lengths (Å) and angles (°) for $3.CH_2Cl_2$ (esd's in parentheses)

N(1)–S(1)	1.584(9)	N(3)–S(3)	1.608(8)
N(1)–Ir(1)	1.983(8)	N(3)–Ir(3)	1.984(8)
N(1)-Au(1)	2.066(7)	N(3)–Au(3)	2.057(7)
Ir(1)-S(2)	2.184(3)	Ir(3)-S(4)	2.203(3)
S(2)–N(2)	1.664(9)	N(4)–S(4)	1.668(9)
N(2)-S(1)	1.561(9)	N(4)–S(3)	1.563(8)
Au(1) - P(1)	2.252(3)	Au(3)–P(3)	2.249(3)
S(1)-N(1)-Ir(1)	119.8(4)	S(3)–N(3)–Ir(3)	119.7(4)
S(1)–N(1)–Au(1)	115.4(4)	S(3)–N(3)–Au(3)	110.8(4)
Ir(1)-N(1)-Au(1)	124.8(4)	Ir(3)–N(3)–Au(3)	129.3(4)
N(1)-Ir(1)-S(2)	86.1(2)	N(3)-Ir(3)-S(4)	85.8(2)
N(2)-S(2)-Ir(1)	107.6(3)	N(4)-S(4)-Ir(3)	107.9(3)
S(1)-N(2)-S(2)	116.3(5)	S(3)-N(4)-S(4)	116.1(5)
N(2)-S(1)-N(1)	110.2(4)	N(4)-S(3)-N(3)	110.4(5)
N(1)-Au(1)-P(1)	172.1(2)	N(3)-Au(3)-P(3)	174.9(2)
N(1)-Au(1)-Au(3)	97.2(2)	N(3)-Au(3)-N(1)	92.9(2)
P(1)-Au(1)-Au(3)	88.45(7)	Au(3)-Au(1)-P(1)	88.13(7)

[AuCl(tht)] (tht = tetrahydrothiophen) with stoichiometric quantities of the phosphine in dichloromethane. ³¹P{¹H} and ¹H NMR spectra (109.4 and 270.0 MHz respectively, d_2 -dichloromethane) were recorded on a JEOL GSX 270 spectrometer, FAB⁺ mass spectra (3-nitrobenzyl alcohol matrix) were carried out by the EPSRC National Mass Spectrometry Service Centre (Swansea).

CAUTION! Perchlorate salts are potentially explosive and should be handled with care.

The general procedure employed is illustrated for 3.

To $[(AuCl)_2(\mu_2-dppm)]$ (41 mg, 0.05 mmol) in dichloromethane (5 cm³) was added silver(1) perchlorate (21 mg, 0.1 mmol) and the solution stirred in the dark for 1.5 h. $[(\eta^5-C_5Me_5)Ir(S_2N_2)]$ (42 mg, 0.1 mmol) was added as a solid in one portion and the mixture stirred for 18 h in the dark. The grey-black mixture was filtered through Celite, which was washed further with dichloromethane (20 cm³). The filtrate was concentrated to *ca*. 1 cm³, vapour diffusion of diethyl ether



Fig. 4 X-Ray crystal structure of $[Au(dppeS-P)Cl]_2(C-H bonds omitted for clarity). Selected bond lengths (Å) and angles (°) (esd's in parentheses): Au(1)–P(1) 2.245(2), Au(1)–Cl(1) 2.292(2), Au(1)–Au(2) 3.2626(5), S(2)–P(2) 1.958(3), Au(2)–P(3) 2.2359(19), Au(2)–Cl(2) 2.2974(17), S(4)–P(4) 1.961(3), P(1)–Au(1)–Cl(1) 174.93(8), P(1)–Au(1)–Au(2) 108.71(6), Cl(1)–Au(1)–Au(2) 75.86(6), P(3)–Au(2)–Cl(2) 172.69(8), P(3)–Au(2)–Au(1) 110.80(5), Cl(2)–Au(2)–Au(1) 76.11(5).$

into this solution afforded 54 mg (59%) of 3 as red crystals. Complexes 5, 6 were crystallised from dichloromethanediethyl ether by solvent diffusion.

1 Red crystals, yield 34%. Found (calc. for $C_{28}H_{30}AuClIr-N_2O_4PS_2$): C 34.2 (34.3), H 2.7 (3.1), N 2.7 (2.9)%. δ_P : 31.7 (s). δ_H : 7.70–7.57 (m, 18H, Ph), 1.51 (s, 15H, C₅Me₅). FAB⁺ MS: m/z 879, $[M^+ - ClO_4]$.

2 Purple crystals, yield 68%. Found (calc. for $C_{23}H_{20}AuCl-CoN_2O_4PS_2$): C 35.8 (35.6), H 2.45 (2.6), N 3.6 (3.6)%. δ_P : 30.0 (s). δ_H : 7.80–7.61 (m, 18H, Ph), 6.01 (s, 5H, C₅H₅). FAB⁺ MS: m/z 675, $[M^+ - ClO_4]$.

3 Red crystals, yield 59%. Found (calc. for C₄₅H₅₂Au₂Cl₂Ir₂-N₄O₈P₂S₄): C 29.8 (29.8), H 2.5 (2.9), N 2.8 (3.1)%. $\delta_{\rm P}$: 26.8 (s). $\delta_{\rm H}$: 7.70–7.44 (m, 20H, Ph), 4.43 (t, 2H, J = 15 Hz, CH₂), 2.04 (s, 30H, C₅Me₅). FAB⁺ MS: m/z, 813 [Au₂(dppm)Cl]⁺, 1233 [Au₂(dppm)Cl(S₂N₂)Ir(C₅Me₅)]⁺.

4 Purple crystals, yield 60%. Found (calc. for $C_{35}H_{32}Au_2Cl_2-Co_2N_4O_8P_2S_4$): C 29.8 (29.8), H 2.2 (2.3), N 3.7 (4.0)%. δ_P : 26.4 (s). δ_H : 7.90 (m, 8H, Ph), 7.51 (m, 12H, Ph), 5.86 (s, 1H, C₅H₅), 4.50 (t, 2H, J = 13 Hz, CH₂). FAB⁺ MS: m/z 813 [Au₂(dppm)Cl]⁺, 1030 [Au₂(dppm)Cl(S₂N₂)Co(C₅H₅)]⁺.

5 Red blocks, yield 24%. Found (calc. for $C_{46}H_{52}Au_2Cl_2Ir_2-N_4O_8P_2S_4$): C 31.05 (30.18), H 2.40 (2.97), N 2.89 (3.06)%. δ_P : 27.5 (s). δ_H : 7.73–7.47 (m, 20H, Ph), 3.22 (d, 4H, J = 6 Hz, CH₂), 2.01 (s, 30H, C_5Me_5). FAB⁺ MS: m/z 827 [Au₂(dp-pe)Cl]⁺, 1246 [Au₂(dppe)Cl(S₂N₂)Ir(C₅Me₅)]⁺.

6 Purple crystals, yield 24%. Found (calc. for $C_{36}H_{34}Au_2Cl_2-Co_2N_4O_8P_2S_4$): C 30.24 (30.38), H 2.32 (2.41), N 3.43 (3.94)%. δ_P : 28.7 (s). δ_H : 7.86 (m, 8H, Ph), 7.54 (m, 12H, Ph), 6.04 (s, 10H, C_5H_5), 3.29 (s, 4H, CH₂). FAB⁺ MS: m/z 827 [Au₂(dp-pe)Cl]⁺.

Reaction between $[(AuCl)_2(\mu_2 P^A P)]$ and $[{}^{n}Bu_2Sn_{(S_2N_2)}]_2$; $[(AuCl)_2(\mu_2 P^A P)]$ (0.17 mmol) and $[{}^{n}Bu_2Sn(S_2N_2)]_2$ (0.17 mmol) were refluxed in thf (8 cm³) for four hours. The solvent was evaporated *in vacuo*, the products extracted into dichloromethane (2 cm³) and purified by gel permeation chromatography (Biobeads SX-8, dichloromethane eluant), 1 cm³ fractions collected prior to, and including, the yellow eluate were monitored by ${}^{31}P({}^{1}H$ } NMR spectroscopy. Fractions giving singlets in their ${}^{31}P\{{}^{1}H\}$ NMR spectra were discarded, those containing a pair of doublets were combined. From the reaction using $[(AuCl)_2(\mu_2-dppe)]$, colourless needles of $[Au(dppeS-P)Cl]_2$ were grown by layering dichloromethane solutions of the appropriate fractions with hexane.

[Au(dppmS-P)Cl]₂ δ_{P} : 36.7 (d, J = 16), 16.9 (d, J = 16 Hz). FAB⁺ MS: m/z 613, [Au(dppmS)]⁺.

[Au(dppeS-P)Cl]₂ $\delta_{\rm P}$: 43.9 (d, J = 63), 32.5 (d, J = 63 Hz). FAB⁺ MS: m/z 627, [Au(dppeS)]⁺.

X-ray crystallography

Single crystal diffraction studies on 1, $3 \cdot \text{CH}_2\text{Cl}_2$ and [Au(dppeS-*P*)Cl]₂ were performed on a Bruker SMART CCD diffractometer at 125 K with graphite monochromated Mo-K_{\alpha} radiation ($\lambda = 0.71073$ Å). Details of data collections and structural refinements are given in Table 3. The structures were solved by direct methods, non-hydrogen atoms were refined with anisotropic displacement parameters, hydrogen atoms bound to carbon were idealised and fixed (C-H 0.95 Å). Structural refinements were by the full-matrix least-squares method on F^2 using the program SHELXTL.³² Crystallographic data (excluding structure factors) for the structures reported in this paper have been deposited with the Cambridge Crystallographic Database Centre.[†]

[†] CCDC reference numbers 213795–213798. See http://www.rsc.org/ suppdata/nj/b3/b306847a/ for crystallographic data in .cif or other electronic format.

Table 3 Details of X-ray data collections and structural refinements for 1, 3.CH ₂ Cl ₂ and [Au(dppeS-	P)Cl] ₂
--	--------------------

	1	$3.CH_2Cl_2$	$[Au(dppeS-P)Cl]_2$
Empirical Formula	C ₂₈ H ₃₀ AuClIrN ₂ O ₄ PS ₂	C46H54Au2Cl4Ir2N4O8P2S4	C52H48Au2Cl2P4S2
M	978.25	1901.24	1325.74
T/K	125(2)	125(2)	125(2)
$\lambda/Å$	0.71073	0.71073	0.71073
Crystal system	Monoclinic	Orthorhombic	Triclinic
Space group	P2(1)/c	Pca2(1)	$P\overline{1}$
a/Å	13.0057(17)	32.363(8)	11.0993(8)
b/Å	11.2301(15)	10.367(3)	13.0681(10)
c/Å	21.098(3)	16.838(4)	19.3948(14)
$\alpha/^{\circ}$	90	90	84.735(1)
β/°	98.213(2)	90	76.483(1)
γ/°	90	90	64.900(1)
$U/Å^3, Z$	3049.9(7), 4	5649(3), 4	2476.9(3), 2
$D_{\rm c}/{\rm Mg}~{\rm m}^{-3}$	2.130	2.235	1.778
μ/mm^{-1}	9.477	10.321	6.273
F(000)	1856	3584	1288
Crystal size/mm	0.1 imes 0.1 imes 0.1	$0.25 \times 0.15 \times 0.06$	0.18 imes 0.1 imes 0.1
θ range/°	1.58-23.29	1.96-23.33	2.02-23.36
Reflections collected	14834	23 332	12636
Independent reflections (R_{int})	4361 (0.0470)	7349 (0.0473)	7038 (0.0410)
Goodness-of-fit on F^2	0.955	0.952	0.900
<i>R</i> 1, <i>wR</i> 2	0.0233, 0.0464	0.0259, 0.0632	0.0364, 0.0713

Acknowledgements

We are grateful to the EPSRC (S. M. A., P. B.) and the University of St. Andrews (H. L. M.) for funding.

References

- 1 R. T. Oakley, Prog. Inorg. Chem., 1988, 36, 299
- 2 T. Torroba, J. Prakt. Chem., 1999, 341, 99.
- 3 P. F. Kelly, A. M. Z. Slawin, D. J. Williams and J. D. Woollins, Chem. Soc. Rev., 1992, 245.
- 4 T. Chivers and F. Edelmann, Polyhedron, 1986, 5, 1661.
- 5 S. M. Aucott, A. M. Z. Slawin and J. D. Woollins, Phosphorus,
- Sulfur Silicon Relat. Elem., 2001, 168, 559.
 S. M. Aucott, A. M. Z. Slawin and J. D. Woollins, Can. J. Chem., 2002, 80, 1481.
- 7 R. Jones, P. F. Kelly, D. J. Williams and J. D. Woollins, J. Chem. Soc., Dalton Trans., 1988, 803.
- 8 R. Jones, C. P. Warrens, D. J. Williams and J. D. Woollins, J. Chem. Soc., Dalton Trans., 1987, 907.
- 9 R. Jones, P. F. Kelly, D. J. Williams and J. D. Woollins, J. Chem. Soc., Chem. Commun., 1985, 1325.
- 10 H. W. Roesky, Adv. Inorg. Chem. Radiochem., 1979, 22, 245.
- 11 D. Hanssgen, M. Jansen, W. Assenmacher and H. Salz, J. Organomet. Chem., 1993, 445, 61.
- 12 J. Weiss, Angew. Chem., Int. Ed. Engl., 1984, 23, 225.
- 13 R. Jones, P. F. Kelly, C. P. Warrens, D. J. Williams and J. D. Woollins, J. Chem. Soc., Chem. Commun., 1986, 711.
- 14 H. Schmidbaur, R. Herr, G. Muller and J. Riede, Organometallics, 1985, 4, 1208.

- 15 R. Uson, A. Laguna, M. Laguna, M. N. Fraile, P. G. Jones and G. M. Sheldrick, J. Chem. Soc., Dalton Trans., 1986, 291.
- 16 H. Schmidbaur, W. Graf and G. Muller, Angew. Chem., Int. Ed. Engl., 1988, 27, 417.
- 17 H. R. C. Jaw, M. M. Savas, R. D. Rogers and W. R. Mason, *Inorg. Chem.*, 1989, 28, 1028.
- 18 L. S. Liou, C. P. Liu and J. C. Wang, Acta. Cryst. C, 1994, 50, 538.
- 19 H. Schmidbaur, A. Wohlleben, U. Schubert, A. Frank and G. Huttner, *Chem. Ber.*, 1977, **110**, 2751.
- 20 P. Lange, A. Schier and H. Schmidbaur, Z. Naturforsch., 1997, 52B, 769.
- 21 H. Schmidbaur, K. Dziwok, A. Grohmann and G. Muller, *Chem. Ber.*, 1989, **122**, 893.
- 22 P. G. Jones and E. Bembenek, Acta Cryst. C, 1996, 52, 2396.
- 23 F. Edelmann, J. Organomet. Chem., 1982, 228, C47.
- 24 C. Hollatz, A. Schier, J. Riede and H. Schmidbaur, J. Chem. Soc., Dalton Trans., 1999, 111.
- 25 K. K. Banger, R. P. Banham, A. K. Brisdon, W. I. Cross, G. Damant, S. Parsons, R. G. Pritchard and A. Sousa-Pedrares, *J. Chem. Soc., Dalton Trans.*, 1999, 427.
- 26 S. Attar, W. H. Bearden, N. W. Alcock, E. C. Alyea and J. H. Nelson, *Inorg. Chem.*, 1990, **29**, 425.
- 27 A. Stutzer, P. Bissinger and H. Schmidbaur, *Chem. Ber.*, 1992, **125**, 367.
- 28 A. Stutzer, P. Bissinger and H. Schmidbaur, Z. Naturforsch., 1992, 47B, 1261.
- 29 P. M. Van Calcar, M. M. Olmstead and A. L. Balch, *Inorg. Chem.*, 1997, 36, 5231.
- 30 D. V. Toronto, B. Weissbart, D. S. Tinti and A. L. Balch, *Inorg. Chem.*, 1996, 35, 2484.
- 31 B. C. Tzeng, K. K. Cheung and C. M. Che, J. Chem. Soc., Chem. Commun., 1996, 1681.
- 32 G. M. Sheldrick, SHELXTL, Bruker AXS, Madison, WI, 1999.