

Tetrahedron Letters 40 (1999) 7497-7500

TETRAHEDRON LETTERS

Diastereo- and enantioselective hydrodimerization of β-monosubstituted acrylic acid amides induced by chiral samarium(II) complexes

Takashi Kikukawa, Takeshi Hanamoto and Junji Inanaga *

Institute for Fundamental Research of Organic Chemistry (IFOC), Kyushu University, Hakozaki, Higashi-ku, Fukuoka 812-8581, Japan

Received 15 July 1999; revised 11 August 1999; accepted 12 August 1999

Abstract

The chiral samarium(II) complex prepared from SmI₂, (*R*)-BINOL, and an achiral tertiary amine promoted the reductive homo-coupling reaction of β -monosubstituted acrylic acid amides to give the corresponding 3,4-*trans*-disubstituted adipamides with high enantioselectivities (up to 85% ee). © 1999 Elsevier Science Ltd. All rights reserved.

Although remarkable progress has been made in the field of asymmetric synthesis, the stereocontrol of radical reactions, especially intermolecular ones, still remain one of the most difficult problems in modern organic synthesis. Recently, some successful trials have been made in the field of free radicals.¹ However, the chiralities are, in most cases, involved in the substrates or reagents² (so-called substrate-or reagent-control) and the reactions which proceed under catalyst-control or ligand-control have not been extensively developed.³ Different from free radicals, conjugated ketyl radicals generated from α , β -unsaturated amides, by treating them with a one-electron transfer reagent such as SmI₂, have anion parts, and therefore, it would be possible to kinetically differentiate the enantiotopic face of the ketyl radicals for the subsequent reaction with the aid of their counter cations coordinated with appropriate chiral ligands as illustrated in Fig. 1. We now report the ligand-controlled enantioselective hydrodimerization of acrylamide derivatives which proceed through the formation of the conjugated ketyl radicals (Scheme 1).⁴

The reaction was carried out as follows: To a cold solution of (R)-BINOL and an amine in THF was added a THF solution of SmI_2 at $-78^{\circ}C$ and the mixture was stirred for 0.5 h at the same temperature, and then N,N-dibenzylcrotonamide **1a** was added. As shown in Table 1, the chiral complexes having the molar ratio of $SmI_2:(R)$ -BINOL:amine=1:2:4 afforded the product **2a** (Scheme 2) with higher enantioselectivities than the corresponding 1:1:2 complexes (cf. entries 1 versus 2, and 3 versus 4).^{5,6} No *meso* isomer was produced as previously reported.⁷

^{*} Corresponding author. E-mail: inanaga@ms.ifoc.kyushu-u.ac.jp

^{0040-4039/99/\$ -} see front matter © 1999 Elsevier Science Ltd. All rights reserved. *P11:* \$0040-4039(99)01551-8



Figure 1. Kinetic discrimination of the enantiotopic face of conjugated ketyl radicals by chiral ligands



Table 1

Effect of molar ratios and amines on the enantioselectivity of the hydrodimerization of 1a

Entry			Product (2a)	
	Chiral Ligand (eq)		Yield (%)	ee (%) ^{a)}
1 ^{b)}	(R)-BINOL (2)	<i>i</i> -Pr ₂ NEt (4)	34	55 (+)
2	(R)-BINOL (4)	<i>i</i> -Pr ₂ NEt (8)	20	69 (+)
3 ^{b)}	(R)-BINOL (2)	TMEDA (4)	38	33 (+)
4	(R)-BINOL (4)	TMEDA (8)	2 9	71 (+)
5	(R)-BINOL (4)	Et ₃ N (8)	6	60 (+)
6	(R)-BINOL (4)	(-)-sparteine (8)	14	50 (+)
7	(S)-BINOL (4)	(-)-sparteine (8)	32	61 (-)
8	(R)-BINOL (4)	N-methylpiperidine (8)	35	67 (+)
9	(R)-BINOL (4)	2,2,6,6-tetramethylpiperidine (8)	34	66 (+)
10	(R)-BINOL (4)	cis-1,2,6-trimethylpiperidine (8)	20	44 (+)

a) Determined by HPLC using Chiral PAK AD. The sign of the optical rotation is shown in parenthesis. b) Reaction time: 4 h.



Scheme 2.

A variety of amines as a component of the chiral samarium complexes were also examined for the reaction of 1a.⁸ As can be seen from Table 1, the use of diisopropylethylamine (entry 2), TMEDA (entry 4), *N*-methylpiperidine (entry 8) and 2,2,6,6-tetramethylpiperidine (entry 9) afforded relatively high ees. When (-)-sparteine was used in combination with (*R*)-BINOL or (*S*)-BINOL, where a matching or mismatching effect can be anticipated, a small difference was observed in the product's ee indicating that (*S*)-BINOL/(-)-sparteine is a matching pair (entry 6 versus 7). In order to clarify the scope and limitation of the present method, the reactions of other substrates were examined. The reactions of 1a-1g were

carried out by using SmI₂ (8 equiv.),⁹ (R)-BINOL (16 equiv.), and TMEDA (32 equiv.) in THF at -78° C for 4 h (Scheme 3) and the results are summarized in Table 2. As expected, high enantioselectivities (up to 85% ee) were obtained when primary alkyl-substituted acrylic amides **1a–1d** were used as substrates. As the β -substituent becomes bulky, the yield of the coupling product decreased (entries 1–4), and in the case of **1e** (R=*i*-Pr) and **1f** (R=*t*-Bu), only the saturated amides **3e** and **3f** were almost quantitatively produced. The reaction of *N*,*N*-diphenylcrotonamide (**1g**) afforded the coupling product **2g** as a mixture of diastereomers (entry 7). The high reactivity of **1g**, and hence the high concentration of the conjugated ketyl radical intermediate, might be responsible for the formation of the *meso* isomer. It is also interesting to note that the sense of enantioselection is reversed (entry 1 versus 7).



 Table 2

 Effect of the substituents of substrates on the enantioselectivity

Entry	Substrate		Product / Yield (%)		2 a -g		
	R	R'		2a-g	3a-g	dl : meso	ee (%) ^{a)}
1	Me	Bn	(1a)	70	20	di oniy	71 (+)
2	Et	Bn	(1b)	45	42	di only	82 (+)
3	<i>n-</i> Pr	Bn	(1c)	35	46	di only	82 (+)
4	BnCH ₂	Bn	(1d)	20	52	dl only	85 (+) ^{b)}
5	<i>i</i> -Pr	Bn	(1e)	_	95	-	_
6	t-Bu	Bn	(1f)		99	_	—
7	Me	Ph	(1g)	55	44	63:37	44 (-)

a) Determined by HPLC using Chiral PAK AD unless otherwise noted. The sign of the optical rotation is shown in parenthesis. b) Determined by HPLC using Chiral PAK OT(+).

A possible nine-membered chelate transition state which leads to the enantioselective C–C bond formation is shown in Fig. 2, in which the conjugated ketyl radical and the ligated crotonamide are arranged in a *cis*-relationship on the chiral coordination sphere of samarium.



Figure 2. A possible transition state for the enantioselective hydrodimerization of crotonamide 1a

In conclusion, the use of 2 equiv. of enantiopure BINOL in combination with achiral tertiary amines for the SmI₂-promoted reductive homo-coupling reaction of β -substituted acrylic acid amides brought about a high degree of asymmetric inductions (up to 85% ee). This is the first successful example of the completely diastereoselective and highly enantioselective C–C bond formation which proceeds through conjugated radical intermediates. Although the chemical yields are not always high, the method would be useful for the preparation of optically active 3,4-*trans*-disubstituted adipic acid derivatives, because they can be obtained in one step with high ees and the chiral source (BINOL) can be quantitatively recovered in an enantiopure state by aqueous alkaline extraction.

Acknowledgements

This work was supported by a Grant-in-Aid for Scientific Research on Priority Areas from the Ministry of Education, Science, Sports, and Culture, Japan.

References

- For reviews, see: (a) Curran, D. P.; Porter, N. A.; Giese, B. Stereochemistry of Radical Reactions; VCH: Weinheim, 1995;
 (b) Giese, B. Angew. Chem., Int. Ed. Engl. 1989, 28, 969; (c) Porter, N. A.; Giese, B.; Curran, D. P. Acc. Chem. Res. 1991, 24, 296; (d) Smadia, W. Synlett 1994, 1.
- For recent examples of the substrate-control, see: (a) Stack, J. G.; Curran, D. P.; Rebek Jr., J.; Ballester, P. J. Am. Chem. Soc. 1991, 113, 5918; (b) Porter, N. A.; Rosensteine, I. J.; Breyer, R. A.; Bruhnke, J. D.; Wu, W. X.; McPhail, A. T. J. Am. Chem. Soc. 1992, 114, 7664; (c) Yamamoto, Y.; Onuki, S.; Yumoto, M.; Asao, N. J. Am. Chem. Soc. 1994, 116, 421; (d) Sibi, M. P.; Jasperse, C. P.; Ji, J. Am. Chem. Soc. 1995, 117, 10779; (e) Fukuzawa, S.; Seki, K.; Tatsukawa, M.; Mutoh, K. J. Am. Chem. Soc. 1997, 119, 1482. For the reagent-control, see: Curran, D. P.; Nanni, D. Tetrahedron: Asymmetry 1996, 7, 2417.
- For recent examples of the catalyst-control, see: (a) Urabe, H.; Yamashita, K.; Suzuki, K.; Kobayashi, K.; Sato, F. J. Org. Chem. 1995, 60, 3576; (b) Murakata, M.; Tsutsui, H.; Hoshino, O. J. Chem. Soc., Chem. Commun. 1995, 481; (c) Wu, J. H.; Radinov, R.; Porter, N. A. J. Am. Chem. Soc. 1995, 117, 11029; (d) Sibi, M. P.; Ji, J. J. Am. Chem. Soc. 1996, 118, 9200; (e) Sibi, M. P.; Ji, J. J. Org. Chem. 1997, 62, 3800; (f) Murakata, M.; Jono, T.; Mizuno, Y.; Hoshino, O. J. Am. Chem. Soc. 1997, 119, 11713. For the ligand control, see: Mikami, K.; Yamaoka, M. Tetrahedron Lett. 1998, 39, 4501.
- 4. A part of this work was presented at the 16th Meeting of the Rare Earth Society of Japan, Osaka, May, 1999.
- 5. Other BINOL derivatives such as 3,3'-diphenyl-BINOL, 6,6'-diphenyl-BINOL, and H₈-BINOL gave less satisfactory results. Also, the use of dilithium binaphthoxide (4 equiv.) in the presence of *t*-butanol in place of the BINOL-amine system afforded **2a** with low ee (6%).
- 6. Ln(OTf)₃:BINOL:amine=1:1:2 system has successfully been used for the asymmetric Diels-Alder reaction: Kobayashi, S.; Ishitani, H. J. Am. Chem. Soc. **1994**, 116, 4083.
- 7. (a) Inanaga, J.; Handa, Y.; Tabuchi, T.; Otsubo, K.; Yamaguchi, M.; Hanamoto, T. Tetrahedron Lett. 1991, 32, 6557; (b) Kanemasa, S.; Yamamoto, H.; Kobayashi, S. Tetrahedron Lett. 1996, 37, 8505.
- 8. No reaction took place at -78° C in the absence of amines.
- 9. As the chiral samarium complex gradually decomposed even at -78°C, an excess amount of the reagent was used.