

## Journal Pre-proofs

Research paper

Facile Synthesis of Heterobimetallic  $[\text{Fe}^{\text{II}}(\mu\text{-diphosphine})\text{Ru}^{\text{II}}]$  and Homobimetallic  $[\text{Fe}^{\text{II}}(\mu\text{-diphosphine})\text{Fe}^{\text{II}}]$  Complexes and their in vitro Cytotoxic Activity on Cisplatin-resistant Cancer Cells

Danya BenYosef, Dario Romano, Mouna Hadiji, Paul J. Dyson, Burgert Blom

PII: S0020-1693(20)30851-3  
DOI: <https://doi.org/10.1016/j.ica.2020.119731>  
Reference: ICA 119731


To appear in: *Inorganica Chimica Acta*

Received Date: 30 April 2020  
Accepted Date: 1 May 2020

Please cite this article as: D. BenYosef, D. Romano, M. Hadiji, P.J. Dyson, B. Blom, Facile Synthesis of Heterobimetallic  $[\text{Fe}^{\text{II}}(\mu\text{-diphosphine})\text{Ru}^{\text{II}}]$  and Homobimetallic  $[\text{Fe}^{\text{II}}(\mu\text{-diphosphine})\text{Fe}^{\text{II}}]$  Complexes and their in vitro Cytotoxic Activity on Cisplatin-resistant Cancer Cells, *Inorganica Chimica Acta* (2020), doi: <https://doi.org/10.1016/j.ica.2020.119731>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier B.V.



# Facile Synthesis of Heterobimetallic [Fe<sup>II</sup>( $\mu$ -diphosphine)Ru<sup>II</sup>] and Homobimetallic [Fe<sup>II</sup>( $\mu$ -diphosphine)Fe<sup>II</sup>] Complexes and their *in vitro* Cytotoxic Activity on Cisplatin-resistant Cancer Cells

Danya BenYosef,<sup>a</sup> Dario Romano,<sup>b,c</sup> Mouna Hadiji,<sup>d</sup> Paul J. Dyson,<sup>d</sup> Burgert Blom<sup>a,\*</sup>

<sup>a</sup>Maastricht Science Programme, Faculty of Science and Engineering, Maastricht University, Kapoenstraat 2, PO Box 616, 6200 MD, Maastricht, The Netherlands

<sup>b</sup>Department of Biobased Materials, Faculty of Science and Engineering, Maastricht University, Brightlands Chemelot Campus, Urmonderbaan 22, Geleen, 6167 RD, The Netherlands

<sup>c</sup>Aachen-Maastricht Institute for Biobased Materials, Brightlands Chemelot Campus, Urmonderbaan 22, Geleen, 6167 RD, The Netherlands

<sup>d</sup>Institut des Sciences et Ingénierie Chimiques, Ecole Polytechnique Fédérale de Lausanne (EPFL), CH1015 Lausanne, Switzerland

## Abstract


A series of heterobimetallic [Fe,Ru] and homobimetallic [Fe,Fe] complexes, featuring  $\mu$ -diphosphine bridges between the two metal centres, are reported along with their *in vitro* cytotoxicity activity on the A2780cisR cell-line. The known starting material [CpFe(CO)<sub>2</sub>(CH<sub>3</sub>)] (Cp =  $\eta^5$ -C<sub>5</sub>H<sub>5</sub>) (**1**) was reacted with dppe (1,2-bis(diphenylphosphino)ethane) forming the known mononuclear  $\eta^1$ -dppe complex: [CpFe(CO)(COCH<sub>3</sub>)( $\eta^1$ -dppe)] (**2**) and the known dinuclear complex: and [[CpFe(CO)(COCH<sub>3</sub>)<sub>2</sub>( $\eta^1$ -dppe)] (**3**). Dimer cleavage reactions between complex **2** and two ruthenium arene complexes (( $\eta^6$ -*p*-cymene)RuCl<sub>2</sub>)<sub>2</sub> and [[( $\eta^6$ -benzene)RuCl<sub>2</sub>]<sub>2</sub>] formed two new heterobimetallic complexes: [CpFe(CO)(COCH<sub>3</sub>)( $\mu$ -dppe)Ru( $\eta^6$ -cymene)Cl<sub>2</sub>] (**4**) and [CpFe(CO)(COCH<sub>3</sub>)( $\mu$ -dppe)Ru( $\eta^6$ -C<sub>6</sub>H<sub>6</sub>)Cl<sub>2</sub>] (**5**). A homobimetallic system, [[CpFe(CO)(I)]<sub>2</sub>( $\eta^1$ -dppb)] (**6**) was obtained by the facile reaction between of [CpFe(CO)<sub>2</sub>I] and 1,4-bis(diphenylphosphino)butane (dppb). Complex **6** was methylated using MeLi to generate the more lipophilic complex [[CpFe(CO)(CH<sub>3</sub>)<sub>2</sub>( $\eta^1$ -dppb)] (**7**). All of the complexes were fully characterized by spectroscopy (<sup>1</sup>H, <sup>13</sup>C {<sup>1</sup>H}, <sup>31</sup>P {<sup>1</sup>H} NMR; FTIR, UV-Vis), and high resolution mass spectrometry (HRMS). Density functional theory calculations (DFT) (Level of theory B3LYP, basis set for H, C, P, O, Cl is 6-31+G(d,p) and for Ru, Fe, I is DGDZVP) on complexes **5** and **6** are also reported. An excellent agreement between the DFT calculated infra-red (IR) spectra of the optimised geometries of **5** and **6** was found with the experimentally determined spectra. *In vitro* cytotoxicity studies of all complexes was carried out on A2780cisR (cisplatin resistant ovarian cancer cell-line) and compared to cisplatin as a positive control and RAPTA-C as a negative control. Although limited to only one cell-line, the results

show that the heterobimetallic complexes **4** and **5** are the most cytotoxic ( $IC_{50} = 4.9 \pm 0.4 \mu\text{M}$  and  $6.5 \pm 0.1 \mu\text{M}$  respectively), whilst the dinuclear  $[\text{Fe},\text{Fe}]$  complexes exhibit no, or very low cytotoxicity.

**Keywords:** Heterobimetallic complexes, Homobimetallic complexes, Anti-cancer activity,  $\mu$ -Phosphine Bridged Complexes

\*Corresponding author email: [burgert.blom@maastrichtuniversity.nl](mailto:burgert.blom@maastrichtuniversity.nl)

### TOC Graphic



### Highlights:

- Synthesis and characterisation of novel heterobimetallic iron-ruthenium and homobimetallic iron-iron complexes and their *in vitro* cytotoxicity on A2780cisR.
- First cytotoxic investigations of diphosphine bridged ( $\mu$ ) diiron complexes.
- DFT studies of the homo and heteronuclear complexes.
- Comparison of DFT calculated and experimentally determined IR spectra.

## Introduction

In 2020 the WHO estimates the number of new cancer cases to rise to 15 million and currently the disease is responsible for 12 % of mortality world-wide.<sup>1</sup> Chemotherapeutic interventions including metal based drugs, are widely employed for treatment,<sup>2</sup> to induce cell apoptosis in cancer cells.<sup>3,4</sup> Cisplatin, found to exhibit anticancer properties in the 1960s by Rosenberg, heralded a new era in cancer chemotherapy.<sup>5</sup> The well-studied mechanism of action of this drug involves impairing cell function by binding to two guanine bases in DNA distorting DNA and inhibiting replication.<sup>6-8</sup> DNA.platinum adducts are recognized by certain proteins which either initiate DNA repair mechanisms, or initiate apoptotic cell death. While cisplatin is a highly effective drug, it causes nephrotoxicity and neurotoxicity and other side effects including hair loss, nausea and vomiting.<sup>9</sup> Other limitations of cisplatin include intrinsic drug resistance or resistance after exposure to some types of tumours, as well as low solubility in aqueous media.<sup>10</sup> Thus, platinum analogues have been developed and studied in order to circumvent these limitations.<sup>11-15</sup> In addition, complexes based on other metals such as gold,<sup>16,17</sup> titanium,<sup>16</sup> ruthenium<sup>18-28</sup> and iron have also been reported to exhibit anti-cancer properties.<sup>29-32</sup> One area that has attracted the attention is the synthesis heterobimetallic systems where potential cooperative behaviour can lead to the potentiation of cytotoxic activity. Systems containing gold(I) and platinum(II) centres, for example, display improved antiproliferative activity, as well as overcome drug resistance issues seen in cisplatin.<sup>34</sup> A study by Fernandez-Gallardo *et al.* shows that ruthenium-gold heterobimetallic systems reveal higher selectivity and cytotoxicity towards renal and colon cancer cell lines than their mononuclear analogs.<sup>35</sup> Serra *et al.* elucidated the enhanced redox properties that iron-ruthenium heterobimetallic systems have for catalytic applications, which served as inspiration for our own work.<sup>33</sup>

We have focussed on the development of heterobimetallic complexes and explored their use in anti-cancer applications. Our initial focus was arene-ruthenium(II) half-sandwich complexes with one or two trichlorostannyl ligands  $\sigma$ -bound to the Ru(II) centre. The compounds showed poor cytotoxicity due to low solubility in aqueous media, which could be overcome in ionic compounds

or by increasing solubility in aqueous media.<sup>36,37</sup> Ruthenium(II) arene complexes bearing a  $\sigma$ -bound germyl group were also prepared and showed low to negligible cytotoxic activity, which was postulated to be due to rapid aquation kinetics.<sup>38</sup> We have also focussed on homo and heterobimetallic systems bearing two group 8 metals and their respective cytotoxicity (Figure 1). In this regard [Fe,Ru] and [Ru,Ru] complexes bridged with  $\mu$ -dppm (dppm = 1,1-bis(diphenylphosphino)methane)) were synthesized with the expectation that the presence of iron in the bimetallic compound may increase drug uptake.<sup>39a</sup> The study revealed that a heteronuclear [Fe,Ru] system was indeed substantially more cytotoxic than cisplatin, on A2780 and A2780cisR cell-lines, while a comparable dinuclear [Ru,Ru] system showed a decrease in cytotoxicity.

In the present study, we report on our latest findings in this area, in which we explored the effects of increasing the lipophilicity at the iron centre by replacing the iodo ligand with an acyl group and explored the effect of increasing the spacer length between the Fe and Ru centres to observe changes in cytotoxicity. Studies by Hartinger, for example have shown that spacer length plays a role in cytotoxic action.<sup>39c</sup> Moreover, the preparation and cytotoxic testing of homobimetallic  $\mu$ -diphosphine [Fe,Fe] complexes is also reported (Figure 1).





Figure 1. Heterobimetallic and homobimetallic complexes reported which exhibit promising cytotoxic profiles and cancer cell selectivity and the compounds reported in the present study.


## Results and Discussion

Our strategy to prepare heterobimetallic [Fe,Ru] complexes is similar to the one we employed previously,<sup>39a</sup> requiring a suitable mononuclear iron complex, bearing a  $\kappa^1$  metal bound bisphosphine ligand with a pendant uncoordinated phosphorus atom of the type  $L_nFe \leftarrow :P-P:$ . The pendant uncoordinated phosphine could then, in a second synthetic step, cleave Ru dimer complexes of the type  $[RuCl_2(\eta^6\text{-arene})]_2$  (arene = benzene and *p*-cymene),<sup>40</sup> to afford the desired dinuclear complexes. We sought complexes bearing a  $\kappa^1$  dppe (dppe = 1,1-bis(diphenylphosphino)ethane)) ligand, as previously we employed dppm (dppe = 1,1-bis(diphenylphosphino)methane)),<sup>39a</sup> and wanted to increase the spacer length between the Fe and Ru centres. Moreover, we were also interested in obtaining an iron centre which was more lipophilic than in our earlier studies, since we envisioned higher cellular uptake and potential efficacy. A suitable agent for these purposes was the known complex  $[CpFe(CO)(COCH_3)(\eta^5\text{-}C_5H_5)]$  (Cp =  $\eta^5\text{-}C_5H_5$ ) (**2**).<sup>43,44</sup> Complex **2** is accessible *via* a facile migratory insertion route using dppe starting from the known complex  $[CpFe(CO)_2(CH_3)]$  **1**.<sup>41,42</sup> Complexes  $[CpFe(CO)(COCH_3)(\eta^5\text{-}C_5H_5)]$  **2** and  $[[CpFe(CO)(COCH_3)]_2(\mu\text{-dppe})]$  **3**,<sup>45</sup> the latter which exists as a mixture diastereoisomers, are obtained from this reaction (Scheme 1).<sup>43,44</sup> While compounds **2** and **3** are known, their cytotoxicity has not been determined, and we decided to include them in our *in vitro* cytotoxicity screening studies (see below).



Scheme 1. Reaction scheme for the synthesis of products  $[\text{CpFe}(\text{CO})(\text{COCH}_3)(\mu\text{-dppe})]$  **2** and  $[[\text{CpFe}(\text{CO})(\text{COCH}_3)]_2(\mu\text{-dppe})]$  **3**. Complexes **2** and **3** possess asymmetric centres (Fe). For complex **2** this exists as a mixture of enantiomers (*rac*-**2**) (only one is depicted), while for complex **3** three stereoisomers are possible, i.e. the *meso* form and a set of enantiomers (only one is depicted in Scheme 1).<sup>46</sup>

Employing complex **2** as starting material (which is chiral and exists as a mixture of enantiomers: *rac*-**2**) the reaction thereof with the Ru dimer complexes  $[\text{RuCl}_2(\eta^6\text{-}p\text{-cymene})]_2$  and  $[\text{RuCl}_2(\eta^6\text{-C}_6\text{H}_6)]_2$  in a 2:1 ratio afforded the heterobimetallic complexes  $[\text{CpFe}(\text{CO})(\text{COCH}_3)(\mu\text{-dppe})\text{Ru}(\eta^6\text{-}p\text{-cymene})\text{Cl}_2]$  **4** and  $[\text{CpFe}(\text{CO})(\text{COCH}_3)(\mu\text{-dppe})\text{Ru}(\eta^6\text{-C}_6\text{H}_6)\text{Cl}_2]$  **5**, respectively (Scheme 2) in a straightforward manner. The  $\mu\text{-dppe}$  bridged  $[\text{Fe},\text{Ru}]$  complexes with no Fe-Ru bond are somewhat rare in the literature.<sup>39b-d</sup> Complex **4** was isolated as a clay red/orange powder in an excellent yield of 85%, while complex **5** required column chromatographic purification using an eluent mixture of 1:1 *n*-hexane/ethyl acetate. The latter afforded a light brown powder with a suboptimal yield of 34%. Both complexes exhibit high decomposition temperatures: **4** at 191 °C, and complex **5** at 130 °C.




Scheme 2. Reaction scheme for the synthesis of complex  $[\text{CpFe}(\text{CO})(\text{COCH}_3)(\mu\text{-dppe})\text{Ru}(\eta^6\text{-}p\text{-cymene})\text{Cl}_2]$  **4** and  $[\text{CpFe}(\text{CO})(\text{COCH}_3)(\mu\text{-dppe})\text{Ru}(\eta^6\text{-C}_6\text{H}_6)\text{Cl}_2]$  **5**. The iron centre in both complexes is asymmetric and complexes **4** and **5** exist as a racemic mixture (*rac*-**4** and *rac*-**5**). Only one stereoisomer is depicted for clarity in both cases.

The full spectroscopic characterization consisting of multinuclear NMR ( $^1\text{H}$ ,  $^{31}\text{P}\{^1\text{H}\}$ ,  $^{13}\text{C}\{^1\text{H}\}$ ) FTIR, and ESI-MS, was carried out for both complexes. In complex **4**, a series of unresolved multiplet resonances in the range  $\delta = 7.84$  to  $7.05$  ppm correspond to the 20 aromatic protons in dppe. Four distinct doublets are observed from  $\delta = 5.24$  to  $4.95$  ppm, corresponding to the 4 inequivalent protons in the *p*-cymene ring, due to the asymmetry at the iron centre. The singlet resonance for  $\eta^5\text{-C}_5\text{H}_5$  is observed at  $\delta = 4.39$  ppm, and the multiplet signals that follow from  $\delta = 2.70$  to  $2.45$  ppm correspond to the inequivalent protons in the dppe bridge. The consecutive singlets represent the methyl groups in the compound: the  $\text{COCH}_3$  resonates at  $\delta = 2.25$  ppm and the  $\text{CH}_3$  from the *p*-cymene is seen at  $\delta = 1.84$  ppm. Finally, the isopropyl methyl groups from the *p*-cymene substituent appear as 2 distinct doublet resonances at  $\delta = 0.88$  and  $0.80$  ppm respectively confirming the diastereotopic nature of the *iPr* groups. The aromatic signals for complex **5** appear at similar ppm values to **4**. A singlet is observed at  $\delta = 5.32$  for complex **5**, and corresponds to 6 equivalent protons from the benzene ( $\eta^6\text{-C}_6\text{H}_6$ ) substituent. Complex **5** shows the  $\eta^5\text{-C}_5\text{H}_5$  resonance at  $\delta = 4.39$  ppm, as well as similar ppm values for the protons in the dppe bridge as with **4**. There is only one singlet resonance for the only methyl group in the compound ( $\text{COCH}_3$ ), the latter which is seen at  $\delta = 2.35$  ppm. The  $^{31}\text{P}\{^1\text{H}\}$  NMR spectra for complexes **4** and **5** are very similar and each contain two doublet resonances. A low-field shifted doublet signal for both complexes is seen at  $\delta = 74.3$  ppm, representing the  $\text{Fe-PPh}_2$  bond, while the second high-field shifted signal, also seen in both complexes, appears at  $\delta = 25.1$  ppm which corresponds to the  $\text{Ru-PPh}_2$  bond. The  $^{13}\text{C}\{^1\text{H}\}$  NMR data for complex **4** shows a series of unresolvable multiplets from  $\delta = 135.5$  to  $126.7$  ppm for the carbon atoms in the aromatic substituents of dppe. Singlets which appear at  $\delta = 107.4$  and  $93.4$  ppm correspond to two C atoms on *p*-cymene ring, while the remaining carbon atoms within the *p*-cymene ring are represented by two sets doublets: one doublet appears at  $\delta = 89.9$  ppm due to  $^2\text{J}(\text{C},\text{P})$  coupling. The singlets at  $\delta = 83.6$  ppm pertains to the equivalent carbons in the  $\eta^5\text{-C}_5\text{H}_5$  group, and the doublet at  $\delta = 50.9$  ppm the methyl carbon from  $\text{COCH}_3$  coupling to phosphorus atom.

Complex **5** shows similar  $^{13}\text{C}\{^1\text{H}\}$  NMR features as **4**. The spectrum displays a doublet resonance at  $\delta = 219.4$  ppm with a  $^2\text{J}(\text{C},\text{P})$  coupling constant of 30 Hz, for the  $\text{Fe-CO}$ . A singlet at  $\delta = 87.6$  ppm corresponds to the benzene substituent and the signal at  $\delta = 83.7$  ppm corresponds to  $\eta^5\text{-C}_5\text{H}_5$ . The FTIR spectra for complexes **4** and **5** reveal the strong stretching vibrations for both the  $\text{Fe-}$

(CO) and the Fe-(C=O)CH<sub>3</sub>: For complex **4**, these are visible at  $\nu = 1901$  and  $1581\text{ cm}^{-1}$  respectively, while for complex **5**, the stretching vibrations appear at  $1897$  and  $1597\text{ cm}^{-1}$ . Both complexes were subjected to high resolution ESI-MS (+) and exhibit signals corresponding the radical cation  $[M]^{\cdot+}$  and  $[M+Na]^+$ , both with matching isotope patterns confirming their constitution. In order to test the stability of the complexes in (DMSO, see below),  $^{31}\text{P}\{^1\text{H}\}$  NMR spectra were recorded of complex **5** dissolved in DMSO-*d*<sub>6</sub> (containing water) after different time intervals.<sup>47</sup> The sample was tested each hour during the first 7 hours, after 24 hours, and finally after 72 hours. When observing the spectra, it is evident that there are no changes in this time period.

We next targeted the homobimetallic [Fe,Fe] complex  $[[\text{CpFe}(\text{CO})(\text{I})]_2(\text{dppb})]$  (complex **6**), bearing the longer tether (dppb as opposed to dppe). We decided to use dppb to explore the effect of a longer spacer between the two metal centres, and to exclude formation of a  $\kappa^2$ -phosphine product. Surprisingly, homobimetallic complexes of the type  $[\text{Fe}(\mu\text{-phosphane})\text{Fe}]$ , although highly prevalent in the literature and used *inter alia* as iron hydrogenase models, have altogether escaped investigation as potential anti-cancer agents, and herein we report the first such study (see below).<sup>48</sup> The complex  $[\text{CpFe}(\text{CO})_2\text{I}]$  and dppb (dppb = 1,1-bis(diphenylphosphino)butane) was dissolved in toluene and refluxed at  $85\text{ }^\circ\text{C}$  for 7 hours. The crude reaction mixture was purified by column chromatography using a 3:2 eluent mixture of dichloromethane/*n*-hexane which exhibited three separable bands. The third band corresponded to  $[[\text{CpFe}(\text{CO})(\text{I})]_2(\text{dppb})]$ , which was isolated as a light green powder in a 37 % yield (Scheme 3) as a mixture of stereoisomers.




Scheme 3. Reaction scheme for the synthesis of complex **6**  $[[\text{CpFe}(\text{CO})(\text{I})]_2(\text{dppb})] \cdot 0.5$  hexane

Complex **6** exists as a mixture of three stereoisomers: a pair of enantiomers and a *meso* form akin to complex **3**. The complex was also subjected to full spectroscopic characterisation. The  $^1\text{H}$  NMR spectrum reveals multiplet resonances from  $\delta = 7.71$  to  $7.35$  ppm correspond to the 20 protons of the phenyl rings of dppb, and a singlet at  $\delta = 4.32$  ppm with an integration of 10 protons corresponds to two equivalent  $\eta^5\text{-C}_5\text{H}_5$  groups in the dinuclear compound. Two multiplet resonances at  $\delta = 2.37$  and  $0.95$  ppm respectively represent the protons of the dppb bridge. The remaining signals correspond to half a molecule of hexane included in the solid. In the  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum, two singlets appear at  $\delta = 61.7$  ppm and  $\delta = 61.6$  ppm, akin to complex **3**; indicating the diastereoisomers in the *meso* form and pair of enantiomers in D,L form. The FTIR confirms the presence of one CO stretching vibration with a medium strong intensity at  $1932\text{ cm}^{-1}$ . The ESI-MS spectra of **6** show a complex fragmentation pattern including the  $[\text{M}]^+$  signal at  $977.9068\text{ m/z}$  and a  $[\text{M}+\text{Na}]^+$  signal at  $1000.9033\text{ m/z}$ , which can be compared well to the theoretical values ( $[\text{M}]^+ 977.9129$ ; for  $[\text{M}+\text{Na}]^+ 1000.9028$ ) confirming constitution.  $^{31}\text{P}\{^1\text{H}\}$  NMR spectra were also recorded of complex **6** in wet  $\text{DMSO-}d_6$  after different time intervals, showing stability.<sup>47</sup> The compound, as with the  $[\text{Fe},\text{Ru}]$  complexes shows stability over this time period (see SI for the spectra).

The related complex **7**,  $[[\text{CpFe}(\text{CO})(\text{CH}_3)]_2(\text{dppb})] \cdot 0.5$  hexane, could readily be isolated upon reaction of complex **6** with MeLi (Scheme 4). After column chromatographic purification using

$\text{CH}_2\text{Cl}_2$ , the compound was successfully isolated as a vermilion powder with a yield of 46.5%. The complex is thermally robust and onset of colour change starts at 134 °C with full decomposition at 146 °C.



Scheme 4. Reaction scheme for the synthesis of complex 7  $[(\text{CpFe}(\text{CO})(\text{CH}_3)_2(\text{dppb}))] \cdot 0.5$  hexane

The  $^1\text{H}$  NMR reveals the Fe- $\text{CH}_3$  resonance at  $\delta = -0.29$  ppm as a doublet of doublets due to coupling to P. In contrast to the precursor **6** the  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum shows only one singlet signal at  $\delta = 75.5$  ppm for the Fe- $\text{PPh}_2$  although two signals are expected due to the diastereoisomerism possibly due to isochronicity. In the FTIR spectrum, the CO stretching vibration appears at  $\nu = 1892$   $\text{cm}^{-1}$ , shifted by 40 wavenumbers from its precursor, **6**, due to enhanced electron density at the iron centre afforded by the methyl group.

Complexes **2-7** were subjected to *in vitro* cytotoxicity studies on the cisplatin resistant ovarian cancer cell line A2780cisR and compared to cisplatin as positive control and RAPTA-C as negative control (Table 1). In the series, the heterobimetallic [Fe,Ru] complexes **4** and **5** exhibit promising cytotoxic activity and are the most active in the series, with the cymene analogue **4** being slightly more cytotoxic than the benzene analogue **5**. This is in keeping with our earlier findings of related complexes bearing a dppm bridge between the Fe and Ru centres, i.e.  $[(\eta^6\text{-C}_6\text{H}_6)\text{RuCl}_2(\mu\text{-dppm})\text{Fe}(\text{CO})\text{I}(\eta^5\text{-C}_5\text{H}_5)]$  ( $\text{IC}_{50} = 1.5$   $\mu\text{M}$  on A2780cisR) and  $[(\eta^6\text{-p-cymene})\text{RuCl}_2(\mu\text{-dppm})\text{Fe}(\text{CO})\text{I}(\eta^5\text{-C}_5\text{H}_5)]$  ( $\text{IC}_{50} = 1.2$   $\mu\text{M}$  on A2780cisR).<sup>39a</sup>

Interestingly, the bimetallic [Fe,Fe] complexes **3** and **6** are inactive with  $\text{IC}_{50}$  values  $> 100$   $\mu\text{M}$ , while the mononuclear iron complex **2** exhibits moderate activity. Complex **7**, the methylated analogue of **6**, exhibits some, albeit it low, cytotoxicity. These latter results suggest that increasing the lipophilicity of the ligand at the iron centre (I in **6** vs  $\text{CH}_3$  in **7**) does enhance cytotoxicity, but overall the dinuclear [Fe,Fe] complexes are rather inactive.

Complex	IC <sub>50</sub> (μM)
<b>2</b>	8.5 ± 0.1
<b>3</b>	>100
<b>4</b>	4.9 ± 0.4
<b>5</b>	6.5 ± 0.1
<b>6</b>	> 100
<b>7</b>	61 ± 3
Cisplatin	35 ± 3
RAPTA-C	> 200

Table 1. IC<sub>50</sub> (μM) values of complexes **2-7** on the A2780cisR cell-line after 72 h exposure to the compound. The values represent the mean between four replicates and ± the std. deviation.

Structural, electronic and spectroscopic characteristics for complexes **5** and **6** were calculated using density function theory methods (DFT). For all the calculations, the level of theory employed was B3LYP with the basis set 6-31+G(d,p) for H, C, O, P and Cl atoms; while for I, Ru and Fe atoms DGDZVP basis set was used. Since we were unable to obtain suitable single crystals for X-ray diffraction analysis of the complexes, despite several attempts and crystallisation techniques, we obtained structural information of the complexes *via* DFT methods. Moreover, the calculated IR spectra of the optimised structures (in the gas phase), were compared to those obtained by experiment (solid state IR). These showed an excellent agreement, in both **5** and **6**, suggesting reasonable structural models for both complexes using DFT methods. Full identification and assignment of the vibrational modes in the IR spectra was also possible for **5** (see SI Fig. 5.5) and **6** (see SI Fig. 6.6) in light of this. Figure 2 shows the optimised structures of complex **5** and **6** with some key metrical parameters obtained from DFT.




Figure 2. DFT optimised structure of complex **5** (left). Selected bond lengths [ $\text{\AA}$ ]: Fe-P2 2.260, Ru-P1 2.387, Fe-C1 1.960, Fe-C2 1.749, Ru-Cl1 2.440, Ru-Cl2 2.466. Fe-P2-P1-Ru dihedral angle:  $145^\circ$ . DFT optimised structure of complex **6** (right). Selected bond lengths [ $\text{\AA}$ ] Fe1-I2/Fe2-I2 2.696, Fe1-P1/Fe2-P2 2.330, Fe1-C<sub>carbonyl</sub>/Fe2-C<sub>carbonyl</sub> = 2.696. Fe1-P1-P2-Fe2 dihedral angle:  $163^\circ$ .

The most striking structural feature of both complexes is the zig-zag arrangement of the heavy atoms, with the two metal centres being somewhat *anti* with respect to each other. In complex **5** a Fe-P-P-Ru dihedral angle of  $145^\circ$  is observed, while complex **6** exhibited an Fe1-P1-P2-Fe2 dihedral angle of  $163^\circ$  (Figure 3).




Figure 3. Perspective of **5** (left) and **6** (right) viewed down the P-P vector. All atoms except the Fe, Ru and P are omitted for clarity.

The electronic nature of complexes **5** and **6** was also studied. Figure 4 shows the highest occupied molecular orbital (HOMO) and the lowest unoccupied molecular orbital (LUMO) with the corresponding molecular orbitals' energies of the two complexes.




Figure 4. LUMO (top) and HOMO (bottom) of the optimized structures of complex **5** (left) and **6** (right). The energies of the LUMOs (top) and HOMOs (bottom) are expressed in hartree. Blue = iron, red = oxygen, grey = carbon, orange = phosphorus, green = chlorine,

Clear differences can readily be observed for the localizations of the HOMO and LUMO when comparing complexes **5** and **6**. For complex **5**, due to its asymmetrical nature, the HOMO and the LUMO are asymmetrically localized on the complex. In particular, the HOMO of complex **5** is delocalized on the iron metal and the ligands (Cp, CO, one phosphorous of the dppe and acetyl), while the LUMO is delocalized on the ruthenium terminus over the ligands (arene and chlorides) excluding the phosphorus atoms of the bridging dppe. This is in contrast to our earlier studies of related heterobimetallic [Fe,Ru] complexes, where *both* HOMO and LUMO were located on the iron terminus of the complex in  $[(\eta^6\text{-Arene})\text{RuCl}_2(\mu\text{-dppe})\text{Fe}(\text{CO})\text{I}(\eta^5\text{-C}_5\text{H}_5)]$  (Ar = C<sub>6</sub>H<sub>6</sub> and *p*-

cymene, dpmm = 1,1-bis(diphenylphosphinomethane)).<sup>39a</sup> The symmetric complex **6** shows the calculated HOMO and LUMO symmetrically localized on both metal centres with their corresponding ligands.

## Conclusions

A series of heterobimetallic [Fe,Ru] and homobimetallic [Fe,Fe] complexes, bridged with a dppe ligand, were prepared and fully characterised spectroscopically and their structures, electronic nature and IR spectra investigated by DFT methods, the latter showing excellent agreement with experiment. The DFT study revealed that for the heterobimetallic complexes, the HOMO and LUMO are not both located on the iron terminus of the complex, in contrast to earlier findings.<sup>39a</sup> An *in vitro* study of all dinuclear complexes was carried out on the A2780cisR cisplatin resistant ovarian cancer cell-line, and compared to the known mononuclear iron complex [CpFe(CO)(COCH<sub>3</sub>)(dppe)] **2**. While limited to only one-cell line, it can be seen that the heterobimetallic [Fe,Ru] complexes exhibit promising cytotoxicity profiles, while the corresponding [Fe,Fe] complexes exhibit very low activity; and the mononuclear complex **2** only moderate activity. Hence such homodinuclear [Fe,Fe] complexes, appear not be suitable candidates for further pursuit, whereas the [Fe,Ru] complexes have promising cytotoxicity.<sup>39a</sup> The increased spacer length, however, seems to slightly reduce cytotoxic action in the latter with the previously reported [Fe,Ru] complexes bearing a dpmm bridge being somewhat more cytotoxic.

## Experimental Section

All reactions, unless stated otherwise, were performed under a nitrogen atmosphere using Schlenk procedures and/or glove-box manipulations. All solvents were degassed by bubbling nitrogen gas for several minutes through them and dried by passage over activated alumina. The starting materials were obtained from commercial suppliers and used as received. Dicarbonylcyclopentadienyliodoiron(II) ([CpFe(CO)<sub>2</sub>I]) (97%), anhydrous tetrahydrofuran (99.9%), methyllithium (1.6 M in diethyl ether), dppe, dppb (99%) and silica gel (60 Å pore size, 70-230 mesh, 63-200 µm) were purchased from Sigma-Aldrich/Merck. Dichloromethane (stab. Amylene), as well as *n*-hexane and diethyl ether (stab. BHT), were obtained from Biosolve

Chemicals. Ethyl acetate (99.5%) was purchased from VWR, as well as chloroform-*d* +0.03% TMS and dimethyl sulfoxide-*d*<sub>6</sub> (99.80%). Dry toluene (99.85%) was obtained from Acros Organics. Dichloro(*p*-cymene)ruthenium(II) dimer ( $[(p\text{-cymene})\text{RuCl}_2]_2$ ) (98%) was purchased from STREM, while benzeneruthenium(II)chloride dimer ( $[(\text{benzene})\text{RuCl}_2]_2$ ) (95%) was purchased from TCI. The known complexes **2** and **3** were prepared by modified literature procedures and can be found in the SI.<sup>43-45</sup> The apparatus used to measure the melting points was a Stuart SMP10 and are uncorrected. NMR experiments at room temperature were recorded using a Bruker Ultrashield 300 MHz/54 mm magnet system. Chemical shifts were observed relative to the residual solvent peak. The <sup>1</sup>H and <sup>13</sup>C {<sup>1</sup>H} NMR chemical shifts were observed relative to tetramethylsilane (TMS), while for <sup>31</sup>P {<sup>1</sup>H} NMR, they were measured relative to phosphoric acid (85%). The software for spectral analysis was Bruker Topspin 4.0.6. FTIR spectra were retrieved using a MIRacle 10 Shimadzu, with a single reflection ATR accessory. Per sample, 64 scans were collected over a range of 400-4000 cm<sup>-1</sup>, with a spectral resolution of 1 cm<sup>-1</sup>; these were analysed using the IR solution software. UV-Vis spectra were obtained using a UV-188 spectrophotometer (Shimadzu) and quartz suprasil precision cells (100- QS, 10 mm light path, from Hellma Analytics). Finally, high resolution mass spectrometry (HR-ESI-MS) was recorded using a Bruker SolariX XR FT-ICR-MS in the Maastricht MultiModal Molecular Imaging Institute (M4I). The samples were dissolved in CHCl<sub>2</sub>-CH<sub>3</sub>CN- CH<sub>3</sub>COOH (49.95 – 49.95 – 0.1 %) and then diluted 100 times. The MS parameters were set in ionization + mode, with a capillary of 4.4 kV, end plate offset of -800V, a nebulizer at 1bar, dry gas at 4 L/min, a dry temperature of 200 °C, and a mass range of 300-1300 m/z. The resolution of all measurements was above 100000. Only the relevant signals are quoted in the experimental section. All isotope patterns were checked using online software; the isotope line with the highest intensity was reported for each sample.

### **Cytotoxicity Tests**

Human ovarian carcinoma (A2780cisR) cell lines were obtained from the European Collection of Cell Cultures Penicillin streptomycin, RPMI 1640 GlutaMAX (where RPMI = Roswell Park Memorial Institute), and DMEM GlutaMAX media (where DMEM = Dulbecco's modified Eagle medium) were obtained from Life Technologies, and fetal bovine serum (FBS) was obtained from Sigma. The cells were cultured in RPMI 1640 GlutaMAX (A2780cisR) and DMEM GlutaMAX (HEK-293) media containing 10% heat-inactivated FBS and 1% penicillin streptomycin at 37 °C

and CO<sub>2</sub> (5%). The A2780cisR cell line was routinely treated with cisplatin (2 μM) in the media to maintain cisplatin resistance. The cytotoxicity was determined using the 3-(4,5-dimethyl 2-thiazolyl)-2,5-diphenyl-2H-tetrazolium bromide (MTT) assay.<sup>49</sup> Cells were seeded in flat-bottomed 96-well plates as a suspension in a prepared medium (100 μL aliquots and approximately 4300 cells/well) and pre-incubated for 24 h. Stock solutions of compounds were prepared in DMSO and were diluted in medium. The solutions were sequentially diluted to give a final DMSO concentration of 0.5% and a final compound concentration range (0–200 μM). Cisplatin and RAPTA-C were tested as a positive (0–100 μM) and negative (200 μM) controls respectively. The compounds were added to the preincubated 96-well plates in 100 μL aliquots, and the plates were incubated for a further 72 h. MTT (20 μL, 5 mg/mL in Dulbecco's phosphate buffered saline) was added to the cells, and the plates were incubated for a further 4 h. The culture medium was aspirated and the purple formazan crystals, formed by the mitochondrial dehydrogenase activity of vital cells, were dissolved in DMSO (100 μL/well). The absorbance of the resulting solutions, directly proportional to the number of surviving cells, was quantified at 590 nm using a SpectroMax M5e multimode microplate reader (using SoftMax Pro software, version 6.2.2). The percentage of surviving cells was calculated from the absorbance of wells corresponding to the untreated control cells. The reported IC<sub>50</sub> values are based on the means from two independent experiments, each comprising four tests per concentration level.

### Density Functional Theory Calculations

DFT calculations were performed to model the complexes **5** and **6**. Gaussian09 software package was used. For all the calculations, the level of theory used for all calculations is B3LYP with the basis set 6-31+G(d,p) for H, C, O, P and Cl atoms; while for I, Ru and Fe atoms DGDZVP basis set was used. Geometry optimizations were calculated without any constrains. All the optimized geometries show not imaginary frequency. Energies and infrared spectra were calculated on the optimized structures.<sup>50</sup>

### Synthesis of [CpFe(CO)(COCH<sub>3</sub>)(η<sup>6</sup>-p-cymene)Ru(η<sup>6</sup>-p-cymene)Cl<sub>2</sub>] (**4**)

0.048 g of [CpFe(CO)(COCH<sub>3</sub>)(η<sup>6</sup>-p-cymene)] (**2**) (0.081 mmol, 1 eq.) and 0.024 g of [(η<sup>6</sup>-p-cymene)RuCl<sub>2</sub>]<sub>2</sub> (0.040 mmol, 0.5 eq) were added to a 100 mL round bottom Schlenk flask. The reactants were dissolved in approximately 20 mL of dichloromethane; the solution was left to stir

at room temperature for 2 hours. The solvent was evaporated under reduced pressure, the residue washed with hexane, and the hexane washings discarded. The resulting solid was dried *in vacuo* affording a clay red/orange powder (0.0616g, 0.068 mmol, 85%) with a decomposition temperature of 191 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>, 298 K): δ 7.84-7.05 (20H, m, C-H dppe), 5.24 (1H, d, <sup>3</sup>J(H,H)= 6.20 Hz, *p*-cymene C-H<sup>D</sup>), 5.20 (1H, d, <sup>3</sup>J(H,H)= 6.20 Hz, *p*-cymene C-H<sup>C</sup>), 5.13 (1H, d, <sup>3</sup>J(H,H)= 6.20 Hz, *p*-cymene C-H<sup>B</sup>), 4.95 (1H, d, <sup>3</sup>J(H,H)= 6.20 Hz, *p*-cymene C-H<sup>A</sup>), 4.39 (5H, s, η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>), 2.70 (1H, m, CHMe<sub>2</sub>), 2.49 (1H, m, PCH<sup>C</sup> bridge), 2.47 (1H, m, PCH<sup>B</sup> bridge), 2.45 (1H, m, PCH<sup>A</sup> bridge), 2.25 (3H, s, COCH<sub>3</sub>), 1.84 (3H, s, *p*-cymene CH<sub>3</sub>), 0.88 (3H, d, <sup>3</sup>J(H,H)= 7.02 Hz, *p*-cymene CH(CH<sub>3</sub>)<sup>B</sup>), 0.80 (3H, d, <sup>3</sup>J(H,H)= 7.02 Hz, *p*-cymene CH(CH<sub>3</sub>)<sup>A</sup>) ppm; <sup>31</sup>P{<sup>1</sup>H} NMR (CDCl<sub>3</sub>, 298 K): δ 74.3 (d, <sup>3</sup>J(P,P)= 34 Hz, Fe-PPh<sub>2</sub>), 25.1 (d, <sup>3</sup>J(P,P)= 34 Hz, Ru-PPh<sub>2</sub>) ppm; <sup>13</sup>C {<sup>1</sup>H} NMR (CDCl<sub>3</sub>, 298 K): δ 135.5-126.7 (m, dppe), 107.4 (s, *p*-cymene C<sup>1</sup> or 4-H), 93.4 (s, *p*-cymene C<sup>1</sup> or 4-H), 89.9 (d, <sup>2</sup>J(C,P)= 4.7 Hz, *p*-cymene C<sup>A</sup>-H), 88.6 (d, <sup>2</sup>J(C,P)= 3.8 Hz, *p*-cymene C<sup>B</sup>-H), 84.8 (d, <sup>2</sup>J(C,P)= 6.0 Hz, *p*-cymene C<sup>C</sup>-H), 84.1 (d, <sup>2</sup>J(C,P)= 5.7 Hz, *p*-cymene C<sup>C</sup>-H), 83.6 (s, η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>), 50.9 (d, <sup>3</sup>J(C,P)= 5.7 Hz, COCH<sub>3</sub>), 28.9 (s, *p*-cymene CH(CH<sub>3</sub>)<sub>2</sub>), 23.9 (m, dppe), 21.6 (s, *p*-cymene *iPr* CH<sub>3</sub><sup>A</sup>), 20.6 (s, *p*-cymene *iPr* CH<sub>3</sub><sup>B</sup>), 19.7 (d, <sup>3</sup>J(C,P)= 28.8 Hz, dppe), 16.3 (s, *p*-cymene CH<sub>3</sub>) ppm; FTIR (cm<sup>-1</sup>): ν= 3059 (vw), 1948 (w), 1901 (m, CO stretch), 1886 (m), 1581 (m, CO stretch), 1481 (w), 1435 (m), 1321 (w), 1261 (vw), 1178 (w), 1093 (w), 1060 (w), 999 (w), 906 (w), 889 (w), 819 (w), 748 (m), 717 (m), 694 (ms) cm<sup>-1</sup>; ESI-MS: *m/z* Calcd. For [M]<sup>+</sup> 896.0745, found 896.0702. Calcd. For [M+Na]<sup>+</sup> 919.0635, found 919.0607.

### Synthesis of [CpFe(CO)(COCH<sub>3</sub>)(<sup>1</sup>-dppe)Ru(η<sup>6</sup>-C<sub>6</sub>H<sub>6</sub>)Cl<sub>2</sub>] (5)

0.100 g of [CpFe(CO)(COCH<sub>3</sub>)(<sup>1</sup>-dppe)] (2) (0.169 mmol, 1 eq.) and 0.042 g of [(η<sup>6</sup>-benzene)RuCl<sub>2</sub>]<sub>2</sub> (0.084 mmol, 0.5 eq.) were added to a 100 mL Schlenk flask. The reactants were dissolved in approximately 20 mL of dichloromethane; the solution was left to stir at room temperature for 2 hours. The solvent was evaporated under reduced pressure, and then column chromatography was performed in which a 1:1 *n*-hexane/ethyl acetate eluent was utilized. The purification afforded a light brown/red powder (0.0477 g, 0.056 mmol, 34%) with a decomposition temperature of 130 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>, 298 K): δ 7.73- 7.14 (20H, m, C-H dppe), 5.32 (6H, s, η<sup>6</sup>-C<sub>6</sub>H<sub>6</sub>), 4.39 (5H, s, η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>), 2.70 (1H, m, PCH<sup>D</sup> bridge), 2.46 (1H, m, PCH<sup>C</sup> bridge), 2.40 (1H, m, PCH<sup>B</sup> bridge), 2.35 (3H, s, COCH<sub>3</sub>), 2.15 (1H, m, PCH<sup>A</sup> bridge) ppm;

$^{31}\text{P}\{^1\text{H}\}$  NMR ( $\text{CDCl}_3$ , 298 K):  $\delta$  74.3 (d,  $^3\text{J}(\text{P},\text{P})= 38.5$  Hz, Fe- $\text{PPh}_2$ ), 25.1 (d,  $^3\text{J}(\text{P},\text{P})= 38.5$  Hz, Ru- $\text{PPh}_2$ ) ppm;  $^{13}\text{C}\{^1\text{H}\}$  NMR ( $\text{CDCl}_3$ , 298 K):  $\delta$  219.4 (d,  $^2\text{J}(\text{C},\text{P})= 30$  Hz, Fe-CO), 135.6-127.0 (m, dppe), 87.6 (d,  $^2\text{J}(\text{C},\text{P})= 3.5$  Hz,  $\eta^6\text{-C}_6\text{H}_6$ ) 83.7 (s,  $\eta^5\text{-C}_5\text{H}_5$ ), 50.7 (d,  $^3\text{J}(\text{C},\text{P})= 5$  Hz,  $\text{COCH}_3$ ), 24.9 (d,  $^1\text{J}(\text{C},\text{P})= 27$  Hz,  $\text{C}^{\text{B}}\text{H}_2\text{PPh}_2$  bridge), 21.1 (d,  $^1\text{J}(\text{C},\text{P})= 27$  Hz,  $\text{C}^{\text{A}}\text{H}_2\text{PPh}_2$  bridge) ppm; FTIR:  $\nu= 3072$  (vw), 2962 (vw), 2897 (vw), 2474 (vw), 2312 (vw), 1897 (ms, CO stretch), 1734 (vw), 1597 (m, CO stretch) 1500 (vw), 1481 (w), 1433 (mw), 1411 (w), 1373 (vw), 1319 (vw), 1259 (w), 1176 (w), 1095 (s), 1055 (s), 1028 (ms), 1014 (ms), 898 (w), 887 (w), 842 (w), 821 (ms), 800 (ms), 746 (ms), 725 (ms), 717 (s), 692 (s), 621 (w), 611 (w), 584 (w), 574 (w), 561 (w), 414 (w)  $\text{cm}^{-1}$  (Full assignment of all stretching vibrations can be found in SI, secured by DFT); ESI-MS:  $m/z$  Calcd. For  $[\text{M}]^+$  840.0117, found 840.0085. Calcd. For  $[\text{M}+\text{Na}]^+$  863.0015, found 862.9983.

### Synthesis of $[[\text{CpFe}(\text{CO})(\text{I})]_2(\text{dppb})] \cdot 0.5$ hexane (6)

0.100 g of  $[\text{CpFe}(\text{CO})_2\text{I}]$  (0.329 mmol) and 0.121 g of 1,4-Bis(diphenylphosphino)butane (dppb) (0.284 mmol) were added to a 100 mL round bottom Schlenk flask. The reactants were dissolved in 30 mL of dry toluene. The reaction was left to stir under a reflux at 85 °C for 7 hours. The solution was then cooled to room temperature and filtered under gravity. The remaining mixture was dried *in vacuo* and was purified using column chromatography with an eluent consisting of 3:2 dichloromethane/*n*-hexane. The first band was a light brown colour, which was discarded. The second band was light green, which was contaminated with starting dppb. The desired product,  $[[\text{CpFe}(\text{CO})(\text{I})]_2(\text{dppb})]$ , was concentrated in the third dark green band. The latter was dried *in vacuo*, which afforded a light green powder with a yield of 0.101 g (0.103 mmol, 37%) and a decomposition temperature of 81-85 °C.  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ , 298 K):  $\delta$  7.71- 7.35 (20H, m, C-H dppb), 4.32 (10H, s,  $\eta^5\text{-C}_5\text{H}_5$ ), 2.37 (4H, m,  $\text{PCH}^{\text{A}}$  bridge), 1.25 (5H, br s,  $\text{CH}_2$  hexane), 0.95 (4H, m,  $\text{PCH}^{\text{B}}$  bridge), 0.87 (1H, m,  $\text{CH}_3$  hexane) ppm (assigned by HSQC/HMBC experiments);  $^{31}\text{P}\{^1\text{H}\}$  NMR ( $\text{CDCl}_3$ , 298 K):  $\delta$  61.7 (s, Fe- $\text{PPh}_2$ , meso stereoisomer and/or D,L form), 61.6 (s, Fe- $\text{PPh}_2$ , meso stereoisomer and/or D,L form) ppm;  $^{13}\text{C}\{^1\text{H}\}$  NMR ( $\text{CDCl}_3$ , 298 K):  $\delta$  220.7 (s, Fe-CO), 133.1-131.7(m, C-H dppb), 130.4 (s, C quaternary dppb), 128.6- 128.2 (m, C-H dppb), 82.4 (s,  $\eta^5\text{-C}_5\text{H}_5$ ), 33.3 (d,  $^1\text{J}(\text{C},\text{P})= 17.5$  Hz,  $\text{C}^{\text{A}}\text{H}_2\text{PPh}_2$  bridge), 32.9 (d,  $^1\text{J}(\text{C},\text{P})= 17.5$  Hz,  $\text{C}^{\text{B}}\text{H}_2\text{PPh}_2$  bridge), 31.6 (s,  $\text{CH}_2(3)$  hexane), 25.2 (s,  $\text{C}^{\text{A}}\text{H}_2$  bridge), 25.1 (s,  $\text{C}^{\text{B}}\text{H}_2$  bridge), 22.7 (s,  $\text{CH}_2(2)$  hexane), 14.1 (s,  $\text{CH}_3$  hexane) ppm; FTIR ( $\text{cm}^{-1}$ ):  $\nu= 2953$  (w), 2924 (w), 2854

(w), 1932 (ms, **CO** stretch), 1587 (vw), 1571 (vw), 1479 (w), 1433 (w), 1359 (vw), 1307 (vw), 1259 (vw), 1180 (w), 1157 (w), 1093 (w), 1070 (w), 1026 (w), 999 (w), 839 (w), 821 (w), 740 (m), 719 (w), 694 (s), 617 (vw), 596 (w), 569 (w), 553 (w), 536 (w), 447 (vw), 418 (vw), 412 (vw)  $\text{cm}^{-1}$  (Full assignment of all stretching vibrations can be found in SI, secured by DFT); UV-Vis (nm)/ dichloromethane  $\lambda_{\text{max}}$ :  $\lambda$  615 (minor), 436 (major); ESI-MS:  $m/z$  Calcd. For  $[\text{M}]^+$  977.9135, found 977.9068. Calcd. For  $[\text{M}+\text{Na}]^+$  1000.9033, found 1000.8967.

### Synthesis of $[[\text{CpFe}(\text{CO})(\text{CH}_3)_2(\text{dppb})] \cdot 0.5 \text{ hexane}$ (7)

0.243 g of  $[[\text{CpFe}(\text{CO})(\text{I})_2(\text{dppb})]$  (0.248 mmol) were added to a 100 mL round bottom flask, containing 50 mL of degassed toluene. The dark green solution was cooled to approximately  $-80^\circ\text{C}$ , and MeLi (0.341 mL, 0.546 mmol, 2.2 eq) was carefully added in a dropwise fashion, turning the solution brown. The mixture was removed from the cooling bath and allowed to warm to ambient temperature, and then was left to stir for 2 hours, which darkened the colour of the brown solution. The solvent was evaporated under reduced pressure. The air-stable dark brown solid was purified *via* column chromatography using  $\text{CH}_2\text{Cl}_2$  as the eluent, and then washed with hexane. The latter yielded 0.087 g of a vermilion-coloured powder (0.115 mmol, 46.5%) with a decomposition temperature of  $134\text{-}146^\circ\text{C}$ .  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ , 298 K):  $\delta$  7.62- 7.29 (20H, m, C-H dppb), 4.12 (10H, s,  $\eta^5\text{-C}_5\text{H}_5$ ), 1.94 (4H, m, bridge), 1.25 (4H, s,  $\text{CH}_2$  hexane), 1.16 (2H, m, bridge), 0.92 (2H, m, bridge), 0.88 (1H, m,  $\text{CH}_3$  hexane), -0.29 (6H, dd,  $^3J(\text{H,P})= 8.0 \text{ Hz}$  & 6.40 Hz,  $\text{CH}_3$ ) ppm;  $^{31}\text{P}$   $\{^1\text{H}\}$  NMR ( $\text{CDCl}_3$ , 298 K):  $\delta$  75.5 (s, Fe-PPh<sub>2</sub>) ppm; FTIR ( $\text{cm}^{-1}$ ):  $\nu=$  3059 (vw), 2924 (w), 2870 (w), 1892 (s, **CO** stretch), 1718 (vw), 1684 (vw), 1653 (vw), 1647 (vw), 1586 (vw), 1571 (vw), 1479 (w), 1458 (w), 1432 (m), 1360 (vw), 1329 (vw), 1306 (w), 1291 (vw), 1260 (w), 1181 (w), 1170 (w), 1157 (w), 1093 (m), 1070 (m), 1058 (m), 1027 (m), 1011 (m), 999 (m), 891 (vw), 840 (m), 818 (m), 801 (m), 743 (m), 738 (m), 723 (m), 693 (vs), 618 (vw), 603 (w), 593 (w), 576 (w), 545 (vw), 535 (vw), 462 (vw)  $\text{cm}^{-1}$ . ESI-MS:  $m/z$  Calcd. For  $[\text{M}]^+$  754.1515, found 754.1524 (17 %). Calcd. For  $[\text{M-CpFe}(\text{CO})(\text{CH}_3)]^+$  590.1591, found 590.1558 (100 %).

### Acknowledgement

We thank the Maastricht Science Programme (MSP) and Maastricht University (Faculty of Science and Engineering) for financial support of this research. We also thank Maarten Honing, Peilang Han and Darya Hadavi (M4I, FHML (Maastricht MultiModal Molecular Imaging; Faculty Health, Medicine & Life

Sciences, Maastricht University) for measuring the HRMS of all the compounds. Dr. Christian Bahn (MSP) is also thanked for useful discussions on the stereochemistry of the complexes.

### Associated Content

The supporting information is available free of charge. Supporting Information is available containing the synthesis and characterisation of known complexes, Multinuclear NMR, IR (experimental and calculated) and ESI-MS spectra of the compounds reported here and Cartesian coordinates for the complexes studied by DFT methods.

### References

1. (a) WHO. Cancer. Geneva: World Health Organization, September 2018. <https://www.who.int/news-room/fact-sheets/detail/cancer> (accessed Sep 18, 2019). (b) V. Tamizhazhagan, K. Pugazhendy, V. Sakthidasan, C. Jayanthi, S. Rajesh, P. Manikanadan, Social and economic burden of cancer on 2020-minireview, *Cancer Research Journal*, 6 (2018) 10-15. (c) R.L. Siegel, K.D. Miller, A. Jemal, *Cancer statistics, 2020*, CA: A Cancer Journal for Clinicians, 70 (2020) 7-30.
2. V.T. DeVita, E. Chu, A history of cancer chemotherapy, *Cancer research*, 68 (2008) 8643-8653.
3. A. Caley, R. Jones, *The principles of cancer treatment by chemotherapy*, Surgery (Oxford), 30 (2012) 186-190.
4. See as examples: (a) S. Rafique, M. Idrees, A. Nasim, H. Akbar, A. Athar, Transition metal complexes as potential therapeutic agents, *Biotechnology and Molecular Biology Reviews*, 5 (2010) 38-45. (b) P.V. Simpson, N.M. Desai, I. Casari, M. Massi, M. Falasca, Metal-based antitumor compounds: beyond cisplatin, *Future Medicinal Chemistry*, 11 (2019) 119-135. (c) M. Zaki, S. Hairat, E.S. Aazam, Scope of organometallic compounds based on transition metal-arene systems as anticancer agents: starting from the classical paradigm to targeting multiple strategies, *RSC Advances*, 9 (2019) 3239-3278. (d) U. Ndagi, N. Mhlongo, M.E. Soliman, Metal complexes in cancer therapy - an update from drug design perspective, *Drug Design Development and Therapy*, 11 (2017) 599-616. (e) W. Liu, R. Gust, Update on metal N-heterocyclic carbene complexes as potential anti-tumor metallodrugs, *Coordination Chemistry Reviews*, 329 (2016) 191-213.
5. See as examples: (a) B. Rosenberg, *Cisplatin: its history and possible mechanisms of action*, Cisplatin, Elsevier 1980, pp. 9-20. (b) F.M. Muggia, A. Bonetti, J.D. Hoeschele, M. Rozenzweig,

- S.B. Howell, Platinum antitumor complexes: 50 years since barnett rosenberg's discovery, *Journal of Clinical Oncology*, 33 (2015) 4219-4226.
6. See as examples: (a) D. Chen, V. Milacic, M. Frezza, Q.P. Dou, Metal complexes, their cellular targets and potential for cancer therapy, *Current Pharmaceutical Design*, 15 (2009) 777-791. (b) M.S. Hassan, A.M. Morgan, M.M. Mekawy, M.A. Zeineb, Molecular mechanisms of Cisplatin-induced placental toxicity and teratogenicity in rats and the ameliorating role of N-acetyl-cysteine, *International Journal of Biochemistry and Cell-biology*, 115 (2019) 105579. (c) M. Raudenska, J. Balvan, M. Fojtu, J. Gumulec, M. Masarik, Unexpected therapeutic effects of cisplatin, *Metallomics*, 11 (2019) 1182-1199. (d) W. Zeng, Z. Du, Q. Luo, Y. Zhao, Y. Wang, K. Wu, F. Jia, Y. Zhang, F. Wang, Proteomic Strategy for Identification of Proteins Responding to Cisplatin-Damaged DNA, *Analytical Chemistry*, 91 (2019) 6035-6042. (e) D. Sheikh-Hamad, Cisplatin-induced cytotoxicity: is the nucleus relevant? *American Journal of Physiology Renal Physiology*, 295 (2008) F42-F43. (f) D.B. Zamble, S.J. Lippard, Cisplatin and DNA repair in cancer chemotherapy, *Trends in Biochemical Sciences*, 20 (1995) 435-439. (g) J. Kozelka, J.C. Chottard, How does cisplatin alter DNA structure? A molecular mechanics study on double-stranded oligonucleotides, *Biophysical Chemistry*, 35 (1990) 165-1.
  7. M.S. Davies, S.J. Berners-Price, T.W. Hambley, Slowing of cisplatin aquation in the presence of DNA but not in the presence of phosphate: improved understanding of sequence selectivity and the roles of mono-aquated and diaquated species in the binding of cisplatin to DNA, *Inorganic chemistry*, 39 (2000) 5603-5613.
  8. R.A. Alderden, M.D. Hall, T.W. Hambley, The discovery and development of cisplatin, *Journal of Chemical Education*, 83 (2006) 728.
  9. E. Wong, C.M. Giandomenico, Current status of platinum-based antitumor drugs, *Chemical Reviews*, 99 (1999) 2451-2466.
  10. B. Lippert, *Cisplatin: chemistry and biochemistry of a leading anticancer drug*, John Wiley & Sons 1999.
  11. S. Dasari, P.B. Tchounwou, Cisplatin in cancer therapy: molecular mechanisms of action, *European Journal of Pharmacology*, 740 (2014) 364-378.
  12. T.W. Hambley, The influence of structure on the activity and toxicity of Pt anti-cancer drugs, *Coordination Chemistry Reviews*, 166 (1997) 181-223.
  13. N.J. Wheate, S. Walker, G.E. Craig, R. Oun, The status of platinum anticancer drugs in the clinic and in clinical trials, *Dalton Transactions*, 39 (2010) 8113-8127.
  14. R.S. Go, A.A. Adjei, Review of the comparative pharmacology and clinical activity of cisplatin and carboplatin, *Journal of Clinical Oncology*, 17 (1999) 409-409.

15. A. Ahmad, F. Sarkar, Breast cancer metastasis and drug resistance, *Progress and Prospects*, (2013), Springer.
16. See: (a) I. Ott, R. Gust, Non platinum metal complexes as anti-cancer drugs, *Archiv der Pharmazie: An International Journal Pharmaceutical and Medicinal Chemistry*, 340 (2007) 117-126. And further examples: (b) E.Y. Tshuva, M. Miller, Coordination complexes of titanium(IV) for anticancer therapy, *Metal Ions in Life Sciences*, 18 (2018) 219-250. (c) H. Skoupilova, R. Hrstka, M. Bartosik, Titanocenes as Anticancer Agents: Recent Insights, *Med. Chem. (Sharjah, United Arab Emirates)*, 13 (2017) 334-344. (d) P. Beckhove, O. Oberschmidt, A.R. Hanauske, C. Pampillon, V. Schirmacher, N.J. Sweeney, K. Strohfeldt, M. Tacke, Antitumor activity of Titanocene Y against freshly explanted human breast tumor cells and in xenografted MCF-7 tumors in mice, *Anticancer Drugs*, 18 (2007) 311-315. (e) O. Oberschmidt, A.-R. Hanauske, F.-J.K. Rehmann, K. Strohfeldt, N. Sweeney, M. Tacke, Activity of [1,2-di(cyclopentadienyl)-1,2-di(p-N,N-dimethylaminophenyl)-ethanediyl] titanium dichloride against tumor colony-forming units, *Anticancer Drugs*, 16 (2005) 1071-1073.
17. See as selected examples: (a) G. Marcon, S. Carotti, M. Coronello, L. Messori, E. Mini, P. Orioli, T. Mazzei, M.A. Cinellu, G. Minghetti, Gold (III) complexes with bipyridyl ligands: solution chemistry, cytotoxicity, and DNA binding properties, *Journal of Medicinal Chemistry*, 45 (2002) 1672-1677. (b) J.H. Sze, P.V. Raninga, K. Nakamura, M. Casey, K.K. Khanna, S.J. Berners-Price, G. Di Trapani, K.F. Tonissen, Anticancer activity of a Gold(I) phosphine thioredoxin reductase inhibitor in multiple myeloma, *Redox Biology*, 28 (2020) 101310. (c) J.H. Kim, E. Reeder, S. Parkin, S.G. Awuah, Gold(I/III)-Phosphine Complexes as Potent Antiproliferative Agents, *Scientific Reports*, 9 (2019) 1-18. (d) N. Estrada-Ortiz, E. Lopez-Gonzales, B. Woods, S. Stuerup, I.A.M. de Graaf, G.M.M. Groothuis, A. Casini, Ex vivo toxicological evaluation of experimental anticancer gold(I) complexes with lansoprazole-type ligands, *Toxicological Research (Cambridge, U. K.)*, 8 (2019) 885-895. (e) F. Guarra, T. Marzo, M. Ferraroni, F. Papi, C. Bazzicalupi, P. Gratteri, G. Pescitelli, L. Messori, T. Biver, C. Gabbiani, Interaction of a gold(I) dicarbene anticancer drug with human telomeric DNA G-quadruplex: solution and computationally aided X-ray diffraction analysis, *Dalton Transactions*, 47 (2018) 16132-16138. (f) C.I. Yeo, K.K. Ooi, E.R.T. Tiekink, Gold-based medicine: a paradigm shift in anti-cancer therapy, *Molecules*, 23 (2018) 1410-1426. (g) I. Marmol, P. Castellnou, R. Alvarez, M.C. Gimeno, M.J. Rodriguez-Yoldi, E. Cerrada, Alkynyl Gold(I) complexes derived from 3-hydroxyflavones as multi-targeted drugs against colon cancer, *European Journal of Medicinal Chemistry*, 183 (2019) 111661.


18. A.R. Simović, R. Masnikosa, I. Bratsos, E. Alessio, Chemistry and reactivity of ruthenium (II) complexes: DNA/protein binding mode and anticancer activity are related to the complex structure, *Coordination Chemistry Reviews*, 398 (2019) 113011.
19. P. Schluga, C.G. Hartinger, A. Egger, E. Reisner, M. Galanski, M.A. Jakupec, B.K. Keppler, Redox behavior of tumor-inhibiting ruthenium (III) complexes and effects of physiological reductants on their binding to GMP, *Dalton Transactions*, (2006) 1796-1802.
20. E.S. Antonarakis, A. Emadi, Ruthenium-based chemotherapeutics: are they ready for prime time? *Cancer Chemotherapy and Pharmacology*, 66 (2010) 1-9.
21. E. Musgrove, C. Rugg, I. Taylor, D. Hedley, Transferrin receptor expression during exponential and plateau phase growth of human tumour cells in culture, *Journal of Cellular Physiology*, 118 (1984) 6-12.
22. H. Yamada, T. Koike, J.K. Hurst, Water Exchange Rates in the Diruthenium  $\mu$ -Oxo Ion cis, cis- $[(bpy)_2Ru(OH_2)]_2O^{4+}$ , *Journal of the American Chemical Society*, 123 (2001) 12775-12780.
23. G. Sava, S. Pacor, A. Bergamo, M. Cocchietto, G. Mestroni, E. Alessio, Effects of ruthenium complexes on experimental tumors: irrelevance of cytotoxicity for metastasis inhibition, *Chemico-Biological Interactions*, 95 (1995) 109-126.
24. J.M. Rademaker-Lakhai, D. van den Bongard, D. Pluim, J.H. Beijnen, J.H. Schellens, A phase I and pharmacological study with imidazolium-trans-DMSO-imidazole-tetrachlororuthenate, a novel ruthenium anticancer agent, *Clinical Cancer Research*, 10 (2004) 3717-3727.
25. A. Bergamo, R. Gagliardi, V. Scarcia, A. Furlani, E. Alessio, G. Mestroni, G. Sava, In vitro cell cycle arrest, in vivo action on solid metastasizing tumors, and host toxicity of the antimetastatic drug NAMI-A and cisplatin, *Journal of Pharmacology and Experimental Therapeutics*, 289 (1999) 559-564.
26. A. Bergamo, C. Gaiddon, J. Schellens, J. Beijnen, G. Sava, Approaching tumour therapy beyond platinum drugs: status of the art and perspectives of ruthenium drug candidates, *Journal of Inorganic Biochemistry*, 106 (2012) 90-99.
27. A.F. Peacock, P.J. Sadler, Medicinal organometallic chemistry: designing metal arene complexes as anticancer agents, *Chemistry—An Asian Journal*, 3 (2008) 1890-1899.
28. B.S. Murray, M.V. Babak, C.G. Hartinger, P.J. Dyson, The development of RAPTA compounds for the treatment of tumors, *Coordination Chemistry Reviews*, 306 (2016) 86-114.
29. S. Inoue, S. Kawanishi, Hydroxyl radical production and human DNA damage induced by ferric nitrilotriacetate and hydrogen peroxide, *Cancer Research*, 47 (1987) 6522-6527.
30. S.V. Torti, F.M. Torti, Iron and cancer: more ore to be mined, *Nature Reviews Cancer*, 13 (2013) 342.

31. E. Hillard, A. Vessières, L. Thouin, G. Jaouen, C. Amatore, Ferrocene-Mediated Proton-Coupled Electron Transfer in a Series of Ferrocifen-Type Breast-Cancer Drug Candidates, *Angewandte Chemie International Edition*, 45 (2006) 285-290.
32. E. Wong, C.M. Giandomenico, Current status of platinum-based antitumor drugs, *Chemical Reviews*, 99 (1999) 2451-2466.
33. D. Serra, K.A. Abboud, C.R. Hilliard, L. McElwee-White, Electronic Interactions in Iron-and Ruthenium-Containing Heterobimetallic Complexes: Structural and Spectroscopic Investigations, *Organometallics*, 26 (2007) 3085-3093.
34. M. Wenzel, E. Bigaeva, P. Richard, P. Le Gendre, M. Picquet, A. Casini, E. Bodio, New heteronuclear gold (I)–platinum (II) complexes with cytotoxic properties: Are two metals better than one? *Journal of Inorganic Biochemistry*, 141 (2014) 10-16.
35. J. Fernández-Gallardo, B.T. Elie, M. Sanaú, M. Contel, Versatile synthesis of cationic N-heterocyclic carbene–gold (I) complexes containing a second ancillary ligand. Design of heterobimetallic ruthenium–gold anticancer agents, *Chemical Communications*, 52 (2016) 3155-3158.
36. O. Renier, C. Deacon-Price, J. Peters, K. Nurekeyeva, C. Russon, S. Dyson, S. Ngubane, J. Baumgartner, P. Dyson, T. Riedel, Synthesis and In Vitro (Anticancer) Evaluation of  $\eta^6$ -Arene Ruthenium Complexes Bearing Stannyl Ligands, *Inorganics*, 5 (2017) 44.
37. M. Gras, B. Therrien, G. Süß-Fink, A. Casini, F. Edafe, P.J. Dyson, Anticancer activity of new organo-ruthenium, rhodium and iridium complexes containing the 2-(pyridine-2-yl)thiazole N,N-chelating ligand, *Journal of Organometallic Chemistry*, 695 (2010) 1119-1125.
38. C. Deacon-Price, D. Romano, T. Riedel, P.J. Dyson, B. Blom, Synthesis, characterisation and cytotoxicity studies of ruthenium arene complexes bearing trichlorogermyl ligands, *Inorganica Chimica Acta*, 484 (2019) 513-520.
39. (a) B. Herry, L.K. Batchelor, B. Roufosse, D. Romano, J. Baumgartner, M. Borzova, T. Reifensahl, T. Collins, A. Benamrane, J. Weggelaar, M.C. Correia, P.J. Dyson, B. Blom, Heterobimetallic Ru( $\mu$ -dppm)Fe and homobimetallic Ru( $\mu$ -dppm)Ru complexes as potential anti-cancer agents, *Journal of Organometallic Chemistry*, (2019) 120934. For  $\mu$ -dppm (or similar) bridged Fe,Ru complexes see: (b) N. Nawar, A.K. Smith, Syntheses of ruthenium-containing heterometallic complexes by use of tridentate phosphine ligands, *Journal of Organometallic Chemistry*, 493 (1995) 239-242. (c) A.W. Coleman, D.F. Jones, P.H. Dixneuf, C. Brisson, J.J. Bonnet, G. Lavigne, Dehalogenation of binuclear arene-ruthenium complexes: a new route to homonuclear triruthenium and heteronuclear ruthenium-iron cluster complexes containing chelating phosphorus ligands. Crystal structure of Ru<sub>3</sub>(CO)<sub>10</sub>(Ph<sub>2</sub>PCH<sub>2</sub>PPh<sub>2</sub>), *Inorganic Chemistry*, 23 (1984) 952-956. (d) P.A.

- Dolby, M.M. Harding, N. Newar, A.K. Smith, Syntheses of iron-containing heterometallic complexes using tridentate phosphine ligands; the crystal structure of  $[(OC)_3Fe\{(Ph_2P)_2CHCH_2PPh_2\}Ru_3(CO)_9\{Ph_2PC(CH_2)PPh_2\}]$ , *Journal of the Chemical Society, Dalton Transactions*, (1992) 2939-2942. (e) M.-G. Mendoza-Ferri, C.G. Hartinger, R.E. Eichinger, N. Stolyarova, K. Severin, M.A. Jakupec, A.A. Nazarov, B. K. Keppler, Influence of the spacer length on the in vitro anticancer activity of dinuclear ruthenium-arene compounds, *Organometallics*, 27 (2008), 2405-2407.
40. (a) R.A. Zelonka, M.C. Baird, *Canadian Journal of Chemistry*, 50(18), (1972) 3063-3072. (b) M.A. Bennett, A.K. Smith, *Journal of the Chemical Society Dalton Transactions*, (1974) 233-241.
41. J.K. Pagano, B.J. Ackley, R. Waterman, Evidence for Iron-Catalyzed  $\alpha$ -Phosphinidene Elimination with Phenylphosphine, *Chemistry—A European Journal*, 24 (2018) 2554-2557.
42. For spectra shifts see: A. Munyaneza, O.G. Adeyemi, N.J. Coville, Solventless migratory-insertion reactions of substituted cyclopentadienyl iron complexes induced by electron donor ligands, *Bulletin of the Chemical Society of Ethiopia*, 23(3) (2009) 399-407.
43. L.-S. Luh, L.-K. Liu, Synthesis and X-ray Structure of  $[(\eta^5-C_5H_5)Fe(CO)C(O)Me](\mu-\eta^1: \eta^1-dppe)[(4-exo-MeC_5H_5)Fe(CO)_2]$ , *Organometallics* 14 (1995) 1514-1517.
44. H.B. Kwon, Reaction of Methyl- $Fe(\eta^5-C_5H_5)(CO)_2$  with 1,2-Bis(diphenylphosphino)ethane, *Journal of the Korean Chemical Society*, 45(1) (2001) 90-91.
45. Similar migratory insertion reactions promoted by diphosphines generating bimetallic diiron systems have been reported: M. Kumar, A. Metta-Magana, H.K. Sharma, K.H. Pannell, Phosphine induced migratory CO insertion into the Fe-CH<sub>2</sub> bond of the organometallic polymer- $[(\eta^5-C_5H_4)Fe(CO)_2CH_2SiMe_2]_n$ -and characterization of model iron complexes, *Dalton Transactions*, 39 (2010) 7125-7131.
46. U. Knof, A. von Zelewsky, Predetermined Chirality at Metal Centres, *Angewandte Chemie International Edition*, 38(3) (1999) 302-322.
47. M. Timm, L. Saaby, L. Moesby, E.W. Hansen, Considerations regarding use of solvents in in vitro cell based assays, *Cytotechnology*, 65(5) (2013) 887-894.
48. A plethora of  $\mu$ -phosphane bridged diiron complexes exist. See as selected examples: (a) Q.-L. Li, S. Lu, R.-F. Zhang, D. Zhao, C.-L. Ma, Substitution reactions of diiron diselenolato complex with bisphosphine ligands, *Polyhedron*, 160 (2019) 255-260. (b) A. Rahaman, S. Ghosh, S. Basak-Modi, A.F. Abdel-Magied, S.E. Kabir, M. Haukka, M.G. Richmond, G.C. Lisensky, E. Nordlander, G. Hogarth, Chalcogenide-capped triiron clusters  $[Fe_3(CO)_9(\mu_3-E)_2]$ ,  $[Fe_3(CO)_7(\mu_3-CO)(\mu_3-E)(\mu-dppm)]$  and  $[Fe_3(CO)_7(\mu_3-E)_2(\mu-dppm)]$  (E = S, Se) as proton-reduction catalysts, *Journal of Organometallic Chemistry*, 880 (2019) 213-222. (c) S. Ghosh, B.E. Sanchez, I. Richards, M.N.

- Haque, K.B. Holt, M.G. Richmond, G. Hogarth, Biomimetics of the [FeFe]-hydrogenase enzyme: Identification of kinetically favoured apical-basal  $[\text{Fe}_2(\text{CO})_4(\mu\text{-H})\{\kappa^2\text{-Ph}_2\text{PC}(\text{Me}_2)\text{PPh}_2\}(\mu\text{-pdt})]^+$  as a proton-reduction catalyst, *Journal of Organometallic Chemistry*, 812 (2016) 247-258. (d) M. Kaiser, G. Knoer, Synthesis, Characterization, and Reactivity of Functionalized Trinuclear Iron-Sulfur Clusters - A New Class of Bioinspired Hydrogenase Models, *European Journal of Inorganic Chemistry*, 2015 (2015) 4199-4206. (e) F.I. Adam, G. Hogarth, S.E. Kabir, I. Richards, Models of the iron-only hydrogenase: Synthesis and protonation of bridge and chelate complexes  $[\text{Fe}_2(\text{CO})_4\{\text{Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2\}(\mu\text{-pdt})]$  ( $n = 2-4$ ) - evidence for a terminal hydride intermediate, *Comptes Rendus Chimie*, 11 (2008) 890-905. (f) S. Ezzaher, J.-F. Capon, F. Gloaguen, F.Y. Petillon, P. Schollhammer, J. Talarmin, Electron-Transfer-Catalyzed Rearrangement of Unsymmetrically Substituted Diiron Dithiolate Complexes Related to the Active Site of the [FeFe]-Hydrogenases, *Inorganic Chemistry*, 46 (2007) 9863-9872. (g) Z.-F. Meng, H.-L. Gao, J.-X. Wang, J.-Y. Shang, C.-G. Li, Phenyl-functionalized diiron propanediselenolato complexes containing the chelated or bridged 1,3-bis(diphenylphosphine)propane ligand, *Journal of Coordination Chemistry*, 71 (2018) 2953-2964. (h) G.-R. Xu, L. Liu, H.-L. Gao, J.-Y. Shang, C.-G. Li, Phenyl-functionalized diiron propane diselenolato complexes containing intramolecular bridging diphosphine ligands, *Journal of Coordination Chemistry*, 70 (2017) 2684-2694.
49. T. Mosmann, Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays, *Journal of Immunological Methods*, 65 (1983) 55-63.
50. Gaussian 09, Revision A.02, M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, G. A. Petersson, H. Nakatsuji, X. Li, M. Caricato, A. Marenich, J. Bloino, B. G. Janesko, R. Gomperts, B. Mennucci, H. P. Hratchian, J. V. Ortiz, A. F. Izmaylov, J. L. Sonnenberg, D. Williams-Young, F. Ding, F. Lipparini, F. Egidi, J. Goings, B. Peng, A. Petrone, T. Henderson, D. Ranasinghe, V. G. Zakrzewski, J. Gao, N. Rega, G. Zheng, W. Liang, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, K. Throssell, J. A. Montgomery, Jr., J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, T. Keith, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, J. M. Millam, M. Klene, C. Adamo, R. Cammi, J. W. Ochterski, R. L. Martin, K. Morokuma, O. Farkas, J. B. Foresman, and D. J. Fox, Gaussian, Inc., Wallingford CT, 2016.

Journal Pre-proofs



Journal Pre-proofs