


This article was downloaded by: [University of Connecticut]

On: 28 October 2014, At: 18:59

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954

Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

Synthesis of Highly Functionalized Phosphorus Ylids Containing Heterocyclic Rings and the Related Iminophosphoranes

Hojatollah Khabazzadeh^a, Kazem Saidi^a, Hassan Sheibani^a & Mohammad Reza Islami^a

^a Department of Chemistry, Shahid Bahonar University of Kerman, Kerman, Iran

Published online: 16 Jul 2007.

To cite this article: Hojatollah Khabazzadeh, Kazem Saidi, Hassan Sheibani & Mohammad Reza Islami (2007) Synthesis of Highly Functionalized Phosphorus Ylids Containing Heterocyclic Rings and the Related Iminophosphoranes, *Phosphorus, Sulfur, and Silicon and the Related Elements*, 182:9, 2163-2170, DOI: [10.1080/10426500701407482](https://doi.org/10.1080/10426500701407482)

To link to this article: <http://dx.doi.org/10.1080/10426500701407482>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with

primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Synthesis of Highly Functionalized Phosphorus Ylides Containing Heterocyclic Rings and the Related Iminophosphoranes

Hojatollah Khabazzadeh

Kazem Saidi

Hassan Sheibani

Mohammad Reza Islami

Department of Chemistry, Shahid Bahonar University of Kerman,
Kerman, Iran

Crystalline phosphorus ylides are obtained in excellent yields from the 1:1:1 addition reaction between triphenylphosphine, dimethyl acetylenedicarboxylate (DMAD) and NH and SH acids, such as 2-amino-4-phenyl thiazole, 2-amino-5-(3-chloro-benzyl) thiaziazole, 3-amino-2-methyl quinazolin-4-one and 3-amino-2-mercapto quinazolin-4-one. These stabilized phosphoranes undergo a smooth intramolecular reaction in boiling toluene to produce aryliminophosphoranes in excellent yields.

Keywords Dimethyl acetylenedicarboxylate; phosphorus ylide; quinazolin-4-one; thiazole

INTRODUCTION

In the realm of ylide compounds and their value for a variety of industrial, biological and chemical synthetic uses, phosphorus ylides are playing increasingly pivotal roles.^{1,2} For a reaction to be synthetically applicable, should have readily available starting materials, high yields and furthermore the product must be free of contaminants so as to avoid the need for extensive purification.³ The chemistry of ylide compounds has provided chemists with exceptionally fertile ground for the design and development of new bond formation.⁴ Since the pioneering work by Staudinger and Meyre in 1919,⁵ on the preparation of iminophosphoranes, compounds of general structure $R_3P=NR$, the

Received November 18, 2006; accepted March 31, 2007.

The authors express appreciation to the Shahid Bahonar University of Kerman Faculty Research Committee for its support of this investigation.

Address correspondence to Kazem Saidi, Department of Chemistry, Shahid Bahonar University of Kerman, 76169, Kerman, Iran. E-mail: saidik@mail.uk.ac.ir

synthetic application of phosphorus ylides to the preparation of various organic products has increased enormously during the last three decades.⁶ Several methods for the preparation of iminophosphoranes have been developed.^{7–9}

RESULTS AND DISCUSSION

As part of our continuing interest in this area, we have recently reported synthesis of highly functionalized phosphorus ylides.¹⁰

Now, in this investigation we wish to report synthesis of stable phosphorus ylides from thiazoles and quinazolinones **1a–d** (Scheme 1).

The ylides **2a, b** were converted to iminophosphoranes **3a, b**, when the reaction mixture was refluxed in toluene (Scheme 2).


It is reasonable to assume that phosphorus ylide **2** results from the initial addition of triphenylphosphine to DMAD and subsequent protonation of the (1:1) adduct. Then the positively charged ion is attacked by conjugate base of **1** to form stable phosphorus ylide **2** (Scheme 3).




The proposed structures of compounds **2a–d** were supported by their elemental analyses and spectroscopic properties, such as ¹H NMR, ¹³C NMR, IR and mass spectral data.

There are two absorptions at the IR spectra in the region of carbonyl groups. One of the carbonyl groups reveals a strong band at 1640 cm⁻¹ which is conjugated with ylide moiety and the nonconjugated carbonyl group appears at about 1750 cm⁻¹.

Since the rotation about the partial double bond in E and Z isomers is slow on the NMR time scale at ambient temperature (Scheme 4); therefore, the presence of a mixture of two geometrical isomers E and Z for compounds **2a–d** was proven by their ¹³C NMR & ¹H NMR absorption. For example ¹H NMR and ¹³C NMR spectra of these compounds show four singlets for methyl hydrogens and four absorptions for carbonyl groups respectively, which are indicative of the presence of two isomers. Presence of the ³¹P-nucleus helps in assignment of compounds **2a–d** by long range spin-spin coupling constants of ³¹P with ¹H and ¹³C nuclei.

The ¹H NMR spectrum of **2a** shows four different sharp lines at $\delta = 3.15, 3.61, 3.71, \text{ and } 3.80$ ppm, which are due to the methoxy protons, along with signals for methine protons at $\delta = 4.45$ and 4.70 ppm which appear as two multiplets respectively for minor and major isomers. Two fairly broad signals at $\delta = 6.25$ and 6.87 ppm are related to protons of N-H groups. The mass spectra of compounds **2a–d** did not reveal molecular ion peak. Peaks at m/z 262 and 183 which are correspond to Ph_3P^+ and $\text{C}_{12}\text{H}_8\text{P}^+$ were observed. The ¹³C NMR of compound **2a** is in good agreement with the proposed structure of this compound. The ¹H




1,2	Z	Yield(%) of 2
a		90
b		88
c		85
d		87


SCHEME 1

NMR and ^{13}C NMR spectra of compounds **2a**–**d** are similar to those of **2a** except for the signals from the aromatic rings which appear in the corresponding chemical shifts.


Compounds **2a** and **2b** undergo a smooth and clean reaction in boiling toluene to produce iminophosphoranes **3a** and **3b**. All of the attempts to convert **2c** to its related iminophosphorane were mostly unsuccessful.



SCHEME 2



SCHEME 3



SCHEME 4

This transformation was suggested to proceed by an initial N-H proton shift onto the ylide carbon of the **2a** and forms the phosphorus betaine **4** and was followed by intramolecular ring closure to produce azaphosphetane **5** which can be in equilibrium with betaine **4**. The only product formed under these conditions was iminophosphorane **3** (Scheme 5).

The ^1H NMR of **3a** displays proton of thiazole ring at 6.41 and aromatic protons at 7.25–7.89 ppm. The ^{13}C NMR of this compound reveals

White powder, m.p. = 140–143°C, yield = 1.04 g (90%); IR (KBr) (ν_{\max} , cm^{-1}): 3404 (NH), 1741 (C=O), 1641 (C=O). MS, m/z (%): 436 (M^+ - $\text{C}_6\text{H}_8\text{O}_4$, 79), 333 (40), 262 (PPh₃, 88), 183 (95), 113 (100). Anal. Calcd. for $\text{C}_{33}\text{H}_{29}\text{N}_2\text{O}_4\text{PS}$: C, 68.26; H, 5.03; N, 4.82%. Found: 68.01; H, 4.95, N, 4.51%.

Major isomer (Z) (63%), ^1H NMR: δ 3.15 and 3.71 (6H, s, 2OCH₃), 4.70 (1H, m, P=C-CH), 6.56 (1H, s, thiazole ring), 6.87 (1H, br s, NH), 7.21–7.77 (40H, m, arom).* ^{13}C NMR: δ 43.29 (d, $^1\text{J}_{\text{PC}}$ = 128.1 Hz, P=C)*, 49.03 and 52.24 (2OCH₃), 57.58 (d, $^2\text{J}_{\text{PC}}$ = 15.6 Hz, P=C-CH)*, 100.73 (CH)*, 126.13 (2CH)*, 126.73 (d, $^1\text{J}_{\text{PC}}$ = 97.9 Hz, C^{ipso})*, 128.06 (2CH)*, 128.43 (CH)*, 128.62 (d, $^3\text{J}_{\text{PC}}$ = 12.4 Hz, C^{meta})*, 131.98 (C^{para})*, 133.78 (d, $^2\text{J}_{\text{PC}}$ = 9.6 Hz, C^{ortho})*, 150.91 (C)*, 167.94 (C)*, 170.16 (d, $^2\text{J}_{\text{PC}}$ = 14.6 Hz, C=O)*, 173.99 (C=O)*.

Minor isomer (E) (37%), ^1H NMR: δ 3.61 and 3.80 (6H, s, 2OCH₃), 4.52 (1H, m, P=C-CH), 6.25 (1H, s br, NH), 6.68 (1H, s, thiazole ring). ^{13}C NMR: δ 50.18 and 52.24 (2OCH₃).

Dimethyl 2-[5-(3-chloro-benzyl)-[1, 3, 4] thiadiazol-2-yl] amino]-3-(triphenylphosphoranylidine)-butanedioate (2b)

White powder, m.p. = 120–124°C, yield = 1.05 g (88%); IR (KBr) (ν_{\max} , cm^{-1}): 3379 (NH), 1765 (C=O), 1620 (C=O). MS, m/z (%): 517 (M^+ - $\text{C}_6\text{H}_8\text{O}_4$, 32), 288 (100), 262 (PPh₃, 25), 183 (47).

Major isomer (E) (77%), ^1H NMR: δ 3.13 and 3.67 (6H, s, 2OCH₃), 4.21 (2H, m, CH₂)*, 4.48 (1H, m, P=C-CH)*, 6.88 (1H, br s, NH)*, 7.21–7.68 (38H, m, arom)*. ^{13}C NMR: δ 38.65 (CH₂), 42.89 (d, $^1\text{J}_{\text{PC}}$ = 126.7 Hz, P=C)*, 49.12 and 52.32 (2OCH₃), 57.84 (d, $^2\text{J}_{\text{PC}}$ = 16.1 Hz, P=C-CH)*, 126.26 (d, $^1\text{J}_{\text{PC}}$ = 92.6 Hz, C^{ipso})*, 127.27 (CH), 127.77 (CH), 128.67 (d, $^3\text{J}_{\text{PC}}$ = 12.3 Hz, C^{meta})*, 129.11 (CH), 129.81 (CH), 132.23 (C^{para})*, 133.72 (C)*, 133.75 (d, $^2\text{J}_{\text{PC}}$ = 9.9 Hz, C^{ortho})*, 134.31 (C), 150.51 (C)*, 165.45 (C)*, 170.29 (d, $^2\text{J}_{\text{PC}}$ = 12.5 Hz, C=O)*, 173.03 (C=O)*.

Minor isomer (Z) (23%), ^1H NMR: δ 3.56 and 3.80 (6H, s, 2OCH₃). ^{13}C NMR: δ 38.34 (CH₂), 50.20 and 52.28 (2OCH₃), 128.43 (CH), 128.53 (CH), 132.02 (CH), 132.10 (CH), 132.80 (C), 133.20 (C).

Dimethyl 2-[(2-methyl-4-oxo-3(4H)-quinazoliny)] amino]-3-(triphenylphosphoranylidine)-butanedioate (2c)

White powder, m.p. = 174–177°C, yield = 0.985 g (85%); IR (KBr) (ν_{\max} , cm^{-1}): 3305 (NH), 1765 (C=O), 1691 (C=O), 1641 (C=N). MS, m/z (%): 405 (M^+ - $\text{C}_9\text{H}_8\text{N}_3\text{O}$, 100), 262 (PPh₃, 18), 183 (25). Anal. Calcd. for $\text{C}_{33}\text{H}_{30}\text{N}_3\text{O}_5\text{P}$: C, 68.39; H, 5.22; N, 7.25 %. Found: 68.07; H, 5.18, N, 7.25%.

*For two geometrical isomers.

Major isomer (*Z*) (73%), $^1\text{H NMR}$: δ 2.59 (1H, br s, NH), 3.20 (3H, s, CH_3), 3.66 and 3.76 (6H, s, 2OCH_3), 3.80 (1H, d, $^3\text{J}_{\text{PH}} = 13.42$ Hz, $\text{P}=\text{C}-\text{CH}^*$), 7.28–8.05 (38H, m, arom)*. $^{13}\text{C NMR}$: δ 22.01 (CH_3)*, 42.08 (d, $^1\text{J}_{\text{PC}} = 133.9$ Hz, $\text{P}=\text{C}$), 49.09 and 50.26 (2OCH_3), 60.62 (d, $^2\text{J}_{\text{PC}} = 19.5$ Hz, $\text{P}=\text{C}-\text{CH}$), 125.94 (d, $^1\text{J}_{\text{PC}} = 91.8$ Hz, C^{ipso}), 128.53 (d, $^3\text{J}_{\text{PC}} = 11.9$ Hz, C^{meta}), 131.79 (C^{para})*, 131.95 (CH)*, 132.02 (CH)*, 132.11 (C)*, 132.94 (C)*, 133.29 (CH)*, 133.45 (d, $^2\text{J}_{\text{PC}} = 9.6$ Hz, C^{ortho}), 134.13 (CH)*, 147.50 (C)*, 165.25 ($\text{C}=\text{O}$)*, 169.60 (d, $^2\text{J}_{\text{PC}} = 14.46$ Hz, $\text{C}=\text{O}$), 173.03 ($\text{C}=\text{O}$).

Minor isomer (*E*) (27%), $^1\text{H NMR}$: δ 2.47 (1H, br s, NH), 2.70 (3H, s, CH_3), 3.80 and 3.82 (6H, s, 2OCH_3). $^{13}\text{C NMR}$: δ 39.86 (d, $^1\text{J}_{\text{PC}} = 141.0$ Hz, $\text{P}=\text{C}$), 50.09 and 52.25 (2OCH_3), 61.62 (d, $^2\text{J}_{\text{PC}} = 18.6$ Hz, $\text{P}=\text{C}-\text{CH}$), 126.58 (d, $^1\text{J}_{\text{PC}} = 95.5$ Hz, C^{ipso}), 128.39 (d, $^3\text{J}_{\text{PC}} = 11.8$ Hz, C^{meta}), 133.38 (d, $^2\text{J}_{\text{PC}} = 9.5$ Hz, C^{ortho}), 170.69 (d, $^2\text{J}_{\text{PC}} = 16.5$ Hz, $\text{C}=\text{O}$).

Dimethyl 2-[(3-amino-4-oxo-3, 4-dihydro-2-quinazoliny)] sulfanyl]-3-(triphenylphosphora-nylidine)-butanedioate (*2d*)

White powder, m.p. = 157–160°C, yield = 1.04 g (87%); IR (KBr) (ν_{max} , cm^{-1}): 3330 and 3203 (NH_2), 1746 ($\text{C}=\text{O}$), 1682 ($\text{C}=\text{O}$), 1640 ($\text{C}=\text{O}$). MS, m/z (%): 320 ($\text{M}^+ - \text{Ph}_3\text{P}=\text{O}$, 15), 301 (20), 288 (85), 277 (46), 262 (PPh_3 , 67), 183 (100), 108 (39), 77 (44), 51 (40).

Major isomer (*Z*) (60%), $^1\text{H NMR}$: δ 3.09 and 3.77 (6H, s, 2OCH_3), 4.39 (2H, s, NH_2), 4.59 (1H, d, $^3\text{J}_{\text{PH}} = 15.6$ Hz, $\text{P}=\text{C}-\text{CH}^*$), 7.05–8.07 (38H, m, arom)*. $^{13}\text{C NMR}$: δ 40.82 (d, $^1\text{J}_{\text{PC}} = 128.1$ Hz, $\text{P}=\text{C}$), 49.08 and 52.11 (2OCH_3), 62.67 (d, $^2\text{J}_{\text{PC}} = 15.1$ Hz, $\text{P}=\text{C}-\text{CH}$), 122.08 (CH)*, 122.80 (CH)*, 124.93 (CH)*, 126.37 (d, $^1\text{J}_{\text{PC}} = 93.4$ Hz, C^{ipso}), 126.48 (CH)*, 128.48 (d, $^3\text{J}_{\text{PC}} = 11.6$ Hz, C^{meta})*, 131.90 (d, $^4\text{J}_{\text{PC}} = 2.0$ Hz, C^{para})*, 133.66 (d, $^2\text{J}_{\text{PC}} = 10.4$ Hz, C^{ortho})*, 134.02 (C)*, 134.49 (C)*, 150.25 (C), 161.88 ($\text{C}=\text{O}$), 170.10 (d, $^2\text{J}_{\text{PC}} = 14.2$ Hz, $\text{C}=\text{O}$), 174.96 (d, $^3\text{J}_{\text{PC}} = 7.0$ Hz, $\text{C}=\text{O}$).

Minor isomer (*E*) (40%), $^1\text{H NMR}$: δ 3.5 and 3.74 (6H, s, 2OCH_3), 4.57 (2H, s, NH_2). $^{13}\text{C NMR}$: δ 42.08 (d, $^1\text{J}_{\text{PC}} = 135.2$ Hz, $\text{P}=\text{C}$), 50.22 and 52.00 (2OCH_3), 62.44 (d, $^2\text{J}_{\text{PC}} = 14.1$ Hz, $\text{P}=\text{C}-\text{CH}$), 150.048 (C), 162.019 ($\text{C}=\text{O}$), 170.54 (d, $^2\text{J}_{\text{PC}} = 16.3$ Hz, $\text{C}=\text{O}$), 175.14 (d, $^3\text{J}_{\text{PC}} = 6.8$ Hz, $\text{C}=\text{O}$).

***N*-(4-Phenyl-1, 3-thiazol-2-yl) triphenyliminophosphorane (*3a*)**

The process for the preparation of **3a** is described as an example. A mixture of **2a** (0.52 g, 1.0 mmol) in toluene (10 mL) was refluxed for 8 h. The solvent was removed under reduced pressure. Dimethyl fumarate sublimated under reduced pressure and the residue recrystallised from 1:1 hexane–ethyl acetate as colorless crystals.

White crystals, m.p. = 162–166°C, yield = 0.41 g (95%), IR (KBr) (ν_{\max} , cm^{-1}): 1591, 1492, 1443, 1319, 1244. MS, m/z (%): 436 (M^+ , 100), 262 (PPh_3 , 40), 183 (82), 108 (53). Anal. Calcd. for $\text{C}_{27}\text{H}_{21}\text{N}_2\text{PS}$: C, 74.29; H, 4.85; N, 6.42 %. Found: 73.97; H, 4.58, N, 6.07%.

^1H NMR: δ 6.41–7.89 (21H, m, arom). ^{13}C NMR: δ 103.29 (C), 125.28 (CH), 125.75 (2CH), 126.77 (C), 128.17 (2CH), 128.56 (d, $^3\text{J}_{\text{PC}} = 12.3$ Hz, C^{meta}), 128.62 (d, $^1\text{J}_{\text{PC}} = 101.8$ Hz, C^{ipso}), 132.21 (d, $^4\text{J}_{\text{PC}} = 2.1$ Hz, C^{para}), 133.22 (d, $^2\text{J}_{\text{PC}} = 10.1$ Hz, C^{ortho}), 137.85 (C), 149.68 (C).

***N*-(5-(3-Chloro-benzyl)-[1, 3, 4] thiadiazol-2-yl) triphenyliminophosphorane (3b)**

White crystals, m.p. = 138–140°C, yield = 0.42 g (93%); IR (KBr) (ν_{\max} , cm^{-1}): 1594, 1581, 1481, 1426, 1320, 1230. MS, m/z (%): 519 ($\text{M}+2$, 12), 519 (M^+ , 30), 282 (100), 183(40). Anal. Calcd. for $\text{C}_{27}\text{H}_{21}\text{ClN}_3\text{PS}$: C, 66.73; H, 4.36; N, 8.65 %. Found: 66.42; H, 4.12, N, 8.46%.

^1H NMR: δ 4.28 (2H, s, CH_2), 7.20–7.81 (19H, m, arom). ^{13}C NMR: δ 37.54 (CH_2), 127.26 (d, $^1\text{J}_{\text{PC}} = 91.2$ Hz, C^{ipso}), 127.32 (CH), 127.62 (CH), 128.82 (d, $^3\text{J}_{\text{PC}} = 12.6$ Hz, C^{meta}), 129.14 (CH), 129.74 (CH), 132.65 (C^{para}), 133.12 (d, $^2\text{J}_{\text{PC}} = 10.2$ Hz, C^{ortho}), 133.37 (C), 134.21 (C), 138.89 (C), 152.14 (C).

REFERENCES

- [1] I. Yavari, M. R. Islami, H. R. Bijanzadeh, *Tetrahedron* **55**, 5547 (1999).
- [2] Y. Shen, *Acc. Chem. Res.*, **31**, 584 (1998).
- [3] Z. Hassani, M. R. Islami, H. Sheibani, M. Kalantari, and K. Saidi, *Arkivoc*, (i), 89 (2006).
- [4] V. P. Balema, J. W. Wiench, M. Pruski, and V. K. Pecharsky, *Chem. Commun.*, **7**, 724 (2002).
- [5] H. Staudinger and J. Meyer, *Helv. Chim. Acta*, **2**, 635 (1919).
- [6] B. E. Maryano and A. B. Reitz, *Chem. Rev.*, **89**, 863 (1989).
- [7] M. W. Dsing, S. Y. Yang, and J. Zhu, *Synthesis*, **1**, 75 (2004).
- [8] N. I. Gusar, *Russ. Chem. Rev.*, **60**, 146 (1991).
- [9] P. M. Feresneda, and P. Molina, *Synlett* **1**, 1 (2004).
- [10] A. Ahmadi, K. Saidi, H. Khabazzadeh, H. Sheibani, and A. Mollahosseini, *Phosphorus, Sulfur, and Silicon, and the Related Elements*, **182**, 1225 (2007).
- [11] A. Geronikaki and G. Theophilidis, *Eur. J. Med. Chem.*, **27**, 1 (1992).
- [12] S. Turner, *J. Med. Chem.*, **31**, 902 (1988).
- [13] R. S. Atkinson, *Tetrahedron*, **45**, 2875 (1989).
- [14] S. N. Pandeya, D. Sriram, D. Nath, and E. De Clercq, *Pharm. Acta Helv.*, **74**, 11 (1999).