


This article was downloaded by: [Universite Laval]

On: 10 September 2013, At: 01:01

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954

Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

### Studies on Reactions of Pyrimidine Compounds. 2<sup>1</sup>. Microwave-Assisted Synthesis of 1,2,3,4-Tetrahydro-2-Thioxopyrimidine Derivatives

Esvet Akbaş<sup>a</sup> & Furgan Aslanoğlu<sup>a</sup>

<sup>a</sup> Department of Chemistry, Faculty of Arts and Sciences, University of Yuzuncu Yil, Kampus, Van, Turkey

Published online: 20 Feb 2008.

To cite this article: Esvet Akbaş & Furgan Aslanoğlu (2007) Studies on Reactions of Pyrimidine Compounds. 2<sup>1</sup>. Microwave-Assisted Synthesis of 1,2,3,4-Tetrahydro-2-Thioxopyrimidine Derivatives, *Phosphorus, Sulfur, and Silicon and the Related Elements*, 183:1, 82-89, DOI: [10.1080/10426500701557021](https://doi.org/10.1080/10426500701557021)

To link to this article: <http://dx.doi.org/10.1080/10426500701557021>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any

losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

## Studies on Reactions of Pyrimidine Compounds. 2<sup>1</sup>. Microwave-Assisted Synthesis of 1,2,3,4-Tetrahydro-2-Thioxopyrimidine Derivatives

**Esvet Akbaş and Furgan Aslanoglu**

Department of Chemistry, Faculty of Arts and Sciences,  
University of Yuzuncu Yil, Kampus, Van, Turkey

*The 5-benzoyl-4,6-diphenyl-1,2,3,4-tetrahydro-2-thioxopyrimidine (1a) and 5-benzoyl-4-(2-chlorophenyl)-6-phenyl-1,2,3,4-tetrahydro-2-thioxopyrimidine (1b) were synthesized using the Biginelli three-component cyclocondensation reaction of a  $\beta$ -diketone, arylaldehyde, and thiourea under microwave irradiation. Thiazolopyrimidine (2), (3) and pyrimido[2,3-b]thiazine (4) derivatives were obtained by a simple one-pot condensation reaction of starting compound 1b and 2-bromopropionic acid, bromoacetic acid and 3-bromopropionic acid, respectively. These reactions performed under microwave irradiation and conventional conditions.*

*This study is suggested a simple and efficient route for the preparation of 1,2,3,4-tetrahydro-2-thioxopyrimidine and derivatives.*

**Keywords** Microwave-Assisted; Biginelli reaction; thioxopyrimidine; synthesis; cyclocondensation; multicomponent reaction

## INTRODUCTION

Recently, there has been focused interest in the three-component cyclocondensation reaction of  $\beta$ -diketone, arylaldehyde, and (thio)urea under Brönsted acid catalysis that was firstly reported by Pietro Biginelli in 1893.<sup>2</sup> The reaction products are dihydropyrimidine derivatives, which display various types of biological and agricultural activities<sup>3</sup> such as analgesic,<sup>3a</sup> antipyretic,<sup>3b</sup> antihypertensive,<sup>3c</sup> anti-inflammatory,<sup>3d</sup>

Received 16 March 2007; accepted 5 June 2007.

This work was supported by the Scientific and Technical Research Council of Turkey (TBAG 2392 103T136). The authors would like to thank Dr. Musa Turker from Department of Biology, Faculty of Arts and Sciences, Yuzuncu Yil University for his linguistic support.

Address correspondence to Esvet Akbaş, Department of Chemistry, Faculty of Arts and Sciences, University of Yuzuncu Yil, Zeve Campus, 65080, Van, Turkey. E-mail: esvakbas@hotmail.com

pesticides,<sup>3e</sup> herbicides,<sup>3f</sup> plant growth regulators,<sup>3g</sup> and as modulators for the transport of calcium ions across the cell membrane.<sup>4</sup>

In the last few years there has been an increased interest in the use of microwave heating in organic synthesis,<sup>5</sup> and it forms now the basis of a number of commercial systems. Some interesting features of this method are the rapid reaction rates, simplicity, solvent-free conditions, and the ease of work-up after the reaction, and better selectivity. Also, microwave irradiation generates rapid intense heating of polar substances, which result in the reduction of reaction time compared to conventional heating.<sup>6</sup>

At present continuing with our studies on the synthesis of 1,2,3,4-tetrahydropyrimidine derivatives using conventional heating and microwave irradiation. The purpose of that work is to extend the Biginelli reactions in order to synthesize some 1,2,3,4-tetrahydro-2-thioxopyrimidine derivatives. The reported method is suggested that a simple and efficient route for the preparation of 1,2,3,4-tetrahydro-2-thioxopyrimidine derivatives.


## RESULTS AND DISCUSSION

In the present study, we achieved the synthesis of the pyrimidine derivatives via conventional heating as well as microwave irradiation experiments. The synthesis of derivative **1a** via conventional heating method was published in our previous work.<sup>7</sup> The derivative is also available in the product list of Aurora Fine Chemicals.<sup>8</sup> The reaction was performed 1,3-diphenyl-1,3-propanedione, benzaldehyde, and thiourea in glacial acetic acid containing a few drops of concentrated hydrochloric acid. The mixture was heated under reflux condition for 8 h. The reaction was finalized in good yield (93%). In a similar way, the compound **1b** was synthesized using 2-chlorobenzaldehyde instead of benzaldehyde (Scheme 1). Yield of product following recrystallization from 1-butanol was of the order of 90%.

When 1,3-diketone, aryl aldehyde, thiourea, and a few drops of concentrated HCl were reacted in glacial acetic acid under microwave irradiation for 5 min obtained the same final products (**1a,b**).

<sup>1</sup>H and <sup>13</sup>C NMR spectra of the isolated compounds **1a**<sup>7,8</sup> and **1b** confirm the expected structures. In <sup>1</sup>H NMR spectra of compound **1b**, the singlets at 10.6 and 9.7 ppm are due to the resonance of the two NH groups of the pyrimidine ring. The doublet and multiplet at 5.8 (*J* = 2.6) and 7.0–7.6 ppm are assigned to C<sub>4</sub>H and to the aryl protons, respectively.


On the other hand, the reactions of thioxopyrimidine derivatives having the NH and C=S group in the suitable position with various bromo



SCHEME 1

compounds were convenient methods to build the thiazolopyrimidine and pyrimido[2,3-b]thiazine derivatives.<sup>9,10</sup> Thus, the compounds **1a** and **1b** were cyclized with various bromo compounds to the thiazolopyrimidine (**2,3**) and pyrimido[2,3-b]thiazine (**4**) derivatives (Scheme 2), in approximately 35–48% yields. (The reactions of **1a** were described Aslanoglu et al.<sup>7</sup>).

The reactions of **1b** and 2-bromopropionic acid as a cyclocondensation reagent in dioxane<sup>10</sup> under reflux gave compound **2**. The IR spectrum of **2** showed absorption band at 1730, 1646  $\text{cm}^{-1}$  because of two carbonyl absorption bands. In  $^1\text{H}$  NMR spectra, the singlet at 6.6 ppm is due to



SCHEME 2

**TABLE I Comparative Study of the Synthesis of 1a, 1b, 2, 3, 4**

Compound	Medium	Yield (%)		Reaction time/min	
		MW	CONV.	MW	CONV.
1a	Glacial acetic acid	75	93	5	480
1b	Glacial acetic acid	80	90	5	480
2	Dioxane	48	35	5	120
3	Glacial acetic acid	44	41	5	60
4	Acetic anhydride CH <sub>3</sub> COONa	30	48	5	120
	Glacial acetic acid				

the resonance of the C<sub>5</sub>H proton. The quintet and doublet at 4.1 and 1.5 ppm ( $J = 7.2$  Hz) are assigned to C<sub>2</sub>H and to the methyl protons, respectively.

The reaction of **1b** with bromoacetic acid lead to 6-benzoyl-5-(2-chlorophenyl)-7-phenyl-3-oxo-2,3-dihydro-5*H*-thiazolo[3,2-*a*]pyrimidine (**3**), whereas reaction with 3-bromopropionic acid afforded 7-benzoyl-6-(2-chlorophenyl)-8-phenyl-4-oxo-2,3-dihydro-6*H*-pyrimido[2,3-*b*]thiazine (**4**). In the IR spectra of compounds **3** and **4** absence of the absorption at IR 3389 and 3146 cm<sup>-1</sup>, the characteristic absorption of NH group of starting material, is a good evidence of the expected reactions. The IR spectrum data of **3** and **4** are in agreement with similar published data.<sup>9,10</sup> The <sup>1</sup>H NMR data of the compounds **3** and **4** are in accord with the expected reaction product (See Experimental Section for details).

The compounds **2–4** were also prepared under microwave irradiation. In that way, we isolated the corresponding thioxopyrimidine derivatives, in good yields within a few minutes. The reaction times are shortened from 480 min to 5 min (Table I).

## EXPERIMENTAL

Solvents were dried by refluxing with the appropriate drying agents and distilled before use. Melting points were determined on a Barnstead Electrothermal 9200 apparatus and are uncorrected. Microanalyses were performed on LECO CHNS 932 Elemental Analyzer. The IR spectra were obtained in as potassium bromide pellets using a Mattson 1000 FTIR spectrometer. The <sup>1</sup>H and <sup>13</sup>C NMR spectra were recorded on Bruker Avance DPX-200 spectrometers, using TMS as an internal standard. All experiments were followed by TLC using DC Alufolien Kieselgel 60 F 254 Merck and Camag TLC lamp (254/366 nm). The

microwave induced reactions for the synthesis of **1a,b** were carried out in a Sinbo SMO 3606 domestic oven with a 230 V–50 Hz power source, 700W output, and 2450 MHz operating frequency.

### Conventional Synthesis— Preparation of the Starting Compounds 1,2,3,4-Tetrahydro-2-thioxopyrimidines (**1a,b**)

A mixture of 1,3-diphenyl-1,3-propanedione (0.3584 g, 1.6 mmol), aryl aldehyde (1.1 mmol), thiourea (0.084 g, 1.1 mmol) and 20 ml of glacial acetic acid containing a few drops concentrated hydrochloric acid was heated under reflux for 8 h. The solution was allowed several hours to yield 0.36 g (90%) of product **1b**, mp 198–199°C (1-butanol). IR (KBr): 3389 and 3146  $\text{cm}^{-1}$  (2NH), 2961  $\text{cm}^{-1}$  ( $\text{C}_4\text{H}$ ), 1625 ( $\text{C}=\text{O}$ ) and 1270  $\text{cm}^{-1}$  ( $\text{C}=\text{S}$ )<sup>11</sup>.  $^1\text{H}$ -NMR ( $\text{DMSO}-d_6$ ):  $\delta$  10.6 (s, 1H, NH), 9.7 (s, 1H, NH), 7.0–7.6 (m, 14H, Harom.), 5.8 (d,  $J = 2.6$ , 1H,  $\text{C}_4\text{H}$ ).  $^{13}\text{C}$  NMR ( $\text{DMSO}-d_6$ ):  $\delta = 196.1$  ( $\text{C}=\text{O}$ ), 176.5 ( $\text{C}=\text{S}$ ),<sup>12</sup> 145.3, 141.4, 140.2, 134.1, 133.6, 133.1, 131.6, 131.5, 131.4, 131.3, 131.1, 130.2, 129.5, 129.3, 129.2, 111.1, 55.6 ppm. Anal. Calcd. for  $\text{C}_{23}\text{H}_{17}\text{ClN}_2\text{OS}$  (404): C, 68.22; H, 4.23; N, 6.92. Found: C, 68.13; H, 4.16; N, 6.79.

The Compound **1a** was yield 93% which have been described in Aslanoglu et al.<sup>7</sup>

### 6-Benzoyl-5-(2-chlorophenyl)-2-methyl-7-phenyl-3-oxo-2,3-dihydro-5H-thiazolo[3,2-a] pyrimidine (**2**)

A mixture of **1b** (0.404 g, 1 mmol) and 2-bromopropionic acid (0.1 ml, 1 mmol) in dioxane (5 ml) were refluxed for 2 h. The reaction mixture was cooled and the precipitate filtered off and then washed with water. The crude product was recrystallized from ethanol. Yield 0.16 g (35%). M.p. 199–200°C, IR(KBr): 3059  $\text{cm}^{-1}$  ( $\text{C}_5\text{H}$ ), 1730, 1646  $\text{cm}^{-1}$  ( $\text{C}=\text{O}$ ),  $^1\text{H}$ -NMR ( $\text{DMSO}-d_6$ ):  $\delta$  6.9–7.5 (m, 14H, Harom.), 6.6 (s, 1H,  $-\text{C}_5\text{H}$ ), 4.1 (q, 1H,  $-\text{C}_2\text{H}$ ,  $J = 7.2$  Hz), 1.5 (d, 3H,  $-\text{CH}_3$ ,  $J = 7.2$  Hz). Anal. Calcd. for  $\text{C}_{26}\text{H}_{19}\text{ClN}_2\text{O}_2\text{S}$  (458). C, 68.04; H, 4.17; N, 6.10. Found: C, 68.10; H, 4.19; N, 6.06.

### 6-Benzoyl-5-(2-chlorophenyl)-7-phenyl-3-oxo-2,3-dihydro-5H-thiazolo[3,2-a]pyrimidine(**3**)

A mixture of **1b** (0.404 g, 1mmol), bromoacetic acid (0.153 g, 1.1 mmol), anhydrous sodium acetate (2 mmol), acetic anhydride (1.2 ml) in acetic acid (20 ml) was heated under reflux for 1 h. The residue was treated with water (100 ml) and the precipitate filtered off and the formed crude product was recrystallized from ethanol. Compound **3** was obtained in yield 0.18 g (41%). M.p. 200–201°C, IR(KBr): 3058  $\text{cm}^{-1}$  ( $\text{C}_5\text{H}$ ), 1734, 1638  $\text{cm}^{-1}$  ( $\text{C}=\text{O}$ ).  $^1\text{H}$ -NMR ( $\text{CDCl}_3$ ):  $\delta$  6.9–7.6 (m, 14H, Harom.), 6.4

(s, 1H,  $-\text{C}_5\text{H}$ ), 3.8–4.0 (q, 2H,  $\text{CH}_2$ ,  $J = 8.7$  Hz, AB system). Anal. Calcd. for  $\text{C}_{25}\text{H}_{17}\text{ClN}_2\text{O}_2\text{S}$  (444). C, 67.49; H, 3.85; N, 6.30. Found: C, 67.52; H, 3.81; N, 6.36.

### **7-Benzoyl-6-(2-chlorophenyl)-8-phenyl-4-oxo-2,3-dihydro-6H-pyrimido[2,3-b]thiazine (4)**

A mixture of **1b** (0.404 g, 1 mmol), 3-bromopropionic acid (0.168 g, 1.1 mmol), anhydrous sodium acetate (2 mmol), acetic anhydride (2 ml) in acetic acid (20 ml) was heated under reflux for 2 h. The residue was treated with water (100 ml) and the precipitate filtered off; the formed crude product was recrystallized from 2-propanol. The compound **4** was obtained in yield 0.22 g (48%). M.p. 257–258°C, IR(KBr): 3059  $\text{cm}^{-1}$  ( $\text{C}_6\text{H}$ ), 1707, 1638  $\text{cm}^{-1}$  ( $\text{C}=\text{O}$ ),  $^{13}\text{C}$ -NMR ( $\text{CDCl}_3$ -DMSO- $d_6$ ):  $\delta = 195.4$  ( $\text{C}=\text{O}$ , benzoyl), 167.5 ( $\text{C}=\text{O}$ ), 154.4, 145.4, 137.1, 136.9, 136.0, 132.0, 131.6, 130.5, 129.8, 129.3, 128.7, 128.5, 128.3, 127.3, 127.2, 126.9, 117.7, 52.7, 35.4, 21.4 ppm. Anal. Calcd. for  $\text{C}_{26}\text{H}_{19}\text{ClN}_2\text{O}_2\text{S}$  (458). C, 68.04; H, 4.17; N, 6.10. Found: C, 68.49; H, 4.76; N, 6.45.

### **Microwave Mediated Synthesis—Preparation of the Starting Compounds 1,2,3,4-Tetrahydro-2-thioxopyrimidines (1a,b)**

A mixture of 1,3-diphenyl-1,3-propanedione (1.6 mmol), aryl aldehyde (1.1 mmol), thiourea (1.1 mmol), and concentrated hydrochloric acid (0.2 ml) with glacial acetic acid (10 ml) were placed in a 25-ml glass beaker and stirred at room temperature for 5 min with a magnetic stirrer. The beaker was placed inside a larger container filled with boiling chips and then was inserted into the microwave oven. The mixture was then subjected to microwave irradiation, 5 min for the compound **1a** and **1b**.

### **5-Benzoyl-4,6-diphenyl-1,2,3,4-tetrahydro-2-thioxopyrimidine (1a)**

The reaction mixture was cooled and the crude products were filtered and recrystallized from acetic acid to give the pure compound **1a**, yield 75%.

### **5-Benzoyl-4-(2-chlorophenyl)-6-phenyl-1,2,3,4-tetrahydro-2-thioxopyrimidine (1b)**

The crude products were filtered and recrystallized from 1-butanol to give **1b**, yield 80%.

### **6-Benzoyl-5-(2-chlorophenyl)-2-methyl-7-phenyl-3-oxo-2,3-dihydro-5H-thiazolo[3,2-a]pyrimidine (2)**

A mixture of **1b** (0.404 g, 1 mmol) and 2-bromopropionic acid (0.1 ml, 1 mmol) in dioxane (2 ml) were stirred at room temperature for 5 min



with a magnetic stirrer. The mixture was irradiated for 5 min in microwave oven. The reaction mixture was cooled and then washed with ethanol and the crude products were filtered. The crude product was recrystallized from ethanol. Yield 0.22 g (48%).

**6-Benzoyl-5-(2-chlorophenyl)-7-phenyl-3-oxo-2,3-dihydro-5H-thiazolo[3,2-a]pyrimidine(3)**

A mixture of **1b** (0.404 g, 1 mmol), bromoacetic acid (0.153 g, 1.1 mmol), anhydrous sodium acetate (2 mmol), acetic anhydride (1.2 ml) in acetic acid (15 ml) were stirred at room temperature for 5 min with a magnetic stirrer. The mixture was irradiated for 5 min in a microwave oven. The reaction mixture was cooled and the residue was treated with water (100 ml); the precipitate filtered off and the formed crude product was recrystallized from ethanol. Yield 0.196 g (44%).

**7-Benzoyl-6-(2-chlorophenyl)-8-phenyl-4-oxo-2,3-dihydro-6H-pyrimido[2,3-b]thiazine (4)**

A mixture of **1b** (0.404 g, 1 mmol), 3-bromopropionic acid (0.168 g, 1.1 mmol), anhydrous sodium acetate (2 mmol), acetic anhydride (2 ml) in acetic acid (15 ml) were stirred at room temperature for 5 min with a magnetic stirrer. The mixture was irradiated for 5 min in microwave oven. The reaction mixture was cooled and the crude products were filtered and washed with water and the formed crude product was recrystallized from 2-propanol. Yield 0.134 g (30%).

## CONCLUSION

In conclusion, we have described a simple and good yielding preparation for thioxopyrimidines by conventional heating and microwave irradiation via Biginelli three-component cyclocondensation reaction. Our investigations will continue on this subject, and the results will be published when our studies are complete.

## REFERENCES

- [1] For part 1 see [7]
- [2] C. O. Kappe, *Molecules*, **3**, 1– 9 (1998); and references cited therein: P. Biginelli, *Gazz. Chim. Ital.*, **23**, 360 (1893).
- [3] N. A. Hassan, *Molecules*, **5**, 827 (2000); references cited therein: (a) M. Pemmsin, C. Lnu-Due, F. Hoguet, C. Gaultier, and J. Narcisse, *Eur. J. Chem.*, **23**, 543 (1988); (b) P. A. S. Smith and R. O. Kan, *J. Org. Chem.*, **29**, 2261 (1964); (c) A. Cannito, M. Pemmsin, C. Lnu-Due, F. Hoguet, C. Gaultier, and J. Narcisse, *Eur. J. Chem.*, **25**, 635 (1990); (d) S. Nega, J. Aionso, A. Diazj, and F. Junquere, *J. Heterocycl. Chem.*, **27**, 269 (1990); (e) S. Tetsuo, T. Mikio, H. Hidetoshi, H. Daijiro, and I. Akira, *Jpn.*

- Kokai Tokyo Koho JP, **62**, 132, 884 (1987); *Chem. Abstr.*, **107**, 198350h (1987); (f) P. K. Chakaravorty, W. J. Grelnee, K. Dooseap, N. B. Mantlo, and A. A. Patchett, A.P.C.T. Int. Appl. WO, 92.20.687.156 (1992); *Chem. Abstr.*, **118**, 213104d (1993); (g) C. J. Shishoo and K. S. Jain, *J. Heterocycl. Chem.*, **29**, 883 (1992).
- [4] C. O. Kappe, *Tetrahedron*, **49**, 6937 (1993).
- [5] (a) M. Larhed, C. Moberg, and A. Hallberg, *Acc. Chem. Res.*, **35**, 717 (2002); (b) C. O. Kappe, *Angew. Chem. Int. Ed.*, **43**, 6250 (2004).
- [6] A. Mobinikhaledi and N. Forughifar, *Phosphorus, Sulfur, and Silicon*, **181**, 2653–2658 (2006); references cited therein: (a) R. S. Varma, *Green. Chem.*, **1**, 43 (1999); (b) S. Caddick, *Tetrahedron*, **51**, 1043 (1995); (c) R. Laurent, A. Laporterie, J. Dubac, J. Berlan, S. Lefeuvre, and M. Audhuy, *J. Org. Chem.*, **57**, 7099 (1992); (d) D. M. P. Mingos and D. R. Baghurst, *Chem. Soc. Rev.*, **20**, 1 (1991).
- [7] F. Aslanoğlu, E. Akbaş, M. Sönmez, and B. Anil, *Phosphorus, Sulfur, and Silicon*, **182**, 1589–1597 (2007).
- [8] Aurora Fine Chemicals Product List, **ck-0179** (2005).
- [9] C. O. Kappe and P. Roschger, *J. Heterocyclic Chem.*, **26**, 55 (1989).
- [10] A. Mobinikhaledi, N. Forughifar, and F. Goodarzi, *Phosphorus, Sulfur, and Silicon*, **178**, 2539–2543 (2003).
- [11] A. A. Aly, *Phosphorus, Sulfur, and Silicon*, **181**, 1285–1298 (2006).
- [12] K. Z. Beresneviciute, Z. Beresnevicius, G. Mikulskiene, J. Kihlberg, and J. Broddefalk, *Magnetic Resonance in Chem.*, **35**, 553–555 (1997).