


ChemComm

Chemical Communications

www.rsc.org/chemcomm



ISSN 1359-7345



COMMUNICATION

Akira Onoda, Takashi Hayashi *et al.*

H₂O₂-dependent substrate oxidation by an engineered diiron site in a bacterial hemerythrin

H₂O₂-dependent substrate oxidation by an engineered diiron site in a bacterial hemerythrin†

Yasunori Okamoto,^a Akira Onoda,^{*a} Hiroshi Sugimoto,^b Yu Takano,^c Shun Hirota,^d Donald M. Kurtz Jr.,^e Yoshitsugu Shiro^b and Takashi Hayashi^{*a}

Cite this: *Chem. Commun.*, 2014, 50, 3421

Received 22nd October 2013,
Accepted 9th December 2013

DOI: 10.1039/c3cc48108e

www.rsc.org/chemcomm

The O₂-binding carboxylate-bridged diiron site in DcrH-Hr was engineered in an effort to perform the H₂O₂-dependent oxidation of external substrates. A His residue was introduced near the diiron site in place of a conserved residue, Ile119. The I119H variant promotes the oxidation of guaiacol and 1,4-cyclohexadiene upon addition of H₂O₂.

O₂ is utilized in several non-heme diiron-carboxylate proteins performing essential biological processes ranging from transport of O₂ (hemerythrin, Hr),^{1,2} hydroxylation of C–H bonds (methane monooxygenase hydroxylase),³ generation of a tyrosyl radical for DNA synthesis (ribonucleotide reductase),⁴ and reduction of unsaturated fatty acids ($\Delta 9$ desaturase).⁵ Despite these functional differences, all of these proteins have a carboxylate-bridged diiron active site within a four-helix bundle protein folding motif.^{2a,6} The coordination sphere and the environment surrounding the diiron site in these proteins have apparently evolved to optimize a variety of reactions with O₂.

Engineering of the first or the second coordination sphere of the diiron site⁷ constitutes one approach to understanding the effects of the protein environment, which distinguish reversible O₂ binding from O₂ activation. As a target protein for engineering the coordination sphere, we have selected DcrH-Hr,⁸ which contains the histidine-rich coordination sphere and exhibits spectroscopic properties characteristic of all Hrs, but contains a larger substrate-accessible

tunnel within the four-helix bundle compared to classical Hrs.^{9,10} As for classical Hrs, the diiron site of DcrH-Hr reversibly binds O₂ but shows more rapid autoxidation. Recently, we succeeded in altering the first coordination sphere of the diiron site in DcrH-Hr by replacing residue Ile119 located close to the exogenous ligand binding site with a Glu residue.¹¹ The I119E variant showed altered properties of exogenous ligand binding and redox behavior in the wild type DcrH-Hr (WT). Herein, we report another engineered variant DcrH-Hr, I119H, of the same residue, in which alteration of the coordination sphere generates the capability of H₂O₂-dependent oxidation of exogenous substrates.

I119H was expressed from *Escherichia coli* and purified as a holo-form containing approximately 1.6 iron atoms per protein, as determined by ferrozine analysis.¹² The diferric form (met-I119H) exhibited two absorption maxima at 325 and 375 nm, which are well-established to arise from oxo to ferric LMCT transitions in the oxo-dicarboxylate-bridged met-Hrs.^{1b,c,8} Resonance Raman (rR) spectra of I119H gave a characteristic peak at 499 cm^{−1} for the symmetric stretching Fe–O_{oxo}–Fe vibrational mode.⁸ The met-I119H protein was crystallized, and the structure of the oxo-bridged diiron site¹³ was determined by X-ray crystallographic analysis at a resolution of 1.9 Å. The overall structure of the met-I119H is similar to that of the met form of WT (met-WT) (RMSD value, 0.324).¹⁰ Surprisingly, H119 coordinates to Fe1 in the crystal structure in place of the conserved H118 ligand residue in WT (Fig. 1). The coordination geometry of other His residues and carboxylate ligands is almost identical to that of met-WT (Fig. S1, ESI†). A computational study of the I119H diiron site yielded alternative structures in which either H118 or H119 coordinates to the Fe1 (Fig. 2).¹⁴ It is, thus, expected that either H118 or H119 coordination would be stable in I119H in solution.

The exogenous ligand binding properties of the met-I119H diiron site were investigated using azide, which is known to coordinate to Fe2 in place of chloride in met-WT.⁹ The azide adduct of met-WT exhibits a UV-vis absorption at 443 nm,⁸ while that of I119H showed a shifted absorption maximum at 425 nm, indicating a change in conformation or orientation of the bound azide ligand (Fig. S2, ESI†). The rR spectrum of I119H showed the Fe–N₃ stretching vibration at 361 cm^{−1}, which was confirmed by ¹⁴N₃[−]/¹⁵N₃[−] isotope

^a Department of Applied Chemistry, Graduate School of Engineering, Osaka University, Suita, Osaka 565-0871, Japan.

E-mail: onoda@chem.eng.osaka-u.ac.jp, thayashi@chem.eng.osaka-u.ac.jp

^b Biometal Science Laboratory, RIKEN SPring-8 Center, Sayo, Hyogo 679-5148, Japan

^c Institute for Protein Research, Osaka University, Suita, Osaka 565-0871, Japan

^d Graduate School of Materials Science, Nara Institute of Science and Technology, Ikoma 630-0192, Japan

^e Department of Chemistry, University of Texas at San Antonio, San Antonio, Texas 78249, USA

† Electronic supplementary information (ESI) available: Information on materials, instrumentation, experimental details and additional data on preparation of proteins, crystal structure analysis, resonance Raman and FTIR spectroscopy, reaction of reduced I119H with O₂, consumption of H₂O₂, and oxidation reactions of guaiacol and 1,4-cyclohexadiene. The atomic coordinates and structure factors (PDB code 3WHN) have been deposited into the Protein Data Bank, <http://www.rcsb.org/>. See DOI: 10.1039/c3cc48108e






Fig. 1 Crystal structures of (a) met-WT (PDB code: 3AGT) and (b) met-I119H (iron in brown, oxygen in red, nitrogen in blue, carbon in white, sulfur in yellow, and chloride in green). The $2F_{\text{obs}} - F_{\text{calc}}$ electron density in light blue grid (contoured at 1.5σ) around the diiron site. The structure drawings were generated in PyMOL.¹⁵




Fig. 2 Depiction of computationally optimized structures of I119H diiron site models (a) where H118 coordinates to Fe1 and (b) where H119 coordinates to Fe1 (iron in brown, carbon in white, oxygen in red, nitrogen in blue, chloride in green, and sulfur in yellow).




Fig. 3 (a) Resonance Raman spectra of the azide adduct of (i) met-WT with Na^{14}N_3 , (ii) met-I119H with Na^{14}N_3 , (iii) met-I119H with Na^{15}N_3 , and (iv) the difference spectrum $^{15}\text{N}_3$ minus $^{14}\text{N}_3$ adducts of met-I119H. Peaks are labeled according to their vibrational assignments, as described in the text. (b) FTIR spectra of the azide adduct of (i) met-WT at 5 K, (ii) met-I119H at 5 K, and (iii) met-I119H at 25 °C.

labeling experiments (Fig. 3a). The Fe–N₃ stretching vibration of I119H is downshifted compared to that of the WT at 372 cm^{-1} .⁸ This finding suggests weaker Fe–N₃ bonding in I119H than that observed in WT.

Additional structural details of the bound azide ligand were obtained from FTIR spectroscopy in the region of the asymmetric stretch of the coordinated azide $\nu_{\text{as}}(\text{NNN})$. The azide adduct of met-WT showed a $\nu_{\text{as}}(\text{NNN})$ band at 2050 cm^{-1} (Fig. 3b). The isotope labeling experiment using $^{15}\text{NN}_2$ indicates that the terminal nitrogen of the azide ligand binds in a η^1 fashion, as expected from the X-ray crystal structure of the met-WT azide adduct.^{1,9,16} The azide adduct of met-I119H exhibits two $\nu_{\text{as}}(\text{NNN})$ bands at 2050 and 2060 cm^{-1} at 5 K (Fig. 3b). The newly observed band, 10 cm^{-1} -higher than that of met-WT, suggests a resonance form shifted farther from the symmetric $[\text{Fe}^{3+}\text{--N}^{\text{--}}\text{=N}^{\text{+}}\text{=N}^{\text{--}}]$ towards the asymmetric $[\text{Fe}^{3+}\text{--N}^{2-}\text{--N}^{\text{+}}\text{=N}]$. The increased asymmetry of the two N–N bonds could be stabilized by a hydrogen bond interaction with the 1N atom of the coordinated azide, possibly with the displaced His119 ligand.¹⁷ No such hydrogen bonding interaction is observed in the structure of the met-WT azide adduct.⁹ The two $\nu_{\text{as}}(\text{NNN})$ bands observed at 5 K coalesced at 25 °C (Fig. 3b). Based on this observation together with the crystal structure and calculated optimized structures, we propose two conformations for I119H: H118off/H119on and H118on/H119off. In the latter case, the H119 residue is capable of forming a hydrogen bonding interaction with an exogenous ligand.

We evaluated the reactivity of I119H toward H_2O_2 as a potential exogenous ligand, which, like azide, could form a hydrogen bond with the pendant H119 residue. The met-WT and met-I119H in 50 mM HEPES (pH 7.0) were individually treated with 50 eq. of H_2O_2 from an aqueous solution, and the amount of H_2O_2 consumed was determined by the iodometry method. Disproportionation of H_2O_2 into O_2 and H_2O has been reported to occur upon reaction of H_2O_2 with a synthetic Hr model complex.¹⁸ We found that the met-WT reacted with H_2O_2 within 10 min with a concomitant evolution of O_2 , which was confirmed by gas chromatography (Fig. S4, ESI†). In addition, the deoxy-WT was oxidized to the met form by H_2O_2 (Fig. S5, ESI†). These results indicate that the disproportionation reaction of H_2O_2 could proceed *via* a mechanism analogous to that proposed for the Hr model complex (Scheme S1, ESI†). The consumption rate of H_2O_2 by I119H was significantly slower than that of WT (Fig. 4a).

Motivated by the differing reactivity of WT and I119H toward H_2O_2 , we tested the promotion of oxidation of external substrates by these proteins. Fig. 4b demonstrates that I119H promotes oxidation of guaiacol by H_2O_2 , as shown by the clear increase in absorption at 470 nm assigned to the oxidized guaiacol product, whereas, under the same conditions, WT does not. Similarly, I119H promotes the oxidation of 1,4-cyclohexadiene by H_2O_2 , producing benzene as a product determined by GC-MS analysis, whereas WT showed no such reactivity (Fig. S6, ESI†). These findings support the generation of an oxidizing intermediate from the altered diiron site in I119H which cannot be generated in WT. According to the reaction analysis, the oxidized product yields for both the guaiacol and cyclohexadiene reactions are approximately 0.26 mol product per mol of a diiron site under the conditions as






Fig. 4 (a) Consumption of H₂O₂ in the presence of met-I119H (solid line) and met-WT (dashed line); [protein] = 100 μ M, [H₂O₂] = 5 mM. All experiments were carried out in 50 mM HEPES (pH 7.0) at 25 $^{\circ}$ C. (b) Oxidation reaction of guaiacol by met-I119H (solid line) and met-WT (dashed line) at 25 $^{\circ}$ C. The absorption of the guaiacol product at 470 nm was monitored; [protein] = 100 μ M, [guaiacol] = 5 mM, [H₂O₂] = 5 mM.

described in Fig. 4b. During the substrate oxidation reaction, H₂O₂ is also expected to be consumed by its concurrent disproportionation. We confirmed that the oxidation of guaiacol could be continued upon the addition of H₂O₂ to the initial reaction solution (Fig. S7, ESI[†]). The key for the generation of the oxidizing intermediate could be a hydrogen bonding interaction between a hydroperoxide coordinating in a fashion similar to that in oxyHr and the displaced H119 ligand. Previous studies have invoked hydrogen bonding to a hydroperoxide bound to Fe(III) as promoting formation of a high-valent iron-oxo species.¹⁹ Support for such a hydrogen bonding interaction comes from the azide ligation experiment monitored by FTIR spectroscopy. An additional contribution could involve dynamic coordination of H118 and H119, which transiently forms a five-coordinated Fe1 site *via* a conformational transition during the reaction. This transient coordination site may allow formation of a μ -1,2-peroxo-diferric species, which is the most commonly proposed precursor to high-valent iron-oxo species in non-heme diiron-carboxylate enzymes.²⁰

In conclusion, we have engineered the diiron site of DcrH-Hr to promote the H₂O₂-dependent oxidation of exogenous substrates. A His residue introduced proximal to the diiron site will participate in a hydrogen bonding interaction with the exogenous H₂O₂ ligand, thereby, activating the peroxide. To the best of our knowledge, this is the first example that successfully converts an O₂-binding non-heme diiron-carboxylate protein to an oxidatively active one. Efforts to characterize the oxidatively active species of the I119H variant and the further engineering of the diiron site are in progress.

This work was financially supported by Grants-in-Aid for Scientific Research ((C), JSPS KAKENHI Grant Number 205590020, and Innovative Areas "Molecular Activation", area 2204, MEXT KAKENHI Grant Number 221050130). Y.O. appreciates support from the Research Fellowship of JSPS and Global COE Program "Global Education and Research Center for Bio-environmental Chemistry" of Osaka University. D. M. K., Jr. acknowledges support from the National Institutes of Health (grant R01 GM040388).

The computations were performed at the Research Center for Computational Science, Okazaki, Japan.

Notes and references

- (a) D. M. Kurtz Jr., D. F. Shriver and I. M. Klotz, *Coord. Chem. Rev.*, 1977, **24**, 145; (b) P. C. Wilkins and R. G. Wilkins, *Coord. Chem. Rev.*, 1987, **79**, 195; (c) R. E. Stenkamp, *Chem. Rev.*, 1994, **94**, 715.
- (a) D. M. Kurtz Jr., *JBC*, *J. Biol. Inorg. Chem.*, 1997, **2**, 159; (b) D. M. Kurtz Jr., *Dioxygen-binding Proteins*, in *Comprehensive Coordination Chemistry II*, ed. J. A. McCleverty and T. J. Meyer, Elsevier, Oxford, U.K., 2004, vol. 8, p. 229.
- (a) B. J. Wallar and J. D. Lipscomb, *Chem. Rev.*, 1996, **96**, 2625; (b) C. E. Tinberg and S. J. Lippard, *Acc. Chem. Res.*, 2011, **44**, 280.
- (a) C. Krebs, J. M. Bollinger and S. J. Booker, *Curr. Opin. Chem. Biol.*, 2011, **15**, 291; (b) A. B. Tomter, G. Zoppellaro, N. H. Andersen, H. Hersleth, M. Hammerstad, Å. K. Røhr, G. K. Sandvik, K. R. Strand, G. E. Nilsson, C. E. Bell III, A. Barra, E. Blasco, L. L. Pape, E. I. Solomon and K. K. Andersson, *Coord. Chem. Rev.*, 2013, **257**, 3.
- B. G. Fox, K. S. Lyle and C. E. Rogge, *Acc. Chem. Res.*, 2004, **37**, 421.
- (a) P. Nordlund and H. Eklund, *Curr. Opin. Struct. Biol.*, 1995, **5**, 758; (b) M. H. Sazinsky and S. J. Lippard, *Acc. Chem. Res.*, 2006, **39**, 558.
- (a) J. Xiong, R. S. Phillips, D. M. Kurtz Jr., S. Jin, J. Ai and J. Sanders-Loehr, *Biochemistry*, 2000, **39**, 8526; (b) C. S. C. Farmer, D. M. Kurtz Jr., R. S. Phillips, J. Ai and J. Sanders-Loehr, *J. Biol. Chem.*, 2000, **275**, 17043; (c) C. S. Farmer, D. M. Kurtz Jr., Z. J. Liu, B. C. Wang, J. Rose, J. Ai and J. Sanders-Loehr, *JBC*, *J. Biol. Inorg. Chem.*, 2001, **6**, 418; (d) M. Faiella, C. Andreozzi, R. T. M. de Rosales, V. Pavone, O. Maglio, F. Natri, W. F. DeGrado and A. Lombardi, *Nat. Chem. Biol.*, 2009, **5**, 882; (e) A. J. Reig, M. M. Pires, R. A. Snyder, Y. Wu, H. Jo, D. W. Kulp, S. E. Butch, J. R. Calhoun, T. G. Szyperski, E. I. Solomon and W. F. DeGrado, *Nat. Chem.*, 2012, **4**, 900.
- J. Xiong, D. M. Kurtz Jr., J. Ai and J. Sanders-Loehr, *Biochemistry*, 2000, **39**, 5117.
- C. E. Isaza, R. Silaghi-Dumitrescu, R. B. Iyer, D. M. Kurtz Jr. and M. K. Chan, *Biochemistry*, 2006, **45**, 9023.
- A. Onoda, Y. Okamoto, H. Sugimoto, Y. Shiro and T. Hayashi, *Inorg. Chem.*, 2011, **50**, 4892.
- Y. Okamoto, A. Onoda, H. Sugimoto, Y. Takano, S. Hirota, D. M. Kurtz Jr., Y. Shiro and T. Hayashi, *Inorg. Chem.*, 2013, **52**, 13014.
- L. L. Stookey, *Anal. Chem.*, 1970, **42**, 779.
- The diferric form of the I119H variant is oxidized by O₂ to the diferric form (Fig. S3, ESI[†]). Although this fact suggests the binding of O₂, the oxy form was not observed due to more rapid autoxidation and slower O₂ binding relative to those of WT DcrH-Hr.
- The total energy of the first coordination sphere in the H118on/H119off and H118off/H119on models differs only 0.67 kcal mol⁻¹. In the H118on/H119off model, the hydrogen bonding interaction between H119 and chloride was formed. The interaction energy between H118 and H119 in the H118on/H119off model and that between H118 and M120 in the H118off/H119on model is 5.88 and 1.84 kcal mol⁻¹, respectively.
- W. L. DeLano, *The PyMOL Molecular Graphics System*, DeLano Scientific, San Carlos, CA, 2008.
- The peak at 2043 cm⁻¹ was assigned as $\nu_{as}(\text{NNN})$ of a free azide. The azide (¹⁵NN₂) adduct of met-WT gave two $\nu_{as}(\text{NNN})$ peaks at 2033 and 2045 cm⁻¹.
- S. Lu, M. H. Sazinsky, J. W. Whittaker, S. J. Lippard and P. Moënnel-Loccoz, *J. Am. Chem. Soc.*, 2005, **127**, 4148.
- B. Maurer, J. Crane, J. Schuler, K. Wieghardt and B. Nuber, *Angew. Chem., Int. Ed.*, 1993, **32**, 289.
- (a) J. T. Groves, *Cytochrome P450*, Springer US, Boston, MA, 2005, p. 1; (b) A. R. McDonald and L. Que, Jr., *Coord. Chem. Rev.*, 2013, **257**, 414.
- (a) A. Ghosh, F. Tiago de Oliveira, T. Yano, T. Nishioaka, E. S. Beach, I. Kinoshita, E. Münck, A. D. Ryabov, C. P. Horwitz and T. J. Collins, *J. Am. Chem. Soc.*, 2005, **127**, 2505; (b) M. Kodera, M. Itoh, K. Kano, T. Funabiki and M. Reglier, *Angew. Chem., Int. Ed.*, 2005, **44**, 7104; (c) G. Xue, A. T. Fiedler, M. Martinho, E. Münck and L. Que, Jr., *Proc. Natl. Acad. Sci. U. S. A.*, 2008, **105**, 20615; (d) G. Xue, D. Wang, R. De Hont, A. T. Fiedler, X. Shan, E. Münck and L. Que, Jr., *Proc. Natl. Acad. Sci. U. S. A.*, 2007, **104**, 20713; (e) M. Kodera, Y. Kawahara, Y. Hitomi, T. Nomura, T. Ogura and Y. Kobayashi, *J. Am. Chem. Soc.*, 2012, **134**, 13236.

