

Metal-template synthesis and co-ordination properties of a palladium complex containing a novel and stable imidazole-substituted phosphine C–P bidentate chelate

Huifang Lang, Jagadese J. Vittal and Pak-Hing Leung*

Department of Chemistry, National University of Singapore, Kent Ridge 119260, Singapore


An organopalladium complex promoted Diels–Alder reaction between 1-phenyl-3,4-dimethylphosphole and 1-vinylimidazole gave a novel imidazole-substituted phosphanorbornene bidentate ligand which co-ordinated to the palladium template *via* the C² carbon atom of the imidazole group and the bridgehead phosphorus donor atom.

Transition-metal complexes containing imidazole and its derivatives play an important role in bioinorganic chemistry.¹ These compounds are frequently considered as models in the development of metal-based enzymes and proteins. In terms of co-ordination chemistry, imidazole may be considered as an ambidentate ligand. For instance, it has been well established that the imidazole unit co-ordinates to transition-metal ions, such as palladium(II) and platinum(II), predominantly *via* one of its nitrogen atoms. In some rare cases, however, the cyclic unit may also co-ordinate as a carbene or an amidine ligand *via* its C² carbon atom. This interesting mode of organometallic bonding has been observed in a small number of Cr⁰, Fe⁰, Ru^{II} and Ru^{III} complexes.² Apart from their biological interest, the availability of stable C-bound imidazole heavy-metal complexes is important for the study of the *trans* influence³ and other related phenomena that are pertinent to the design and development of efficient support ligands for homogenous catalysis.⁴ Indeed, it has been reported that palladium complexes containing C-bound imidazole ligands are efficient catalysts for cross-coupling reactions.⁵ In general, however, palladium complexes containing monodentate ligands are kinetically labile. We believe that the development of a new class of stable C-bound imidazole complexes may significantly influence the design of catalysts in homogenous catalysis. Here we report the palladium-template synthesis of the first imidazole-substituted tertiary phosphine bidentate ligand in which the C² carbon of the imidazole unit is involved in metal chelation.

In the absence of the transition-metal ion, no Diels–Alder reaction was observed between 1-phenyl-3,4-dimethylphosphole (DMPP) and 1-vinylimidazole. Upon co-ordination to palladium, however, DMPP is activated towards the [4 + 2] cycloaddition reaction. Thus, when [PdCl₂(DMPP)₂] or the organopalladium complex **1**⁶ was treated with the dienophile in 1,2-dichloroethane at 84 °C, the *exo*-cycloadduct **2** was obtained as the sole product (Scheme 1).[†] When the template complex **1** was used, the cycloaddition reaction was completed in 30 d. Interestingly, during the course of heating, the N³ nitrogen of the imidazole unit underwent a parallel N-alkylation reaction with 1,2-dichloroethane and produced HCl as the side product.⁷ The HCl thus generated led to further chemoselective cleavage of the orthometallated benzylamine ligand from the palladium template and thus facilitate the formation of the dichloro complex **2** as the final product.⁶ The

molecular structure and the co-ordination chemistry of **2** have been determined by X-ray structural analysis (Fig. 1).[‡] The study reveals that the reaction of **1** with 1-vinylimidazole in 1,2-dichloroethane has resulted in the removal of the benzylamine chelate and the imidazole substituted *exo*-phosphanorbornene ligand created co-ordinates to palladium as a bidentate chelate *via* the bridgehead phosphorus atom and the C² carbon atom of the imidazole group. The palladium atom is in a slightly distorted square-planar geometry with the bond angles in the ranges 84.8(1)–94.8(2) and 167.0(1)–172.2(2)°. Due to the aromaticity of the imidazole ring, all C–C and C–N bonds [1.350(8)–1.383(6) Å] within the five-membered ring are noticeably shorter than the two attached C–N bonds [1.473(7)–1.453(7) Å]. The Pd(1)–C(11) distance of 1.993(5) Å is similar to the Pd–C bonds observed in other reported complexes containing the orthometallated benzylamine [2.004(11) Å] and naphthylamine units (2.006 Å) which experience a similar *trans* electronic influence from a chloro ligand.⁸ The Pd(1)–P(1) distance of 2.207(1) Å is also within the normal range observed for this class of phosphanorbornene complex. The Pd···Cl(3) distance is 3.622(2) Å indicating that there is no interaction between the two heavy atoms.

Interestingly, when [PdCl₂(DMPP)₂] was used for the Diels–Alder reaction, a much lower reaction rate was observed. The ³¹P NMR studies indicated that 80% of [PdCl₂(DMPP)₂]



Scheme 1

[†] Preparation of complex **2**. A solution of the organopalladium complex **1** (0.46 g, 0.97 mmol) in 1,2-dichloroethane (40 cm³) was treated with silver perchlorate (0.2 g, 0.97 mmol) in water (1 cm³) for 30 min. The resulting mixture was filtered through a layer of Celite to remove silver chloride and the organic layer was dried over anhydrous MgSO₄. The dried solution was treated with 1-vinylimidazole (0.37 g, 3.87 mmol) and the reaction mixture was then stirred at 84 °C for 30 d. The solution was washed with water (50 cm³) and then dried over MgSO₄. The solvent was removed under reduced pressure to give a yellow residue. Upon crystallization from acetonitrile–diethyl ether, the dichloro complex **2** was obtained as yellow prisms (0.21 g, 40%), m.p. 256–258 °C (decomp.). ³¹P NMR (CDCl₃): δ 105.3 (s).

[‡] Crystal data for **2**: C₁₉H₂₂Cl₂N₂PPd, *M* = 522.11, orthorhombic, space group *Pna*2₁, *a* = 14.9398(1), *b* = 16.5294(1), *c* = 8.6207(1) Å, *U* = 2128.85(3) Å³, *Z* = 4, *D*_c = 1.629 g cm^{−3}, *T* = 293 K, μ(Mo–Kα) = 13.29 cm^{−1}, *F*(000) = 1048, *R*₁ = 0.0408, *wR*₂ = 0.0885 for 4214 independent observed reflections [*I* > 2σ(*I*), 1.84 ≤ 2θ ≤ 29.31°] and 236 parameters. The Flack parameter was refined to −0.02(4). CCDC reference number 186/1018.

* E-Mail: chmlph@nus.edu.sg




Fig. 1 Molecular structure and co-ordination chemistry of complex **2**. Selected bond lengths (Å) and angles (°): Pd(1)–P(1) 2.207(1), Pd(1)–C(11) 1.993(5), Pd(1)–Cl(1) 2.391(2), Pd(1)–Cl(2) 2.358(2), P(1)–C(1) 1.859(6), P(1)–C(4) 1.842(5), N(1)–C(6) 1.473(7), N(1)–C(11) 1.375(7), N(1)–C(9) 1.383(6), C(9)–C(10) 1.350(8), C(10)–N(2) 1.381(7), N(2)–C(11) 1.357(7), N(2)–C(12) 1.453(7), C(13)–Cl(3) 1.781(9); P(1)–Pd–Cl(1) 167.0(1), P(1)–Pd–Cl(2) 84.8(1), P(1)–Pd–C(11) 87.8(2), C(11)–Pd–Cl(1) 94.8(2), C(11)–Pd–Cl(2) 172.2(2), Cl(1)–Pd–Cl(2) 92.0(1), C(1)–P(1)–C(4) 81.8(3), C(6)–N(1)–C(11) 124.0(4), N(1)–C(9)–C(10) 106.3, C(9)–C(10)–N(2) 107.7(4), C(10)–N(2)–C(11) 110.7(4), N(2)–C(11)–N(1) 104.4(4)

remains unchanged after the complex was treated with 1-vinylimidazole for 30 d under similar reaction conditions. Nevertheless, a small quantity of **2** was observed in the ^{31}P NMR spectra of the reaction mixture. Clearly, the co-ordinated cyclic diene in

the template complex **1** receives a higher degree of activation as compared with its counterparts in $[\text{PdCl}_2(\text{DMPP})_2]$. We are currently investigating the optical resolution and the catalytical properties of transition-metal complexes containing the imidazole-substituted phosphine ligand.

Acknowledgements

We are grateful to the National University of Singapore for support of this research (Grant No. RP972667) and research scholarships (to H. F. L.).

References

- 1 R. J. Sundberg and R. B. Martin, *Chem. Rev.*, 1974, **74**, 471; M. Grebl and B. Krebs, *Inorg. Chem.*, 1994, **33**, 3877.
- 2 F. A. Cotton and G. Wilkinson, in *Advanced Inorganic Chemistry*, John Wiley & Son, New York, 5th edn., 1988, ch. 10.
- 3 M. F. Tweedle and H. Taube, *Inorg. Chem.*, 1982, **21**, 3361.
- 4 K. H. Weiss, in *Transition Metal Carbene Complexes*, Verlag Chemie, Weinheim, 1983, p. 227.
- 5 W. A. Herrmann and C. Kocher, *Angew. Chem., Int. Ed. Engl.*, 1997, **36**, 2162.
- 6 S. Selvaratnam, P. H. Leung, A. J. P. White and D. J. Williams, *J. Organomet. Chem.*, 1997, **542**, 61.
- 7 C. A. Ghilardi, S. Midollini, S. Moneti, A. Orlandini and J. A. Ramirez, *J. Chem. Soc., Chem. Commun.*, 1989, 304.
- 8 J. Albert, J. Granell and J. Sales, *Organometallics*, 1995, **14**, 1393; S. Y. Siah, P. H. Leung and K. F. Mok, *J. Chem. Soc., Chem. Commun.*, 1995, 1747.

Received 20th May 1998; Communication 8/03820A