



A Journal of the Gesellschaft Deutscher Chemiker

Angewandte Chemie

GDCh

International Edition

www.angewandte.org

Accepted Article

Title: Iron in a cage: fixation of a Fe(II)tpy₂ complex by fourfold interlinking

Authors: Thomas Brandl, Sven Johannsen, Daniel Häussinger, Nithin Suryadevara, Alessandro Prescimone, Stefan Bernhard, Manuel Gruber, Mario Ruben, Richard Berndt, and Marcel Mayor

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: *Angew. Chem. Int. Ed.* 10.1002/anie.202006340

Link to VoR: <https://doi.org/10.1002/anie.202006340>

COMMUNICATION

Iron in a cage: fixation of a Fe(II)tpy₂ complex by fourfold interlinking

Thomas Brandl,^[a] Sven Johannsen,^[b] Daniel Häussinger,^[a] Nithin Suryadevara,^[c] Alessandro Prescimone,^[a] Stefan Bernhard,^[c] Manuel Gruber,^[b] Mario Ruben,^[d,e] Richard Berndt^[b] and Marcel Mayor*^[a,d,f]

Dedicated to our joint friend and tutor Jean-Marie Lehn

Abstract: The coordination sphere of the Fe(II) terpyridine complex **1** is rigidified by fourfold interlinking of both terpyridine ligands. Profiting from an octa-aldehyde precursor complex, the ideal dimensions of the interlinking structures are determined by reversible *Schiff*-base formation, before irreversible *Wittig* olefination provided the rigidified complex. Reversed-phase HPLC enables the isolation of the all-*trans* isomer of the Fe(II) terpyridine complex **1**, which is fully characterized. While temperature independent low-spin states were recorded with superconducting quantum interference device (SQUID) measurements for both, the open precursor **8** and the interlinked complex **1**, evidence of the increased rigidity of the ligand sphere in **1** was provided by proton T₂ relaxations in NMR experiments. The ligand sphere fixation in the macrocyclized complex **1** even reaches a level resisting substantial deformation upon deposition on a Au(111) surface, as demonstrated by its pristine form in a low temperature ultra-high vacuum scanning tunneling microscope experiment. The observed two level fluctuation of the tunneling current at elevated bias might even point at spin-switching in the deposited complex.

Switches are essential components in electronic devices and driven by the ongoing miniaturization, molecular switches move into the focus of interest.^[1–3] While various structures triggered by light, heat, or (electro) chemical potential have been reported,^[4–6] molecular switches responding on the applied potential and/or mechanical pressure are less frequent, in spite of their better miniaturization potential.

Iron bis-terpyridine (Fe(II)tpy₂) complexes are particularly appealing as potential switching motives due to their coordination geometry sensitive spin crossover (SCO) properties. And indeed, single molecule junctions with tailor-made Fe(II)tpy₂ complexes

investigated at low temperature displayed bistability features pointing at SCO events triggered by the distortion of the ligand sphere. Examples are the mechanical spin-state manipulation of a homoleptic Fe(II)tpy₂ complex in a mechanically controlled break junction (MCBJ),^[7] or electric field (*E*-field) sensing junctions with heteroleptic Fe(II)tpy₂ complexes with push-pull systems integrated in one tpy-ligand.^[8] In the latter case, the reliability of the *E*-field triggered switching was recently improved by separating the active Fe(II) complex from the electrodes.^[9]

While these MCBJ based single molecule experiments showing the potential of Fe(II)tpy₂ complexes as switchable subunits were encouraging, scanning tunneling microscopy (STM) may provide insights into the arrangement of the molecule at the electrode surface and its electron transport. In a first attempt, the arrangement of both tpy-ligands around the central Fe²⁺ ion was stabilized by macrocyclization, providing chiral Fe(II)tpy₂ complexes.^[10] Upon deposition on Au(111) in an ultra-high vacuum (UHV) STM set-up however, the complex disintegrated as clearly documented by the macrocycle spread on the metal surface (Figure SI22).^[11]

Here we report the next generation of Fe(II)tpy₂ complexes with rigidified coordination sphere. Fourfold interlinking of both tpy-ligands freezes their spatial arrangement further improving the stability of the central Fe(II)tpy₂ complex in **1** (Figure 1).




Figure 1. Fourfold interlinked Fe(II) terpyridine complexes **1** and **2**. While **1** was isolated and characterized, **2** was observed by mass spectrometry and served to identify linkers with dimensions suitable to bridge the two aldehydes.

- [a] Department of Chemistry, University of Basel, St. Johannis-Ring 19, 4056 Basel, Switzerland, E-mail: marcel.mayor@unibas.ch
Homepage: <https://www.chemie.unibas.ch/~mayor/>
- [b] Institut für Experimentelle und Angewandte Physik, Christian-Albrechts-Universität zu Kiel, Leibnizstr. 19, 24098 Kiel, Germany
- [c] Department of Chemistry, Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA
- [d] Institute of Nanotechnology, Karlsruhe Institute of Technology (KIT), Hermann-von-Helmholtz-Platz 1, 76344 Eggenstein-Leopoldshafen, Germany, E-mail: marcel.mayor@kit.edu
- [e] Institut de Physique et Chimie des Matériaux de Strasbourg (IPCMS), CNRS-Université de Strasbourg, 23, rue de Loess, BP 43, 67034 Strasbourg cedex 2, France
- [f] Lehn Institute of Functional Materials, School of Chemistry, Sun Yat-Sen University, Guangzhou 510275, China

Supporting information for this article is given via a link at the end of the document.

COMMUNICATION

And indeed, the coordination sphere of the isolated target complex **1** was considerably reinforced as displayed by *Carr-Purcell-Meiboom-Gill*-NMR experiments. The improved structural integrity of **1** even enabled its investigation by UHV STM experiments.

To explore a fourfold interlinking strategy, the octa-aldehyde complex **8** (Scheme 1) moved into the focus of interest. The symmetry of the compound facilitates its synthesis and the exposed aldehyde functions enables a variety of potential interlinking reactions, like e.g. the *McMurry* reaction,^[12] carbonyl-olefin metathesis,^[13] *Schiff* base condensation^[14] or *Wittig* olefination.^[15]

And indeed, as displayed in Scheme 1, the octa-aldehyde Fe(II)tpy₂ complex **8** was obtained in five steps. First the aldehyde groups of commercially available 2-bromoisophthalaldehyde **3** were protected by refluxing **3** in toluene in the presence of ethylene glycol and catalytic amounts of *p*-toluenesulfonic acid. By removing the water with a *Dean-Stark* apparatus the reaction was driven to completion and the bis-dioxolan species **4** was isolated in quantitative yield. The boronic acid pinacol ester **5** was obtained in 87% yield by exposing **4** to standard *Miyaura* borylation conditions. Thus **4** was kept for 18 h at 100 °C in 1,4-dioxane with bis(pinacolato)diboron (Bpin)₂ as borane source, bis(triphenylphosphine)palladium(II) dichloride (Pd(PPh₃)₂Cl₂) as catalyst and potassium acetate (KOAc) as base. The required 4,4''-dichloro-2,2':6',2''-terpyridine building block was assembled following our already published protocol.^[16] The catalytic system reported by *Guram* and coworkers^[17] enabled the twofold *Suzuki-Miyaura* cross-coupling reaction with **5**. Using potassium carbonate (K₂CO₃) as base and bis(di-tert-butyl(4-dimethylamino)pehnyl)phosphine) dichloropalladium (II) (Pd(amphos)Cl₂) as catalyst with a toluene/water solvent mixture in a sealed tube at 116 °C, provided the desired tpy-ligand **6** in 75% isolated yield after column chromatography (CC). To liberate the aldehydes prior to Fe²⁺ coordination, two reported protocols were combined.^[18,19] The tetra-aldehyde tpy ligand **7** was obtained in 85% yield by treating **6** with pyridinium *p*-toluenesulfonate as proton source in a mixture of acetone and water at 100 °C in a microwave reactor. The homoleptic Fe(II) complex **8** was assembled by the addition of iron (II) chloride (FeCl₂) to a solution of the ligand **7** in a dichloromethane (DCM) and methanol (MeOH) solvent mixture. An immediate color change to deep purple indicated the complex formation. After solvent removal and precipitation from a NH₄PF₆ water solution, the octa-aldehyde complex **8** was isolated in 94% yield.

All new compounds were fully characterized by ¹H- and ¹³C-NMR spectroscopy, and high resolution mass spectrometry. Single crystals suitable for x-ray diffraction of the complex **8** were obtained by diffusing diethyl ether vapor into an acetonitrile solution of **8**. The solid state structure (Figure 2) not only corroborated the identity of the complex, but also gave insight into the intramolecular distances between the aldehyde groups. The inter-aldehyde O-O spacing varied between 3.60 and 7.15 Å.





Figure 2. Solid state structure of complex **8** with rotation ellipsoids at 50% probability. a) top view along a N-Fe-N axis and b) side view. Hydrogen atoms and the PF₆⁻ counter ions were omitted for clarity. Color code: N: blue, Fe: orange, C: gray, O: red.

To explore both, the feasibility of the fourfold interlinking strategy and the ideal dimension of the interlinking units, the *Schiff* base condensation was selected as test reaction. The reversibility of the imine bond should enable to investigate the thermodynamic stability and accessibility of the rigidified complex.

Initial attempts using ethylene diamine as linker (MM2 calculated N-N distance: 3.80 Å) failed, the focus moved towards aromatic diamines, hoping that the increased delocalization of the imine π-systems additionally favors the interlinking. And indeed,



Scheme 1. Synthesis of the Fe(II)tpy cage complexes **2** and **1**. Reagents and conditions: a) ethylene glycol, *p*TsOH.H₂O, toluene, reflux, 12 h, 100%; (Bpin)₂, KOAc, Pd(PPh₃)₂Cl₂, 1,4-dioxane, 100 °C, 18 h, 87%; c) K₂CO₃, Pd(amphos)Cl₂, toluene/water (5:1), 116 °C, 18 h, 75%; d) PPTS, acetone/water (5:1), microwave (MW), 80 °C, 3 h, 85%; e) 1.) FeCl₂, DCM/MeOH (1:1), RT, 1 h, 2.) NH₄PF₆, water, 12 h, 94%; f) *m*-phenylenediamine, *p*TsOH.H₂O, DCM to DCE, reflux, 10 d, n.d.; g) (1,3-phenylenebis(methylene))bis(triphenylphosphonium) dibromide, DBU, DMF, -20 °C, 12 h, 4%.

COMMUNICATION

with *meta*-phenylenediamine (MM2 calculated N-N distance: 4.5 Å) the formation of an eightfold condensation product with **8** was observed among a variety of possible condensation products by direct injection electrospray mass spectrometry (DI-ESI-MS, Figure SI18). While reaction conditions favoring the formation of the fourfold interlinked complex **2** could be found (8 equivalents of the diamine in refluxing 1,2-dichloroethane (83.5 °C) for 10 days, Figure SI19), all attempts to isolate the fourfold interlinked structure **2** failed, pointing at the lability of the imine bonds. Also numerous efforts to reduce the imine linkers to more robust amine bonds employing a variety of reducing agents were not successful.

Thus, in spite the fact that the mass signal in the DI-ESI-MS remains the only indication of the existence of the fourfold interlinked complex **2**, the experiments were indicative to identify the dimension of the ideal interlinking structure.

In order to improve the stability of the fourfold interlinked complex the strategy was to replace the reversible and hydrolysis sensitive imine-bonds of **2** by the olefins in **1**. Like that the interlinking structure would be of comparable dimension and by using *Wittig* chemistry, **1** should be accessible from the same octa-aldehyde precursor Fe(II)tpy₂ complex **8**. And indeed, treating **8** with 10 equivalents of (1,3-phenylenebis(methylene))bis(triphenylphosphonium) dibromide in -20 °C cold dimethylformamide using 1,8-diazabicyclo[5.4.0]undec-7-ene as a base resulted in a crude reaction mixture with the mass of the expected fourfold interlinked complex **1** (1330 m/z) as main signal recorded in a matrix-assisted laser desorption ionization mass spectrometer (MALDI-MS). However, even though the interlinking reaction proceeded smoothly, the purification remained challenging due to the lack of conformational selectivity in the double bonds formed by the *Wittig* reaction. With the *D*_{2h} symmetry of **1**, 64 different isomers due to *cis/trans* configurations of the eight double bonds are possible in theory. Even though not all of them are likely to be formed, the uninterpretable NMR spectrum of the crude reaction mixture points at several isomers being present. Thus the crude material was subjected to reversed-phase HPLC and after two runs, the all-*trans* isomer could be separated in 4% yield (Figure SI23). The similar MALDI-MS signals of all isolated peaks confirmed the hypothesis of various olefin isomers being formed during the interlinking *Wittig* reaction. The ¹H-NMR spectrum of the all-*trans* isomer of **1** is displayed in Figure 3. The rather simple spectrum corroborates the identity of the highly symmetric complex, with the 16.4 Hz coupling constants of both alkene protons (magenta in Fig. 3) documenting the *trans*-connection of the double bonds. The all-*trans* isomer of **1** was further characterized by ¹H-, ¹³C-, COSY-, ROESY-, HSQC- and HMBC-NMR spectroscopy, UV-Vis spectroscopy, and high-resolution electrospray ionization mass spectrometry (HR-ESI-MS, see supporting information).

Of particular interest was the rigidity gained by the fourfold interlinking in the all-*trans* isomer of **1**, which was analyzed by the comparison of the spin-spin relaxation times *T*₂ for the protons of **1** (all-*trans* isomer) and the octa-aldehyde precursor complex **8**. In these complexes of comparable molecular weight, the spin-spin relaxation correlates with the intramolecular mobility of the spin and is thus a sensitive probe for the flexibility of the spin bearing protons. Using a *Carr-Purcell-Meiboom-Gill* (CPMG) spin echo pulse sequence, *T*₂ was measured for all aryl protons (Fig. SI15). And indeed, the recorded values display for every proton a decreased *T*₂ by a factor of 1.20 to 1.75 for the all-*trans* isomer of **1** compared to **8** (table SI3), pointing at a substantial rigidification




Figure 3. ¹H-NMR of the all-*trans* isomer of **1** with assignment of the signals.

of the ligand sphere upon fourfold interlinking. The experiment even allows to localize the mobility of individual subunits in both complexes. While the protons of the central pyridine ring **A** are the most static part, the *ortho*-positioned **B** rings are more mobile in both compounds. Interestingly, the protons of the **C** and **D** rings as subunits of the linker structure of **1** (all-*trans* isomer) displayed flexibilities in between the values recorded for **A** and **B**, suggesting a tight fixation of the interlinking bridges. In contrast, the **C** ring of **8** displayed an up to three-fold *T*₂ increase, as expected for the higher degree of freedom of a rotating subunit. Another observation pointing at the rigidity of **1** is the increased uniformity of the values recorded for *T*₂ (max/min: 1.52) compared with the ones observed for **8** (max/min: 3.04). While the molecular weight increase from **8** to **1** is with 20.9 % too small to rationalize the recorded differences in *T*₂, an aggregation of **1** to oligomers might be an alternative explanation for the observation. To disqualify this hypothesis, the hydrodynamic radii of both complexes were determined by diffusion ordered NMR spectroscopy (Fig. SI16 and SI17). The observed difference of 5.2 % in diffusion coefficients corresponds to a mass increase of 16.4 % of **1** compared to **8**, corroborating that both complexes were analyzed as dissolved monomeric complexes.

Density functional theory (DFT) calculations were performed to obtain a better understanding of the electronic structure of **1** and to gauge the ring strain energy of the large ring system surrounding the [Fe(tpy)₂]²⁺ core of the complex. DFT calculations were completed for compound **1** using a B3LYP functional and a 6-31G (d,p) basis set using the Gaussian 09 suite (Figure 4). An unrestricted geometry optimization revealed that **1** optimized to retain the *D*_{2d} symmetry commonly observed for octahedral bis-2,2':6',2''-terpyridine complexes. The flexibility of the surrounding ring structure allowed for a slight bending; the direction of this deformation was identical for the ring's quarter-segments diagonally opposed to each other. The electronic structure retained some of the parent complex' features: The low energy pyridinic π* orbitals were also observed for the terpyridine moiety of **1** and covered the orbitals from LUMO to LUMO+7; at higher energies the π* orbitals of the ring structure started mixing in. The HOMO of the parent [Fe(tpy)₂]²⁺ is located on the Fe d orbitals and is the origin of the complex' well-studied MLCT transition. In contrast, compound **1**'s calculated HOMO is exclusively located on the ring's large π system and the first metal contributions do not reveal itself until reaching the 1.5 eV more stable HOMO-8 and the degenerate pair HOMO-9 and HOMO-10. The reaction

COMMUNICATION

enthalpy of an isodesmic reaction separating the peripheral ring and the metal complex moiety was estimated by DFT calculations and revealed that the overall ring strain energy for **1** is around 16 kcal/mol. The strain is distributed over 4 ring systems formed by more than 70 atoms making it negligible. Details of these calculations are documented in the Supporting Information (Fig. SI24).




Figure 4. Frontier molecular orbital diagram of **1** generated from DFT calculations (B3LYP/6-31G (d,p)). The rendered 98% surfaces of selected orbitals are also depicted. The first filled orbital with a significant metal contribution is HOMO-8, which is in stark contrast to the metal-centred HOMO observed for the parent $[\text{Fe}(\text{tpy})_2]^{2+}$ complex.

The rigidity of **1** (all-*trans* isomer) was also observed in its magnetic behavior recorded in a superconducting quantum interference device (SQUID) magnetometer. In a temperature range between 5 K and 365 K, complex **1** remained in its low spin state. While the observation supports the claim of a fixed coordination sphere in the all-*trans* isomer of **1**, it does not qualify to account for an increased rigidity by fourfold interlinking, as also the open octa-aldehyde **8** remains in its low-spin state in this temperature range.

To our delight, the structural integrity of **1** was even preserved upon deposition on a gold surface in a STM experiment. Employing an electrospray setup with in line mass selection^[20] to deposit **1** on an Au(111) surface (Fig. SI21) enabled its deposition and subsequent investigation of the sample with a STM at 4.4 K under UHV conditions. The electrospray deposition ensured the absence of counter ions on the surface. A STM constant-current topograph of a typical molecule is displayed in Figure 5a. The adsorbate most probably correspond to intact **1**, although a degree of distortion may be present. The geometry of **1** on Au(111) is most likely the one represented in the side view (Fig. 5b), exposing three dominant extrusions into vacuum (two phenyl groups of interlinking subunits and the central pyridine moiety of

a ligand). In agreement with this simple model, the topograph exhibits three protrusions with submolecular dimensions. The distance between the outermost protrusions (1.6 nm) approximately matches the phenyl to phenyl distance of **1** (see scale bars in Fig. 5 a) and b). The apparent height of the molecular adsorbates (Fig. 5a) is with about 550 pm considerably higher than the 180 pm observed for the flattened terpyridine ligands of the first generation of $\text{Fe}(\text{II})\text{tpy}_2$ complexes disintegrating upon deposition on Au(111).^[11] This observation is in agreement with a limited adsorption-induced distortion of **1**, further documenting the structural integrity gained by fourfold interlinking of the tpy ligands. The charge of **1** cannot be unambiguously clarified from the present data.




Figure 5. a) Constant-current STM topograph of **1** on Au(111). Current $I = 5$ pA and sample voltage $V = -0.5$ V. (b) Side-view model of **1**. Orange, blue, dark gray and light gray spheres represent iron, nitrogen, carbon and hydrogen atoms, respectively. (c) Time series of the tunneling current measured at $V = 0.9$ V with the tip position frozen over the center of **1**.

While spin switching of individual complexes has been evidenced on partially insulating layers,^[21–26] the switching on metal surfaces remains a challenge with a few exceptions.^[27,28] The strong molecule-substrate interaction on metal surfaces often affects the integrity and functionality of adsorbed complexes.^[11,26] The fixation of the ligands by fourfold interlinking not only increases the structural integrity of **1**, but also is expected to preserve the functionality of the $\text{Fe}(\text{II})\text{tpy}_2$ complex caged in the center of the macrocycle upon adsorption, which was not the case for less tightly fixed $\text{Fe}(\text{II})\text{tpy}_2$ model complexes (Figure SI22). And indeed, a two-level fluctuation of the tunneling current was recorded at slightly elevated sample voltages (e.g., $V = 0.9$ V in Fig. 5c). Even though the finding is consistent with structural and electronic changes expected upon spin transitions in spin crossover complexes, an unambiguous identification of the underlying switching mechanism will require further experiments, most likely in higher magnetic fields.

In summary, the tight fixation of the coordination sphere of a $\text{Fe}(\text{II})\text{tpy}_2$ complex by fourfold interlinking of the suitably decorated tpy ligands is reported. Profiting from the octa-aldehyde precursor **8**, the ideal dimensions of the interlinking

COMMUNICATION

subunits was determined by Schiff base condensation, while the caged iron complex **1** was obtained by Wittig olefination. The all-*trans* isomer of **1** was isolated by HPLC and fully characterized by NMR spectroscopy, UV-Vis spectroscopy and mass spectrometry. The rigidity of the ligand sphere was analyzed by CPMG-NMR experiments and increased the integrity of the caged complex to a level enabling to preserve its structure and functionality upon adsorption on metal surfaces. Low-temperature STM experiments evidenced preserved switching functionality of the complex.

The approach holds perspectives not only for spin-switch functionality, but also to couple the spin of the complexes to more reactive ferromagnetic substrates,^[29] of relevance for molecular spintronics. Currently, we are exploring both, the structural diversity of caged complexes as well as their peripheral functionalization.

Acknowledgements

The authors gratefully acknowledge financial support by the Swiss National Science Foundation (SNF, grant numbers: 200020-178808). This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant no. 766726. M. M. acknowledges support by the 111 project (90002-18011002). S.B. acknowledges support by the US NSF (CHE-1764353). We thank Thomas Knaak for providing some of the data shown in Fig. SI22.

Keywords: Fe terpyridine • Macrocyclic ligands • Wittig reactions • Cage compounds • Hexadentate ligand


- [1] A. R. Pease, J. O. Jeppesen, J. F. Stoddart, Y. Luo, C. P. Collier, J. R. Heath, *Acc. Chem. Res.* **2001**, *34*, 433–444.
- [2] A. S. Blum, J. G. Kushmerick, D. P. Long, C. H. Patterson, J. C. Yang, J. C. Henderson, Y. Yao, J. M. Tour, R. Shashidhar, B. R. Ratna, *Nat. Mater.* **2005**, *4*, 167–172.
- [3] S. Y. Quek, M. Kamenetska, M. L. Steigerwald, H. J. Choi, S. G. Louie, M. S. Hybertsen, J. B. Neaton, L. Venkataraman, *Nat. Nanotechnol.* **2009**, *4*, 230–234.
- [4] D. Čavlović, M. Juriček, *Chim. Int. J. Chem.* **2019**, *73*, 313–316.
- [5] G. Ke, C. Duan, F. Huang, X. Guo, *InfoMat* **2019**, inf2.12068.
- [6] J. L. Zhang, J. Q. Zhong, J. D. Lin, W. P. Hu, K. Wu, G. Q. Xu, A. T. S. Wee, W. Chen, *Chem. Soc. Rev.* **2015**, *44*, 2998–3022.
- [7] R. Frisenda, G. D. Harzmann, J. A. Celis Gil, J. M. Thijssen, M. Mayor, H. S. J. van der Zant, *Nano Lett.* **2016**, *16*, 4733–4737.
- [8] G. D. Harzmann, R. Frisenda, H. S. J. van der Zant, M. Mayor, *Angew. Chem. Int. Ed.* **2015**, *54*, 13425–13430.
- [9] T. Brandl, M. El Abbassi, D. Stefani, R. Frisenda, G. D. Harzmann, H. S. J. van der Zant, M. Mayor, *Eur. J. Org. Chem.* **2019**, *2019*, 5334–5343.
- [10] T. Brandl, V. Hoffmann, A. Pannwitz, D. Häussinger, M. Neuburger, O. Fuhr, S. Bernhard, O. S. Wenger, M. Mayor, *Chem. Sci.* **2018**, *9*, 3837–3843.
- [11] T. Knaak, C. González, Y. J. Dappe, G. D. Harzmann, T. Brandl, M. Mayor, R. Berndt, M. Gruber, *J. Phys. Chem. C* **2019**, *123*, 4178–4185.
- [12] J. E. McMurry, M. P. Fleming, *J. Am. Chem. Soc.* **1974**, *96*, 4708–4709.
- [13] M. R. Becker, R. B. Watson, C. S. Schindler, *Chem. Soc. Rev.* **2018**, *47*, 7867–7881.
- [14] H. Schiff, *Ann. Chem. Pharm.* **1864**, *131*, 118–119.
- [15] B. E. Maryanoff, A. B. Reitz, *Chem. Rev.* **1989**, *89*, 863–927.
- [16] G. D. Harzmann, M. Neuburger, M. Mayor, *Eur. J. Inorg. Chem.* **2013**, *2013*, 3334–3347.
- [17] A. S. Guram, A. O. King, J. G. Allen, X. Wang, L. B. Schenkel, J. Chan, E. E. Bunel, M. M. Faul, R. D. Larsen, M. J. Martinelli, et al., *Org. Lett.* **2006**, *8*, 1787–1789.
- [18] Y. He, M. Johansson, O. Sterner, *Synth. Commun.* **2004**, *34*, 4153–4158.
- [19] R. Sterzycki, *Synthesis* **2002**, *1979*, 724–725.
- [20] Chr. Hamann, R. Woltmann, I.-P. Hong, N. Hauptmann, S. Karan, R. Berndt, *Rev. Sci. Instrum.* **2011**, *82*, 033903.
- [21] T. G. Gopakumar, F. Matino, H. Naggert, A. Bannwarth, F. Tuczek, R. Berndt, *Angew. Chem. Int. Ed.* **2012**, *51*, 6262–6266.
- [22] T. Miyamachi, M. Gruber, V. Davesne, M. Bowen, S. Boukari, L. Joly, F. Scheurer, G. Rogez, T. K. Yamada, P. Ohresser, et al., *Nat. Commun.* **2012**, *3*, 938.
- [23] M. Bernien, H. Naggert, L. M. Arruda, L. Kipgen, F. Nickel, J. Miguel, C. F. Hermanns, A. Krüger, D. Krüger, E. Schierle, et al., *ACS Nano* **2015**, *9*, 8960–8966.
- [24] T. Jasper-Toennies, M. Gruber, S. Karan, H. Jacob, F. Tuczek, R. Berndt, *Nano Lett.* **2017**, *17*, 6613–6619.
- [25] L. Kipgen, M. Bernien, S. Ossinger, F. Nickel, A. J. Britton, L. M. Arruda, H. Naggert, C. Luo, C. Lotze, H. Ryll, et al., *Nat. Commun.* **2018**, *9*, 2984.
- [26] S. Rohlf, J. Grunwald, T. Jasper-Toennies, S. Johannsen, F. Diekmann, M. Studniarek, R. Berndt, F. Tuczek, K. Rosnagel, M. Gruber, *J. Phys. Chem. C* **2019**, *123*, 17774–17780.
- [27] K. Bairagi, O. Iasco, A. Bellec, A. Kartsev, D. Li, J. Lagoute, C. Chacon, Y. Girard, S. Rousset, F. Miserque, et al., *Nat. Commun.* **2016**, *7*, 12212.
- [28] A. Köbke, F. Gutzeit, F. Röhricht, A. Schlimm, J. Grunwald, F. Tuczek, M. Studniarek, D. Longo, F. Choueikani, E. Otero, et al., *Nat. Nanotechnol.* **2020**, *15*, 18–21.
- [29] S. Gueddida, M. Gruber, T. Miyamachi, E. Beaupaire, W. Wulfhekel, M. Alouani, *J. Phys. Chem. Lett.* **2016**, *7*, 900–904.

COMMUNICATION

Entry for the Table of Contents

COMMUNICATION

Caging an Fe(II) ion: Like a cube the fourfold interlinked bisterpyridine ligand surrounds the Fe(II) ion. The post-coordinative macrocyclization rigidifies the ligand sphere and guaranties the preservation of the complex's properties.



Thomas Brandl, Sven Johannsen, Daniel Häussinger, Nithin Suryadevara, Alessandro Prescimone, Stefan Bernhard, Manuel Gruber, Mario Ruben, Richard Berndt and Marcel Mayor

Page No. – Page No.

Iron in a cage: fixation of a Fe(II)tpy₂ complex by fourfold interlinking