# The Kinetics of the Reaction of $[Ru(CI)H(PPh_3)_3]$ with Various Olefins

Giuseppe Alibrandi and Brian E. Mann'

Department of Chemistry, The University of Sheffield, Sheffield S3 7HF, UK

A detailed kinetic investigation of the reaction of  $[Ru(CI)H(PPh_3)_3]$  with cycloheptatriene, cyclohepta-1,3-diene, cyclooctatetraene, penta-1,4-diene, cycloocta-1,5-diene and dimethyl maleate, has been carried out spectrophotometrically in  $CH_2CI_2$  at 10 °C. It is shown that the major mechanism is *via* dissociation of PPh<sub>3</sub> to give  $[Ru(CI)H(PPh_3)_2]$  which then reacts with the olefin. There is also a second mechanism involving direct attack of the olefin on  $[Ru(CI)H(PPh_3)_3]$ .

The complex [Ru(Cl)H(PPh<sub>3</sub>)<sub>3</sub>] is a very valuable hydrogenation catalyst, but its mechanism of action is far from being understood.<sup>1</sup> It is a very difficult system to study due to its great reactivity resulting in such rapid hydrogen consumption that hydrogen starvation can occur in the solution.<sup>2</sup> This is in contrast with [RhCl(PPh<sub>3</sub>)<sub>3</sub>] which has been thoroughly investigated.<sup>3</sup> In the original paper by Wilkinson and coworkers<sup>4</sup> describing the preparation and initial kinetic investigations of the catalytic activity of [Ru(Cl)H(PPh<sub>3</sub>)<sub>3</sub>], it was tentatively concluded that the active species is [Ru(Cl)H(PPh<sub>3</sub>)<sub>2</sub>], but this was, in part, by analogy with the [RhH(CO)(PPh<sub>3</sub>)<sub>3</sub>] system and by showing inhibition by PPh<sub>3</sub>.<sup>4</sup> Support for this suggestion came from the isolation of  $[Ru(Cl)H(PPh_3)_2]$  by treating  $[Ru(C_6H_4PPh_2)(PPh_3)Cl]$ with hydrogen<sup>5</sup> and  $[RuBr_3(PPh_3)_2]$  reacts with H<sub>2</sub> to give  $[Ru(Br)H(PPh_3)_2]$ .<sup>6</sup> More recently, it has been established that  $[Ru(Cl)H(PPh_3)_3]$  undergoes dissociative PPh<sub>3</sub> exchange. The complex  $[{Ru(Cl)H(PPh_3)_2}_2]$  is an effective hydrogenation catalyst in dimethylacetamide, and the kinetics for its action as a hydrogenation catalyst for acrylamide were determined.<sup>8</sup> The mechanism involves a [Ru(Cl)H(PPh<sub>3</sub>)<sub>2</sub>] species. Hydrogenation catalysed by [RuCl<sub>2</sub>(PPh<sub>3</sub>)<sub>3</sub>] has also been studied.<sup>9</sup> It was found that the mechanism involves first the formation of [Ru(Cl)H(PPh<sub>3</sub>)<sub>3</sub>] which is the active catalyst. Inhibition by PPh<sub>3</sub> is found and attributed to the formation of  $[Ru(Cl)H(PPh_3)_2].$ 

The complex  $[RuH_4(PPh_3)_3]$  is also a very effective hydrogenation catalyst and its mechanism for the hydrogenation of cyclohexanone has been thoroughly investigated.<sup>10</sup> The mechanism was shown to involve the initial loss of H<sub>2</sub> to give  $[RuH_2(PPh_3)_3]$ , which rapidly co-ordinates the cyclohexanone to give  $[RuH_2(PPh_3)_3(C_6H_{10}O)]$ . This complex reacts with H<sub>2</sub> to give  $[RuH_2(PPh_3)_3(C_6H_{11}OH)]$ , and finally the cyclohexanol is displaced by H<sub>2</sub>, to give  $[RuH_4(PPh_3)_3]$  and cyclohexanol. Although  $[Ru(Cl)H(PPh_3)_3]$  could, in the presence of H<sub>2</sub>, yield  $[RuH_4(PPh_3)_3]$  which would then be the catalyst, this is unlikely, and there is no evidence for this reaction. It is more probable that the hydrogenation catalysis by  $[Ru(Cl)H(PPh_3)_3]$  involves species which still have chloride co-ordinated to the ruthenium.

#### **Results and Discussion**

The Kinetics of the Reaction of  $[Ru(Cl)H(PPh_3)_3]$  and Cycloheptatriene.—Previous work had shown that the reaction of  $[Ru(Cl)H(PPh_3)_3]$  with cycloheptatriene proceeds smoothly to give  $[Ru(\eta^5-C_7H_9)Cl(PPh_3)_2]$ .<sup>11a</sup> The reaction is easy to follow by electronic spectroscopy as the starting material,  $[Ru(Cl)H(PPh_3)_3]$ , is purple, and the product,  $[Ru(\eta^5-C_7H_9)Cl(PPh_3)_3]$   $C_7H_9$ )Cl(PPh\_3)<sub>2</sub>], is yellow-brown. The reaction proceeds at a convenient rate at 10 °C, and was therefore studied at this temperature. The reaction was always carried out with at least a ten-fold excess of cycloheptatriene over [Ru(Cl)H(PPh\_3)\_3], so that pseudo-first-order kinetics could be used. Triphenyl-phosphine is liberated during the reaction, but is at very low concentration, as the concentration of [Ru(Cl)H(PPh\_3)\_3] is only *ca*. 10<sup>-5</sup> mol dm<sup>-3</sup>. Examination of the PPh<sub>3</sub> dependence of the rate constant shows that at this concentration the effect on the rate constant is negligible, see below. The rate constants obtained spectrophotometrically for various concentrations of cycloheptatriene and PPh<sub>3</sub> are collected in Table 1.

Examination of the data in Table 1 shows that the reaction rate is enhanced linearly, but only slightly by the addition of cycloheptatriene, and there is a non-zero intercept, see Fig. 1. The reaction rate is inhibited by the addition of  $PPh_3$ , see Fig. 2.

These data are consistent with the mechanism given in Scheme 1. It is necessary to propose two different ways for the reaction to proceed. The dominant mechanism is the initial loss of PPh<sub>3</sub> to give [Ru(Cl)H(PPh<sub>3</sub>)<sub>2</sub>], which then can either react with PPh<sub>3</sub> to give [Ru(Cl)H(PPh<sub>3</sub>)<sub>2</sub>] or with cycloheptatriene to give [Ru( $\eta^5$ -C<sub>7</sub>H<sub>9</sub>)Cl(PPh<sub>3</sub>)<sub>2</sub>]. The pre-dissociation of PPh<sub>3</sub> has been suggested previously <sup>4</sup> and then proven by magnetization transfer measurements.<sup>7</sup> The rate of this mechanism is set by the rate of PPh<sub>3</sub> dissociation to give [Ru(Cl)H(PPh<sub>3</sub>)<sub>2</sub>], and is independent of the concentration of C<sub>7</sub>H<sub>8</sub>, but is slowed by the addition of PPh<sub>3</sub>. The second slower mechanism is the direct attack of cycloheptatriene on [Ru(Cl)H(PPh<sub>3</sub>)<sub>3</sub>] to give

Table 1 Rate constants for the reaction of  $[Ru(Cl)H(PPh_3)_3]$  with cycloheptatriene in  $CH_2Cl_2$  at 283.2 K

$[C_7H_8]/mol  dm^{-3}$	$[PPh_3]/mol dm^{-3}$	$10^4 k_{ m obs}/{ m s}^{-1}$
0.0289	0.0	4.09
0.0482	0.0	4.15
0.0964	0.0	4.38
0.241	0.0	4.79
0.578	0.0	5.60
0.771	0.0	6.06
0.0482	0.003	3.40
0.0482	0.01	2.68
0.0482	0.02	1.92
0.0482	0.035	1.58
0.0482	0.05	1.30
0.0482	0.075	1.05
0.241	0.01	4.22
0.241	0.05	3.03
0.241	0.10	2.31



**Fig. 1** A plot of the rate of the reaction of  $[Ru(Cl)H(PPh_3)_3]$  with various olefins, as a function of  $C_7H_8$  concentration, in the absence of added PPh\_3 in CH\_2Cl\_2 at 283.2 K; cyclohepta-1,3-diene ( $\blacklozenge$ ), cycloocta-1,5-diene ( $\diamondsuit$ ), penta-1,4-diene ( $\blacksquare$ ), cyclooctatetraene ( $\Box$ ), cycloheptatriene ( $\bigcirc$ ), dimethyl maleate ( $\blacklozenge$ )



**Fig. 2** A plot of the rate of the reaction of  $[Ru(Cl)H(PPh_3)_3]$  with  $C_7H_8$ , as a function of PPh<sub>3</sub> concentration in  $CH_2Cl_2$  at 283.2 K;  $[C_7H_8] = 0.0482$  ( $\bigcirc$ ) or 0.241 mol dm<sup>-3</sup> ( $\triangle$ ). The fitted lines were calculated using equation (1) with  $k_1 = 4.05$  ( $\pm 0.06$ ) × 10<sup>-4</sup> s<sup>-1</sup>,  $k_{-1}/k_3 = 2.9$  ( $\pm 0.2$ ),  $k_2 = 2.61$  ( $\pm 0.16$ ) × 10<sup>-4</sup> dm<sup>3</sup> mol<sup>-1</sup> s<sup>-1</sup>

 $[Ru(\eta^5-C_7H_9)Cl(PPh_3)_2]$  and depends on the concentration of  $C_7H_8$ .

Analysis of the mechanism in Scheme 1 gives the rate law given in equation (1).

$$k_{\rm obs} = \frac{k_1 [C_7 H_8]}{(k_{-1}/k_3) [PPh_3] + [C_7 H_8]} + k_2 [C_7 H_8] \quad (1)$$

Without PPh<sub>3</sub> added to the solution, the term  $(k_{-1}/k_3)$ [PPh<sub>3</sub>] can be approximated to zero and the rate-law is reduced to equation (2) as illustrated in Fig. 1. In the presence of PPh<sub>3</sub>, all

$$k_{\rm obs} = k_1 + k_2 [C_7 H_8] \tag{2}$$

terms operate. The values of  $k_{obs}$ , [PPh<sub>3</sub>], and [C<sub>7</sub>H<sub>8</sub>] were fitted to equation (1) using a non-linear, least-squares curve fitting program, and a good fit was obtained yielding the kinetic constants:  $k_1 = 4.05 (\pm 0.06) \times 10^{-4} \text{ s}^{-1}$ ,  $k_{-1}/k_3 = 2.9 (\pm 0.2)$ ,  $k_2 = 2.61 (\pm 0.16) \times 10^{-4} \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ , see Fig. 2.

In the absence of added PPh<sub>3</sub>, the concentration of liberated PPh<sub>3</sub> is negligible, and equation (2) applies. Using this analysis, the rates in Table 2 were obtained, which for cycloheptatriene are not significantly different from those obtained using the full equation (1).

A further check on the validity of this mechanism comes from repeating the measurements with other dienes and trienes. It is predicted that  $k_1$  should be independent of the olefin chosen.

**Table 2** The values for  $k_1$  and  $k_2$  derived using equation (2) for a variety of olefins  $[C_7H_8 = cycloheptatriene, C_7H_{10} = cyclohepta-1,3-diene, C_8H_8 = cyclooctatetraene, 1,4-C_5H_8 = penta-1,4-diene, 1,5-C_8H_{12} = cycloocta-1,5-diene, C_2H_2(CO_2Me)_2 = dimethyl maleate]$ 

Olefin	$10^4 k_1 / \mathrm{s}^{-1}$	$10^4 k_2 / s^{-1}$
$C_7H_8$	4.08	2.61
$C_7 H_{10}$	3.91	0.00
$C_2H_2(CO_2Me)_2$	6.20	7.38
$1,5-C_8H_{12}$	3.85	0.30
$1,4-C_5H_8$	4.14	0.31
C <sub>8</sub> H <sub>8</sub>	4.07	1.03



Scheme 1 The proposed mechanism for the reaction of  $[Ru(Cl)-H(PPh_3)_3]$  with cycloheptatriene



Scheme 2 The reaction of  $[Ru(Cl)H(PPh_3)_3]$  with cyclohepta-1,3diene

The Kinetics of the Reaction of  $[Ru(Cl)H(PPh_3)_3]$  and Cyclohepta-1,3-diene.—Previous work has shown that this is not a single reaction, but consists of two steps.<sup>11a</sup> The initial reaction gives the expected product,  $[Ru(\eta^3-C_7H_9)Cl(PPh_3)_2]$ , but there is then a hydrogen-transfer reaction between free cyclohepta-1,3-diene and  $[Ru(\eta^3-C_7H_{11})Cl(PPh_3)_2]$  to yield  $[Ru(\eta^5-C_7H_9)Cl(PPh_3)_2]$  and presumably cycloheptene, see Scheme 2.\*

The second reaction proceeds at a rate comparable with the first, which made the full analysis of the rate data from electronic spectroscopy subject to error. However reasonable rates could be obtained from the initial gradient, see Tables 2 and 3.

The Kinetics of the Reaction of  $[Ru(Cl)H(PPh_3)_3]$  and Cyclooctatetraene.—The reaction between  $[Ru(Cl)H(PPh_3)_3]$  and cyclooctatetraene has been shown to yield a mixture of three products, see Scheme 3.<sup>12</sup>

The reaction proceeds cleanly as a single step reaction, yielding the rate data in Table 2.

The Kinetics of the Reaction of  $[Ru(Cl)H(PPh_3)_3]$  and Penta-1,4-diene.—The reaction between  $[Ru(Cl)H(PPh_3)_3]$  and penta-1,4-diene has been shown to involve sequential reactions, yielding initially  $[Ru(\eta^3-C_5H_9)Cl(PPh_3)_2]$ , then  $[Ru(\eta^5-C_5H_9)Cl(PPh_3)_2]$ 

<sup>\*</sup> Note added at proof: We now believe that  $H_2$  is liberated. Only traces of cycloheptene are detected and  $H_2$  reverses the reaction.<sup>11b</sup>



Scheme 3 The reaction of [Ru(Cl)H(PPh<sub>3</sub>)<sub>3</sub>] with cyclooctatetraene



Scheme 4 The reaction of [Ru(Cl)H(PPh<sub>3</sub>)<sub>3</sub>] with penta-1,4-diene

 $C_5H_7)Cl(PPh_3)_2]$  and finally  $[Ru(\eta^5\text{-}C_5H_5)Cl(PPh_3)_2],$  see Scheme 4.13.\*

The kinetics of the consumption of  $[Ru(Cl)H(PPh_3)_3]$  were determined, see Fig. 1 and Tables 2 and 3.

The Kinetics of the Reaction of  $[Ru(Cl)H(PPh_3)_3]$  and Cycloocta-1,5-diene.—The reaction does not produce olefin insertion into the Ru–H bond, but stops at the diene complex, see Scheme 5.<sup>4,14</sup>

This reaction is analogous to that previously reported between  $[Ru(Cl)H(PPh_3)_3]$  and bicyclo[2.2.1]hepta-2,5-diene, which yields I as the product.<sup>4,15</sup> In view of the similarity of the kinetics with those observed for the other dienes, cycloheptatriene, and cyclooctatetraene, it is probable that the first stages of the reaction are analogous to those given in Scheme 1, but the reaction stops without insertion of the double bond into the Ru–H bond.

The kinetics of the consumption of  $[Ru(Cl)H(PPh_3)_3]$  were determined, see Fig. 1 and Tables 2 and 3.

The Kinetics of the Reaction of  $[Ru(Cl)H(PPh_3)_3]$  and Dimethyl Maleate.—Previous investigations had shown that  $[Ru(Cl)H(PPh_3)_3]$  reacts with dimethyl maleate to give  $[Ru{CH(CO_2Me)CH_2CO_2Me}Cl(PPh_3)_2]$  and PPh<sub>3</sub>. Only a single ruthenium containing product was detected by <sup>1</sup>H and <sup>31</sup>P NMR spectroscopy, but the stereochemistry was not established.<sup>16</sup>

The kinetics of the consumption of  $[Ru(Cl)H(PPh_3)_3]$  were determined, see Fig. 1 and Tables 2 and 3.

Comparison of the Rate Data.—Previous work has shown the inhibition of the catalytic hydrogenation using [Ru(Cl)H-(PPh<sub>3</sub>)<sub>3</sub>] by PPh<sub>3</sub> and proposed the formation of [Ru(Cl)H-(PPh<sub>3</sub>)<sub>2</sub>] as a key step of the mechanism.<sup>4</sup> The rate data presented here confirm the suggestion that the dissociation of PPh<sub>3</sub> is the rate-determining step. Examination of the rate data in Table 2 shows that  $k_1$  is the same for cycloheptatriene, cyclohepta-1,3-diene, cyclooctatetraene, cyclopenta-1,4-diene and cycloocta-1,5-diene, as is required for the dissociation of PPh<sub>3</sub> from [Ru(Cl)H(PPh<sub>3</sub>)<sub>3</sub>]. Only when dimethyl maleate is used,  $k_1$  is somewhat increased, and this is attributed to a pre-coordination of the ester group to the ruthenium before PPh<sub>3</sub> 953

**Table 3** Rate constants for the reaction of  $[Ru(Cl)H(PPh_3)_3]$  with various olefins in CH<sub>2</sub>Cl<sub>2</sub> at 283.2 K; olefins as in Table 2

Olefin	[Olefin]/ mol dm <sup>-3</sup>	$10^4 k_{obs}/s^{-1}$
$C_7H_{10}$	0.046	3.92
- /10	0.138	3.88
	0.230	3.90
	0.346	3.82
	0.461	4.21
	0.599	3.72
	0.737	3.98
	0.922	3.88
$C_8H_8$	0.0444	4.15
0 0	0.133	4.10
	0.222	4.43
	0.444	4.42
	0.666	4.80
$1,4-C_5H_8$	0.0483	4.28
	0.242	4.05
	0.725	4.42
$1,5-C_8H_{12}$	0.122	3.83
	0.305	4.03
	0.610	4.00
$C_2H_2(CO_2Me)_2$	0.0399	5.98
	0.120	7.53
	0.199	7.78
	0.299	8.58
	0.399	9.18
	0.519	9.95
	0.638	10.63
	0.798	12.22





Scheme 5 The reaction of  $[Ru(Cl)H(PPh_3)_3]$  with cycloocta-1,5-diene



dissociation with a large enough formation constant so that under the reaction conditions, the complex is in effect fully formed. Once  $[Ru(Cl)H(PPh_3)_2]$  is formed it has the choice of either reacting with PPh<sub>3</sub> to give back the starting material, and hence the inhibition by PPh<sub>3</sub>, or reacting with the olefin to give the product(s).

There is also a minor direct reaction of the olefin with  $[Ru(Cl)H(PPh_3)_3]$ . The reaction with the olefin is very dependent on the olefin and presumably depends on the ability of the olefin to co-ordinate to the ruthenium and then to insert into the Ru–H bond. Examination of the data for  $k_2$  in Table 2 shows that the rate decreases in the order dimethyl maleate > cycloheptatriene > cyclooctatetraene > cycloocta-1,5-diene  $\approx$  penta-1,4-diene > cyclohepta-1,3-diene.

The parallel between these observations and the mechanism proven for the hydrogenation of cyclohexene catalysed by  $[RhCl(PPh_3)_3]$  is noteworthy.<sup>3</sup> In both cases the major pathway involves fourteen-electron intermediates, namely  $[Ru(Cl)H(PPh_3)_2]$  and  $[RhCl(PPh_3)_2]$ , but a slower pathway also exists involving species containing three PPh<sub>3</sub> ligands.

### Experimental

The complex  $[Ru(Cl)H(PPh_3)_3]$  was prepared by the literature method.<sup>17</sup> Dichloromethane was freshly distilled over CaH<sub>2</sub>.

<sup>\*</sup> Note added at proof: We now believe that the formation of  $[Ru(\eta^{5}-C_{5}H_{7})Cl(PPh_{3})_{2}]$  and  $[Ru(\eta^{5}-C_{5}H_{5})Cl(PPh_{3})_{2}]$  involves the loss of  $H_{2}$ .<sup>11b</sup>

The kinetics were followed spectrophotometrically using a Perkin Elmer Lambda 5 spectrophotometer by repetitive scanning in the range 350 to 650 nm. The reactions were carried out in a silica cell sealed with a subaseal in the thermostatted cell compartment with a temperature accuracy of  $\pm 0.05$  °C. The use of at least a 10-fold excess of olefin over complex always ensured pseudo-first order conditions. The rate constants were obtained from a non-linear least-squares fit of the experimental  $A_t$ , the absorbance at time t, versus time data to the expression  $A_t = A_{\infty} + (A_0 - A_{\infty})\exp(-k_{obs}t)$ , with  $A_0$ , the absorbance at time t = 0,  $A_{\infty}$  and  $k_{obs}$  as the parameters to be optimized.

## Acknowledgements

We wish to thank the Consiglio Nationale, Italy for financial support of G. A.

## References

- B. R. James, *Homogeneous hydrogenation*, Wiley, New York, 1973,
   p. 83; *Adv. Organomet. Chem.*, 1979, 17, 319; F. H. Jardine, *Prog. Inorg. Chem.*, 1983, 31, 265.
- 2 F. H. Jardine, Prog. Inorg. Chem., 1983, 31, 327.
- 3 J. Halpern, in Organotransition Metal Chemistry, eds. Y. Ishida and M. Tsusui, Plenum, New York, 1975, p. 109.

- 4 P. S. Hallman, B. R. McGarvey and G. Wilkinson, J. Chem. Soc. A, 1968, 3143.
- 5 B. R. James, L. D. Markham and D. K. W. Wang, J. Chem. Soc., Chem. Commun., 1974, 439.
- 6 B. R. James, A. D. Rattray and D. K. W. Wang, J. Chem. Soc., Chem. Commun., 1976, 792.
- 7 Ref. 27 of A. M. Stolzenberg and E. L. Muetterties, Organometallics, 1985, 4, 1739.
- 8 B. R. James and D. K. W. Wang, J. Chem. Soc., Chem. Commun., 1977, 5500. 9 S. R. Patil, D. N. San and P. V. Chaudhari, J. M. J. C. J. 1000, 10
- 9 S. R. Patil, D. N. Sen and R. V. Chaudhari, *J. Mol. Catal.*, 1983, **19**, 233; 1984, **23**, 51.
- 10 D. E. Linn, jun. and J. Halpern, J. Am. Chem. Soc., 1987, 109, 2969.
- 11 (a) M. Grassi, B. E. Mann, P. Manning and C. M. Spencer, J. Organomet. Chem., 1986, 307, C55; (b) B. E. Mann and G. A. Sun, unpublished results.
- 12 G. Alibrandi and B. E. Mann, J. Chem. Soc., Dalton Trans., 1992, 1439.
- 13 B. E. Mann, P. W. Manning and C. M. Spencer, J. Organomet. Chem., 1986, 312, C64.
- 14 G. Alibrandi, B. E. Mann and P. Manning, unpublished work.
- 15 T. W. Dekleva, A. M. Joshi, I. S. Thorburn, B. R. James, S. V. James, S. V. Evans and J. Trotter, *Isr. J. Chem.*, 1990, **30**, 343.
- 16 B. E. Mann and P. Manning, unpublished work.
- 17 R. A. Schunn and E. R. Wonchoba, Inorg. Synth., 1972, 13, 131.

Received 15th October 1993; Paper 3/06185J