



One pot synthesis of aryl substituted aurones

Céu M. Sousa^{a,*}, Jérôme Berthet^b, Stephanie Delbaere^b, Paulo J. Coelho^a

^a Centro de Química – Vila Real, Universidade de Trás-os-Montes e Alto Douro, Quinta dos Prados, 5001-801 Vila Real, Portugal

^b Université Lille Nord de France, CNRS UMR 8516, UDSL, Faculté des Sciences Pharmaceutiques et Biologiques, F-59006 Lille, France

ARTICLE INFO

Article history:

Received 28 February 2011

Received in revised form

27 May 2011

Accepted 31 May 2011

Available online 6 June 2011

Keywords:

Aurone

Benzofuranone

Dyes

2',6'-Dihydroxyacetophenone

Benzophenone

9-Fluorenone


ABSTRACT

An efficient one-pot synthesis of new 4-hydroxy-2-(diarylmethylene)benzofuran-3(2H)-ones dyes (aryl substituted aurones) from 2',6'-dihydroxyacetophenone, potassium *tert*-butoxide and aromatic ketones under thermal conditions is described.

© 2011 Elsevier Ltd. All rights reserved.

1. Introduction

Aurones (2-benzylidenebenzofuran-3(2H)-ones) are natural occurring yellow dyes found especially in plants and in some marine organisms, usually in a hydroxylated form. Both *E* and *Z* isomers can be found in nature, although the latter is thermodynamically more stable and thus much more abundant. Besides their contribution to the pigmentation of flowers and fruits [1] aurones exhibit a wide variety of biological activities: they have been described as antifungal agents [2–4], antibacterial agents [5–9], insect antifeedant agents [10], antioxidants [11–14], as inhibitors of tyrosinase [15], iodothyronine-deiodinase [16], acetylcholinesterase [17], anticancer [18–20] and as possessing anti-inflammatory properties [10]. These compounds are usually prepared by three main methods: condensation between benzofuranones and benzaldehydes [10,15,17,18,21,22], oxidative cyclisation of 2'-hydroxychalcones [11] or by catalysed cyclisation of *ortho*-(1-hydroxyprop-2-ynyl)phenols or *ortho*-hydroxyaryl arylethynyl ketones [23–25]. In this paper we describe a fast and easy one-pot synthesis of new aryl substituted aurones starting from commercially available materials under thermal conditions.




2. Results and discussion


2'-Hydroxychalcones can be easily prepared, under mild conditions, by aldol condensation between 2'-hydroxyacetophenone and different benzaldehydes in the presence of base. This reaction is very general and these compounds are valuable intermediates in the synthesis of flavones [26–28] and aurones [11] (Scheme 1).

The reaction of 2'-hydroxyacetophenone with benzophenone in the presence of *t*-BuOK in toluene at reflux, gives the expected aryl substituted 2'-hydroxychalcone an useful compound that can be converted in acid medium to the 2,2-diphenylchroman-1-one, an important precursor for the synthesis photochromic chromenes [29,30]. In an attempt to prepare 2',6'-dihydroxy-3-phenylchalcone we tried the reaction of 2',6'-dihydroxyacetophenone with benzophenone in the presence of *t*-BuOK and using toluene as solvent, but no reaction was detected after 8 h at reflux. Changing the solvent to xylene led to the same result while when the reaction was performed in high boiling point solvents like DMSO, ethyleneglycol or nitrobenzene only degradation products were observed (Scheme 2).

* Corresponding author. Tel.: +351 259 350284; fax: +351 259 350480.
E-mail address: pcoelho@utad.pt (C.M. Sousa).



Scheme 1. Synthesis of flavones and aurones from 2'-hydroxychalcones.




Scheme 2. Reaction of 2'-hydroxyacetophenone and 2',6'-dihydroxyacetophenone with benzophenone.

However, heating a mixture of 2',6'-dihydroxyacetophenone, potassium *tert*-butoxide and benzophenone in the presence of a very small amount of DMSO at 200 °C for 2 min affords a red mass that upon acid hydrolysis originated a deep yellow dye.

The ^1H NMR spectrum of this compound showed the expected aromatic signals but the absence of any signal that could be attributed to ethylenic protons, and the presence of only one phenolic proton pointed to a different structure. Mass spectrometry confirmed the molecular formula of the product as $\text{C}_{21}\text{H}_{14}\text{O}_3$ which indicates a difference of two hydrogen atoms from the predicted chalcone structure.

The structure of this new yellow dye was unambiguously established using 2D NMR experiments as being an aurone derivative (Scheme 3). ^1H – ^1H scalar correlations were measured in COSY experiment (Fig. 1) from H-5 at 6.60 ppm up to H-7 at 6.76 ppm via H-6 at 7.51 ppm and for the two phenyl groups between H-3'/7' at 7.60 ppm and H-4'/6' at 7.44 ppm, and between H-9'/13' at 7.37 ppm and H-10'/12' at 7.50 ppm.

The HMBC spectrum (Fig. 2) evidences long-range C–H correlations between carbon C-4 at 156.9 ppm and aromatic protons H-5, H-6 and H-7, and the phenolic function at 8.06 ppm. The furanone quaternary carbons C-7a and C-3 were evidenced at 163.8 ppm and 184.8 ppm, respectively. Carbon C-7a is long-range correlated with H-5, H-6 and H-7, while carbon C-3 is correlated with H-5 and H-7. The carbon C-1' at 133.0 ppm is long-range correlated with H-3'/7' and H-9'/13'.



Scheme 3. Synthesis of 4-hydroxy-2-(diphenylmethylene)benzofuran-3(2H)-one dye **1a**.

The dye **1a** was obtained in 10% yield after column chromatography. Heating the reaction mixture for more than 2 min, or at higher temperature leads to more degradation while heating at lower temperature, leaves too much unreactive material. The reaction also works using *t*-butanol or DMF as solvent but with lower yields (7 and 8%, respectively) and the amount of solvent must be very low. If several mL are used only degradation products are detected. The best results were obtained using 60 mg of DMSO for 0.5 mmol of 2',6'-dihydroxyacetophenone. Heating the reagents in an inert atmosphere (N_2) gave the same result.




Fig. 1. ^1H – ^1H COSY of **1a** in CDCl_3 .





Fig. 2. ^1H – ^{13}C HMBC of **1a** in CDCl_3 .

The mechanism of this reaction may involve the intermediate formation of the chalcone by aldol condensation followed by intramolecular nucleophilic addition to the conjugated double bond leading to a carbanion stabilised by the two aromatic substituents, and finally dehydrogenation under thermal conditions leading to the conjugated aurone **1a** (Scheme 4).

Starting from fluorenone or 2-hydroxyfluorenone we obtained new 4-hydroxy-2-(diarylmethylene)benzofuran-3(2H)-ones dyes (aryl substituted aurones) **1b** and **1c** in 9 and 8% yield, respectively (Table 1). Although the yields of this transformation are rather low (8–10%), this reaction gives rise in one-pot to rather complex molecules that are not easy to prepare by other methods. To our knowledge there is only one report concerning the synthesis of 2-(diarylmethylene)-3-benzofuranones, in 4 steps from commercially available compounds [24].

The complete assignments of all proton and carbon resonances for dyes **1a–c** is reported in Table 2. For aurone **1c** the higher chemical shift of proton H-3' (8.89 ppm), relative to proton H-13' (8.36 ppm), suggests the position of the former near the carbonyl group. To unambiguously establish the configuration of the double bond of this compound, a 1D-NOESY experiment was performed (Fig. 3). Since a strong dipolar correlation between H-13' at 8.36 ppm and H-7 at 6.98 ppm and H-12' at 7.15 ppm was observed, proton H-13' must be on the same side of H-7 and therefore the double bond has an *E* configuration.

These dyes show a broad absorption band in the UV–Vis spectrum with a maximum between 398–435 nm and molar absorption coefficients (ϵ) of 0.9×10^4 to $1.5 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$ in CH_2Cl_2 .



Scheme 4. Proposed mechanism for the formation of 4-hydroxy-2-(diphenylmethylene)benzofuran-3(2H)-one dye **1a**.

Table 1
Synthesis of 4-hydroxy-2-(diarylmethylene)benzofuran-3(2H)-ones dyes.

Ketone	Product	Yield (%)	λ_{max} (nm) (ϵ)	λ_{max} in basic medium (nm)
	 1a	10	398 (9×10^3)	462
	 1b	9	435 (1.5×10^4)	519
	 1c	8	433 (4.6×10^4)	500

Table 2
NMR spectral data for dyes **1a–c**.

	Dye 1a (CDCl ₃)		1b in CDCl ₃		1c in DMSO-d ₆	
	¹ H	¹³ C	¹ H	¹³ C	¹ H	¹³ C
2	—	143.6	—	146.3	—	147.0
3	—	184.8	—	186.3	—	182.2
3a	—	110.8	—	110.4	—	110.6
4	—	156.9	—	157.3	—	157.9
5	6.60	109.3	6.69	109.9	6.69	111.1
6	7.51	139.0	7.59	139.4	7.61	139.3
7	6.76	103.2	6.88	103.3	6.98	102.8
7a	—	163.8	—	163.7	—	165.0
1'	—	133.0	—	129.4	—	125.6
2'	—	137.1	—	135.3	—	136.5
3'	7.60	131.4	9.33	128.0	8.89	115.5
4'	7.44	128.3	7.37	128.0	—	158.5
5'	7.44	129.6	7.44	130.8	6.86	117.5
6'	7.44	128.3	7.68	119.9	7.65	121.5
7'	7.60	131.4	—	141.3	—	132.5
8'	—	135.4	—	137.0	—	136.7
9'	7.37	130.5	—	142.2	—	142.0
10'	7.50	128.3	7.68	119.9	7.69	119.6
11'	7.50	129.1	7.44	130.8	7.41	130.9
12'	7.50	128.3	7.37	128.0	7.29	127.0
13'	7.37	130.5	8.46	129.5	8.36	129.0
OH	8.06	—	8.32	—	11.26 (C-4) 9.71 (C-4')	—
J in Hz						
3'–4'	7.8		7.6			
4'–5'			7.4			
5'–6'			7.2		8.6	
6'–7'	7.8					
9'–10'	7.8					
10'–11'			7.2		7.4	
11'–12'			7.4		7.7	
12'–13'	7.8		8.1		7.8	
5–6	8.0		8.3		8.2	
6–7	8.0		8.2		8.2	
3'–5'					2.2	




Fig. 4. UV/Vis spectra of compound **1b** (4.3×10^{-5} M, CH₂Cl₂) before and after the addition of base (Bu₄NOH).

which are in the range of those reported in the literature for aurones [31]. Since these aurones have phenolic groups their absorption spectra is pH dependent. The addition of base (Bu₄NOH) leads, as expected, to a strong bathochromic shift of the λ_{max} (64–84 nm) due to the formation of the phenolate anion (Fig. 4).

3. Experimental

The reactions were monitored by thin-layer chromatography on aluminum plates precoated with Merck silica gel 60 F254 (0.25 mm). Melting point were determined in capillary tubes and are uncorrected. The new compounds were determined to be >95% pure by ¹H NMR spectroscopy. UV–Vis spectra were recorded on a CARY 50 Varian spectrophotometer. IR spectra were obtained on a Perkin–Elmer FTIR 1600 spectrometer using KBr disks (wave-numbers in cm^{−1}). Electronic impact mass spectra were measured




Fig. 3. a) ¹H NMR spectrum of **1c** and b) 1D NOESY spectrum with an irradiation of H-13' to observe dipolar correlation.

on a AutoSpecE spectrometer. The ^1H and ^{13}C NMR spectra were recorded at 298 K in CDCl_3 or DMSO using a Bruker Avance-500 spectrometer. Chemical shifts (δ) were reported in ppm and coupling constants (J) in Hz. Heteronuclear ^1H – ^{13}C HSQC and HMBC experiments were carried out using standard procedures.

3.1. 4-Hydroxy-2-(diphenylmethylene)benzofuran-3(2H)-one **1a**

2',6'-dihydroxyacetophenone (76 mg, 0.5 mmol), potassium *tert*-butoxide (168 mg, 1.5 mmol), benzophenone (182 mg, 1.0 mmol) and DMSO (60 mg) were mixed to get an homogeneous mass and heated in an oil bath at 200 °C for 2 min. The red-dark mass was hydrolysed with HCl (30 mL, 5%) and the mixture extracted with diethyl ether (3×20 mL). The yellow-orange organic phase was extracted with NaOH (10%, 20 mL) and the aqueous phase discarded. 20 mL of HCl (10%) were added and the mixture shaken vigorously. Finally the organic phase was separated, dried over anhydrous sodium sulphate and evaporated to give an yellow oil that was purified by column chromatography on SiO_2 (2–5% EtOAc/Petroleum ether) to afford dye **1a** as an yellow solid (16 mg, 10% yield). mp 117–119 °C. IR: 3157, 3064, 1674, 1614, 1580, 1458, 1300, 1184, 1048. For ^1H NMR and ^{13}C NMR data see Table 1. MS: m/z (%): 314 (M^+ , 62), 313 (100), 237 (5), 165 (15). Exact mass for $\text{C}_{21}\text{H}_{14}\text{O}_3$: 314.0947; Found 314.0943.

3.2. 2-(9H-Fluoren-9-ylidene)-4-hydroxybenzofuran-3(2H)-one **1b**

Prepared from 9-fluorenone using the same procedure. The compound **1b** was isolated as a yellow solid (14 mg, 9% yield). mp 245–246. IR: 3337, 3106, 3050, 1680, 1636, 1610, 1585, 1467, 1448, 1347, 1318, 1140, 1045. For ^1H NMR and ^{13}C NMR data see Table 1. MS: m/z (%): 312 (M^+ , 6.3), 311 (9), 256 (9), 213 (13), 98 (100). Exact mass for $\text{C}_{21}\text{H}_{12}\text{O}_3$: 312.0786; Found 312.0780.

3.3. (E)-4-Hydroxy-2-(2-hydroxy-9H-fluoren-9-ylidene)benzofuran-3(2H)-one **1c**

Prepared from 2-hydroxyfluorenone using 2.5 mmol of potassium *tert*-butoxide (5 eq). The compound **1c** was isolated as an orange solid (15 mg, 8% yield). mp 220–222. IR: 3470, 3060, 1674, 1639, 1617, 1581, 1453, 1443, 1302, 1263, 1135, 1048. For ^1H NMR and ^{13}C NMR data see Table 1. MS: m/z (%): Exact mass for $\text{C}_{21}\text{H}_{12}\text{O}_4$: 328.0736; Found 328.0726.

Acknowledgements

To FCT (Portugal's Foundation for Science and Technology) and FEDER for financial support through the research unit Centro de Química-Vila Real (POCTI-SFA-3-616) and project PTDC/QUI/66012/2006. The 500 MHz NMR facilities were funded by the Région Nord-Pas de Calais (France), the Ministère de la Jeunesse de l'Éducation Nationale et de la Recherche (MJENR) and the Fonds Européens de Développement Régional (FEDER).

Appendix. Supplementary data

Supplementary data associated with this article can be found in the online version, at doi:10.1016/j.dyepig.2011.05.026.

References

- [1] Boumendjel A. Aurones: a subclass of flavones with promising biological potential. *Curr Med Chem* 2003;10:2621–30.
- [2] Brooks CJ, Watson DG. Phytoalexins. *Nat Prod Rep* 1985;2(5):427–59.
- [3] Keyser O, Kiderlen AF, Folkers U, Kolodziej H. In vitro leishmanicidal activity of aurones. *Planta Med* 1999;65(4):316–9.
- [4] Pare PW, Dmitrieva N, Mabry TJ. Phytoalexin aurone induced in *Cephalocereus senilis* liquid suspension-culture. *Phytochemistry* 1991;30:1133–5.
- [5] Ferreira EO, Salvador MJ, Pral EMF, Alfieri SC, Ito IY, Dias DA. A new hepta-substituted (E)-aurone glucoside and other aromatic compounds of *Gomphrena agrestis* with biological activity. *Z Naturforsch C Biosci* 2004;59(7–8):499–505.
- [6] Kayser O, Kiderlen AF. Leishmanicidal activity of aurones. *Tokai J Exp Clin Med* 1999;23(6):423–6.
- [7] Pires JR, Saito C, Gomes SL, Giesbrecht AM, Amaral AT. Investigation of 5-nitrofuran derivatives: Synthesis, antibacterial activity, and quantitative structure-activity relationships. *J Med Chem* 2001;44(22):3673–81.
- [8] Thomas MG, Lawson C, Allanson NM, Leslie BW, Bottomley JR, McBride A, et al. A series of 2(Z)-2-benzylidene-6,7-dihydroxybenzofuran-3[2H]-ones as inhibitors of chorismate synthase. *Bioorg Med Chem Lett* 2003;13(3):423–6.
- [9] Hadj-esfandiari N, Navidpour L, Shadnia H, Amini M, Samadi N, Faramarzi M, et al. Synthesis, antibacterial activity, and quantitative structure-activity relationships of new (Z)-2-(nitroimidazolylmethylene)-3(2H)-benzofuranone derivatives. *Bioorg Med Chem Lett* 2007;17(22):6354–63.
- [10] Moritomo M, Fukumoto H, Nozoe T, Hagiwara A, Komai K. Synthesis and insect antifeedant activity of aurones against *Spodoptera litura* Larvae. *J Agr Food Chem* 2007;55(3):700–5.
- [11] Venkateswarlu S, Panchagnula GK, Gottumukkala AL, Subbaraju GV. Synthesis, Structural revision, and biological activities of 4'-chloroaurone, a metabolite of marine brown alga *Spatoglossum variable*. *Tetrahedron* 2007;63(29):6909–14.
- [12] Venkateswarlu S, Panchagnula GK, Subbaraju GV. Synthesis and antioxidative activity of 3',4',6,7-tetrahydroxyaurone, a metabolite of *Bidens frondosa*. *Biosci Biotech Biochem* 2004;68(10):2183–5.
- [13] Detsi A, Majdalani M, Kontogiorgis CA, Hadjipavlou-Litina D, Kefalas P. Natural and synthetic 2'-hydroxy-chalcones and aurones: synthesis, characterization and evaluation of the antioxidant and soybean lipoxygenase inhibitory activity. *Bioorg Med Chem* 2009;17(23):8073–85.
- [14] Barros A, Silva A, Alkorta L, Elguero J. Synthesis, experimental and theoretical NMR study of 2'-hydroxychalcones bearing a nitro substituent on their B ring. *Tetrahedron* 2004;60(31):6513–21.
- [15] Okombi S, Rival D, Bonnet S, Mariotte A, Perrier E, Boumendjel A. Discovery of benzylidenebenzofuran-3(2H)-one (aurones) as potent inhibitors of human tyrosinase. *J Med Chem* 2006;49:329–33.
- [16] Aufmkolk M, Koerhle J, Hesch RD, Cody V. Inhibition of rat liver iodothyronine deiodinase. Interaction of aurones with the iodothyronine ligand-binding site. *J Biol Chem* 1986;261:11623–30.
- [17] Sheng R, Xu Y, Hu C, Zhang J, Lin X, Li J, et al. Design, synthesis and AChE inhibitory activity of indanone and aurone derivatives. *Eur J Med Chem* 2009;44(1):7–17.
- [18] Sim HM, Lee CY, Ee PLR, Go ML. Dimethoxyaurones: potent inhibitors of abcg2 (breast cancer resistance protein). *Eur J Pharm Sci* 2008;35:293–306.
- [19] Lawrence NJ, Rennison D, McGown AT, Hadfield JA. The total synthesis of an aurone isolated from *Uvaria hamiltonii*: aurones and flavones as anticancer agents. *Bioorg Med Chem Lett* 2003;13(21):3759–63.
- [20] Huang W, Liu MZ, Li Y, Tan Y, Yang GF. Design, syntheses, and antitumor activity of novel chromone and aurone derivatives. *Bioorg Med Chem* 2007;15(15):5191–7.
- [a] Bandgar BP, Patil SA, Korbad BL, Biradar SC, Nile SN, Khobragade CN. Synthesis and biological evaluation of a novel series of 2,2-bisaminomethylated aurone analogues as anti-inflammatory and antimicrobial agents. *Eur J Med Chem* 2010;45(7):3223–7.
- [21] Hadjeri M, Beney C, Boumendjel A. Recent advances in the synthesis of conveniently substituted flavones, quinolones, chalcones and aurones: potential biologically active molecules. *Curr Org Chem* 2003;7:679–89.
- [22] Varma RS, Varma M. Alumina-mediated condensation - a simple synthesis of aurones. *Tetrahedron Lett* 1992;33(40):5937–40.
- [23] Harkat H, Blanc A, Weibel J, Pale P. versatile and expeditious synthesis of aurones via Au^{I} -catalyzed cyclization. *J Org Chem* 2008;73(4):1620–3.
- [24] Lin CF, Lu WD, Wang IW, Wu MJ. Synthesis of 2-(diarylmethylene)-3-benzofuranones promoted via palladium-catalyzed reactions of aryl iodides with 3-aryl-1-(2-*tert*-butyldimethylsilyloxy)phenyl-2-propyn-1-ones. *Synlett* 2003;13:2057–61.
- [25] Kerr PJ, Pyke SM, Ward AD. Synthesis, and cyclization to aurones and flavones, of alkoxy-substituted aryl arylalkynyl ketones. *Aust J Chem* 2008;61:350–8.
- [26] Menezes MJ, Manjrekar S, Pai V, Patre RE, Tilve SG. A facile microwave assisted synthesis of flavones. *Indian J Chem B* 2009;48B:1311–4.
- [27] Larsen L, Yoon DH, Weavers RT. Synthesis of a range of polyhydroxy 8-aryl flavones. *Synthetic Commun* 2009;39:2935–48.
- [28] Cabrera M, Simoens M, Falchi G, Lavaggi ML, Piro OE, Castellano EE, et al. Synthetic chalcones, flavanones, and flavones as antitumoral agents: biological evaluation and structure-activity relationships. *Bioorg Med Chem* 2007;15(10):3356–67.
- [29] Wawzonek S, Nagler RC, Carlson LJ. The reaction of grignard reagents with 4-hydroxy- and 4-ethoxycoumarin. *J Am Chem Soc* 1954;76:1080.
- [30] Cottam J, Livingstone R, Walshaw M, Bartle KD, Jones DW. Dimerisation of 2,2-diphenyl- and 2,2-dimethyl-chromen. 2. Reaction between 3-(2-hydroxyphenyl)-1,3-diphenylpropan-1-one and 3-(2-hydroxyphenyl)-1,1-diphenylprop-2-en-1-ol. *J Chem Soc*; 1965:5261–5.
- [31] Yugai GA, Mostoslavskii MA, Paramonov VD. Absorption spectra of photoinduced E isomers of aurones, thioindogenides, and selenoindogenides. *Theor Exp Chem* 1977;12:549–51.