New Approaches to the Synthesis of 4-(2-Aminophenyl)-1,3-dihydro-2*H*imidazol-2-ones and 3-Ureidoindoles and a Study of Their Interconversion

by Antonín Klásek*^a), Antonín Lyčka^b), Michal Holčapek^c), and Ignác Hoza^d)

^a) Department of Chemistry, Faculty of Technology, Tomas Bata University, CZ-76272 Zlín (e-mail: klasek@ft.utb.cz)

^b) Research Institute for Organic Syntheses, Rybitví 296, CZ-53354 Pardubice 20

^c) Department of Analytical Chemistry, Faculty of Chemical Technology, University of Pardubice, CZ-53210 Pardubice

d) Department of Food Engineering, Faculty of Technology, Tomas Bata University, CZ-76272 Zlín

Dedicated to Professor Vojeslav Štěrba on the occasion of his 85th birthday

3-Alkyl/aryl-3-ureido-1*H*,3*H*-quinoline-2,4-diones (**2**) and 3a-alkyl/aryl-9b-hydroxy-3,3a,5,9b-tetrahydro-1*H*-imidazo[4,5-*c*]quinoline-2,4-diones (**3**) react in boiling concentrated HCl to give 5-alkyl/aryl-4-(2-aminophenyl)-1,3-dihydro-2*H*-imidazol-2-ones (**6**). The same compounds were prepared by the same procedure from 2-alkyl/aryl-3-ureido-1*H*-indoles (**4**), which were obtained from the reaction of 3alkyl/aryl-3-aminoquinoline-2,4(1*H*,3*H*)-diones (**1**) with 1,3-diphenylurea or by the transformation of 3a-alkyl/aryl-9b-hydroxy-3,3a,5,9b-tetrahydro-1*H*-imidazo[4,5-*c*]quinoline-2,4-diones (**3**) and 5-alkyl/ aryl-4-(2-aminophenyl)-1,3-dihydro-2*H*-imidazol-2-ones (**6**) in boiling AcOH. The latter were converted into 1,3-bis[2-(2-oxo-2,3-dihydro-1*H*-imidazol-4-yl)phenyl]ureas (**5**) by treatment with triphosgene. All compounds were characterized by ¹H- and ¹³C-NMR and IR spectroscopy, as well as atmospheric pressure chemical-ionisation mass spectra.

1. Introduction. – In our laboratory, much attention has been paid to study the reactivity of 3-alkyl/aryl-3-hydroxy-quinoline-2,4(1H,3H)-diones and their 3-amino analogues $\mathbf{1}$ [1–9]. Both types of compounds readily rearrange into other heterocyclic systems. In the case of the 3-hydroxy derivatives, thermally induced rearrangement produces dioxindoles or 3,1-benzoxazin-2-ones [1][2], rearrangement in an alkaline environment produces dioxindoles and 2-hydroxyindoxyls [3]. 3-Amino derivatives rearrange, when treated with urea in boiling AcOH, to yield imidazo[1,5-c]quinazoline-3,5-diones [4] or 3-ureidoindolin-2-ones [5]. The reaction of 3-aminoquinoline-2,4diones 1 with nitrourea in dioxane was successful in preparing anticipated reaction intermediates of this rearrangement [6][7], whereas the reaction with isocyanates yielded 3'-substituted 3-ureidoquinoline-2,4-diones 2 or their cyclic isomers 3 [8]. These compounds rearrange in boiling AcOH to give either mixtures of 3ureidoindoles 4 and 1,3-disubstituted ureas 5, or merely 1,3-disubstituted ureas 5 (Scheme 1) [9]. Refluxing compounds 4 in conc. HCl changes the indole system into imidazolinone, and compounds 6 prepared in this manner provide compounds 5 by a reaction with triphosgene [9].

The discovery that the addition products of 3-aminoquinoline-2,4-diones and isocyanates (2 and 3) rearrange to indole and imidazolinone derivatives, opens up a

© 2008 Verlag Helvetica Chimica Acta AG, Zürich



path for the preparation of new compounds of type **4**, **5**, and **6**, which may be interesting structures for studying biological activity. For example, 2-phenyl-3-semicarbazidoin-doles, which have a structure similar to **4**, display antimicrobial and antihistaminic activity [10][11]; interesting biological activity is described for many imidazolin-2-ones [12], and a number of 1,3-diarylureas act as anti-cytokinine agents [13].

Transformation of compounds 2 and/or 3 to compounds 4 occurs only in rather low yield, and the main or even sole product in most cases are 1,3-disubstituted ureas 5 [9]. Another difficulty is, that mixtures of 4 and 5 can be separated only with great difficulty. For that reason, we tried to find reaction conditions where 3-ureidoindoles 4 or imidazolinones 6 arise, while the formation of compounds 5 is limited. This paper describes the results achieved in this matter so far.

2. Results and Discussion. – Our first aim was to find reaction conditions that would enable us to prepare 3-ureidoindoles **4** directly from α -amino ketones **1**. In one of our earlier papers [6], we described that 3-aminoquinoline-2,4-diones **1** in boiling AcOH yield products of molecular rearrangement not only in the reaction with urea but also in reactions with monosubstituted or 1,1-disubstituted ureas (butylurea, phenylurea, 1,1-dibenzylurea). Under the given reaction conditions, monosubstituted or 1,1-disubstituted ureas split into the respective amine and isocyanic acid, which reacts with 3-aminoquinoline-2,4-diones **1** giving rise to intermediates analogous to **2** and **3**

 $(R^3 = H)$, subject of molecular rearrangement. This fact led to the idea that, under similar reaction conditions, 1,3-disubstituted ureas could produce the corresponding amine and isocyanate that would react with 1 to yield intermediates 2 and/or 3, which subsequently rearrange to the products 4. This assumption was confirmed, and from the reaction of 3-aminoquinolones 1f, 1h, and 1l with 1,3-diphenylurea in boiling AcOH (*Scheme 2*) the rearranged compounds 4f, 4h, and 4l were isolated as the main products (*Table 1, Method A*).



As a side product, acetanilide was isolated. Additionally, in one single case, the diarylurea 51 was obtained in negligible quantities. Other side products, which exhibited a molecular peak of even m/z values in the mass spectra were obtained in two cases (starting from 1f and 1l). Their molecular peaks split into two primary fragments, which differ by a value of m/z 26. Such splitting of the molecular ion is typical for 1.3disubstituted ureas [9] and, therefore, we propose structure 7 for the side products of the reaction of compounds 1f and 1l with 1,3-diphenylurea (Scheme 2). This proposal was confirmed by synthesizing 7 from 6 and phenyl isocyanate. The formation of compounds 7f and 7l may be explained by a reaction mechanism analogous to that of the transformation of compounds 2 and 3 into 4 and 5 via an intermediate A [9] (Scheme 2). The reaction of amines 1 with diphenylurea (Table 1, Method A) gives generally lower yields of 4 than the reaction of 2 or 3, which are transformed to 4 in boiling AcOH [9]; but the formation of 1,3-diarylureas is limited in every case. However, the disadvantage of the direct transformation of 1 into 4 (Method A) is its limitation to 3-ureidoindoles 4 that bear a Ph group in position 1' ($R^3 = Ph$). When compounds 1 were reacted with 1,3-dibutylurea, we merely obtained inseparable mixtures of products. A further disadvantage of the direct transformation of 1 to 4 is the difficult chromatographic separation of 4 from the reaction mixture containing considerable quantities of unreacted 1,3-diphenylurea and acetanilide. A considerable proportion of compounds displaying a polar character also arose, and these could not

Entry	Starti	ing com	pound		Method	Reaction time [h]	Products (Yield [%])
		\mathbb{R}^1	\mathbb{R}^2	R ³			
1	2a	Bu	Н	Bu	Ε	1+1.5	4a (68)
2	2a	Bu	Н	Bu	В	1.5	4a (18), 5a (67)
3	3a	Bu	Н	Bu	В	1.5	4a (31), 5a (53)
4	6a	Bu	Н	Bu	D	3	4a (45)
5	6a	Bu	Н	Bu	D	1.5	4a (30)
6	6b	Bu	Н	Ph	D	1	4b (31)
7	3c	Ph	Н	Bu	Ε	1 + 1.5	4c (82)
8	6c	Ph	Н	Bu	D	1	4c (67)
9	2d	Ph	Н	Ph	Ε	1 + 1.5	4d (65)
10	6d	Ph	Н	Ph	D	1	4d (65)
11	3e	Bu	Me	Bu	Ε	1 + 1.5	4e (42), 6e (31)
12	6e	Bu	Me	Bu	D	2	4e (28)
13	1f	Bu	Me	Ph	Α	1.5	4f (21), 7f (8)
14	1f	Bu	Me	Ph	Α	2	4f (29), 7f (6)
15	6f	Bu	Me	Ph	D	1	4f (76)
16	6g	Bu	Bu	Bu	D	3	4g (86)
17	1ĥ	Bu	Bu	Ph	Α	4.5	4h (33)
18	3h	Bu	Bu	Ph	В	0.5	4h (36), 5h (28)
19	6h	Bu	Bu	Ph	D	4	4h (56)
20	6i	Bu	PhCH ₂	Bu	D	3.5	4i (53)
21	6i	Bu	PhCH ₂	Ph	D	2.5	4i (36)
22	3k	Ph	Bu	Bu	Ε	1 + 1.5	4k (67)
23	6k	Ph	Bu	Bu	D	1	4k (43), 8k (16), 6k (7)
24	11	Ph	Bu	Ph	Α	6	41 (47), 51 (3), 71 (8)
25	61	Ph	Bu	Ph	D	3	4I (58)
26	3m	Ph	PhCH ₂	Bu	Ε	1 + 1.5	4m (53), 6m (12), 8m (9)
27	3m	Ph	PhCH ₂	Bu	В	0.5	4m (4), 5m (51), 6m (12)
28	6m	Ph	PhCH ₂	Bu	D	5	4m (14), 6m (25), 8m (21)
29	3n	Ph	PhCH	Ph	В	0.5	4n (4), 5n (34), 6n (32)
30	3n	Ph	PhCH ₂	Ph	Ε	1 + 1.5	4n (77)
31	6n	Ph	PhCH	Ph	D	1.5	4n (21), 8n (16), 6n (32)
32	6n	Ph	PhCH-	Ph	D	3	4n (16) $8n$ (25) $6n$ (32)

Table 1. Preparation of 3-Ureidoindoles 4 from Compounds 1 (Method A), 2 (Methods B and E), 3 (Methods B and E), and 6 (Method D)

be separated by chromatography. For this reason, we abandoned this path for preparing 3-ureidoindoles 4 and sought for other possibilities.

We propose a reaction mechanism with a rearrangement of compounds 2 and 3 in boiling AcOH to 4 and 5 (or 5 only) via the formation of an intermediate A' (Scheme 3) [9]. This intermediate partly hydrolyses to give 6, which is transformed to 4 as indicated in Scheme 4. However, at the same time, A' reacts with 6 to yield the disubstituted urea 5. It can be assumed that, in the presence of excess H₂O, compounds 4 should arise as the sole products. Therefore, we carried out the rearrangement of compounds 2a and 3a, 3h, 3m, and 3n in aqueous AcOH (Method B). However, the results of these experiments (Table 1, Entries 2, 3, 18, 27, and 29) show that even under these reaction conditions, intermediate A' reacts with the amino group of 6, and mixtures of 4 and 5





arise. In cases where $R^1 = Ph$, compounds **6** are obtained as well (*Table 1, Entries 27* and 29). This finding is in accordance with the fact that, in these cases, the tendency for intramolecular addition of the amino group in **6** is lowered [9]. Because the use of *Method B* does not lead to satisfactory results, these experiments were discontinued.

Compounds 6 are also possible sources for the preparation of 3-ureidoindoles 4. As already described [9], compounds 6 are unstable in polar solvents, and some of them were converted back to compounds 4 during NMR spectra collection in $(D_6)DMSO$. This conversion is particularly easy when $R^2 = H$. However, there is a question how 6

can be easily prepared. We have described the transformation of 3-ureidoindoles 4a, 4b, 4f, 4h, 4i, 4j, and 4l to imidazolones 6a, 6b, 6f, 6h, 6i, 6j, and 6l (*Scheme 1*) by boiling compounds 4 in HCl [9]. Compounds 4c, 4d, 4e, 4g, 4k, 4m, and 4n have not been available earlier, because only 1,3-disubstituted ureas 5 were isolated from the corresponding reaction mixtures after reacting the corresponding compounds 2 or 3 with urea in AcOH. However, the easy transformation of 3-ureidoindoles 4 into imidazolinones 6 led us to reason that compounds 6 could also be formed by direct transformation of ureidoquinolinediones 2 or their cyclic isomers 3 in strongly acidic aqueous environments. This assumption was fully confirmed, and the corresponding dihydroimidazolones 6 were isolated in high yields from the reactions of 2a-2d or 3c, 3e-3n with conc. HCl (*Table 2, Method C*). Acid-catalyzed rearrangement of 2 or 3 (*Scheme 3*) gives rise to carbocation B, followed by the opening of the indolone ring under formation of intermediate A'. The latter immediately hydrolyses in a strongly acidic aqueous environment to give dihydroimidazolone 6 and, therefore, formation of disubstituted ureas 5 cannot take place.

Table 2. Preparation of Dihydroimidazolones 6, from Compounds 2, 3, 4, and 8 under Reflux in conc. HCl (Method C)^a)^b)

Entry	Starting compound	Reaction time [h]	Product (Yield [%])	Entry	Starting compound	Reaction time [h]	Product (Yield [%])
1	2a	2	6a (80)	16	4g	1	6g (83)
2	2a	1	6a (54)	17	3h	0.5	6h (66)
3	3a	4	6a (76)	18	3h	2	6h (52)
4	2b	1	6b (74)	19	3i	2	6i (72)
5	2c	1	6c (79)	20	3j	2	6j (50)
6	3c	1	6c (78)	21	3k	1	6k (85)
7	4c	1	6c (68)	22	4k	1	6k (75)
8	2d	0.5	6d (83)	23	8k	1	6k (65)
9	4d	2.5	6d (79)	24	31	1.5	61 (78)
10	3e	2	6e (75)	25	3m	1	6m (72)
11	4 e	0.5	6e (80)	26	4m	1	6m (79)
12	3f	1.5	6f (83)	27	8m	1	6m (49)
13	3f	2	6f (73)	28	3n	0.5	6n (73)
14	3f	3	6f (63)	29	4n	2.5	6n (54)
15	3g	1	6g (92)	30	8n	1	6n (53)

^a) For substituents, see *Table 1*. ^b) Conversion of compounds **4a**, **4b**, **4f**, **4h**, **4i**, **4j**, and **4l** to the corresponding compounds **6** has been described in [9].

Now we found that reconversion of **6** into ureidoindoles **4** may be successfully executed in boiling AcOH (*Table 1, Method D*). In a weakly acidic environment, compounds **6** undergo an intramolecular nucleophilic addition of the amino group to form intermediate **C**. The intermediate's imidazolidinone ring is opened, and ureidoindoles **4** are formed (*Scheme 4*).

The anticipated intermediates C are aza-analogues of hexahydropyrrolo[2,3b]indole alkaloids, which are cyclic analogues of N-acylated derivatives of tryptophane. It is well-known, that these alkaloids and their synthetic analogues arise from Nacylated tryptophane derivatives in a strongly acidic environment but, under weakly acidic conditions, they reconvert into acyclic tautomers [14-18]. The transformation of **6** to **4** via the intermediate **C** bears a certain distant analogy to the conversions of hexahydropyrrolo[2,3-b]indoles. Despite all efforts, we did not succeed in our earlier work [9] and have not yet succeeded in isolating or trapping anticipated intermediates **C**. In a majority of cases, heating **6** in AcOH to reflux merely provided the corresponding compounds **4** (*Table 1, Method D*). However, the reaction of **6k**, **6m**, and **6n** in AcOH gave, apart from the corresponding products **4k**, **4m**, and **4n** and unreacted **6k**, **6m**, and **6n**, other side products (**8k**, **8m**, and **8n**). These compounds revert to **6** in boiling conc. HCl (*Table 2, Method C*). Hence, **6k**, **6m**, and **6n**, which differ from other compounds **6** by the presence of a Ph group at C(5) of the imidazolone nucleus, obviously convert only slowly into indole derivatives **4** by heating in AcOH, and a competitive transformation into acetylated compounds **8** occurs (*Scheme 4*). The structure of **8** was confirmed by its synthesis from **6** and Ac₂O. The reason why this transformation from **6** to **8** does not take place in the case of **6l**, which bears a Ph group at C(5) like **6k**, **6m**, and **6n**, is an issue we failed to resolve.

Rearrangement of compounds 2 or 3 in conc. HCl (*Table 2, Method C*) offers high yields of imidazolinones 6 (mostly in a 74–92% range). In a number of cases, however, the following transformation of 6 into ureidoindoles 4 (*Table 1, Method D*) shows substantially lower yields (28-86%). If the target products are compounds 4, it is much more favorable to prepare them from compounds 2 or 3 by performing a two-stage reaction without isolating an intermediate (*Scheme 3, Table 1, Method E*). Compounds 4a, 4c, 4d, 4k, and 4n were prepared from 2a, 3c, 2d, 3k, and 3n in this manner with yields of 65-82% (*Table 1*). In the cases of 3e and 3m, compounds 6 were obtained additionally to 4 (*Table 1*).

In our previous work [9], we described the preparation of **5a**, **5f**, **5h**, **5i**, **5j**, and **5l** by a reaction of the corresponding dihydroimidazolones **6** with triphosgene (bis(trichloromethyl) carbonate). The other compounds of type **5** could not be prepared because ureidoindoles **4** were not available from the reactions of **2** and/or **3** in AcOH, thus dihydroimidazolones **6** could not be prepared either. But now, having in hand the relevant compounds **6** from the reaction of **2** and/or **3** in conc. HCl, we attempted to prepare the remaining **5c**, **5d**, **5e**, **5g**, **5k**, **5m**, and **5n**. As in our previous work [9], we found that compounds **6** react with triphosgene at room temperature in CH₂Cl₂ very slowly, even though this procedure has been currently employed by other groups [19][20]. The disubstituted ureas **5e**, **5g**, **5k**, **5m**, and **5n** were obtained by performing the reaction in boiling benzene and in the presence of a catalytic quantity of 4-(dimethylamino)pyridine (*Scheme 5*). Preparation failed only in cases where **6** is not substituted at C(1) (R² = H). Similar to our previous work [9], where compound **5a** was prepared in only 6% yield and preparation of **5b** was not possible, we obtained **5c** in only 12% yield and failed to prepare **5d**.

In conclusion, we would like to emphasise that the elaborated methods enable us to prepare 3-ureidoindoles 4 and/or dihydroimidazolones 6 from 3-aminoquinolinediones 1 *via* the intermediates 2 or 3 in high yields. The unprecedented mutual transformation of 4 and 6 is not merely of theoretical importance but enables, through a simple procedure, a targeted preparation of new types of compounds suitable for biological testing as well as for studying further transformations, particularly in the case of dihydroimidazolones 6 bearing a reactive aniline fragment.





This study was supported by the *Ministry of Education, Youth and Sports* of the Czech Republic (Grant No. MSM 7088352101) and *Czech Science Foundation* (grant No. 203/07/0320). *M. H.* acknowledges the support of grant projects MSM 0021627502 (the *Ministry of Education, Youth and Sports* of the Czech Republic) and 203/05/2106 (*Czech Science Foundation*). The authors thank Mrs. *H. Geržová* (Faculty of Technology, Tomas Bata University in Zlín), and Mr. *R. Jirásko* (University of Pardubice) for technical help.

Experimental Part

1. General. Reactions, as well as the course of separation and also the purity of substances, were monitored by TLC (elution systems benzene/AcOEt, 4:1 (solvent system S3), CHCl₃/EtOH, 9:1 (S4) and/or 19:1 (S5), and CHCl₃/AcOEt, 7:3 (S6)) on Alugram[®] SIL G/UV₂₅₄ foils (Macherey-Nagel). Column chromatography (CC) was carried out on silica gel (Merck, grade 60, 70-230 mesh), using CHCl₃, and then successive mixtures of CHCl₃/EtOH (99:1 to 8:2, solvent system S1) or benzene, and then successive mixtures of benzene/AcOEt (99:1 to 8:2, solvent system S2). M.p.: Kofler block or Gallencamp apparatus. IR (KBr) Spectra: Mattson 3000 spectrophotometer. NMR Spectra were recorded on a *Bruker Avance* spectrometer (500.13 MHz for ¹H and 125.76 MHz for ¹³C) in (D_6)DMSO. ¹H and ¹³C chemical shifts are given on the δ scale (in ppm) and are referenced to internal TMS. All 2D experiments (gradient-selected (gs)-COSY, NOESY, gs-TOCSY, gs-HMQC, gs-HMBC) were performed using the manufacturer's software. ¹H-NMR Spectra were assigned using gs-COSY. Protonated C-atoms were assigned by gs-HMQC. Quaternary C-atoms were assigned by gs-HMBC. The positive-ion atmospheric pressure chemical ionization (APCI) mass spectra were measured on an ion trap analyzer Esquire 3000 (Bruker Daltonics) within the mass range m/z 50-500. Samples were dissolved in MeCN and analyzed by direct infusion at the flow rate of 50 µl/min. The ion-source temp. was 350°, the APCI probe temp. was 300° , the flow rate and the pressure of N₂ were 4 l/min and 45 psi, resp. For MS/MS measurements, the isolation width of precursor ions was 4 m/z and the collision amplitude was 1 V. Elemental analyses (C, H, N): EA 1108 Elemental Analyzer (Fisons Instrument).

Starting compounds **2** and **3** were prepared from the corresponding compounds **1** according to the protocol described in [8]. Yields of products **4**, **6**, and **8** are given in *Tables 1* and *2*.

2. General Procedures for the Preparation of (1H-Indol-3-yl)ureas **4** and 4-(2-Acetamidophenyl)-1,3dihydroimidazol-2-ones **8**. Method A. A soln. of **1** (0.5 mmol) and 1,3-diphenylurea (117 mg, 0.55 mmol) in AcOH (2 ml) was heated to reflux for the time given in Table 1. After cooling, the precipitated crystals were filtered off with suction and recrystallized from an appropriate solvent. In cases when crystallization failed, the mixtures were evaporated *in vacuo* to dryness, and the residue was crystallized from an appropriate solvent. In some cases, mother liquors were chromatographed on silica gel using solvent systems S1 or S2.

Method B. A soln. of **2** or **3** (0.5 mmol) in aq. AcOH (70% v/v, 6 ml) was heated to reflux for the time given in *Table 1*. After cooling, the solvent was evaporated *in vacuo* to dryness, and the residue was crystallized from an appropriate solvent. In some cases, mother liquors were worked up by CC on silica gel.

Method D. A soln. of **6** (0.5 mmol) in AcOH (6 ml) was heated to reflux for the time given in *Table 1*. Following treatment of the reaction mixture, workup was carried out as described in *Method B*.

Method E. A soln. of compound **2** or **3** (0.5 mmol) in conc. HCl (5 ml) was heated to reflux for 1 h. After cooling, the soln. was evaporated *in vacuo* to dryness, the residue was dissolved in pyridine (5 ml), anh. K_2CO_3 (350 mg) was added, and the mixture was stirred for 1 h. Insoluble material was filtered off, the filtrate was evaporated do dryness, dissolved in AcOH (5 ml), and the mixture was heated to reflux for 1.5 h. After cooling, the solvent was evaporated *in vacuo* to dryness, and the residue was crystallized from an appropriate solvent. In some cases, mother liquors were separated by CC on silica gel.

2.1. *1-Butyl-1-(2-butyl-1H-indol-3-yl)urea* (**4a**). Prepared from **2a** (*Methods B* (besides **5a**) and *E*), **3a** (*Method B*), and **6a** (*Method D*). Colorless crystals. M.p. $136-141^{\circ}$ (benzene/hexane), identical in all respects to authentic **4a** of m.p. $135-142^{\circ}$, prepared from **2a** and **3a** [9].

2.2. 1-(2-Butyl-1H-indol-3-yl)-1-phenylurea (**4b**). Prepared from **6b** (*Method D*). Colorless crystals. M.p. 205–208° (AcOEt), identical in all respects to authentic **4b** of m.p. 204–208°, prepared from **2b** [9].

2.3. *1-Butyl-1-(2-phenyl-1*H-*indol-3-yl)urea* (**4c**). Prepared from **3c** (*Method E*) and **6c** (*Method D*). Colorless crystals. M.p. 212–215° (benzene). IR: 3445, 3306, 3218, 3195, 3064, 2952, 2932, 2869, 1641, 1581, 1491, 1464, 1453, 1427, 1379, 1356, 1339, 1309, 1279, 1231, 1179, 1036, 1115, 1074, 1028, 1008, 965, 921, 764, 739, 697, 630, 614, 564. ¹H- and ¹³C-NMR: see *Table 3*. APCI-MS (pos.): 308 (28, $[M + H]^+$), 291 (100, $[M + H - NH_3]^+$), 263 (36, $[M + H - NH_3 - CO]^+$). APCI-MS/MS (pos.) of *m*/*z* 308: 291 (100, $[M + H - NH_3]^+$). APCI-MS (neg.): 306 (100, $[M - H]^-$), 261 (18, $[M - H - COOH]^-$), 233 (4, $[M - H - BuNH_2]^-$). APCI-MS/MS (neg.) of *m*/*z* 306: 263 (43, $[M - H - NHCO]^-$), 249 (75, $[M - H - Bu]^-$), 206 (100, $[M - H - Bu - NHCO]^-$). Anal. calc. for C₁₉H₂₁N₃O (307.39): C 74.24, H 6.89, N 13.67; found: C 74.32, H 6.92, N 13.51.

2.4. *1-Phenyl-1-(2-phenyl-1*H-*indol-3-yl)urea* (**4d**). Prepared from **2d** (*Method E*) and **6d** (*Method D*). Colorless crystals. M.p. 250–256° (benzene). IR: 3458, 3394, 3318, 3190, 3061, 1659, 1579, 1492, 1453, 1404, 1340, 1264, 1239, 1198, 1148, 1073, 1030, 829, 763, 741, 693, 637, 616, 564, 519. ¹H- and ¹³C-NMR: see *Table 3*. APCI-MS (pos.): 328 (67, $[M + H]^+$), 311 (55, $[M + H - NH_3]^+$), 283 (100, $[M + H - COOH]^+$), 206 (3, $[M + H - COOH - Ph]^+$). APCI-MS/MS (pos.) of *m/z* 328: 311 (100, $[M + H - NH_3]^+$), 284 (5, $[M + H - NH_2CO]^+$), 209 (6, $[M + H - PhNCO]^+$). APCI-MS (neg.): 326 (100, $[M - H]^-$), 282 (94, $[M - H - NH_2CO]^-$). APCI-MS/MS (neg.) of *m/z* 326: 283 (100, $[M - H - NH_2CO]^-$). Anal. calc. for C₂₁H₁₇N₃O (327.38): C 77.04, H 5.23, N 12.84; found: C 77.14; H 5.36, N 12.69.

2.5. *1-Butyl-1-(2-butyl-1H-indol-3-yl)-3-methylurea* (4e). Prepared from **3e** (*Method E*, besides **6e**) and from **6e** (*Method D*). Colorless crystals. M.p. $127-130^{\circ}$ (hexane). IR: 3348, 3249, 2959, 2930, 2872, 2745, 1613, 1523, 1458, 1409, 1383, 1292, 1227, 1185, 1144, 1124, 1011, 924, 768, 743, 715, 646, 568, 521. ¹H- and ¹³C-NMR: see *Table 3*. APCI-MS (pos.): 302 (100, $[M + H]^+$), 271 (45, $[M + H-MeNH_2]^+$), 243 (50, $[M + H-MeNH_2-CO]^+$), 215 (22, $[M + H-MeNH_2-butene]^+$), 172 (79, $[M + H-MeNH_2-butene-NHCO]^+$). APCI-MS/MS (pos.) of *m/z* 302: 271 (100, $[M + H-MeNH_2]^+$), 243 (72, $[M + H-MeNH_2-CO]^+$), 215 (37, $[M + H-MeNH_2-butene]^+$), 172 (79, $[M + H-MeNH_2-butene-NHCO]^+$). APCI-MS (neg.): 300 (100, $[M - H]^-$), 269 (3, $[M - H-MeNH_2]^-$). APCI-MS/MS (neg.) of *m/z* 300: 243 (100, $[M - H-Bu]^-$), 186 (45, $[M - H-2Bu]^-$), 170 (5, $[M - H-Bu-BuNH_2]^-$). Anal. calc. for $C_{18}H_{27}N_3O$ (301.43): C 71.72, H 9.03, N 13.94; found: C 71.62, H 9.14, N 13.79.

2.6. 1-(2-Butyl-1H-indol-3-yl)-3-methyl-1-phenylurea (4f). Prepared from 1f (Method A) and from 6f (Method D). Colorless crystals. M.p. $208-214^{\circ}$ (AcOEt), identical in all respects to authentic 4f of m.p. $204-208^{\circ}$, prepared from 3f [9]. In a small quantity (8 and 6%, resp.), a side-product with m.p. $229-234^{\circ}$ (benzene/hexane), identical in all respects to synthetic 7f (Section 6.1), was isolated from the reactions of 1f with diphenylurea (Method A).

2.7. *I*,3-*Dibutyl-I-(2-butyl-IH-indol-3-yl)urea* (**4g**). Prepared from **6g** (*Method D*). Yellowish oil. IR: 2958, 2929, 2870, 1708, 1639, 1586, 1517, 1458, 1379, 1330, 1287, 1225, 1178, 1148, 1128, 1014, 924, 845, 767, 743, 624, 522. ¹H- and ¹³C-NMR: see *Table 3*. APCI-MS (pos.): 344 (100, $[M + H]^+$), 271 (13, $[M + H-BuNH_2]^+$), 243 (12, $[M + H-BuNHCOH]^+$), 215 (5, $[M + H-BuNH_2-butene]^+$). APCI-MS/ MS (pos.) of *m*/*z* 344: 271 (85, $[M + H-BuNH_2]^+$), 243 (100, $[M + H-BuNHCOH]^+$), 215 (32, $[M + H-BuNH_2-butene]^+$), 172 (54, $[M + H-BuNH_2-butene-NHCO]^+$). APCI-MS (neg.): 342 (100,

			Table	э 3. ¹ Н-	and ¹³ C-Chem	tical Shi	fts (ð [ppm])	of Com	pounds 4 in ((D ₆)DM	OS.			
Position	4c		4d		4e		4g		4k		4m		4n	
	$\delta_{ m H}$	δ_{C}	$\delta_{ m H}$	δ_{C}	$\delta_{ m H}$	δ_{C}	$\delta_{ m H}$	$\delta_{ m C}$	$\delta_{ m H}$	δ_{C}	$\delta_{ m H}$	δ_{C}	$\delta_{ m H}$	δ_{C}
1	11.62	I	11.75	I	11.10	I	11.13	I	11.64	I	11.66	I	11.78	I
2	I	131.3	I	131.1	I	137.4	I	137.2	ļ	133.0	I	133.3	I	133.8
3	I	114.9	I	114.9	I	113.1	I	112.9	ļ	114.2	I	114.1	I	113.8
3a	I	126.0	I	125.9	I	125.1	I	124.8	ļ	126.0	I	126.3	I	126.0
4	7.40	117.7	7.40	117.6	7.26	116.9	7.26	116.7	7.35	117.6	7.39	117.8	7.40	117.6
5	7.12	119.8	7.12	120.1	7.01	119.2	7.01	119.1	7.10	119.9	7.11	119.9	7.02	120.0
6	7.21	122.2	7.21	123.6	7.09	120.8	7.09	120.7	7.20	122.4	7.21	122.4	7.23	123.5
7	7.47	111.9	7.47	112.0	7.35	111.4	7.35	111.4	7.47	112.0	7.47	112.0	7.52	112.0
7a	I	134.9	I	134.9	I	134.5	I	134.4	I	135.0	I	135.0	I	135.0
NHCONH	I	158.5	I	157.2	I	158.4	I	157.6	I	157.7	I	158.4	I	156.5
CONHR ²	5.62	I	6.11	I	5.30	I	5.25	I	5.72	I	6.44	I	6.95	Ι
$1'(R^1)$	I	132.6	I	133.2	2.62, 2.59	24.8	2.62, 2.59	24.8	I	131.3	I	133.2	I	131.0
$2'$ (\mathbf{R}^{1})	7.83	126.5	7.88	126.3	1.76, 1.66	31.2	1.74, 1.69	30.4	7.79	126.5	7.78	126.6	7.85	126.8
3' (R ¹)	7.51	128.8	7.52	128.9	1.39	22.3	1.39	22.3	7.52	128.9	7.50	128.6	7.52	128.9
4' (R ¹)	7.41	127.9	7.42	128.1	0.96	14.1	0.96	13.8	7.40	128.0	7.42	128.1	7.42	128.1
$1'(R^2)$	I	I	I	I	2.51	27.4	3.03, 2.91	39.6	3.01, 2.96	39.8	4.21, 4.19	43.6	4.27, 4.21	43.6
2' (R ²)	I	I	I	I	I	I	1.28	32.3	1.26	32.4	I	141.5	I	140.8
3' (R ²)	I	I	I	I	I	I	1.16	19.4	1.13	19.4	7.17	127.0	7.24	126.8
$4'$ (\mathbb{R}^2)	I	I	I	I	I	I	0.83	13.9	0.72	13.9	7.25	128.1	7.20	128.2
5' (R ²)	I	I	I	I	I	I	I	I	I	I	7.21	126.4	7.20	126.3
$1'(R^3)$	3.73, 3.22	48.1	I	143.6	3.59, 3.52	48.2	3.58, 3.51	48.1	3.72, 3.24	48.4	3.76, 3.28	48.6	I	143.7
2' (R ³)	1.36, 1.18	30.8	7.27	124.0	1.39	30.5	1.39	31.0	1.35, 1.26	31.0	1.39, 1.27	31.0	7.30	123.8
3' (R ³)	1.14	19.7	7.21	128.2	1.28	19.8	1.28	19.7	1.13	19.9	1.16	19.9	7.11	128.0
$4'(R^3)$	0.73	13.8	7.24	122.5	0.87	13.9	0.85	13.8	0.80	13.9	0.73	13.9	7.23	122.5

Helvetica Chimica Acta – Vol. 91 (2008)

 $[M - H]^{-}$, 241 (3, $[M - H - PhNHCOH]^{-}$). APCI-MS/MS (neg.) of m/z 342: 285 (19, $[M - H - Bu]^{-}$), 243 (100, $[M - H - PhNCO]^{-}$), 186 (94, $[M - H - PhNCO - Bu]^{-}$), 143 (7, $[M - H - PhNCO - Bu - NHCO]^{-}$). Anal. calc. for C₂₁H₃₃N₃O (343.51): C 73.43, H 9.68, N 12.23; found: C 73.31, H 9.75, N 12.13.

2.8. 3-Butyl-1-(2-butyl-1H-indol-3-yl)-1-phenylurea (4h). Prepared from 1h (Method A), 3h (Method B, besides 5h), and 6h (Method D). Colorless crystals. M.p. $166-169^{\circ}$ (benzene/hexane), identical in all respects to authentic 4h of m.p. $168-171^{\circ}$, prepared from 3h [9].

2.9. 3-Benzyl-1-butyl-1-(2-butyl-1H-indol-3-yl)urea (**4i**). Prepared from **6i** (*Method D*). Colorless crystals. M.p. $135-140^{\circ}$ (benzene/hexane), identical in all respects to authentic **4i** of m.p. $137-142^{\circ}$, prepared from **3i** [9].

2.10. 3-Benzyl-1-(2-butyl-1H-indol-3-yl)-1-phenylurea (4j). Prepared from 6j (Method D). Colorless crystals. M.p. $152-156^{\circ}$ (cyclohexane), identical in all respects to authentic 4j of m.p. $153-157^{\circ}$, prepared from 3j [9].

2.11. *1,3-Dibutyl-1-(2-phenyl-1*H-*indol-3-yl)urea* (**4k**). Prepared from **3k** (*Method E*) and from **6k** (*Method D*, besides **8k** and unreacted **6k**). Colorless crystals. M.p. $150-153^{\circ}$ (benzene/hexane). IR: 3426, 3185, 2956, 2929, 2870, 1632, 1585, 1518, 1455, 1383, 1335, 1285, 1227, 1147, 1113, 1032, 1007, 916, 764, 744, 695, 633, 613, 563. ¹H- and ¹³C-NMR: see *Table 3*. APCI-MS (pos.): 364 (100, $[M + H]^+$), 291 (10, $[M + H-BuNH_2]^+$). APCI-MS/MS (pos.) of *m*/*z* 364: 291 (100, $[M + H-BuNH_2]^+$), 265 (51, $[M + H-BuNCO]^+$), 235 (9, $[M + H-BuNH_2-butene]^+$). APCI-MS (neg.): 362 (100, $[M - H]^-$), 261 (4, $[M - H-BuNHCOH]^-$). APCI-MS/MS (neg.) of *m*/*z* 362: 305 (12, $[M - H-Bu]^-$), 263 (52, $[M - H-BuNCO]^-$), 206 (100, $[M - H-BuNCO-Bu]^-$). Anal. calc. for C₂₃H₂₉N₃O (363.50): C 76.00, H 8.04, N 11.56; found: C 75.88, H 8.15, N 11.39.

2.12. 4-(2-Acetamidophenyl)-1,3-dibutyl-5-phenyl-1,3-dihydro-2H-imidazol-2-one (=N-[2-(1,3-Dibutyl-2-oxo-5-phenyl-2,3-dihydro-1H-imidazol-4-yl)phenyl]acetamide; **8k**). Prepared from **6k** (*Method* D, besides **4k** and unreacted **6k**). Colorless crystals. M.p. 135 – 137° (hexane). IR: 3290, 3074, 2956, 2936, 2871, 1679, 1604, 1580, 1526, 1454, 1443, 1406, 1372, 1296, 1257, 1209, 1115, 1074, 1009, 944, 869, 769, 760, 703, 677, 649, 597, 550, 535. ¹H- and ¹³C-NMR: see *Table 4*. APCI-MS (pos.): 406 (100, $[M + H]^+$), 364 (3, $[M + H - NCO]^+$). APCI-MS/MS (pos.) of *m*/z 406: 364 (100, $[M + H - NCO]^+$), 332 (7, $[M + H - NCO - BuNCO]^+$). APCI-MS (neg.): 404 (100, $[M - H]^-$). APCI-MS/MS (neg.) of *m*/z 404: 362 (5, $[M - H - NCO]^-$), 348 (32, $[M - H - NCO - BuNCO - Bu]^-$). APCI-MS/MS (neg.) of *m*/z 404: 362 (5, $[M - H - NCO]^-$), 206 (8, $[M - H - NCO - BuNCO - Bu]^-$). Anal. calc. for C₂₅H₃₁N₃O₂ (405.53): C 74.04, H 7.70, N 10.36; found: C 74.18, H 7.56, 10.29.

2.13. 3-Butyl-1-phenyl-1-(2-phenyl-1H-indol-3-yl)urea (41). Prepared from 11 (Method A) and from 61 (Method D). Colorless crystals. M.p. $230-232^{\circ}$ (AcOH), identical in all respects to authentic 41 of m.p. $226-230^{\circ}$, prepared from 31 [9]. Two side products were isolated besides 41 from the reaction of 11 with diphenylurea: compound of m.p. $224-229^{\circ}$ (benzene/hexane), identical in all respects to synthetic 71 (Section 6.2) and compound of m.p. $260-265^{\circ}$, identical in all respects to authentic 51, prepared from 31 [9].

2.14. 3-Benzyl-1-butyl-1-(2-phenyl-IH-indol-3-yl)urea (4m). Prepared from 3m (*Method E*, besides 6m and 8m and *Method B*, besides 5m and 6m) and from 6m (*Method D*, besides 6m and 8m). Colorless crystals. M.p. 164–166° (benzene/hexane). IR: 3419, 3061, 2955, 2928, 2969, 1638, 1603, 1584, 1512, 1452, 1418, 1383, 1333, 1296, 1227, 1167, 1113, 1027, 965, 827, 763, 738, 719, 693, 612, 563. ¹H- and ¹³C-NMR: see *Table 3*. APCI-MS (pos.): 398 (100, $[M + H]^+$), 381 (3, $[M + H - NH_3]^+$), 291 (25, $[M + H - NH_3 - MeC_6H_3]^+$). APCI-MS/MS (pos.) of *m*/*z* 398: 381 (33, $[M + H - NH_3]^+$), 291 (100, $[M + H - NH_3 - MeC_6H_3]^+$), 235 (17, $[M + H - NH_3 - MeC_6H_3 - butene]^+$), 193 (4, $[M + H - NH_3 - MeC_6H_3 - butene - NCO]^+$). APCI-MS (neg.): 396 (100, $[M - H]^-$), 261 (9, $[M - H - C_6H_6 - Bu]^-$), 233 (7, $[M - H - C_6H_6 - Bu - CO]^-$). APCI-MS/MS (neg.) of *m*/*z* 396: 339 (17, $[M - H - Bu]^-$), 295 (18, $[M - H - BuNHCOH]^-$), 263 (77, $[M - H - Bu - C_6H_6]^-$), 206 (100, $[M - H - 2 Bu - C_6H_6]^-$). Anal. calc. for $C_{26}H_{27}N_3O$ (397.51): C 78.56, H 6.85, N 10.57; found: C 78.41, H 6.94, N 10.41.

2.15. 4-(2-Acetamidophenyl)-1-benzyl-3-butyl-5-phenyl-1,3-dihydro-2H-imidazol-2-one (= N-[2-(1-Benzyl-3-butyl-2-oxo-5-phenyl-2,3-dihydro-1H-imidazol-4-yl)phenyl]acetamide; **8m**). Prepared from **6m** (Method D, besides **4m** and **6m**). Colorless crystals. M.p. 120–121° (hexane). IR: 3410, 3315, 3063, 3032,

Position	7f		71		8k		8m		8n	
	δ_{H}	$\delta_{\rm C}$	$\delta_{ m H}$	δ_{C}	$\delta_{ m H}$	$\delta_{ m C}$	$\delta_{ m H}$	$\delta_{\rm C}$	$\delta_{ m H}$	δ_{C}
2	_	152.5	-	152.2	-	152.9	-	153.1	_	152.5
4	_	113.9	-	115.8	-	116.5	-	117.3	-	117.0
5	_	122.7	-	123.0	-	121.1	-	121.2	-	122.4
1'	_	118.2	-	117.8	-	121.1	-	121.4	-	120.4
2'	_	138.9	-	138.8	-	137.8	-	137.9	-	137.9
3'	8.06	119.6	7.90	119.6	7.79	123.7	7.76	124.0	7.66	122.9
4'	7.27	129.1	7.21	129.1	7.35	129.4	7.35	129.1	7.17	129.3
5'	6.95	121.8	6.85	121.6	7.12	124.0	7.14	124.2	6.91	123.5
6'	7.11	132.0	7.13	132.5	7.24	132.4	7.26	132.4	7.03	132.1
$1'(\mathbf{R}^{1})$	2.42	22.8	-	128.8	-	129.0	-	129.1	-	128.9
$2'(R^1)$	1.50	30.1	^a)	129.4	7.29	127.2	7.12	126.5	^b)	^b)
3' (R ¹)	1.23	21.6	^a)	128.5	7.34	128.4	7.32	128.4	^b)	^b)
4' (R ¹)	0.74	13.4	^a)	128.5	7.34	127.9	7.24	127.0	^b)	^b)
$1'(R^2)$	3.34	27.7	3.86, 3.60	40.1	3.73, 3.60	40.7°)	4.97, 4.78	44.5	5.02, 4.84	44.7
$2'(R^2)$	_	_	1.52	30.9	1.38	30.8°)	-	138.2	-	137.9
$3'(R^2)$	_	_	1.23	19.4	1.15	19.2°)	7.23	128.2	^b)	^b)
$4'(R^2)$	_	_	0.81	13.5	^d)	13.4°)	7.18	128.4	^b)	^b)
5' (R ²)	_	_	-	_	-	-	7.23	127.9	^b)	^b)
$1'(R^3)$	_	135.4	-	135.2	3.56, 3.31	40.7°)	3.62, 3.32	41.0	-	135.5
$2'(R^3)$	7.12	126.4	a)	126.8	1.38	30.8°)	1.41	30.8	^b)	^b)
$3'(R^3)$	7.23	128.4	a)	128.5	1.15	19.2°)	1.14	19.2	^b)	^b)
$4'(R^3)$	7.16	126.5	a)	126.9	^d)	13.4°)	0.75	13.4	^b)	^b)
C(2')-NHCO	9.14	152.1	9.03	151.9	8.96	168.3	9.10	168.3	9.09	167.6
N <i>H</i> Ph	7.85		7.84							
NHPh or NHMe	-	139.6	-	139.6	1.96	23.6	1.98	23.6	1.86	23.7
	7.44	118.3	7.42	118.5						
	7.29	128.9	7.33	128.9						
	7.02	122.0	7.03	122.0						

Table 4. ¹*H*- and ¹³*C*-Chemical Shifts (δ [ppm]) of Compounds 7 and 8 in (D_6)DMSO

^a) Strong overlap of ¹H resonances (δ (¹H) = 7.16–7.35). ^b) Strong overlap of ¹H and ¹³C resonances of three Ph groups, δ (¹H) = 7.16–7.36 (*m*, 15 H), δ (¹³C) = 129.9, 128.5, 128.4, 127.0, 126.9, 126.7. ^c) ¹³C Resonances of appropriate pairs in two Bu group signals differ maximally in 0.04 ppm. ^d) δ (¹H) = 0.77 or 0.72.

2957, 2930, 2871, 1678, 1606, 1580, 1530, 1497, 1451, 1402, 1368, 1299, 1252, 1209, 1030, 975, 941, 764, 752, 724, 701, 659, 598, 550. ¹H- and ¹³C-NMR: see *Table 4*. APCI-MS (pos.): 440 (100, $[M + H]^+$). APCI-MS/MS (pos.) of *m*/*z* 440: 422 (10, $[M + H - H_2O]^+$), 398 (9, $[M + H - NCO]^+$), 349 (100, $[M + H - MeC_6H_4]^+$), 332 (6, $[M + H - MeC_6H_4 - NH_3]^+$), 307 (21, $[M + H - MeC_6H_4 - NCO]^+$). APCI-MS (neg.): 438 (100, $[M - H]^-$). APCI-MS/MS (neg.) of *m*/*z* 438: 382 (5, $[M - H - Bu]^-$), 347 (43, $[M - H - MeC_6H_5]^+$), 304 (100, $[M - H - MeC_6H_5 - NHCO]^+$), 291 (4, $[M - H - MeC_6H_5 - butene]^+$), 262 (5, $[M - H - MeC_6H_5 - NCO]^+$). Anal. calc. for C₂₈H₂₉N₃O₂ (439.55): C 76.51, H 6.65, N 9.56; found: C 76.35, H 6.69, N 9.39.

2.16. *3-Benzyl-1-phenyl-1-(2-phenyl-1*H-*indol-3-yl)urea* (**4n**). Prepared from **3n** (*Method B*, besides **5n** and **6n** or *Method E*), and from **6n** (*Method D*, besides **8n** and **6n**). Colorless crystals. M.p. 222–224° (benzene/hexane). IR: 3427, 3342, 3061, 3027, 2949, 1669, 1620, 1584, 1512, 1491, 1454, 1374, 1321, 1284, 1244, 1151, 1113, 1075, 1028, 918, 741, 696, 632, 562, 517. ¹H- and ¹³C-NMR: see *Table 3*. APCI-MS

(pos.): 418 (100, $[M + H]^+$), 401 (3, $[M + H - NH_3]^+$), 311 (11, $[M + H - NH_3 - MeC_6H_3]^+$), 283 (51, $[M + H - NH_3 - MeC_6H_3 - CO]^+$). APCI-MS/MS (pos.) of *m*/*z* 418: 401 (52, $[M + H - NH_3]^+$), 375 (8, $[M + H - NHCO]^+$), 311 (100, $[M + H - NH_3 - MeC_6H_3]^+$), 285 (54, $[M + H - C_6H_5CH_2NCO]^+$). APCI-MS (neg.): 416 (100, $[M - H]^-$), 282 (41, $[M - H - Ph - Bu]^-$), 255 (3, $[M - H - Ph - Bu - HCN]^-$). APCI-MS/MS (neg.) of *m*/*z* 416: 283 (100, $[M - H - Ph - butene]^-$). Anal. calc. for $C_{28}H_{23}N_3O$ (417.50): C 80.55, H 5.55, N 10.06; found: C 80.42, H 5.67, N 10.11.

2.17. 4-(2-Acetamidophenyl)-1-benzyl-3,5-diphenyl-1,3-dihydro-2H-imidazol-2-one (= N-[2-(1-Benzyl-2-oxo-3,5-diphenyl-2,3-dihydro-1H-imidazol-4-yl)phenyl]acetamide; **8n**). Prepared from **6n** (Method D, besides **4n** and **6n**). Colorless crystals. M.p. 237–241° (benzene/hexane). IR: 3341, 3335, 3061, 2953, 1677, 1595, 1579, 1520, 1493, 1453, 1383, 1296, 1253, 1231, 1158, 1125, 1042, 1015, 957, 887, 820, 781, 763, 747, 736, 710, 700, 651, 592, 519, 502. ¹H- and ¹³C-NMR: see *Table 4*. APCI-MS (pos.): 460 (100, $[M + H]^+$), 416 (15, $[M + H - NCO]^+$). APCI-MS/MS (pos.) of m/z 460: 442 (50, $[M + H - H_2O]^+$), 418 (32, $[M + H - NCO]^+$), 401 (9, $[M + H - NCO - NH_3]^+$), 369 (100, $[M + H - MeC_6H_4]^+$), 352 (12, $[M + H - MeC_6H_4 - NH_3]^+$), 327 (48, $[M + H - MeC_6H_4 - NCO]^+$), 309 (11, $[M + H - MeC_6H_4 - NCO]^-$). APCI-MS/MS (neg.) of m/z 458: 416 (12, $[M - H - NCO]^-$), 325 (55, $[M - H - C_6H_5CH_2NCO]^-$), 282 (100, $[M - H - C_6H_5CH_2NCO - NCO]^-$). Anal. calc. for $C_{30}H_{25}N_3O_2$ (459.54): C 78.41, H 5.48, N 9.14; found: C 78.29, H 5.62, N 9.09.

3. General Procedure for the Preparation of 4-(2-Aminophenyl)-1,3-dihydro-2H-imidazol-2-ones (6) from Compounds 2, 3, 4, and 8 (Method C). A soln. of respective starting compound 2, 3, 4, and 8 (2.5 mmol) in conc. HCl (25 ml) was heated to reflux for the time given in Table 2. If necessary, AcOH was added to dissolve poorly soluble starting compound. After cooling, the soln. was evaporated *in vacuo* to dryness. The residue was dissolved in pyridine (5 ml), anh. K_2CO_3 (350 mg) was added and the mixture was stirred for 1 h. The insoluble material was filtered off, the filtrate was evaporated to dryness, and the residue was crystallized from an appropriate solvent or separated by CC on silica gel.

3.1. 5-(2-Aminophenyl)-1,4-dibutyl-1,3-dihydro-2H-imidazol-2-one (**6a**). Prepared from **2a** and **3a**. Colorless crystals. M.p. $134-139^{\circ}$ (benzene/cyclohexane), identical in all respects to authentic **6a** of m.p. $135-140^{\circ}$, prepared from **4a** [9].

3.2. 5-(2-Aminophenyl)-4-butyl-1-phenyl-1,3-dihydro-2H-imidazol-2-one (**6b**). Prepared from **2b**. Colorless crystals. M.p. $134-140^{\circ}$ (without recrystallization), identical in all respects to authentic **6b** of m.p. $138-141^{\circ}$, prepared from **4b** [9].

3.3. 5-(2-Aminophenyl)-1-butyl-4-phenyl-1,3-dihydro-2H-imidazol-2-one (**6c**). Prepared from **2c**, **3c**, and **4c**. Colorless crystals. M.p. 167–169° (benzene/hexane). IR: 3457, 3345, 3142, 3057, 2959, 2933, 2871, 1675, 1615, 1604, 1574, 1506, 1490, 1453, 1403, 1376, 1361, 1310, 1264, 1158, 1115, 1077, 1031, 969, 935, 908, 751, 693, 665, 520, 498. ¹H- and ¹³C-NMR: see *Table 5*. APCI-MS (pos.): 308 (100, $[M + H]^+$), 291 (59, $[M + H - NH_3]^+$), 263 (28, $[M + H - COOH]^+$). APCI-MS/MS (pos.) of *m/z* 308: 291 (100, $[M + H - NH_3]^+$), 265 (15, $[M + H - NHCO]^+$), 252 (21, $[M + H - butene]^+$), 235 (22, $[M + H - NH_3 - butene]^+$), 209 (29, $[M + H - BuNCO]^+$). APCI-MS (neg.): 306 (100, $[M - H]^-$), 261 (11, $[M - H - COOH]^-$), 233 (4, $[M - H - BuNH_2]^-$). APCI-MS/MS (neg.) of *m/z* 306: 263 (82, $[M - H - NHCO]^-$), 249 (92, $[M - H - Bu]^-$), 206 (100, $[M - H - Bu - NHCO]^-$). Anal. calc. for C₁₉H₂₁N₃O (307.39): C 74.24, H 6.89, N 13.67; found: C 74.29, H 6.98, N 13.53.

3.4. 5-(2-Aminophenyl)-1,4-diphenyl-1,3-dihydro-2H-imidazol-2-one (**6d**). Prepared from **2d** and **4d**. Colorless crystals. M.p. 297–309° (benzene/dioxane). IR: 3432, 3333, 3134, 3061, 3030, 2978, 2878, 2762, 1674, 1622, 1596, 1492, 1454, 1392, 1309, 1256, 1151, 1072, 1030, 915, 853, 776, 764, 747, 709, 693, 664, 649, 549, 504. ¹H- and ¹³C-NMR: see *Table 5*. APCI-MS (pos.): 328 (100, $[M + H]^+$), 311 (4, $[M + H-NH_3]^+$), 283 (99, $[M + H-COOH]^+$), 208 (3, $[M + H-COOH-C_6H_3]^+$). APCI-MS/MS (pos.) of *m*/*z* 328: 311 (82, $[M + H-NH_3]^+$), 285 (78, $[M + H-NHCO]^+$), 235 (48, $[M + H-NH_3-C_6H_4]^+$), 209 (100, $[M + H-NHCO-C_6H_4]^+$), 182 (51, $[M + H-NHCO-C_6H_4-HCN]^+$). APCI-MS (neg.): 326 (66, $[M - H]^-$), 282 (100, $[M - H-NH_2CO]^-$), 255 (31, $[M - H-NH_2CO - HCN]^-$). APCI-MS/MS (neg.) of *m*/*z* 326: 283 (100, $[M - H-NHCO]^-$). Anal. calc. for C₂₁H₁₇N₃O (327.38): C 77.04, H 5.23, N 12.84; found: C 76.85, H 5.35, N 12.71.

3.5. 4-(2-Aminophenyl)-3,5-dibutyl-1-methyl-1,3-dihydro-2H-imidazol-2-one (6e). Prepared from 3e and 4e. Colorless crystals. M.p. 74 – 78° (hexane). IR: 3401, 3337, 3226, 3028, 2957, 2931, 2866, 1668, 1632,

Position	6c		6d		6e		6g		6k		em (6n	
	$\delta_{ m H}$	δ_{C}	$\delta_{\rm H}$	δ_{C}	$\delta_{ m H}$	δ_{C}	$\delta_{ m H}$	δ_{C}	$\delta_{\rm H}$	δ_{C}	$\delta_{\rm H}$	δ_{C}	$\delta_{ m H}$	δ_{C}
	10.72	I	11.00	I	1	Т	I	I	I	I	1	T	I	Т
2	I	153.5	I	153.0	I	153.1	I	152.9	I	152.9	I	153.2	I	152.7
4	I	117.8	I	117.9	I	115.4	I	115.6	I	117.7	I	118.2	I	118.0
5	I	117.3	I	118.6	I	119.8	I	119.3	I	120.6	I	120.8	I	122.2
1′	Ι	113.2	I	113.3	I	112.8	I	113.0	Ι	112.7	I	112.4	I	112.5
2,	I	148.1	I	148.2	I	148.0	I	147.8	I	148.3	I	148.3	I	148.1
3,	6.82	114.8	6.65	114.5	6.76	114.5	6.76	114.5	6.68	114.5	6.70	114.5	6.50	114.3
4	7.21	130.3	7.34	130.0	7.14	129.6	7.13	129.6	7.07	128.9	7.08	129.9	6.92	128.4
5'	6.55	116.1	6.45	115.8	6.61	115.7	6.62	115.8	6.52	115.7	6.52	115.7	6.37	115.4
6′	7.03	132.1	7.04	132.3	6.96	132.1	6.97	132.1	6.97	132.7	6.98	132.7	6.96	132.8
\mathbf{NH}_2	5.05	I	5.18	I	4.91	I	4.86	I	4.97	I	5.05	I	5.08	I
$1'(R^1)$	I	130.2	I	130.0	2.32, 2.23	22.8	2.33, 2.25	22.7	I	129.8	I	129.4	I	129.1
2' (R ¹)	7.26	124.4	7.28	124.9	1.40	30.6	(q	(q	7.33	128.3	7.11	126.6	с)	с)
3′ (R ¹)	7.22	128.3	7.27	128.4	1.27	21.5	(q	(q	7.33	129.5	7.31	128.4	c)	c)
4' (R ¹)	7.13	126.0	7.22	126.6	0.78	13.5	(p	(p	7.30	127.7	7.25	127.7	с)	с)
$1'(R^2)$	I	I	I	I	3.19	27.2	3.58	40.5	3.67, 3.60	40.7	4.93, 4.78	44.6	4.99, 4.82	44.8
$2'(R^2)$	I	I	I	I	I	I	1.62	(q	1.38	30.9	I	138.4	I	138.0
3′ (R ²)	I	I	I	I	I	I	(q	(q	1.15	19.2	7.23	128.2	с)	с)
$4'(R^2)$	I	I	I	I	I	I	(p	(p	0.76	13.4	7.18	129.5	c)	c)
5' (R ²)	I	I	I	I	I	I	- 1	1	Ι	I	7.23	127.0	c)	c)
$1'(R^3)$	3.44, 3.26	39.9	I	135.5	3.44, 3.27	40.4	3.44, 3.26	40.3	3.55, 3.30	40.5	3.62, 3.34	42.4	- 1	135.6
$2'(R^3)$	1.37	30.9	7.35	127.4	1.31	30.9	(q	(q	1.38	30.9	1.41	31.0	с)	с)
3' (R ³)	1.15	19.3	7.27	128.2	1.09	19.2	(q	(q	1.15	19.2	1.16	19.3	c)	c)
4' (R ³)	0.76	13.4	7.19	126.5	0.72	13.4	(p	(p	0.76	13.4	0.78	13.5	c)	с)
^a) Numbe ^b) Strong e three Ph g ^d) Resonau	ring as indica overlap of ¹ H groups, $\delta(^{1}H)$ nces of three	tread in <i>S</i> resonant = 7.16 - Me grou	cheme I. ces of CI 7.36 $(m, ps: \delta(^{1}H)$	This nuu H_2 group 15 H), δ) = 0.96,	mbering does s, $\delta(^{1}H) = 1.0$ $\delta(^{13}C) = 129.6$ 0.77 and 0.74	(not matrix) 7 - 1.40 (2 C), (2 C),	tch with the $(m, 10 \text{ H}), \delta((m, 10 \text{ H})), \delta((m, 128.4 (2 \text{ C})), 2)) = 13.8, 13.5$	systemat ${}^{13}C) = 3.5$ 128.3 (2 and 13.4.	ic names giv l.1, 30.9, 21.6 C), 128.2 (2	en in th , 19.6, 19 C), 128.	e <i>Exper. Pari</i> .5. ^c) Strong 1, 127.0, 126.	for con overlap 9 (2 C),	ıpounds 6c ε of ¹ H resona 126.8 (2 C).	nd 6d . 1ces of 125.7.

Table 5. ¹*H*- and ¹³*C*-*Chemical Shifts* (δ [ppm]) of *Compounds* **6** in $(D_6)DMSO^a$)

1567, 1493, 1457, 1401, 1370, 1317, 1265, 1152, 1081, 1023, 950, 940, 825, 749, 656, 599, 552. ¹H- and ¹³C-NMR: see *Table 5*. APCI-MS (pos.): 302 (100, $[M + H]^+$), 256 (7, $[M + H - HCOOH]^+$). APCI-MS/MS (pos.) of *m*/*z* 302: 284 (10, $[M + H - H_2O]^+$), 258 (100, $[M + H - HH_2O]^+$), 246 (98, $[M + H - butene]^+$), 201 (85, $[M + H - NH_2CO - Bu]^+$), 189 (78, $[M + H - butene - Bu]^+$), 172 (97, $[M + H - butene - Bu - NH_3]^+$). APCI-MS (neg.): 300 (100, $[M - H]^-$). APCI-MS/MS (neg.) of *m*/*z* 300: 257 (3, $[M - H - NHCO]^-$), 243 (100, $[M - H - Bu]^-$), 199 (13, $[M - H - Bu - NH_2CO]^-$), 186 (15, $[M - H - 2 Bu]^-$). Anal. calc. for $C_{18}H_{27}N_3O$ (301.43): C 71.72, H 9.03, N 13.94; found: C 71.52, H 9.14, N 13.87.

3.6. 4-(2-Aminophenyl)-5-butyl-1-methyl-3-phenyl-1,3-dihydro-2H-imidazol-2-one (6f). Prepared from **3f**. Colorless crystals. M.p. $155-160^{\circ}$ (benzene/hexane), identical in all respects to authentic 6f of m.p. $154-162^{\circ}$, prepared from **4f** [9].

3.7. 4-(2-Aminophenyl)-1,3,5-tributyl-1,3-dihydro-2H-imidazol-2-one (**6**g). Prepared from **3**g and **4**g. Yellowish oil. IR: 3464, 3331, 2957, 2932, 2871, 1676, 1640, 1573, 1494, 1457, 1412, 1368, 1310, 1260, 1157, 1112, 939, 749, 653, 555. ¹H- and ¹³C-NMR: see *Table 5*. APCI-MS (pos.): 344 (100, $[M + H]^+$), 298 (41, $[M + H - HCOOH]^+$), 271 (6, $[M + H - BuNH_2]^+$), 243 (9, $[M + H - BuNHCOH]^+$). APCI-MS/MS (pos.) of *m*/*z* 344: 300 (4, $[M + H - NH_2CO]^+$), 288 (4, $[M + H - butene]^+$), 271 (20, $[M + H - BuNH_2]^+$), 243 (100, $[M + H - BuNHCOH]^+$). APCI-MS (neg.): 342 (100, $[M - H]^-$), 312 (18, $[M - H - CH_2O]^-$), 241 (11, $[M - H - BuNHCOH]^-$). APCI-MS/MS (neg.) of *m*/*z* 342: 285 (12, $[M - H - Bu]^-$), 243 (100, $[M - H - BuNCO]^-$), 186 (92, $[M - H - BuNCO - Bu]^-$). Anal. calc. for C₂₁H₃₃N₃O (343.51): C 73.43, H 9.68, N 12.23; found: C 73.52, H 9.59, N 12.37.

3.8. 4-(2-Aminophenyl)-1,5-dibutyl-3-phenyl-1,3-dihydro-2H-imidazol-2-one (6h). Prepared from **3h**. Colorless crystals. M.p. 115–121° (benzene/hexane), identical in all respects to authentic **3h** of m.p. 115–122°, prepared from **4h** [9].

3.9. 4-(2-Aminophenyl)-1-benzyl-3,5-dibutyl-1,3-dihydro-2H-imidazol-2-one (**6i**). Prepared from **3i**. Colorless crystals. M.p. $98-104^{\circ}$ (hexane), identical in all respects to authentic **6i** of m.p. $98-105^{\circ}$, prepared from **4i** [9].

3.10. 4-(2-Aminophenyl)-1-benzyl-5-butyl-3-phenyl-1,3-dihydro-2H-imidazol-2-one (**6j**). Prepared from **3j**. Colorless crystals. M.p. $87-93^{\circ}$ (hexane), identical in all respects to authentic **6j** of m.p. $88-94^{\circ}$, prepared from **4j** [9].

3.11. 4-(2-Aminophenyl)-1,3-dibutyl-5-phenyl-1,3-dihydro-2H-imidazol-2-one (**6k**). Prepared from **3k**, **4k**, and **8k**. Colorless crystals. M.p. 77–81° (hexane). IR: 3466, 3424, 3336, 3229, 3059, 2953, 2930, 2867, 1663, 1625, 1602, 1573, 1491, 1457, 1406, 1366, 1312, 1262, 1209, 1159, 1114, 1077, 1027, 938, 805, 749, 701, 664, 651, 624, 550, 513. ¹H- and ¹³C-NMR: see *Table 5*. APCI-MS (pos.): 364 (100, $[M + H]^+$), 318 (15, $[M + H - HCOOH]^+$), 263 (10, $[M + H - BuNHCOH]^+$). APCI-MS/MS (pos.) of *m*/*z* 364: 346 (8, $[M + H - H_2O]^+$), 308 (38, $[M + H - butene]^+$), 291 (9, $[M + H - BuNH_2]^+$), 263 (100, $[M + H - BuNHCOH]^+$). APCI-MS (neg.) of *m*/*z* 362: 306 (45, $[M - H - butene]^-$), 263 (100, $[M - H - BuNCO - Bu]^-$). APCI-MS/MS (neg.) of *m*/*z* 362: 306 (45, $[M - H - butene]^-$), 263 (100, $[M - H - BuNCO]^-$), 206 (71, $[M - H - BuNCO - Bu]^-$). Anal. calc. for C₂₃H₂₉N₃O (363.50): C 76.00, H 8.04, N 11.56; found: C 75.87, H 7.87, N 11.42.

3.12. 4-(2-Aminophenyl)-1-butyl-3,5-diphenyl-1,3-dihydro-2H-imidazol-2-one (61). Compound was prepared from 31. Colorless crystals. M.p. $65-73^{\circ}$ (without recrystallization), identical in all respects to authentic compound of m.p. $54-57^{\circ}$, prepared from 41 [9].

3.13. 4-(2-Aminophenyl)-1-benzyl-3-butyl-5-phenyl-1,3-dihydro-2H-imidazol-2-one (**6m**). Prepared from **3m**, **4m**, and **8m**. Colorless crystals. M.p. 114–115° (benzene/hexane). IR: 3451, 3325, 3203, 3063, 3029, 2954, 2932, 2872, 1672, 1622, 1600, 1572, 1489, 1448, 1398, 1356, 1308, 1265, 1155, 1075, 1028, 951, 921, 775, 749, 702, 831, 577, 501. ¹H- and ¹³C-NMR: see *Table 5*. APCI-MS (pos.): 398 (100, $[M + H]^+$), 352 (10, $[M + H - HCOOH]^+$). APCI-MS/MS (pos.) of m/z 398: 381 (25, $[M + H - NH_3]^+$), 355 (25, $[M + H - NHCO]^+$), 307 (100, $[M + H - MeC_6H_4]^+$), 263 (29, $[M + H - MeC_6H_4 - NH_2CO]^+$), 208 (33, $[M + H - MeC_6H_4 - BuNCO]^+$). APCI-MS (neg.): 396 (100, $[M - H]^-$). APCI-MS/MS (neg.) of m/z 396: 340 (31, $[M - H - butene]^-$), 263 (100, $[M - H - butene - C_6H_5]^-$), 219 (15, $[M - H - butene - C_6H_5 - NH_2CO]^-$), 206 (42, $[M - H - butene - C_6H_5Bu]^-$). Anal. calc. for $C_{26}H_{27}N_3O$ (397.51): C 78.56, H 6.85, N 10.57; found: C 78.41, H 6.94, N 10.39.

3.14. 4-(2-Aminophenyl)-1-benzyl-3,5-diphenyl-1,3-dihydro-2H-imidazol-2-one (**6n**). Prepared from **3n**, **4n**, and **8n**. Colorless crystals. M.p. 209–212° (benzene/hexane). IR: 3465, 3366, 3061, 3028, 2926,

1687, 1615, 1600, 1572, 1498, 1453, 1427, 1386, 1370, 1310, 1241, 1157, 1130, 1072, 1026, 953, 917, 824, 784, 763, 745, 701, 660, 587, 544, 520. ¹H- and ¹³C-NMR: see *Table 5*. APCI-MS (pos.): 418 (100, $[M + H]^+$), 373 (16, $[M + H - COOH]^+$). APCI-MS/MS (pos.) of *m*/*z* 418: 401 (11, $[M + H - H_2O]^+$), 375 (12, $[M + H - NHCO]^+$), 327 (100, $[M + H - MeC_6H_4]^+$), 283 (52, $[M + H - MeC_6H_4 - NH_2CO]^+$). APCI-MS (neg.): 416 (100, $[M - H]^-$). APCI-MS/MS (neg.) of *m*/*z* 416: 325 (100, $[M - H - MeC_6H_4]^-$), 283 (48, $[M - H - MeC_6H_4 - NH_2CO]^-$). Anal. calc. for C₂₈H₂₃N₃O (417.50): C 80.55, H 5.55, N 10.06; found: C 80.41, H 5.67, 10.12.

4. General Procedure for the Preparation of 4-(2-Acetamidophenyl)-5-phenyl-1,3-dihydro-2Himidazol-2-ones (8). A mixture of 6 (0.1 mmol), Ac₂O (1 ml), and pyridine (1 ml) was left at r.t. for 4 h, evaporated to dryness *in vacuo*, and the residue was crystallized from an appropriate solvent. The following compounds 8, identical in all respects to those described in *Sect. 3.2*, were prepared: 8k (86% yield), 8m (92 % yield), and 8n (96% yield).

5. General Procedure for the Preparation of 1,3-Bis[2-(2-oxo-2,3-dihydro-1H-imidazol-4-yl)phenyl]ureas (5). Compounds 5c, 5e, 5g, 5k, 5m, and 5n were prepared from the corresponding compounds 6 and triphosgene according to the same procedure as was reported for preparation of compounds 5a, 5f, 5h, 5i, 5j, and 5l [9].

5.1. *1,3-Bis[2-(3-butyl-2-oxo-5-phenyl-2,3-dihydro-1*H-*imidazol-4-yl)phenyl]urea* (**5c**). Prepared from **6c** in 12% yield. Colorless crystals. M.p. 176–186° (AcOEt/hexane), identical in all respects to authentic **6c** of m.p. 173–179°, prepared from **3c** [9].

5.2. 1,3-Bis[2-(3,5-dibutyl-1-methyl-2-oxo-2,3-dihydro-1H-imidazol-4-yl)phenyl]urea (5e). Prepared from 6e in 18% yield. Yellowish oil, identical in all respects to authentic 5e prepared from 3e [9].

5.3. 1,3-Bis[2-(1,3,5-tributyl-2-oxo-2,3-dihydro-IH-imidazol-4-yl)phenyl]urea (5g). Prepared from 6g in 55% yield. Yellowish oil, identical in all respects to authentic 5g prepared from 3g [9].

5.4. 1,3-Bis[2-(1,3-dibutyl-2-oxo-5-phenyl-2,3-dihydro-1H-imidazol-4-yl)phenyl]urea (**5k**). Prepared from **6k** in 65% yield. Colorless crystals. M.p. $173 - 176^{\circ}$ (hexane), identical in all respects to authentic **5k** of m.p. $154 - 164^{\circ}$, prepared from **3k** [9].

5.5. 1,3-Bis[2-(1-benzyl-3-butyl-2-oxo-5-phenyl-2,3-dihydro-1H-imidazol-4-yl)phenyl]urea (5m). Prepared from 6m in 68% yield. Colorless crystals. M.p. $94-98^{\circ}$ (hexane), identical in all respects to authentic 5m of m.p. $95-100^{\circ}$, prepared from 3m [9].

5.6. 1,3-Bis[2-(1-benzyl-2-oxo-3,5-diphenyl-2,3-dihydro-1H-imidazol-4-yl)phenyl]urea (**5n**). Prepared from **6n** in 78% yield. Colorless crystals. M.p. 242–248°, identical in all respects to authentic **5n** of m.p. 227–232°, prepared from **3n** [9].

6. General Procedure for the Preparation of Ureas **7** from Dihydroimidazolones **6**. Phenylisocyanate (0.03 ml, 0.28 mmol) was added under stirring at r.t. to the soln. of **6** (0.25 mmol) in CHCl₃ (2 ml). The mixture was stirred for 2.5 h at r.t., evaporated *in vacuo* to dryness, and the residue was crystallized from benzene/hexane (1:1, v/v).

6.1. I-[2-(5-Butyl-1-methyl-2-oxo-3-phenyl-2,3-dihydro-1H-imidazol-4-yl)phenyl]-3-phenylurea (**7f**). Prepared from **6f** in 91% yield. Colorless crystals. M.p. 235–237° (benzene/hexane). IR: 3335, 3298, 2957, 2932, 2868, 1709, 1670, 1645, 1600, 1584, 1554, 1530, 1500, 1443, 1396, 1294, 1259, 1229, 1198, 889, 753, 684, 633, 535, 506. ¹H- and ¹³C-NMR: see *Table 4*. APCI-MS (pos.): 441 (100, $[M + H]^+$), 348 (38, $[M + H - PhNH_2]^+$), 322 (37, $[M + H - PhNCO]^+$). APCI-MS/MS (pos.) of m/z 441: 397 (13, $[M + H - NH_2CO]^+$), 348 (37, $[M + H - PhNH_2]^+$), 322 (98, $[M + H - PhNCO]^+$), 265 (21, $[M + H - PhNCO - Bu]^+$), 203 (7, $[M + H - 2 PhNCO]^+$), 172 (100, $[M + H - PhNCO - PhNH_2 - Bu]^+$). APCI-MS (neg.): 439 (42, $[M - H]^-$), 346 (9, $[M - H - PhNH_2]^-$), 320 (100, $[M - H - PhNCO]^-$). APCI-MS/MS (neg.) of m/z 439: 320 (100, $[M - H - PhNCO]^-$). Anal. calc. for $C_{27}H_{28}N_4O_2$ (440.54): C 73.61, H 6.41, N 12.72; found: C 73.52, H 6.24, N 12.67.

6.2. 1-[2-(1-Butyl-2-oxo-3,5-diphenyl-2,3-dihydro-1H-imidazol-4-yl)phenyl]-3-phenylurea (71). Prepared from **61** in 79% yield. Colorless crystals. M.p. 221–228° (benzene/hexane). IR: 3342, 3302, 3057, 2957, 2932, 2861, 1713, 1672, 1599, 1582, 1527, 1497, 1446, 1395, 1292, 1258, 1229, 1192, 1094, 755, 695, 656, 505. ¹H- and ¹³C-NMR: see *Table 4*. APCI-MS (pos.): 503 (100, $[M + H]^+$), 410 (28, $[M + H-PhNH_2]^+$), 384 (31, $[M + H-PhNCO]^+$). APCI-MS/MS (pos.) of m/z 503: 410 (100, $[M + H-PhNH_2]^+$), 384 (52, $[M + H-PhNCO]^+$), 354 (9, $[M + H-PhNH_2-butene]^+$), 285 (25, $[M + H-PhNCO]^+$). APCI-MS (neg.): 501 ($[M - H]^-$), 382 ($[M - H-PhNCO]^-$). APCI-MS/

MS (neg.) of m/z 501: 382 (100, $[M - H - PhNCO]^-$). Anal. calc. for $C_{32}H_{30}N_4O_2$ (502.61): C 76.47, H 6.02, N 11.15; found: C 76.57, H 6.19, N 11.02.

REFERENCES

- [1] A. Klásek, K. Kořistek, J. Polis, J. Košmrlj, Tetrahedron 2000, 56, 1551.
- [2] A. Klásek, K. Kořistek, S. Kafka, J. Košmrlj, Heterocycles 2003, 60, 1811.
- [3] S. Kafka, A. Klásek, J. Košmrlj, J. Org. Chem. 2001, 66, 6394.
- [4] A. Klásek, K. Kořistek, A. Lyčka, M. Holčapek, Tetrahedron 2003, 59, 1283.
- [5] A. Klásek, K. Kořistek, A. Lyčka, M. Holčapek, Tetrahedron 2003, 59, 5279.
- [6] A. Klásek, A. Lyčka, M. Holčapek, I. Hoza, Tetrahedron 2004, 60, 9953.
- [7] A. Klásek, A. Lyčka, M. Holčapek, M. Kovář, I. Hoza, J. Heterocycl. Chem. 2006, 43, 1251.
- [8] A. Klásek, A. Lyčka, M. Holčapek, I. Hoza, J. Heterocycl. Chem. 2006, 43, 203.
- [9] A. Klásek, A. Lyčka, M. Holčapek, Tetrahedron 2007, 63, 7059.
- [10] S. Hiremath, N. N. Gondar, M. G. Purohit, Indian J. Chem., Sect. B 1982, 21, 321.
- [11] S. Hiremath, P. Shivayogi, P. S. Badami, M. G. Purohit, Indian J. Chem., Sect. B 1985, 24, 1115.
- [12] S. Wendeborn, T. Winkler, I. Foisy, Tetrahedron Lett. 2000, 41, 6387, and refs. cit. therein.
- [13] G. Wagner, S. Laufer, Med. Res. Rev. 2006, 26, 1.
- [14] G. T. Bourne, D. Crich, J. W. Davies, D. C. Horwell, J. Chem. Soc., Perkin Trans. 1 1991, 1693.
- [15] M. Bruncko, D. Crich, J. Org. Chem. 1994, 59, 5543.
- [16] T. Hino, Yakugaku Zasshi 1996, 116, 566.
- [17] F. J. Sardina, H. Rapoport, Chem. Rev. 1996, 96, 1825.
- [18] K. M. Depew, S. P. Marsden, D. Zatorska, A. Zatorski, W. G. Bornmann, S. J. Danishefsky, J. Am. Chem. Soc. 1999, 121, 11953.
- [19] D. B. Grotjahn, C. Joubran, Tetrahedron: Asymmetry 1995, 6, 745.
- [20] F. Lortie, S. Boileau, L. Bouteiller, Chem. Eur. J. 2003, 9, 3008.

Received August 31, 2007