

Article

## Introduction of Quinolines and Isoquinolines onto Non-activated alpha-C-H Bond of Tertiary Amides through Radical Pathway

Naoki Okugawa, Katsuhiko Moriyama, and Hideo Togo

*J. Org. Chem.*, **Just Accepted Manuscript** • DOI: 10.1021/acs.joc.6b02303 • Publication Date (Web): 07 Dec 2016

Downloaded from <http://pubs.acs.org> on December 7, 2016

### Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.

# Introduction of Quinolines and Isoquinolines onto Non-activated $\alpha$ -C-H Bond of Tertiary Amides through Radical Pathway

Naoki Okugawa<sup>†</sup>, Katsuhiko Moriyama<sup>†‡</sup>, and Hideo Togo<sup>\*†</sup>

<sup>†</sup>Graduate School of Science, Chiba University, Yayoi-cho 1-33, Inage-ku, Chiba  
263-8522, Japan


<sup>‡</sup>Molecular Chirality Research Center, Chiba University, Yayoi-cho 1-33, Inage-ku,  
Chiba 263-8522, Japan

**RECEIVED DATE (to be automatically inserted after your manuscript is accepted  
if required according to the journal that you are submitting your paper to)**

CORRESPONDING AUTHOR FOOTNOTE: E-mail: [togo@faculty.chiba-u.jp](mailto:togo@faculty.chiba-u.jp),

Tel: 81-43-290-2792, Fax: 81-43-290-2792


## ABSTRACT



Treatment of quinolines and isoquinolines with benzoyl peroxide in tertiary amides, such as *N,N*-dimethylacetamide, *N,N*-dimethylpropionamide, and *N*-acetylpyrrolidine, *etc.*, under irradiation with a Hg lamp in the temperature range of 35 °C ~ 40 °C gave C-C-bonded quinolines and isoquinolines bearing amide groups with high regioselectivity in good to moderate yields, respectively, under transition-metal-free conditions.

## INTRODUCTION

Synthetic studies of electron-deficient heteroaromatic bases, such as quinoline, isoquinoline, phenanthridine, *etc.*, bearing amide functional groups are very attractive and important because most of the heteroaromatic bases possess biological activities, *e. g.*, 5-chloro-2-hydroxy-*N*-(quinolin-8'-yl)benzamide **I** is a potent inhibitor of VEGFR-2 kinase,<sup>1a</sup> quinoline derivative **II** bearing an amide-*N*-methylene group at 8-position is a renin inhibitor for the treatment of hypertension,<sup>1b</sup> and 4-hydroxy-8-(methanesulfonylamino)quinoline-2-carboxamide **III** is an inhibitor of human plasma Lp-PLA2.<sup>1c</sup> The heteroaromatic bases also can be used as a luminescent probe, an example of which is phenanthridine derivative **IV** bearing an aza-crown-bonded amide-*N*-methylene group at 6-position is a luminescent probe of adenosine nucleotides, as shown in Fig. 1.<sup>1d</sup> Moreover, the amide groups can be easily transformed into amines by hydrolysis or reduction.



**Fig. 1** Quinolines and Phenanthridine Bearing Amide Group

Today, there are many methods for the introduction of alkyl groups onto electron-deficient heteroaromatic bases, such as pyridine, quinoline, *etc.*, through radical pathways,<sup>2</sup> typically, the decarboxylative alkylation of electron-deficient heteroaromatic bases with alkyl carboxylic acids, AgNO<sub>3</sub>, and K<sub>2</sub>S<sub>2</sub>O<sub>8</sub> under heating conditions (the Minisci reaction),<sup>3</sup> the decarboxylative alkylation of electron-deficient heteroaromatic bases with *N*-acyloxy-2-thiopyridone prepared from alkyl carboxylic acids and *N*-hydroxy-2-thiopyridone, under tungsten-lamp irradiation (the Barton decarboxylation),<sup>4</sup> and the introduction of alkyl groups onto electron-deficient heteroaromatic bases with alkyl halides and tris(trimethylsilyl)silane or


1,1,2,2-tetraphenyldisilane in the presence of AIBN under warming conditions, or with alkyl halides and tetra(trimethylsilyl)silane under irradiation with a Hg lamp.<sup>5</sup> The introduction of crown ether onto an electron-deficient heteroaromatic base by using crown ether and *t*-butyl hydrogen peroxide (TBHP) in the presence of FeSO<sub>4</sub> can be also carried out.<sup>6</sup> Those methods of introducing alkyl groups onto electron-deficient heteroaromatic bases through radical pathways are very attractive and useful because alkylated heteroaromatic bases that cannot be easily obtained by ionic electrophilic substitution (S<sub>E</sub>Ar) onto electron-deficient heteroaromatic bases using the Friedel-Crafts alkylation can be easily prepared. Recently, we reported a simple method for the introduction of ether, cyclic ether, and crown ether groups onto electron-deficient heteroaromatic bases with benzoyl peroxide (BPO) under heating and transition-metal-free conditions.<sup>7</sup> On the other hand, only a few studies of the introduction of a tertiary amide group onto electron-deficient heteroaromatic bases have been reported, *i.e.*, the introduction of an amide group onto lepidine and quinoxaline with H<sub>2</sub>SO<sub>4</sub> and hydroxylamine-*O*-sulfonic acid in the presence of FeSO<sub>4</sub>•7H<sub>2</sub>O in a mixture of formamide or DMF and water (10:1),<sup>8a</sup> the introduction of an amide group onto lepidine and quinoxaline with formamide or *N,N*-dimethylacetamide solvent in the presence of H<sub>2</sub>SO<sub>4</sub>, H<sub>2</sub>O<sub>2</sub>, and TiO<sub>2</sub> under sunlight irradiation,<sup>8b</sup> the preparation of 6-amidophenanthridine from 2-isocyanobiphenyl with amide and *t*-butyl peroxybenzoate (TBPB) at 120 °C,<sup>8c</sup> and the introduction of an amide group at 2-position of benzothiazoles with potassium persulfate in *N,N*-dimethylacetamide at 70 °C.<sup>8d</sup> Additionally, there are a few reports for amidation of arenes or heteroaromatic bases with formamides via formation of •CONH<sub>2</sub> (carbamoyl) radicals.<sup>9</sup> Here, we would like to report the introduction of quinolines and isoquinolines onto the non-activated α-C-H bond of tertiary amides with BPO under irradiation with a Hg lamp without using any transition metals.

## RESULTS AND DISCUSSION

First, a solution of lepidine **1a** (1.0 mmol) and BPO (2.0 mmol) in the presence of trifluoroacetic acid (1.0 mmol) in *N,N*-dimethylacetamide **A** (5 mL) in a 30 mL screw-capped flask was irradiated with a Hg lamp (400 W high pressure) in the temperature range of 35 °C ~ 40 °C to give 2-(*N*-acetyl,*N*-methylamino)methyl-4-methylquinoline **2aA** in 82% yield, as shown in

Table 1 (entry 1). When BPO was replaced with aq. hydrogen peroxide ( $\text{H}_2\text{O}_2$ ), *t*-butyl hydrogen peroxide (TBHP), di-*t*-butyl peroxide (DTBP), *t*-butyl peroxybenzoate (TBPB), and potassium peroxodisulfate ( $\text{K}_2\text{S}_2\text{O}_8$ ) under the same reaction conditions, the yield of compound **2aA** became low or moderate, and lepidine was recovered (entries 2~6, Table 1). When 3 equivalents of  $\text{K}_2\text{S}_2\text{O}_8$  was used under the same conditions as those in entry 6 in 5 mL and 8 mL of *N,N*-dimethylacetamide **A**, product **2aA** was obtained in moderate to low yields (entries 7 and 8, Table 1). This was because the solubility of  $\text{K}_2\text{S}_2\text{O}_8$  in *N,N*-dimethylacetamide **A** was quite low under the present reaction conditions, and a homogeneous solution was not obtained during the irradiation with  $\text{K}_2\text{S}_2\text{O}_8$ . In contrast, a homogeneous solution was maintained during

**Table 1.** Introduction of Lepidine **1a** onto *N,N*-Dimethylacetamide **A** to form **2aA**





Reaction scheme: Lepidine **1a** (1.0 mmol) + *N,N*-dimethylacetamide **A** (X mL)  $\xrightarrow[\text{Hg-}h\nu, 35-40^\circ\text{C}, 24\text{ h}]{\text{oxidant, Brønsted acid}}$  Product **2aA**.

entry	X	oxidant (equiv.)	Brønsted acid (equiv.)	yield (%)
1	5.0	BPO (2.0)	$\text{CF}_3\text{CO}_2\text{H}$ (1.0)	82
2	5.0	$\text{H}_2\text{O}_2$ (2.0)	$\text{CF}_3\text{CO}_2\text{H}$ (1.0)	49
3	5.0	TBHP (2.0)	$\text{CF}_3\text{CO}_2\text{H}$ (1.0)	10
4	5.0	DTBP (2.0)	$\text{CF}_3\text{CO}_2\text{H}$ (1.0)	8
5	5.0	TBPB (2.0)	$\text{CF}_3\text{CO}_2\text{H}$ (1.0)	14
6	5.0	$\text{K}_2\text{S}_2\text{O}_8$ (2.0)	$\text{CF}_3\text{CO}_2\text{H}$ (1.0)	68
7	5.0	$\text{K}_2\text{S}_2\text{O}_8$ (3.0)	$\text{CF}_3\text{CO}_2\text{H}$ (1.0)	56
8	8.0	$\text{K}_2\text{S}_2\text{O}_8$ (3.0)	$\text{CF}_3\text{CO}_2\text{H}$ (1.0)	36
9 <sup>a</sup>	5.0	BPO (2.0)	$\text{CF}_3\text{CO}_2\text{H}$ (1.0)	21
10 <sup>a</sup>	5.0	TBHP (2.0)	$\text{CF}_3\text{CO}_2\text{H}$ (1.0)	36
11	5.0	BPO (2.0)	$\text{TsOH} \cdot \text{H}_2\text{O}$ (1.0)	80
12	5.0	BPO (2.0)	$\text{CH}_3\text{SO}_3\text{H}$ (1.0)	81
13	5.0	BPO (2.0)	$\text{CF}_3\text{SO}_3\text{H}$ (1.0)	78
14	5.0	BPO (2.0)	$\text{CH}_3\text{CO}_2\text{H}$ (1.0)	trace
15	2.0	BPO (2.0)	$\text{CF}_3\text{CO}_2\text{H}$ (1.0)	67
16	4.0	BPO (2.0)	$\text{CF}_3\text{CO}_2\text{H}$ (1.0)	76
17	8.0	BPO (2.0)	$\text{CF}_3\text{CO}_2\text{H}$ (1.0)	87
18	8.0	BPO (2.0)	$\text{CF}_3\text{CO}_2\text{H}$ (1.3)	85
19	8.0	BPO (1.5)	$\text{CF}_3\text{CO}_2\text{H}$ (1.0)	86
20	8.0	BPO (1.7)	$\text{CF}_3\text{CO}_2\text{H}$ (1.0)	88
21	8.0	BPO (2.3)	$\text{CF}_3\text{CO}_2\text{H}$ (1.0)	81
22	8.0	—	$\text{CF}_3\text{CO}_2\text{H}$ (1.0)	14
23	8.0	BPO (1.7)	—	trace
24 <sup>b</sup>	8.0	BPO (1.7)	$\text{CF}_3\text{CO}_2\text{H}$ (1.0)	13
25 <sup>c</sup>	8.0	BPO (1.7)	$\text{CF}_3\text{CO}_2\text{H}$ (1.0)	0

<sup>a</sup>  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$  (0.3 equiv.) was added. <sup>b</sup> Reaction was carried out at 100 °C for 24 h without irradiation with a Hg lamp. <sup>c</sup> Reaction was carried out at 40 °C for 8 h under irradiation with a tungsten lamp.

the irradiation with BPO. The addition of  $\text{FeSO}_4$  as an electron transfer catalyst to a solution containing BPO (similar to the Fenton system) also reduced the yield of compound **2aA** (entry 9, Table 1). Moreover, when the acid was changed to a stronger acid than trifluoroacetic acid, such as toluenesulfonic acid, methanesulfonic acid, and trifluoromethanesulfonic acid, the yield of compound **2aA** almost did not change (entries 11~13, Table 1). When it was changed to a weaker acid than trifluoroacetic acid, such as acetic acid, compound **2aA** was not obtained at all (entry 14, Table 1). Thus, the complete protonation of lepidine to increase its reactivity toward nucleophilic radical species is required. Then, the effect of the amounts of solvent (2.0 mL, 4.0 mL, and 8.0 mL) and BPO (1.5 equiv., 1.7 equiv., and 2.3 equiv.) was studied, and it was found that treatment of lepidine **1a** (1.0 mmol) with BPO (1.7 mmol) in the presence of trifluoroacetic acid (1.0 mmol) in *N,N*-dimethylacetamide **A** (8.0 mL) under irradiation with a Hg lamp for 24 h in the temperature range of 35 °C ~ 40 °C was the best choice, giving compound **2aA** in 88% yield (entry 20 in entries 15 ~21, Table 1). The reason why the yield of **2aA** depends on the amount of *N,N*-dimethylacetamide **A** may be the smooth formation of a carbon-centered amide radical by the reaction of *N,N*-dimethylacetamide **A** with  $\text{PhCO}_2\cdot$  and the suppression of the oxidation of the carbon-centered amide radical to carbo-cation by BPO under diluted conditions. In the absence of BPO under the same reaction conditions, the yield of compound **2aA** was 14% (entry 22, Table 1). In the absence of trifluoroacetic acid under the same reaction conditions, the yield of compound **2aA** was trace (entry 23, Table 1). When the present reaction was carried out under warming conditions at 100 °C for 24 h and under irradiation with a tungsten lamp (300 W) at 40 °C for 8 h, instead of irradiation with a Hg lamp, the yield of compound **2aA** was only 13% and 0%, respectively (entries 24 and 25, Table 1). Then, a gram-scale experiment was performed with 8 mmol of lepidine in a 100 mL screw-capped flask under the same procedure and conditions as those in entry 20, Table 1, and compound **2aA** was obtained in 89% yield, as shown in Table 2. Thus, this method can be used for gram-scale preparation. Based on those results, various quinoline derivatives, such as 4,6-dimethylquinoline **1b**, 4,7-dichloroquinoline **1c**, 2-methylquinoline **1d**, 2,6-dimethylquinoline **1e**, 6-methoxy-2-methylquinoline **1f**, 6-fluoro-2-methylquinoline **1g**, 6-chloro-2-methylquinoline **1h**,

**Table 2.** Introduction of Quinoline, Isoquinoline, and Phenanthridine **1** onto Tertiary Amides **A~E**




<sup>a</sup> Reaction was carried out on a 8.0 mmol scale. <sup>b</sup> Solvent (10.0 mL) was used.

<sup>c</sup> BPO (2.0 equiv.) was used. <sup>d</sup> Reaction was carried out for 18 h.

7-chloro-2-methylquinoline **1i**, and 6-bromo-2-methylquinoline **1j**, were treated with BPO in the presence of trifluoroacetic acid in *N,N*-dimethylacetamide **A** under the same reaction conditions as those in entry 20, Table 1 to provide the corresponding quinoline derivatives **2bA~2jA** bearing a tertiary amide group in good to moderate yields, respectively, as shown in Table 2. When quinoline derivatives **1k** and **1l** bearing an ester or an acetyl group were reacted with BPO in *N,N*-dimethylacetamide under the same reaction conditions, compounds **2kA** and **2lA** were produced in moderate yields, respectively. Similarly, treatment of isoquinoline **1m**, 4-bromoisoquinoline **1n**, and quinoxaline **1o** with BPO in *N,N*-dimethylacetamide **A** under the same reaction conditions gave compounds **2mA**, **2nA**, and **2oA** in good to moderate yields, respectively. The same treatment of phenanthridine **1p**, a tricyclic heteroaromatic base, with BPO in *N,N*-dimethylacetamide **A** generated also the corresponding C-C bonded compound **2pA** in 65% yield. On the other hand, when lepidine was treated with BPO in *N,N*-dimethylpropionamide **B** and 1-methyl-2-pyrrolidone **C** instead of *N,N*-dimethylacetamide **A** under irradiation with a Hg lamp, 4-methyl-2-(*N*-methyl,*N*-propanoylamino)methylquinoline **2aB** and a mixture of 4-methylquinolines **2aC** and **2aC'** bearing a pyrrolidone group at 2-position were obtained in 72% and 73% yields (19% and 54%), respectively. Thus, 1-methyl-2-pyrrolidone has two reaction positions, *i.e.*, the *N*-methyl group and the cyclic *N*-methylene group. These results are consistent with reported results, *i.e.*, the smooth generation of carbon-centered amide radical by the hydrogen-atom abstraction of the *N*-methyl group in *N,N*-dimethylacetamide and two carbon-centered amide radicals by the hydrogen-atom abstraction of the *N*-methyl group and the *N*-methylene

group in 1-methyl-2-pyrrolidone by a cumyloxy radical.<sup>10</sup> However, when 1-ethyl-2-pyrrolidone **D** instead of 1-methyl-2-pyrrolidone **C** was used as solvent with lepidine and BPO under the same irradiation conditions, single product **2aD** was obtained in 62% yield. This result may have been derived from the steric hindrance in the reaction of protonated lepidine salt with the carbon-centered amide radical formed by the hydrogen-atom abstraction of the *N*-ethyl group in 1-ethyl-2-pyrrolidone **D**, and the carbon-centered amide radical formed by the hydrogen-atom abstraction of the *N*-methylene group in 1-ethyl-2-pyrrolidone **D** reacted with protonated lepidine salt to give **2aD** alone. When *N*-acetylpyrrolidine **E** was used as solvent with lepidine and BPO under the same irradiation conditions, 1-acetyl-2-(4'-methylquinolin-2'-yl)pyrrolidine **2aE** was obtained in moderate yield. Similarly, when isoquinoline was treated with BPO in *N,N*-dimethylpropionamide **B**, 1-methyl-2-pyrrolidone **C**, and 1-ethyl-2-pyrrolidone **D** under irradiation with a Hg lamp, the corresponding C-C bonded products **2mB**, a mixture of **2mC** and **2mC'**, and **2mD** were obtained in 77%, 76% (13% and 63%), and 60% yields, respectively. When phenanthridine was treated with BPO in *N,N*-dimethylpropionamide **B** and

**Scheme 1.** Introduction of Quinoline **1q** onto *N,N*-Dimethylacetamide **A**




1-ethyl-2-pyrrolidone **D** under the same reaction conditions, the corresponding phenathridine derivatives **2pB** and **2pD** bearing an amide group at 6-position were obtained in 77% and 70% yields, respectively. When quinoline, which has two reactive positions toward nucleophilic radical species, was treated with BPO (1.0 equiv.) in *N,N*-dimethylacetamide **A** under irradiation with a Hg lamp for 24 h, 2-(*N*-acetyl,*N*-methylamino)methylquinoline **2qA** (20%),

4-(*N*-acetyl,*N*-methylamino)methylquinoline **2qA'** (31%), and 2,4-bis[(*N*-acetyl,*N*-methylamino)methyl]quinoline **2qA''** (24%) were obtained in 75% yield, as shown in Scheme 1 (eq. 1). On the other hand, when quinoline was treated with BPO (2.0 equiv.) in *N,N*-dimethylacetamide under irradiation with a Hg lamp for 24 h, 2,4-bis[(*N*-acetyl,*N*-methylamino)methyl]quinoline **2qA''** was obtained in 84% yield as a single product (eq. 2).


The structure of 1-ethyl-5-(4'-methylquinolin-2'-yl)pyrrolidin-2-one **2aD** was supported by X-ray crystallographic analysis (see Supporting Information).

Finally, the obtained C-C bonded compounds **2** are mainly mixtures of *anti* and *syn* forms with respect to the amide groups. Therefore, C-C bonded compounds **2** were converted into secondary amines **3** bearing a quinoline group and an isoquinoline group in good yields, respectively, by aq. HCl hydrolysis for analysis, as shown in Table 3.


**Table 3.** Hydrolysis of Tertiary Amides **2** to Secondary Amines **3**



<sup>a</sup> Reaction was carried out on a 0.5 mmol scale.

**Scheme 2.** Reduction of Tertiary Amides **2** to Tertiary Amines **4** with NaBH<sub>4</sub> and I<sub>2</sub>

Then, C-C bonded compounds **2aA** and **2aC'** were reduced to the corresponding tertiary amines **4aA** and **4aC'** bearing a quinoline group in 70% and 61% yields, respectively, by treatment with NaBH<sub>4</sub> (2.3 equiv.) and I<sub>2</sub> (1.0 equiv.) in THF for 4 h under refluxing conditions, as shown in Scheme 2 (eqs. 1 and 2).


**Scheme 3.** Introduction of Quinoline **1a** and **1e** onto *N*-Methylacetamide **F**

When *N*-methylacetamide **F** instead of *N,N*-dimethylacetamide **A** was treated with lepidine **1a** and 2,6-dimethylquinoline **1e** in the presence of BPO and trifluoroacetic acid under the same irradiation conditions with a Hg lamp,

2-(*N*-acetylamino)methyl-4-methylquinoline **2aF** and 4-(*N*-acetylamino)methyl-2,6-dimethylquinoline **2eF** were obtained in 62% and 59% yields, respectively, as shown in Scheme 3.

To support the present reaction mechanism, two experiments were carried out. When the reaction of lepidine with BPO (1.7 equiv.) in the presence of 2,2,6,6-tetramethylpiperidine-1-oxyl radical (TEMPO, 3.0 equiv.) and trifluoroacetic acid (1.0 equiv.) in *N,N*-dimethylacetamide **A** was carried out under the same irradiation conditions, C-C bonded compound **2aA** was not obtained at all. Moreover, the reaction of lepidine with BPO (1.7 equiv.) and trifluoroacetic acid (1.0 equiv.) in *N,N*-dimethylacetamide **A** in the presence of *p*-cresol (3.0 equiv.), which functions as a hydrogen atom donor to radical species, under the same irradiation conditions was also retarded to give compound **2aA** in only 14% yield. Thus, it is suggested that the present reaction proceeds via the radical pathway. The proposed reaction mechanism is shown in Scheme 4 with lepidine **1a** in *N,N*-dimethylacetamide **A**.

**Scheme 4.** Plausible Reaction Pathway



Benzoyloxyl radical formed by the homolytic bond cleavage of BPO under irradiation with a Hg lamp abstracts a hydrogen atom from the *N*-methyl group in

*N,N*-dimethylacetamide **A** to form carbon-centered amide radical **IA** and benzoic acid (or abstraction of a hydrogen atom from the *N*-methyl group or the *N*-methylene group in 1-methyl-2-pyrrolidone **C** to form carbon-centered amide radicals **IC** and **IC'**). This nucleophilic carbon-centered amide radical **IA** reacts at the most electrophilic position of protonated lepidine, *i.e.*, 2-position, to form C-C bonded intermediate **IIA**, which further reacts with BPO to give protonated C-C bonded quinoline derivative, together with the generation of benzoic acid and benzoyloxyl radical. Quinoline derivative **2aA** was obtained by neutralization with aq. NaHCO<sub>3</sub> solution. Practically, the smooth generation of carbon-centered amide radical **IA** from *N,N*-dimethylacetamide **A** and two carbon-centered amide radicals adjoined to the nitrogen atom in 1-methyl-2-pyrrolidone **C** by a cumyloxy radical, *i.e.*, an alkoxyl radical, has been reported by Bietti.<sup>10</sup>

## CONCLUSION

The introduction of quinolines, isoquinolines, and phenanthridine onto the non-activated  $\alpha$ -C-H bond of tertiary amides was efficiently carried out by treatment with benzoyl peroxide under irradiation with a Hg lamp via the radical pathway. We believe that the present reaction would be useful because it would enable the preparation of unique quinolines, isoquinolines, and phenanthridines bearing tertiary amide groups.

## EXPERIMENTAL SECTION

**General Information.** <sup>1</sup>H NMR spectra were measured on 400 MHz spectrometers. Chemical shifts were recorded as follows: chemical shift in ppm from internal tetramethylsilane on the  $\delta$  scale, multiplicity (s = singlet; d = doublet; t = triplet; q = quartet; quin = quintet; sext = sextet; m = multiplet; br = broad), coupling constant (Hz), integration, and assignment. <sup>13</sup>C NMR spectra were measured on 100 MHz spectrometers. Chemical shifts were recorded in ppm from the solvent resonance employed as the internal standard (deuteriochloroform at 77.0 ppm). High-resolution mass spectra (HRMS) were measured on orbitrap mass spectrometers. Characteristic peaks in the infrared (IR) spectra were recorded in wave numbers, cm<sup>-1</sup>. Melting points

were uncorrected. Thin-layer chromatography (TLC) was performed using 0.25 mm silica gel plates (60F-254). The products were purified by short column chromatography on neutral silica gel 60 (63–200 mesh).

**Typical Experimental Procedure for Introduction of Lepidine to *N,N*-Dimethylacetamide with BPO under Irradiation with Hg-Lamp:** To a solution of lepidine (**1a**; 1.0 mmol, 143.2 mg) and trifluoroacetic acid (1.0 mmol, 0.077 mL) in *N,N*-dimethylacetamide (**A**, 8.0 mL) in a 30 mL screw-capped flask was added benzoyl peroxide (wetted with *ca.* 25% water, 1.7 mmol, 549.1 mg) at room temperature, and the flask was flushed by Ar gas. Then, the mixture was stirred for 24 h under irradiation with a high-pressure mercury lamp (Hg-lamp, 400W: AHH400S, 1.7 mW/cm<sup>2</sup>) at the range of 35 °C~40 °C. Then, the reaction mixture was cooled to room temperature and quenched with sat. aq. NaHCO<sub>3</sub> (10 mL). The mixture was extracted with AcOEt (20 mL × 5) and the organic layer was washed with brine (50 mL). The obtained organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>. After filtration, the solvent was removed under reduced pressure, and the residue was purified by short column chromatography on neutral silica gel (eluent: AcOEt/MeOH = 20/1) to give the product **2aA** (199.8 mg, 88% yield).

**Typical Experimental Procedure for Hydrolysis of *tertiary* Amides to *secondary* Amines with aq. HCl:** A solution of 2-(*N*-acetyl,*N*-methylamino)methyl-4-methylquinoline (**2aA**; 1.0 mmol, 228.3 mg) in aq. 3 M HCl (3.0 mL) was stirred for 3 h at 100 °C. Then, the reaction mixture was cooled to room temperature and quenched with sat. aq. NaHCO<sub>3</sub> (20 mL). The mixture was extracted with CHCl<sub>3</sub> (20 mL × 5), and the obtained organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>. After filtration, the solvent was removed under reduced pressure to give the product **3aA** (184.0 mg, 99 % yield).

**Typical Experimental Procedure for Reduction of *tertiary* Amides to *tertiary* Amines with NaBH<sub>4</sub> and I<sub>2</sub>:** To a solution of 2-(*N*-acetyl,*N*-methylamino)methyl-4-methylquinoline (**2aA**; 0.5 mmol, 114.2 mg) and NaBH<sub>4</sub> (1.15 mmol, 43.5 mg) in THF (1.5 mL) was added dropwise a solution of I<sub>2</sub> (0.5 mmol, 126.9 mg) in THF (1.0 mL) under Ar atmosphere at 0 °C. Then, the mixture was

1  
2  
3  
4  
5  
6 stirred for 4 h at 70 °C. Then, the reaction mixture was cooled to 0 °C and added to aq.  
7 3 M HCl (3.0 mL) solution. The mixture was stirred for 1 h at 70°C. The reaction  
8 mixture was cooled to room temperature and quenched with sat. aq. NaHCO<sub>3</sub> (20 mL)  
9 solution. Then, the mixture was extracted with CHCl<sub>3</sub> (20 mL × 5), and the  
10 obtained organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>. After filtration, the solvent was  
11 removed under reduced pressure, and the residue was purified by short column  
12 chromatography on neutral silica gel (AcOEt /MeOH= 2/1) to give the product **4aA**  
13 (74.9 mg, 70% yield).  
14  
15  
16  
17  
18  
19  
20

21 For amide products **2aC**, **2aC'**, **2aD**, **2aF**, **2eF**, **2mC**, **2mC'**, **2mD**, and **2pD**, oil or  
22 solid, IR, <sup>1</sup>H-NMR, <sup>13</sup>C-NMR, and HRMS were shown in the following experimental  
23 section, due to the single isomer. However, other amide products **2** are a mixture of  
24 *anti* and *syn* in the amide group, and their <sup>1</sup>H-NMR and <sup>13</sup>C-NMR spectra are rather  
25 complicated. Therefore, for those amide products **2**, oil or solid, IR, and HRMS were  
26 shown in the following experimental section, and copies of <sup>1</sup>H-NMR and <sup>13</sup>C-NMR  
27 were attached in Supporting Information. After acid hydrolysis of those amide  
28 products **2**, each single amine **3** was obtained, and oil or solid, IR, <sup>1</sup>H-NMR, <sup>13</sup>C-NMR,  
29 and HRMS were shown in the following experimental section, and copies of those  
30 <sup>1</sup>H-NMR and <sup>13</sup>C-NMR were attached in Supporting Information.  
31  
32  
33  
34  
35  
36  
37  
38  
39

40 **2-(N-Acetyl,N-methylamino)methyl-4-methylquinoline (2aA)<sup>8a</sup>**: (a mixture of *anti*  
41 and *syn* isomers) 199.8 mg, 88% yield; yellow solid; IR (neat) 2926, 1637, 1601, 1401,  
42 756 cm<sup>-1</sup>; HRMS (ESI) calcd for C<sub>14</sub>H<sub>17</sub>N<sub>2</sub>O [M+H]<sup>+</sup> 229.1335, found 229.1331.  
43  
44

45 **2-(N-Acetyl,N-methylamino)methyl-4,6-dimethylquinoline (2bA)**: (a mixture of *anti*  
46 and *syn* isomers) 188.1 mg, 78% yield; yellow oil; IR (neat) 2921, 1644, 1600, 1398,  
47 824 cm<sup>-1</sup>; HRMS (ESI) calcd for C<sub>15</sub>H<sub>19</sub>N<sub>2</sub>O [M+H]<sup>+</sup> 243.1492, found 243.1489.  
48  
49

50 **2-(N-Acetyl,N-methylamino)methyl-4,7-dichloroquinoline (2cA)**: (a mixture of *anti*  
51 and *syn* isomers) 143.3 mg, 51% yield; white solid; IR (neat) 2927, 1638, 1604, 1396,  
52 819 cm<sup>-1</sup>; HRMS (ESI) calcd for C<sub>13</sub>H<sub>13</sub>Cl<sub>2</sub>N<sub>2</sub>O [M+H]<sup>+</sup> 283.0399, found 283.0398.  
53  
54

55 **4-(N-Acetyl,N-methylamino)methyl-2-methylquinoline (2dA)**: (a mixture of *anti* and  
56 *syn* isomers) 188.0 mg, 82% yield; yellow oil; IR (neat) 2924, 1636, 1604, 1402, 755  
57 cm<sup>-1</sup>; HRMS (ESI) calcd for C<sub>14</sub>H<sub>17</sub>N<sub>2</sub>O [M+H]<sup>+</sup> 229.1335, found 229.1331.  
58  
59  
60

**4-(N-Acetyl,N-methylamino)methyl-2,6-dimethylquinoline (2eA)**: (a mixture of *anti*

and *syn* isomers) 186.2 mg, 77% yield; yellow solid; IR (neat) 2920, 1645, 1603, 1400, 825  $\text{cm}^{-1}$ ; HRMS (ESI) calcd for  $\text{C}_{15}\text{H}_{19}\text{N}_2\text{O}$   $[\text{M}+\text{H}]^+$  243.1492, found 243.1489.

**4-(*N*-Acetyl,*N*-methylamino)methyl-6-methoxy-2-methylquinoline (2fA):** (a mixture of *anti* and *syn* isomers) 166.3 mg, 64% yield; yellow solid; IR (neat) 2933, 1621, 1404, 1232, 727  $\text{cm}^{-1}$ ; HRMS (ESI) calcd for  $\text{C}_{15}\text{H}_{19}\text{N}_2\text{O}_2$   $[\text{M}+\text{H}]^+$  259.1441, found 259.1438.

**4-(*N*-Acetyl,*N*-methylamino)methyl-6-fluoro-2-methylquinoline (2gA):** (a mixture of *anti* and *syn* isomers) 169.3 mg, 69% yield; yellow solid; IR (neat) 2924, 1637, 1609, 1405, 830  $\text{cm}^{-1}$ ; HRMS (ESI) calcd for  $\text{C}_{14}\text{H}_{16}\text{FN}_2\text{O}$   $[\text{M}+\text{H}]^+$  247.1241, found 247.1235.

**4-(*N*-Acetyl,*N*-methylamino)methyl-6-chloro-2-methylquinoline (2hA):** (a mixture of *anti* and *syn* isomers) 202.4 mg, 77% yield; yellow solid; IR (neat) 2924, 1637, 1605, 1401, 830  $\text{cm}^{-1}$ ; HRMS (ESI) calcd for  $\text{C}_{14}\text{H}_{16}\text{ClN}_2\text{O}$   $[\text{M}+\text{H}]^+$  263.0946, found 263.0939.

**4-(*N*-Acetyl,*N*-methylamino)methyl-7-chloro-2-methylquinoline (2iA):** (a mixture of *anti* and *syn* isomers) 191.1 mg, 73% yield; white solid; IR (neat) 2927, 1627, 1606, 1402, 890  $\text{cm}^{-1}$ ; HRMS (ESI) calcd for  $\text{C}_{14}\text{H}_{16}\text{ClN}_2\text{O}$   $[\text{M}+\text{H}]^+$  263.0946, found 263.0943.

**4-(*N*-Acetyl,*N*-methylamino)methyl-6-bromo-2-methylquinoline (2jA):** (a mixture of *anti* and *syn* isomers) 205.8 mg, 67% yield; yellow solid; IR (neat) 2924, 1640, 1607, 1401, 828  $\text{cm}^{-1}$ ; HRMS (ESI) calcd for  $\text{C}_{14}\text{H}_{16}\text{BrN}_2\text{O}$   $[\text{M}+\text{H}]^+$  307.0441, found 307.0434.

**4-(*N*-Acetyl,*N*-methylamino)methyl-6-methoxycarbonyl-2-methylquinoline (2kA):** (a mixture of *anti* and *syn* isomers) 158.7 mg, 55% yield; yellow solid; IR (neat) 2952, 1716, 1644, 1272, 751  $\text{cm}^{-1}$ ; HRMS (ESI) calcd for  $\text{C}_{16}\text{H}_{19}\text{N}_2\text{O}_3$   $[\text{M}+\text{H}]^+$  287.1390, found 287.1386.

**4-(*N*-Acetyl,*N*-methylamino)methyl-6-acetyl-2-methylquinoline (2lA):** (a mixture of *anti* and *syn* isomers) 106.8 mg, 40% yield; yellow solid; IR (neat) 2924, 1671, 1642, 1402, 1263, 841  $\text{cm}^{-1}$ ; HRMS (ESI) calcd for  $\text{C}_{16}\text{H}_{19}\text{N}_2\text{O}_2$   $[\text{M}+\text{H}]^+$  271.1441, found 271.1437.

**1-(*N*-Acetyl,*N*-methylamino)methylisoquinoline (2mA):** (a mixture of *anti* and *syn* isomers) 183.5 mg, 86% yield; white solid; IR (neat) 2927, 1638, 1403, 826, 747  $\text{cm}^{-1}$ ; HRMS (ESI) calcd for  $\text{C}_{13}\text{H}_{15}\text{N}_2\text{O}$   $[\text{M}+\text{H}]^+$  215.1179, found 215.1175.

**1-(*N*-Acetyl,*N*-methylamino)methyl-4-bromoisoquinoline (2nA):** (a mixture of *anti* and *syn* isomers) 151.3 mg, 52% yield; yellow solid; IR (neat) 2927, 1631, 1410, 1255,

763  $\text{cm}^{-1}$ ; HRMS (ESI) calcd for  $\text{C}_{13}\text{H}_{14}\text{BrN}_2\text{O}$   $[\text{M}+\text{H}]^+$  293.0284, found 293.0279.

**2-(N-Acetyl,N-methylamino)methylquinoxaline (2oA)<sup>8b</sup>**: (a mixture of *anti* and *syn* isomers) 97.6 mg, 45% yield; yellow oil; IR (neat) 2931, 1642, 1492, 1400, 756  $\text{cm}^{-1}$ ; HRMS (ESI) calcd for  $\text{C}_{12}\text{H}_{14}\text{N}_3\text{O}$   $[\text{M}+\text{H}]^+$  216.1131, found 216.1128.

**6-(N-Acetyl,N-methylamino)methylphenanthridine (2pA)<sup>8c</sup>**: (a mixture of *anti* and *syn* isomers) 171.6 mg, 65% yield; yellow oil; IR (neat) 2926, 1643, 1486, 1402, 750  $\text{cm}^{-1}$ ; HRMS (ESI) calcd for  $\text{C}_{17}\text{H}_{17}\text{N}_2\text{O}$   $[\text{M}+\text{H}]^+$  265.1335, found 265.1332.

**2-(N-Acetyl,N-methylamino)methylquinoline (2qA)**: (a mixture of *anti* and *syn* isomers) 42.2 mg, 20% yield; yellow oil; IR (neat) 2930, 1634, 1600, 1401, 776  $\text{cm}^{-1}$ ; HRMS (ESI) calcd for  $\text{C}_{13}\text{H}_{15}\text{N}_2\text{O}$   $[\text{M}+\text{H}]^+$  215.1179 found 215.1175

**4-(N-Acetyl,N-methylamino)methylquinoline (2qA')**: (a mixture of *anti* and *syn* isomers) 65.9 mg, 31% yield; yellow oil; IR (neat) 2930, 1643, 1596, 1401, 752  $\text{cm}^{-1}$ ; HRMS (ESI) calcd for  $\text{C}_{13}\text{H}_{15}\text{N}_2\text{O}$   $[\text{M}+\text{H}]^+$  215.1179 found 215.1177.

**2,4-Bis[(N-methyl-N-acetylamino)methyl]quinoline (2qA'')**: (a mixture of *anti* and *syn* isomers) 71.3 mg, 24% yield; yellow oil; IR (neat) 2931, 1633, 1485, 1401, 754  $\text{cm}^{-1}$ ; HRMS (ESI) calcd for  $\text{C}_{17}\text{H}_{22}\text{N}_3\text{O}_2$   $[\text{M}+\text{H}]^+$  300.1707, found 300.1704.

**2-(N-Methyl,N-propanoylamino)methyl-4-methylquinoline (2aB)**: (a mixture of *anti* and *syn* isomers) 174.3 mg, 72% yield; yellow oil; IR (neat) 2936, 1643, 1601, 1402, 756  $\text{cm}^{-1}$ ; HRMS (ESI) calcd for  $\text{C}_{15}\text{H}_{19}\text{N}_2\text{O}$   $[\text{M}+\text{H}]^+$  243.1492, found 243.1487.

**1-[(4'-Methylquinolin-2'-yl)methyl]pyrrolidin-2-one (2aC)**: 44.6 mg, 19% yield; colorless oil; IR (neat) 2923, 1674, 1602, 1421, 757  $\text{cm}^{-1}$ ;  $^1\text{H}$ -NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  2.04 (quin,  $J = 7.6$  Hz, 2H), 2.50 (t,  $J = 7.6$  Hz, 2H), 2.69 (s, 3H), 3.41 (t,  $J = 7.6$  Hz, 2H), 4.73 (s, 2H), 7.22 (s, 1H), 7.56 (t,  $J = 8.4$  Hz, 1H), 7.71 (t,  $J = 8.4$  Hz, 1H), 7.98 (d,  $J = 8.4$  Hz, 1H), 8.05 (d,  $J = 8.4$  Hz, 1H) ppm;  $^{13}\text{C}$ -NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  17.8, 18.8, 30.8, 47.2, 49.2, 120.5, 123.7, 126.2, 127.4, 129.3, 129.5, 145.5, 147.3, 156.8, 175.3 ppm; HRMS (ESI) calcd for  $\text{C}_{15}\text{H}_{17}\text{N}_2\text{O}$   $[\text{M}+\text{H}]^+$  241.1335, found 241.1331.

**1-Methyl-5-(4'-methylquinolin-2'-yl)pyrrolidin-2-one (2aC')**: 129.6 mg, 54% yield; white solid, mp 145-146  $^\circ\text{C}$ ; IR (neat) 2916, 1686, 1601, 1393, 763  $\text{cm}^{-1}$ ;  $^1\text{H}$ -NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  1.98-2.09 (m, 1H), 2.47-2.71 (m, 3H), 2.73 (s, 3H), 2.78 (s, 3H), 4.82 (dd,  $J = 8.6, 5.2$  Hz, 1H), 7.12 (s, 1H), 7.59 (t,  $J = 8.4$  Hz, 1H), 7.74 (t,  $J = 8.4$  Hz, 1H), 8.00 (d,  $J = 8.4$  Hz, 1H), 8.07 (d,  $J = 8.4$  Hz, 1H) ppm;  $^{13}\text{C}$ -NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  18.9, 26.2, 28.6, 30.0, 66.5, 118.0, 123.7, 126.5, 127.5, 129.6 (2C), 146.1, 147.5, 160.6, 175.8 ppm; HRMS (ESI) calcd for  $\text{C}_{15}\text{H}_{17}\text{N}_2\text{O}$   $[\text{M}+\text{H}]^+$  241.1335, found 241.1334.

**1-Ethyl-5-(4'-methylquinolin-2'-yl)pyrrolidin-2-one (2aD):** 157.5 mg, 62% yield; white solid, mp 100-101 °C; IR (neat) 2935, 1685, 1600, 1419, 764 cm<sup>-1</sup>; <sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>) δ 1.05 (t, *J* = 7.2 Hz, 3H), 1.98-2.09 (m, 1H), 2.47-2.62 (m, 2H), 2.63-2.71 (m, 1H), 2.72 (s, 3H), 2.77 (dq, *J* = 14.5, 7.2 Hz, 1H), 3.80 (dq, *J* = 14.5, 7.2 Hz, 1H), 4.96 (dd, *J* = 8.5, 4.8 Hz, 1H), 7.13 (s, 1H), 7.59 (t, *J* = 8.4 Hz, 1H), 7.74 (t, *J* = 8.4 Hz, 1H), 8.00 (d, *J* = 8.4 Hz, 1H), 8.07 (d, *J* = 8.4 Hz, 1H) ppm; <sup>13</sup>C-NMR (100 MHz, CDCl<sub>3</sub>) δ 12.3, 18.9, 26.3, 30.3, 36.0, 63.8, 118.2, 123.7, 126.4, 127.5, 129.57, 129.64, 145.9, 147.5, 160.9, 175.4 ppm; HRMS (ESI) calcd for C<sub>16</sub>H<sub>19</sub>N<sub>2</sub>O [M+H]<sup>+</sup> 255.1492, found 255.1489.

**1-Acetyl-2-(4'-methylquinolin-2'-yl)pyrrolidine (2aE):** (a mixture of *anti* and *syn* isomers) 122.8 mg, 48% yield; white solid; IR (neat) 2968, 1639, 1595, 1410, 758 cm<sup>-1</sup>; HRMS (ESI) calcd for C<sub>16</sub>H<sub>19</sub>N<sub>2</sub>O [M+H]<sup>+</sup> 255.1492, found 255.1490.

**2-(*N*-Acetylamino)methyl-4-methylquinoline (2aF)<sup>8a</sup>:** 132.4 mg, 62% yield; white solid, mp 121-123 °C; IR (neat) 3293, 2920, 1634, 1549, 765 cm<sup>-1</sup>; <sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>) δ 2.15 (s, 3H), 2.70 (s, 3H), 4.69 (d, *J* = 4.3 Hz, 2H), 7.17 (s, 1H), 7.57 (t, *J* = 8.4 Hz, 1H), 7.72 (t, *J* = 8.4 Hz, 1H), 7.99 (d, *J* = 8.4 Hz, 1H), 8.05 (d, *J* = 8.4 Hz, 1H) ppm; <sup>13</sup>C-NMR (100 MHz, CDCl<sub>3</sub>) δ 18.6, 23.2, 44.8, 120.5, 123.8, 126.1, 127.4, 129.1, 129.4, 145.1, 146.9, 155.7, 170.2 ppm; HRMS (ESI) calcd for C<sub>13</sub>H<sub>15</sub>N<sub>2</sub>O [M+H]<sup>+</sup> 215.1179, found 215.1175.

**4-(*N*-Acetylamino)methyl-2,6-dimethylquinoline (2eF):** 134.1 mg, 59% yield; yellow solid, mp 154-155 °C; IR (neat) 3253, 2916, 1628, 1560, 829 cm<sup>-1</sup>; <sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>) δ 2.10 (s, 3H), 2.53 (s, 3H), 2.66 (s, 3H), 4.82 (d, *J* = 5.6 Hz, 2H), 7.12 (s, 1H), 7.51 (d, *J* = 8.5 Hz, 1H), 7.66 (s, 1H), 7.90 (d, *J* = 8.5 Hz, 1H) ppm; <sup>13</sup>C-NMR (100 MHz, CDCl<sub>3</sub>) δ 21.8, 23.1, 25.0, 40.3, 120.7, 121.8, 124.5, 128.8, 131.5, 135.9, 142.3, 146.3, 157.6, 170.1 ppm; HRMS (ESI) calcd for C<sub>14</sub>H<sub>17</sub>N<sub>2</sub>O [M+H]<sup>+</sup> 229.1335, found 229.1332.

**1-(*N*-Acetyl,*N*-methylamino)methylisoquinoline (2mB):** (a mixture of *anti* and *syn* isomers) 175.6 mg, 77% yield; yellow oil; IR (neat) 2936, 1638, 1481, 1404, 747 cm<sup>-1</sup>; HRMS (ESI) calcd for C<sub>14</sub>H<sub>17</sub>N<sub>2</sub>O [M+H]<sup>+</sup> 229.1335, found 229.1334.

**1-[(Isoquinolin-1'-yl)methyl]pyrrolidin-2-one (2mC):** 30.5 mg, 13% yield; white solid, mp 97-99 °C; IR (neat) 2951, 1672, 1256, 834, 747 cm<sup>-1</sup>; <sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>) δ 1.94 (quin, *J* = 7.6 Hz, 2H), 2.45 (t, *J* = 7.6 Hz, 2H), 3.30 (t, *J* = 7.6 Hz, 2H), 5.09 (s, 2H), 7.61-7.67 (m, 2H), 7.71 (t, *J* = 8.0 Hz, 1H), 7.85 (d, *J* = 8.0 Hz, 1H), 8.38

(d,  $J = 8.0$  Hz, 1H), 8.45 (d,  $J = 5.8$  Hz, 1H) ppm;  $^{13}\text{C}$ -NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  17.6, 30.8, 46.9, 47.1, 121.0, 125.4, 126.7, 127.1, 127.9, 130.4, 136.3, 141.5, 155.8, 174.6 ppm; HRMS (ESI) calcd for  $\text{C}_{14}\text{H}_{15}\text{N}_2\text{O}$   $[\text{M}+\text{H}]^+$  227.1179, found 227.1176.

**5-(Isoquinolin-1'-yl)-1-methylpyrrolidin-2-one (2mC')**: 141.7 mg, 63% yield; yellow oil; IR (neat) 2925, 1673, 1395, 825, 747  $\text{cm}^{-1}$ ;  $^1\text{H}$ -NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  2.08-2.20 (m, 1H), 2.44-2.56 (m, 1H), 2.56-2.72 (m, 2H), 2.80 (s, 3H), 5.51 (dd,  $J = 7.9, 4.3$  Hz, 1H), 7.61 (d,  $J = 5.8$  Hz, 1H), 7.65 (t,  $J = 7.9$  Hz, 1H), 7.73 (t,  $J = 7.9$  Hz, 1H), 7.90 (d,  $J = 7.9$  Hz, 1H), 8.13 (d,  $J = 7.9$  Hz, 1H), 8.51 (d,  $J = 5.8$  Hz, 1H) ppm;  $^{13}\text{C}$ -NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  26.0, 28.8, 29.9, 62.0, 120.5, 123.3, 125.9, 127.6, 127.8, 130.0, 136.6, 142.0, 158.0, 175.9 ppm; HRMS (ESI) calcd for  $\text{C}_{14}\text{H}_{15}\text{N}_2\text{O}$   $[\text{M}+\text{H}]^+$  227.1179, found 227.1176.

**1-Ethyl-5-(isoquinolin-1'-yl)pyrrolidin-2-one (2mD)**: 144.3 mg, 60% yield; yellow oil; IR (neat) 2975, 1672, 1416, 1245, 727  $\text{cm}^{-1}$ ;  $^1\text{H}$ -NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  1.01 (t,  $J = 7.3$  Hz, 3H), 2.09-2.19 (m, 1H), 2.45-2.63 (m, 2H), 2.63-2.82 (m, 2H), 3.84 (dq,  $J = 14.7, 7.3$  Hz, 1H), 5.64 (dd,  $J = 7.8, 4.4$  Hz, 1H), 7.61 (d,  $J = 5.7$  Hz, 1H), 7.65 (t,  $J = 7.9$  Hz, 1H), 7.73 (t,  $J = 7.9$  Hz, 1H), 7.90 (d,  $J = 7.9$  Hz, 1H), 8.16 (d,  $J = 7.9$  Hz, 1H), 8.51 (d,  $J = 5.7$  Hz, 1H) ppm;  $^{13}\text{C}$ -NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  12.4, 26.0, 30.2, 35.9, 59.0, 120.4, 123.1, 125.8, 127.5, 127.7, 130.0, 136.5, 141.9, 158.3, 175.4 ppm; HRMS (ESI) calcd for  $\text{C}_{15}\text{H}_{17}\text{N}_2\text{O}$   $[\text{M}+\text{H}]^+$  241.1335, found 241.1332.

**6-(N-Methyl,N-propanoylamino)methylphenanthridine (2pB)**: (a mixture of *anti* and *syn* isomers) 213.1 mg, 77% yield; yellow solid; IR (neat) 2936, 1636, 1462, 1404, 752  $\text{cm}^{-1}$ ; HRMS (ESI) calcd for  $\text{C}_{18}\text{H}_{19}\text{N}_2\text{O}$   $[\text{M}+\text{H}]^+$  279.1492, found 279.1489.

**1-Ethyl-5-(phenanthridin-6'-yl)pyrrolidin-2-one (2pD)**: 204.1 mg, 70% yield; white solid, mp 154-156  $^{\circ}\text{C}$ ; IR (neat) 2974, 1669, 1443, 1272, 762  $\text{cm}^{-1}$ ;  $^1\text{H}$ -NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  1.08 (t,  $J = 7.4$  Hz, 3H), 2.12-2.25 (m, 1H), 2.44-2.58 (m, 1H), 2.60-2.74 (m, 2H), 2.90-3.03 (m, 1H), 3.92 (dq,  $J = 14.7, 7.4$  Hz, 1H), 5.69 (dd,  $J = 8.2, 3.2$  Hz, 1H), 7.64-7.77 (m, 3H), 7.89 (t,  $J = 8.2$  Hz, 1H), 8.15 (d,  $J = 8.0$  Hz, 1H), 8.22 (d,  $J = 8.2$  Hz, 1H), 8.57 (d,  $J = 8.0$  Hz, 1H), 8.72 (d,  $J = 8.2$  Hz, 1H) ppm;  $^{13}\text{C}$ -NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  12.5, 25.8, 30.2, 36.3, 59.5, 121.8, 122.9, 123.6, 123.7, 124.2, 127.1, 127.5, 128.7, 130.3, 130.5, 133.4, 143.1, 157.8, 175.7 ppm; HRMS (ESI) calcd for  $\text{C}_{19}\text{H}_{19}\text{N}_2\text{O}$   $[\text{M}+\text{H}]^+$  291.1492, found 291.1490.

**2-(N-Methylamino)methyl-4-methylquinoline (3aA)**: 184.0 mg, 99% yield; yellow oil; IR (neat) 3307, 2927, 1603, 1445, 755  $\text{cm}^{-1}$ ;  $^1\text{H}$ -NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  2.54 (s,

3H), 2.70 (s, 3H), 4.00 (s, 2H), 7.31 (s, 1H), 7.53 (t,  $J = 8.4$  Hz, 1H), 7.69 (td,  $J = 8.4$ , 1.1 Hz, 1H), 7.97 (dd,  $J = 8.4$ , 1.1 Hz, 1H), 8.06 (d,  $J = 8.4$  Hz, 1H) ppm;  $^{13}\text{C}$ -NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  18.7, 36.4, 57.8, 121.2, 123.6, 125.8, 127.3, 129.1, 129.5, 144.5, 147.5, 159.8 ppm; HRMS (ESI) calcd for  $\text{C}_{12}\text{H}_{15}\text{N}_2$   $[\text{M}+\text{H}]^+$  187.1230, found 187.1225.

**4,6-Dimethyl-2-(*N*-methylamino)methylquinoline (3bA):** yellow oil; IR (neat) 3303, 2920, 1602, 1442, 823  $\text{cm}^{-1}$ ;  $^1\text{H}$ -NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  2.54 (s, 3H), 2.55 (s, 3H), 2.67 (s, 3H), 3.98 (s, 2H), 7.27 (d,  $J = 1.6$  Hz, 1H), 7.52 (dd,  $J = 8.6$ , 1.6 Hz, 1H), 7.72 (s, 1H), 7.94 (d,  $J = 8.6$  Hz, 1H) ppm;  $^{13}\text{C}$ -NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  18.7, 21.8, 36.4, 57.8, 121.2, 122.7, 127.2, 129.1, 131.2, 135.5, 143.8, 146.0, 158.8 ppm; HRMS (ESI) calcd for  $\text{C}_{13}\text{H}_{17}\text{N}_2$   $[\text{M}+\text{H}]^+$  201.1386, found 201.1382.

**7-Chloro-4-hydroxy-2-(*N*-methylamino)methylquinoline (3cA):** 4-Chloro group was converted into OH group in aq. HCl hydrolysis. white solid, mp 177 °C (decomp.); IR (neat) 3285, 3069, 2939, 1494, 816  $\text{cm}^{-1}$ ;  $^1\text{H}$ -NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  2.48 (s, 3H), 3.81 (s, 2H), 6.14 (s, 1H), 7.28 (dd,  $J = 8.6$ , 1.8 Hz, 1H), 7.40 (d,  $J = 1.8$  Hz, 1H), 8.27 (d,  $J = 8.6$  Hz, 1H) ppm;  $^{13}\text{C}$ -NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  36.2, 51.7, 108.0, 117.0, 124.2, 124.3, 127.8, 138.0, 139.8, 150.5, 178.2 ppm; HRMS (ESI) calcd for  $\text{C}_{11}\text{H}_{12}\text{ClN}_2\text{O}$   $[\text{M}+\text{H}]^+$  223.0633, found 223.0633.

**4-(*N*-Methylamino)methyl-2-methylquinoline (3dA):** 187.3 mg, >99% yield; yellow oil; IR (neat) 3296, 2935, 1604, 1445, 726  $\text{cm}^{-1}$ ;  $^1\text{H}$ -NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  2.58 (s, 3H), 2.74 (s, 3H), 4.20 (s, 2H), 7.33 (s, 1H), 7.51 (t,  $J = 8.4$  Hz, 1H), 7.68 (t,  $J = 8.4$  Hz, 1H), 8.01 (d,  $J = 8.4$  Hz, 1H), 8.04 (d,  $J = 8.4$  Hz, 1H) ppm;  $^{13}\text{C}$ -NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  25.4, 36.6, 52.1, 120.6, 122.9, 125.2, 125.6, 129.0, 129.3, 145.2, 147.9, 158.8 ppm; HRMS (ESI) calcd for  $\text{C}_{12}\text{H}_{15}\text{N}_2$   $[\text{M}+\text{H}]^+$  187.1230, found 187.1226.

**2,6-Dimethyl-4-(*N*-methylamino)methylquinoline (3eA):** yellow solid, mp 67-68 °C; IR (neat) 3302, 2919, 1603, 1442, 823  $\text{cm}^{-1}$ ;  $^1\text{H}$ -NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  2.54 (s, 3H), 2.59 (s, 3H), 2.71 (s, 3H), 4.17 (s, 2H), 7.29 (s, 1H), 7.50 (d,  $J = 8.6$  Hz, 1H), 7.75 (s, 1H), 7.93 (d,  $J = 8.6$  Hz, 1H) ppm;  $^{13}\text{C}$ -NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  21.8, 25.2, 36.6, 52.1, 120.6, 121.8, 125.1, 129.0, 131.2, 135.4, 144.5, 146.5, 157.8 ppm; HRMS (ESI) calcd for  $\text{C}_{13}\text{H}_{17}\text{N}_2$   $[\text{M}+\text{H}]^+$  201.1386, found 201.1387.

**6-Methoxy-2-methyl-4-(*N*-methylamino)methylquinoline (3fA):** yellow solid, mp 51-52 °C; IR (neat) 3315, 2924, 1603, 1229, 839  $\text{cm}^{-1}$ ;  $^1\text{H}$ -NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  2.59 (s, 3H), 2.70 (s, 3H), 3.94 (s, 3H), 4.14 (s, 2H), 7.25 (d,  $J = 2.8$  Hz, 1H), 7.30 (s, 1H), 7.34 (dd,  $J = 9.3$ , 2.8 Hz, 1H), 7.95 (d,  $J = 9.3$  Hz, 1H) ppm;  $^{13}\text{C}$ -NMR (100 MHz,

CDCl<sub>3</sub>)  $\delta$  25.0, 36.6, 52.3, 55.5, 101.5, 120.7, 120.9, 126.0, 130.7, 143.86, 143.91, 156.1, 157.1 ppm; HRMS (ESI) calcd for C<sub>13</sub>H<sub>17</sub>N<sub>2</sub>O [M+H]<sup>+</sup> 217.1335, found 217.1332.

**6-Fluoro-2-methyl-4-(N-methylamino)methylquinoline (3gA):** yellow solid, mp 45-46 °C; IR (neat) 3307, 2792, 1515, 1222, 836 cm<sup>-1</sup>; <sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  2.58 (s, 3H), 2.72 (s, 3H), 4.12 (s, 2H), 7.35 (s, 1H), 7.44 (td, *J* = 9.3, 2.7 Hz, 1H), 7.64 (dd, *J* = 9.3, 2.7 Hz, 1H), 8.02 (dd, *J* = 9.3, 5.7 Hz, 1H) ppm; <sup>13</sup>C-NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  25.2, 36.6, 52.3, 106.8 (d, *J*<sub>C,F</sub> = 21.9 Hz), 119.0 (d, *J*<sub>C,F</sub> = 25.8 Hz), 121.3, 126.0 (d, *J*<sub>C,F</sub> = 9.5 Hz), 131.6 (d, *J*<sub>C,F</sub> = 9.5 Hz), 144.8 (d, *J*<sub>C,F</sub> = 5.7 Hz), 145.0, 158.1, 160.0 (d, *J*<sub>C,F</sub> = 246.1 Hz) ppm; HRMS (ESI) calcd for C<sub>12</sub>H<sub>14</sub>FN<sub>2</sub> [M+H]<sup>+</sup> 205.1136, found 205.1132.

**6-Chloro-2-methyl-4-(N-methylamino)methylquinoline (3hA):** yellow solid, mp 94-96 °C; IR (neat) 3319, 2784, 1603, 1444, 764 cm<sup>-1</sup>; <sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  2.57 (s, 3H), 2.72 (s, 3H), 4.13 (s, 2H), 7.35 (s, 1H), 7.60 (dd, *J* = 8.9, 2.3 Hz, 1H), 7.96 (d, *J* = 8.9 Hz, 1H), 7.99 (d, *J* = 2.3 Hz, 1H) ppm; <sup>13</sup>C-NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  25.3, 36.6, 52.1, 121.3, 122.2, 126.0, 129.8, 130.8, 131.4, 144.6, 146.3, 159.2 ppm; HRMS (ESI) calcd for C<sub>12</sub>H<sub>14</sub>ClN<sub>2</sub> [M+H]<sup>+</sup> 221.0840, found 221.0839.

**7-Chloro-2-methyl-4-(N-methylamino)methylquinoline (3iA):** 201.7 mg, 91% yield; yellow oil; IR (neat) 3292, 2791, 1604, 1445, 768 cm<sup>-1</sup>; <sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  2.57 (s, 3H), 2.72 (s, 3H), 4.16 (s, 2H), 7.33 (s, 1H), 7.45 (dd, *J* = 8.8, 2.0 Hz, 1H), 7.97 (d, *J* = 8.8 Hz, 1H), 8.03 (d, *J* = 2.0 Hz, 1H) ppm; <sup>13</sup>C-NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  25.4, 36.6, 52.1, 120.8, 123.7, 124.5, 126.5, 128.2, 134.8, 145.4, 148.5, 160.1 ppm; HRMS (ESI) calcd for C<sub>12</sub>H<sub>14</sub>ClN<sub>2</sub> [M+H]<sup>+</sup> 221.0840, found 221.0839.

**6-Bromo-2-methyl-4-(N-methylamino)methylquinoline (3jA):** yellow solid, mp 123-124 °C; IR (neat) 3315, 2783, 1600, 1443, 785 cm<sup>-1</sup>; <sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  2.58 (s, 3H), 2.71 (s, 3H), 4.14 (s, 2H), 7.35 (s, 1H), 7.73 (dd, *J* = 8.8, 2.0 Hz, 1H), 7.89 (d, *J* = 8.8 Hz, 1H), 8.17 (d, *J* = 2.0 Hz, 1H) ppm; <sup>13</sup>C-NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  25.4, 36.6, 52.0, 119.6, 121.3, 125.5, 126.5, 131.0, 132.4, 144.5, 146.6, 159.3 ppm; HRMS (ESI) calcd for C<sub>12</sub>H<sub>14</sub>BrN<sub>2</sub> [M+H]<sup>+</sup> 265.0335, found 265.0334.

**2-Methyl-6-methoxycarbonyl-4-(N-methylamino)methylquinoline (3kA):** yellow solid, mp 57-59 °C; IR (neat) 3327, 2951, 1703, 1279, 758 cm<sup>-1</sup>; <sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  2.60 (s, 3H), 2.76 (s, 3H), 3.99 (s, 3H), 4.27 (s, 2H), 7.42 (s, 1H), 8.06 (d, *J* = 8.8 Hz, 1H), 8.26 (dd, *J* = 8.8, 1.7 Hz, 1H), 8.78 (d, *J* = 1.7 Hz, 1H) ppm; <sup>13</sup>C-NMR

(100 MHz, CDCl<sub>3</sub>)  $\delta$  25.6, 36.6, 51.9, 52.4, 121.2, 124.5, 126.2, 127.0, 128.7, 129.5, 146.8, 150.0, 161.5, 166.9 ppm; HRMS (ESI) calcd for C<sub>14</sub>H<sub>17</sub>N<sub>2</sub>O<sub>2</sub> [M+H]<sup>+</sup> 245.1285, found 245.1281.

**6-Acetyl-2-methyl-4-(N-methylamino)methylquinoline (3IA):** yellow solid, mp 79-80 °C; IR (neat) 3323, 2925, 1672, 1602, 1260 cm<sup>-1</sup>; <sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  2.61 (s, 3H), 2.74 (s, 3H), 2.76 (s, 3H), 4.27 (s, 2H), 7.42 (s, 1H), 8.08 (d, *J* = 8.8 Hz, 1H), 8.21 (dd, *J* = 8.8, 1.8 Hz, 1H), 8.70 (d, *J* = 1.8 Hz, 1H) ppm; <sup>13</sup>C-NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  25.6, 26.8, 36.7, 52.0, 121.4, 124.6, 125.0, 127.6, 129.7, 133.9, 147.1, 150.0, 161.6, 197.7 ppm; HRMS (ESI) calcd for C<sub>14</sub>H<sub>17</sub>N<sub>2</sub>O [M+H]<sup>+</sup> 229.1335, found 229.1331.

**1-(N-Methylamino)methylisoquinoline (3mA):** 142.7 mg, 83% yield; yellow oil; IR (neat) 3323, 2931, 1562, 822, 743 cm<sup>-1</sup>; <sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  2.60 (s, 3H), 4.38 (s, 2H), 7.56 (d, *J* = 5.8 Hz, 1H), 7.61 (t, *J* = 7.9 Hz, 1H), 7.68 (t, *J* = 7.9 Hz, 1H), 7.83 (d, *J* = 7.9 Hz, 1H), 8.19 (d, *J* = 7.9 Hz, 1H), 8.46 (d, *J* = 5.8 Hz, 1H) ppm; <sup>13</sup>C-NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  36.6, 54.2, 119.9, 124.5, 126.6, 127.2, 127.3, 129.9, 136.0, 141.6, 158.5 ppm; HRMS (ESI) calcd for C<sub>11</sub>H<sub>13</sub>N<sub>2</sub> [M+H]<sup>+</sup> 173.1073, found 173.1071.

**4-Bromo-1-(N-methylamino)methylisoquinoline (3nA):** yellow solid, mp 150 °C (decomp); IR (neat) 3327, 2844, 1255, 924, 755 cm<sup>-1</sup>; <sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  2.59 (s, 3H), 4.35 (s, 2H), 7.68 (t, *J* = 8.1 Hz, 1H), 7.80 (t, *J* = 8.1 Hz, 1H), 8.196 (d, *J* = 8.1 Hz, 1H), 8.198 (d, *J* = 8.1 Hz, 1H), 8.65 (s, 1H) ppm; <sup>13</sup>C-NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  36.6, 54.1, 118.7, 125.0, 126.7, 127.8, 128.2, 131.2, 134.7, 143.3, 158.2 ppm; HRMS (ESI) calcd for C<sub>11</sub>H<sub>12</sub>BrN<sub>2</sub> [M+H]<sup>+</sup> 251.0178, found 251.0181.

**2-(N-Methylamino)methylquinoxaline (3oA)<sup>11a</sup>:** yellow oil; IR (neat) 3318, 2926, 1493, 1126, 758 cm<sup>-1</sup>; <sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  2.58 (s, 3H), 4.14 (s, 2H), 7.72-7.81 (m, 2H), 8.05-8.14 (m, 2H), 8.90 (s, 1H) ppm; <sup>13</sup>C-NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  36.2, 55.2, 129.0, 129.2, 129.4, 130.1, 141.8, 141.9, 145.2, 154.6 ppm; HRMS (ESI) calcd for C<sub>10</sub>H<sub>12</sub>N<sub>3</sub> [M+H]<sup>+</sup> 174.1026, found 174.1022.

**6-(N-Methylamino)methylphenanthridine (3pA):** yellow solid, mp 57-59 °C; IR (neat) 3323, 2930, 1586, 1446, 750 cm<sup>-1</sup>; <sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  2.67 (s, 3H), 4.46 (s, 2H), 7.64 (t, *J* = 8.0 Hz, 1H), 7.67-7.75 (m, 2H), 7.84 (t, *J* = 8.0 Hz, 1H), 8.15 (d, *J* = 8.0 Hz, 1H), 8.26 (d, *J* = 8.0 Hz, 1H), 8.55 (d, *J* = 8.0 Hz, 1H), 8.64 (d, *J* = 8.0 Hz, 1H) ppm; <sup>13</sup>C-NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  36.7, 54.7, 121.9, 122.4, 123.9, 124.8,

125.4, 126.6, 127.4, 128.6, 129.8, 130.4, 132.7, 143.3, 158.2 ppm; HRMS (ESI) calcd for  $C_{15}H_{15}N_2$   $[M+H]^+$  223.1230, found 223.1231.

**2-(*N*-Methylamino)methylquinoline (3qA):** yellow oil; IR (neat) 3303, 2930, 1600, 1504, 747  $cm^{-1}$ ;  $^1H$ -NMR (400 MHz,  $CDCl_3$ )  $\delta$  2.56 (s, 3H), 4.09 (s, 2H), 7.45 (d,  $J$  = 8.4 Hz, 1H), 7.52 (t,  $J$  = 8.0 Hz, 1H), 7.71 (t,  $J$  = 8.0 Hz, 1H), 7.81 (d,  $J$  = 8.0 Hz, 1H), 8.10 (d,  $J$  = 8.0 Hz, 1H), 8.14 (d,  $J$  = 8.4 Hz, 1H) ppm;  $^{13}C$ -NMR (100 MHz,  $CDCl_3$ )  $\delta$  36.4, 57.6, 120.5, 126.1, 127.3, 127.6, 128.8, 129.6, 136.7, 147.5, 160.0 ppm; HRMS (ESI) calcd for  $C_{11}H_{13}N_2$   $[M+H]^+$  173.1073, found 173.1073.

**4-(*N*-Methylamino)methylquinoline (3qA')<sup>11b</sup>:** yellow oil; IR (neat) 3291, 2931, 1593, 1508, 753  $cm^{-1}$ ;  $^1H$ -NMR (400 MHz,  $CDCl_3$ )  $\delta$  2.58 (s, 3H), 4.24 (s, 2H), 7.44 (d,  $J$  = 4.3 Hz, 1H), 7.58 (t,  $J$  = 8.4 Hz, 1H), 7.72 (t,  $J$  = 8.4 Hz, 1H), 8.09 (d,  $J$  = 8.4 Hz, 1H), 8.13 (d,  $J$  = 8.4 Hz, 1H), 8.88 (d,  $J$  = 4.3 Hz, 1H) ppm;  $^{13}C$ -NMR (100 MHz,  $CDCl_3$ )  $\delta$  36.5, 52.1, 119.7, 123.1, 126.5, 127.0, 129.0, 130.1, 145.3, 148.2, 150.3 ppm; HRMS (ESI) calcd for  $C_{11}H_{13}N_2$   $[M+H]^+$  173.1073, found 173.1071.

**2,4-Bis(*N*-methylamino)methylquinoline (3qA''):** 147.3 mg, 84% yield; brown oil; IR (neat) 3285, 2931, 1602, 1445, 755  $cm^{-1}$ ;  $^1H$ -NMR (400 MHz,  $CDCl_3$ )  $\delta$  2.55 (s, 3H), 2.58 (s, 3H), 4.05 (s, 2H), 4.22 (s, 2H), 7.49 (s, 1H), 7.54 (t,  $J$  = 8.1 Hz, 1H), 7.69 (t,  $J$  = 8.1 Hz, 1H), 8.05 (d,  $J$  = 8.1 Hz, 1H), 8.09 (d,  $J$  = 8.1 Hz, 1H) ppm;  $^{13}C$ -NMR (100 MHz,  $CDCl_3$ )  $\delta$  36.4, 36.6, 52.3, 57.9, 119.2, 123.1, 126.0 (2C), 129.1, 129.7, 145.6, 147.9, 160.0 ppm; HRMS (ESI) calcd for  $C_{13}H_{18}N_3$   $[M+H]^+$  216.1495, found 216.1494.

**4-Methyl-2-(pyrrolidin-2'-yl)quinoline (3aE):** 93.4 mg, 88% yield; yellow oil; IR (neat) 3303, 2959, 1603, 1445, 756  $cm^{-1}$ ;  $^1H$ -NMR (400 MHz,  $CDCl_3$ )  $\delta$  1.76-1.86 (m, 1H), 1.86-1.95 (m, 2H), 2.33 (sext,  $J$  = 6.5 Hz, 1H), 2.70 (s, 3H), 3.05-3.13 (m, 1H), 3.25-3.33 (m, 1H), 4.42 (t,  $J$  = 7.5 Hz, 1H), 7.31 (s, 1H), 7.52 (t,  $J$  = 8.3 Hz, 1H), 7.68 (t,  $J$  = 8.3 Hz, 1H), 7.96 (d,  $J$  = 8.3 Hz, 1H), 8.03 (d,  $J$  = 8.3 Hz, 1H) ppm;  $^{13}C$ -NMR (100 MHz,  $CDCl_3$ )  $\delta$  18.8, 26.1, 33.9, 47.4, 63.9, 120.2, 123.6, 125.7, 127.4, 129.0, 129.5, 144.6, 147.1, 163.4 ppm; HRMS (ESI) calcd for  $C_{14}H_{17}N_2$   $[M+H]^+$  213.1386, found 213.1382.

**2-(*N*-Ethyl,*N*-methylamino)methyl-4-methylquinoline (4aA):** 74.9 mg, 70% yield; yellow solid, mp 167 °C (decomp); IR (neat) 2969, 2790, 1603, 1446, 754  $cm^{-1}$ ;  $^1H$ -NMR (400 MHz,  $CDCl_3$ )  $\delta$  1.15 (t,  $J$  = 7.2 Hz, 3H), 2.30 (s, 3H), 2.57 (q,  $J$  = 7.2 Hz, 2H), 2.70 (s, 3H), 3.78 (s, 2H), 7.48 (s, 1H), 7.53 (t,  $J$  = 8.3 Hz, 1H), 7.68 (t,  $J$  = 8.3 Hz, 1H), 7.97 (d,  $J$  = 8.3 Hz, 1H), 8.07 (d,  $J$  = 8.3 Hz, 1H) ppm;  $^{13}C$ -NMR (100 MHz,

CDCl<sub>3</sub>)  $\delta$  12.3, 18.7, 42.1, 51.8, 64.2, 121.7, 123.6, 125.8, 127.4, 129.0, 129.5, 144.5, 147.4, 159.9 ppm; HRMS (ESI) calcd for C<sub>14</sub>H<sub>19</sub>N<sub>2</sub> [M+H]<sup>+</sup> 215.1543, found 215.1539.

**4-Methyl-2-(1'-methylpyrrolidin-2'-yl)quinoline (4aC')**: 68.9 mg, 61% yield, colorless oil; IR (neat) 2941, 2778, 1603, 1447, 757 cm<sup>-1</sup>; <sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  1.82-1.95 (m, 2H), 1.96-2.12 (m, 1H), 2.28 (s, 3H), 2.30-2.45 (m, 2H), 2.71 (s, 3H), 3.30 (t, *J* = 8.4 Hz, 1H), 3.44 (t, *J* = 8.4 Hz, 1H), 7.48 (s, 1H), 7.53 (t, *J* = 8.3 Hz, 1H), 7.68 (t, *J* = 8.3 Hz, 1H), 7.97 (d, *J* = 8.3 Hz, 1H), 8.08 (d, *J* = 8.3 Hz, 1H) ppm; <sup>13</sup>C-NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  18.8, 23.1, 33.8, 40.8, 57.2, 73.4, 119.6, 123.6, 125.7, 127.6, 128.9, 129.6, 144.9, 147.3, 163.6 ppm; HRMS (ESI) calcd for C<sub>15</sub>H<sub>19</sub>N<sub>2</sub> [M+H]<sup>+</sup> 227.1543, found 227.1539.

## ASSOCIATED CONTENTS

## SUPPORTING INFORMATION

The Supporting Information is available free of charge on the ACS Publication website at DOI: <http://pubs.acs.org>.

<sup>1</sup>H NMR and <sup>13</sup>C NMR spectra of all products **2**, **3**, and **4**, and X-ray crystallographic data of **2aD**.

## AUTHOR INFORMATION

E-mail: [togo@faculty.chiba-u.jp](mailto:togo@faculty.chiba-u.jp) (H.T.)

## ACKNOWLEDGEMENT

Financial support in the form of a Grant-in-Aid for Scientific Research (No. 15K05418) from the Ministry of Education, Culture, Sports, Science, and Technology in Japan is acknowledged.

## REFERENCES

- (1) (a) Yang, Y.; Shi, L.; Zhou, Y.; Li, H.; Zhu, Z.; Zhu, H. *Bioorg. Med. Chem. Lett.* **2010**, *20*, 6653-6656. (b) Chen, A.; Aspiotis, R.; Campeau, L.; Cauchon, E.; Chefson, A.; Ducharme, Y.; Falgueyret, J.; Gagné, S.; Han, Y.; Houle, R.; Laliberté, S.; Larouche, G.; Lévesque, J.; McKay, D.; Percival, D. *Bioorg. Med. Chem. Lett.*

- 2011, 21, 7399-7404. (c) Hu, Y.; Lin, E. C. K.; Pham, L. M.; Cajica, J.; Amantea, C. M.; Okerberg, E.; Brown, H. E.; Fraser, A.; Du, L.; Kohno, Y.; Ishiyama, J.; Kozarich, J. W.; Shreder, K. R. *Bioorg. Med. Chem. Lett.* **2013**, 23, 1553-1556. (d) Weitz, E. A.; Chang, J. Y.; Rosenfield, A. H.; Pierre, V. C. *J. Am. Chem. Soc.* **2012**, 134, 16099-16102.
- (2) (a) Giese, B. “*Radicals in Organic Synthesis: Formation of Carbon-Carbon-Bonds*” (Ed.: Baldwin, J. E.), Chap. 5, 210-261, Pergamon Press, 1986. (b) Togo, H. “*Advanced Free Radical Reactions for Organic Synthesis*”, Chap. 5, 157-170, Elsevier, 2004.
- (3) (a) Minisci, F.; Galli, E.; Ceoere, M.; Malatesta, V.; Caronna, T. *Tetrahedron Lett.* **1968**, 5609-5612. (b) Minisci, F.; Galli, R.; Valatesta, V.; Caronna, T. *Tetrahedron* **1970**, 26, 4083-4091. (c) Buratti, W.; Gardini, G. P.; Minisci, F.; Bertini, F.; Galli, G.; Perchinunno, M. *Tetrahedron* **1971**, 27, 3655-3668. (d) Minisci, F. *Synthesis* **1973**, 1-24. (e) Caronna, T.; Citterio, A.; Crolla, T.; Minisci, F. *Tetrahedron Lett.* **1976**, 1731-1732. (f) Caronna, T.; Citterio, A.; Crolla, T.; Minisci, F. *J. Chem. Soc., Perkin Trans 1* **1977**, 865-869. (g) Castaldi, G.; Minisci, F.; Tortelli, V.; Vismara, E. *Tetrahedron Lett.* **1984**, 25, 3897-3900.
- (4) (a) Barton, D. H. R.; Garcia, B.; Togo, H.; Zard, S. Z. *Tetrahedron Lett.* **1986**, 27, 1327-1330. (b) Castagino, E.; Corsano, S.; Barton, D. H. R.; Zard, S. Z. *Tetrahedron Lett.* **1986**, 27, 6337-6338. (c) Togo, H.; Fujii, M.; Ikuma, T.; Yokoyama, M. *Tetrahedron Lett.* **1991**, 32, 3377-3380.
- (5) (a) Togo, H.; Hayashi, K.; Yokoyama, M. *Chem. Lett.* **1991**, 2063-2064. (b) Togo, H.; Hayashi, K.; Yokoyama, M. *Chem. Lett.* **1993**, 641-644. (c) Togo, H.; Hayashi, K.; Yokoyama, M. *Bull. Chem. Soc. Jpn.* **1994**, 67, 2522-2527. (d) Yamazaki, O.; Togo, H.; Matsubayashi, S.; Yokoyama, M. *Tetrahedron Lett.* **1998**, 39, 1921-1924. (e) Yamazaki, O.; Togo, H.; Matsubayashi, S.; Yokoyama, M. *Tetrahedron* **1999**, 55, 3735-3747.
- (6) Zeletchonok, Y. B.; Zorin, V. V.; Klyavlin, M. C. *Tetrahedron Lett.* **1988**, 29, 5037-5038.
- (7) Okugawa, N.; Moriyama, K.; Togo, H. *Eur. J. Org. Chem.* **2015**, 4973-4981.
- (8) (a) Citterio, A.; Gentile, A.; Minisci, F.; Serravalle, M.; Ventura, S. *J. Org. Chem.* **1984**, 49, 3364-3367. (b) Caronna, T.; Gambarotti, C.; Palmisano, L.; Punta, C.;

Recupero, F. *Chem. Commun.* **2003**, 2350-2351. (c) Fang, H.; Zhao, J.; Ni, S.; Mei, H.; Han, J.; Pan, Y. *J. Org. Chem.* **2015**, *80*, 3151-3158. (d) Wang, J.; Li, J.; Huang, J.; Zhu, Q. *J. Org. Chem.* **2016**, *81*, 3017-3022.

- (9) Typical reports; (a) Photochemical amidation of arenes with formamide in acetone; Elad, D. *Tetrahedron Lett.* **1963**, 77-79. (b) Thermal amidation of pyridines with formamides, AgNO<sub>3</sub>, and K<sub>2</sub>S<sub>2</sub>O<sub>8</sub>; Han, W.; Jin, F.; Zhao, Q.; Du, H.; Yao, L. *Synlett* **2016**, 27, 1854-1859.
- (10) Salamone, M.; Milan, M.; DiLabio, G. A.; Bietti, M. *J. Org. Chem.* **2014**, *79*, 7179.
- (11) (a) Butera, J. A.; Spinelli, W.; Anantharaman, V.; Marcopulos, N.; Parsons, R. W.; Moubarak, I. F.; Cullinan, C.; Baglit, J. F. *J. Med. Chem.* **1991**, *34*, 3212-3228. (b) Jansen, K.; Heirbaut, L.; Verkerk, R.; Cheng, J. D.; Joossens, J.; Cos, P.; Maes, L.; Lambeir, A.; De Meester, I.; Augustyns, K.; Veken, P. V. *J. Med. Chem.* **2014**, *57*, 3053-3074.