Journal Pre-Proof

Three-component couplings for the synthesis of pyrroloquinoxalinones by azomethine ylide 1,3-dipolar cycloaddition chemistry

Anthony Choi, Iain Coldham

PII:	S0040-4039(19)30780-4
DOI:	https://doi.org/10.1016/j.tetlet.2019.151023
Reference:	TETL 151023
To appear in:	Tetrahedron Letters
Received Date:	18 July 2019
Revised Date:	2 August 2019
Accepted Date:	7 August 2019



Please cite this article as: Choi, A., Coldham, I., Three-component couplings for the synthesis of pyrroloquinoxalinones by azomethine ylide 1,3-dipolar cycloaddition chemistry, *Tetrahedron Letters* (2019), doi: https://doi.org/10.1016/j.tetlet.2019.151023

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2019 Published by Elsevier Ltd.

Graphical Abstract

To create your abstract, type over the instructions in the template box below. Fonts or abstract dimensions should not be changed or altered.





Tetrahedron Letters journal homepage: www.elsevier.com

Three-component couplings for the synthesis of pyrroloquinoxalinones by azomethine ylide 1,3-dipolar cycloaddition chemistry

Anthony Choi, Iain Coldham*

Department of Chemistry, University of Sheffield, Brook Hill, Sheffield S3 7HF, UK

ARTICLE INFO * Corresponding author. Tel.: +44-114-222-9428; fax: +44-114-222-9346; e-mail: i.coldham@sheffield.ac.uk

Article history: Received Received in revised form Accepted Available online

Keywords: Azomethine ylide 1,3-Dipolar cycloaddition Heterocycle Multicomponent reaction

1-Methyl-3,4-dihydroquinoxalin-2(1*H*)-one was heated with a range of aldehydes to generate intermediate azomethine ylides which underwent [3+2] cycloaddition reactions with *N*-methyl or *N*-phenylmaleimide to give substituted tetrahydropyrroloquinoxalinones. Only one (racemic) stereoisomer was formed in each case and the stereochemical outcome was verified by single crystal X-ray analysis. The products from this multicomponent reaction could be oxidised with DDQ to the pyrroloquinoxalinones.

2019 Elsevier Ltd. All rights reserved.

1. Introduction

Cycloaddition reactions using azomethine ylides have been used widely to synthesise organic molecules containing pyrrolidine and pyrrole rings.¹⁻⁴ There are many ways to prepare the intermediate azomethine ylides and one of the most simple involves the reaction between a secondary amine and an aldehyde. If the amine contains an electron withdrawing group in the α -position, such as an ester, then the resulting iminium ion can be deprotonated to give an azomethine ylide.5-16 As an alternative to an ester, a lactone can be used as a stabilising group to generate azomethine ylides, which has allowed access to a variety of different polycyclic compounds.¹⁷⁻¹⁹ In contrast, electron withdrawing amides or lactams are relatively uncommon as stabilising groups for intermediate azomethine ylides.²⁰⁻²⁵ Sixmembered lactams such as piperazinones are important functional groups which have a large number of applications, especially in medicinal chemistry.²⁶⁻²⁸ However piperazinones and their benzo-fused derivatives (dihydroquinoxalinones, 1) have not, to the best of our knowledge, been used to form azomethine ylides for dipolar cycloaddition reactions. Cycloadducts 2 have an interesting tricyclic structure, where a pyrrolidine is fused with a dihydroquinoxalinone (Scheme 1).









The pyrroloquinoxalinone structure has been synthesised previously using cyclization reactions.^{29–36} However, [3+2] cycloaddition reactions could provide a very efficient route to these compounds, which have the potential for significant biologically activity. Similar structures have been found to have biological relevance as, for example, GABA/benzodiazepine receptor ligands, 5-HT₃ receptor agonists, or vascular smooth muscle relaxants (Fig. 1).^{37–40} Herein, we highlight the use of 3,4-dihydroquinoxalin-2(1*H*)-one **1** in [3+2] cycloaddition reactions to generate a series of novel tetracyclic compounds based on this core ring system.

2. Results and Discussion

We began by synthesising the dihydroquinoxalinone **1** using a known procedure (see supplementary data).⁴¹ With this compound in hand we studied the use of conditions similar to those reported by Siedel and co-workers⁴² to try to effect the desired cycloaddition reaction (Scheme 2). Heating dihydroquinoxalinone **1** with benzaldehyde, 4 Å molecular sieves (MS), benzoic acid and *N*-methylmaleimide in toluene failed to give the desired cycloaddition product and returned only recovered starting materials. Substituting benzoic acid for the stronger acid camphorsulfonic acid (CSA) gave a similar result, but oxidised compound **3** was isolated instead of compound **1**. It was clear that oxidation of the starting material was interfering

with the desired reaction.⁴³ To investigate the oxidation reaction further, dihydroquinoxalinone **1** was heated at reflux in toluene with 10 mol% CSA in the absence of aldehyde or maleimide. Analysis of the reaction mixture after 17 h by ¹H NMR spectroscopy indicated a mixture of dihydroquinoxalinone **1** and its oxidised form **3** in a 1:1.2 ratio. In contrast, by carrying out the same reaction under an argon atmosphere, ¹H NMR spectroscopy of the reaction mixture indicated that no oxidised material **3** had been formed.



Scheme 2. Initial attempted cycloaddition reaction.

We therefore repeated the cycloaddition reaction under an inert atmosphere and were pleased to find that the desired cycloadduct 4 was formed as a single racemic diastereoisomer in moderate yield (Scheme 3). The reaction conditions were optimised as shown in Table 1. Initially increasing the amount of CSA caused an increase in the yield of cycloadduct 4, with a maximum yield of 57% being obtained when 15 mol% of CSA was added (Entry 3). Decreasing the reaction temperature caused a decrease in the yield of cycloadduct 4 (Entry 5). Although the addition of the drying agent MgSO4 did not affect the yield (Entry 6), the use of 4 Å molecular sieves was beneficial (Entry 7). Changing the molar ratio of dihydroquinoxalinone 1 and benzaldehyde gave variable yields of cycloadduct 4 (Entries 8-10) with the best result being obtained when using 1.5 eq. dihydroquinoxalinone 1 and 1 eq. benzaldehyde (Entry 8). However, it was possible to obtain a good yield with 1 eq. of each of component: 1, PhCHO, and N-methylmaleimide (Entry 9). Finally, when the acid was changed from CSA to benzoic acid there was a decrease in the yield (Entry 11).



Scheme 3. Cycloaddition to give tetracycle 4.

Table 1 . Optimization of the cycloaddition in Scheme 3.			
Entry	Conditions	Yield 4 (%)	
1	2 eq. 1, no acid, 110 °C, no drying agent	0	
2	2 eq. 1, 10 mol% CSA, 110 °C, no drying agent	47	
3	2 eq. 1, 15 mol% CSA, 110 °C, no drying agent	57	
4	2 eq. 1, 20 mol% CSA, 110 °C, no drying agent	49	
5	2 eq. 1, 15 mol% CSA, 90 °C, no drying agent	40	
6	2 eq. 1, 15 mol% CSA, 110 °C, MgSO ₄	48	
7	2 eq. 1, 15 mol% CSA, 110 °C, 4 Å MS	63	
8	1.5 eq. 1, 15 mol% CSA, 110 °C, 4 Å MS	69	
9	1 eq. 1, 15 mol% CSA, 110 °C, 4 Å MS	54	
10	1.5 eq. PhCHO, 15 mol% CSA, 110 °C, 4 Å MS	25	
11	1.5 eq. 1, 15 mol% PhCO ₂ H, 110 °C, 4 Å MS	32	

The stereochemistry of compound **4** was confirmed by single crystal X-ray crystallography (Fig. 2), which showed a *trans* configuration between the protons α to the amine nitrogen atom and therefore that the [3+2] cycloaddition occurs with the S-

shaped ylide (assuming a concerted process, although it is possible that the reaction is stepwise). The dipolarophile *N*-methyl maleimide approaches *trans* to the phenyl group (*endo* selectivity).





Using the optimised conditions, we then explored the substrate scope of the reaction. By varying the aldehyde, we were able to obtain a range of different cycloadducts 5-8 (Scheme 4). The reaction was successful with both electron-poor (4-nitro) and electron-rich (4-methoxy) benzaldehydes, although the yield of cycloadduct 6 in the latter case was low. It was pleasing to find that the reaction was successful with a heteroaryl group (thiophenyl) and with the non-aromatic aldehyde cyclohexane-carbaldehyde, which gave the desired cycloadducts 7 and 8 respectively. In all cases for products 5-8 only a single stereoisomer was isolated and each had very similar coupling constants in the ¹H NMR spectra, suggesting that they all have the same relative stereochemistry as determined for adduct 4.



Scheme 4. Exploring the scope of the aldehyde RCHO.

We then attempted to expand the scope of the dipolarophile by testing different electron-poor alkenes. However mostly starting materials and no desired cycloadducts were isolated on heating amine **1** and PhCHO with dimethyl fumarate, methyl *trans*-cinnamate, diethyl acetylenedicarboxylate, or benzylidenemalononitrile. The reaction was successful with *N*-phenylmaleimide to give cycloadduct **9** in 39% yield (Scheme 5).



Scheme 5. Cycloaddition with N-phenylmaleimide.

Finally, the scope of chemistry was extended by oxidation of cycloadducts **4** and **7** using 4,5-dichloro-3,6-dioxo-1,4-

cyclohexadiene-1,2-dicarbonitrile (DDQ). This resulted in the formation of compounds **10** and **11**, respectively, in good yields (Scheme 6).



Scheme 6. Oxidation of the cycloadducts.

3. Conclusion

In conclusion, we have demonstrated a successful and simple protocol for a multicomponent coupling reaction involving azomethine ylide dipolar cycloaddition chemistry from dihydroquinoxalinone **1** and a variety of aldehydes. The scope of the dipolarophile needs further study but the chemistry allows access to novel heterocyclic products as single stereoisomers. The strategy provides a way to prepare libraries of compounds based on the pyrroloquinoxalinone core for biological screening.

4. Acknowledgments

We acknowledge support from the EPSRC and the University of Sheffield. We thank Craig Robertson for the single crystal Xray analysis, Sandra van Meurs and Craig Robertson for help with NMR spectroscopy, and the Faculty of Science mass spectrometry service.

5. References and notes

- 1. Li, J.; Ye, Y.; Zhang, Y. Org. Chem. Front. 2018, 5, 864–892.
- Burrell, A. J. M.; Coldham, I. *Curr. Org. Synth.* 2010, 7, 312–331.
 Pandey, G.; Banerjee, P.; Gadre, S. R. *Chem. Rev.* 2006, *106*,
- 4484–4517.4. Coldham, I.; Hufton, R. *Chem. Rev.* 2005, *105*, 2765–2810.
- Confalone, P. N.; Huie, E. M. J. Am. Chem. Soc. 1984, 106, 7175– 7177.
- Grigg, R.; Duffy, L. M.; Dorrity, M. J.; Malone, J. F.; Rajviroongit, S.; Thornton-Pett, M. *Tetrahedron* 1990, 46, 2213– 2230.
- Coldham, I.; Coles, S. J.; Crapnell, K. M.; Fernàndez, J.-C.; Haxell, T. F. N.; Hursthouse, M. B.; Moseley, J. D.; Treacy, A. B. *Chem. Commun.* 1999, 1757–8.
- Coldham, I.; Crapnell, K. M.; Moseley, J. D.; Rabot, R. J. Chem. Soc., Perkin Trans. 1 2001, 1758–1763.
- Coldham, I.; Crapnell, K. M.; Fernàndez, J.-C.; Moseley, J. D.; Rabot, R. J. Org. Chem. 2002, 67, 6181–6187.
- Coldham, I.; Dobson, B. C.; Franklin, A. I.; Fletcher, S. R. *Tetrahedron Lett.* 2007, 48, 873–875.
- Coldham, I.; Dobson, B. C.; Fletcher, S. R.; Franklin, A. I. Eur. J. Org. Chem. 2007, 2676–2686.
- 12. Burrell, A. J. M.; Watson, L.; Martin, N. G.; Oram, N.; Coldham, I. Org. Biomol. Chem. 2010, 8, 4530–4532.
- Coldham, I.; Burrell, A. J. M.; Guerrand, H. D. S.; Watson, L.; Martin, N. G.; Oram, N. *Beilstein J. Org. Chem.* 2012, *8*, 107– 111.
- Sutariya, T. R.; Labana, B. M.; Parmar, N. J.; Kant, R.; Gupta, V. K.; Plata, G. B.; Padrón, J. M. New J. Chem. 2015, 39, 2657–2668.

- Pathak, R.; Dobson, B. C.; Ghosh, N.; Ageel, K. A.; Alshawish, M. R.; Saruengkhanphasit, R.; Coldham, I. Org. Biomol. Chem. 2015, 13, 3331–3340.
- 16. Yamada, R.; Adachi, Y.; Yokoshima, S.; Fukuyama, T. *Angew. Chem. Int. Ed.* **2016**, *55*, 6067–6070.
- Anslow, A. S.; Harwood, L. M.; Phillips, H.; Watkin, D.; Wong, L. F. *Tetrahedron: Asymmetry* 1991, *2*, 1343–1358.
- Williams, R. M.; Zhai, W.; Aldous, D. J.; Aldous, S. C. J. Org. Chem. 1992, 57, 6527–6532.
- Lo, M. M.-C.; Neumann, C. S.; Nagayama, S.; Perlstein, E. O.; Schreiber, S. L. J. Am. Chem. Soc. 2004, 126, 16077–16086.
- Ollis, W. D.; Stanforth, S. P.; Ramsden, C. A. J. Chem. Soc. Perkin Trans. 1 1989, 965–968.
- Garner, P.; Ho, W. B.; Grandhee, S. K.; Youngs, W. J.; Kennedy, V. O. J. Org. Chem. 1991, 56, 5893–5903.
- 22. Peyronel, J.-F.; Grisoni, S.; Carboni, B.; Courgeon, T.; Carrié, R. *Tetrahedron* **1994**, *50*, 189–198.
- 23. Trávníček, M.; Pospíšil, J.; Potáček, M. Collect. Czech. Chem. Commun. 1999, 64, 1993–2006.
- 24. Pospíšil, J.; Trávníček, M.; Potáček, M. Arkivoc 2001, ii, 146-162.
- 25. Choi, A.; Morley, R. M.; Coldham, I. *Beilstein J. Org. Chem.* **2019**, *15*, 1480–1484.
- Kakarla, R.; Liu, J.; Naduthambi, D.; Chang, W.; Mosley, R. T.; Bao, D.; Steuer, H. M. M.; Keilman, M.; Bansal, S.; Lam, A. M.; Seibel, W.; Neilson, S.; Furman, P. A.; Sofia, M. J. *J. Med. Chem.* 2014, *57*, 2136–2160.
- Zhuo, C.-X.; Zhang, X.; You, S.-L. ACS Catal. 2016, 6, 5307– 5310.
- Sun, A. W.; Hess, S. N.; Stoltz, B. M. Chem. Sci. 2019, 10, 788– 792.
- Taylor, E. C.; McKillop, A.; Ross, R. E. J. Am. Chem. Soc. 1965, 87, 1990–1995.
- Kasatkina, S. O.; Stepanova, E. E.; Dmitriev, M. V; Maslivets, A. N. Russ. J. Org. Chem. 2018, 54, 1055–1060.
- 31. Chicharro, R.; de Castro, S.; Reino, J. L.; Arán, V. J. *Eur. J. Org. Chem.* **2003**, 2314–2326.
- 32. Li, D.; Ollevier, T. Eur. J. Org. Chem. 2019, 1273-1280.
- Luo, X.; Chenard, E.; Martens, P.; Cheng, Y.-X.; Tomaszewski, M. J. Org. Lett. 2010, 12, 3574–3577.
- 34. Xu, L.; Jiang, Y.; Ma, D. Synlett 2010, 2285–2288.
- Jamieson, C.; Congreve, M. S.; Emiabata-Smith, D. F.; Ley, S. V.; Scicinski, J. J. Org. Process Res. Dev. 2002, 6, 823–825.
- Tanimori, S.; Nishimura, T.; Kirihata, M. Bioorg. Med. Chem. Lett. 2009, 19, 4119–4121.
- TenBrink, R. E.; Im, W. B.; Sethy, V. H.; Tang, A. H.; Carter, D. B. J. Med. Chem. 1994, 37, 758–768.
- Tang, A. H.; Franklin, S. R.; Himes, C. S.; Ho, P. M. J. Pharmacol. Exp. Ther. 1991, 259, 248–254.
- Campiani, G.; Morelli, E.; Gemma, S.; Nacci, V.; Butini, S.; Hamon, M.; Novellino, E.; Greco, G.; Cagnotto, A.; Goegan, M.; Cervo, L.; Valle, F. D.; Fracasso, C.; Caccia, S.; Mennini, T. J. Med. Chem. 1999, 42, 4362–4379.
- Abou-Gharbia, M.; Freed, M. E.; McCaully, R. J.; Silver, P. J.; Wendt, R. L. J. Med. Chem. 1984, 27, 1743–1746.
- Wang, F.; Hu, B.-L.; Liu, L.; Tu, H.-Y.; Zhang, X.-G. J. Org. Chem. 2017, 82, 11247–11252.
- 42. Mantelingu, K.; Lin, Y.; Seidel, D. Org. Lett. 2014, 16, 5910– 5913.
- For oxidation of 3,4-dihydroquinoxalin-2(1*H*)-ones see, for example, Akula, P. S.; Hong, B.-C.; Lee, G.-H. *RSC Adv.* 2018, *8*, 19580–19584.

7. Supplementary Material

Experimental procedures, spectroscopic and X-ray data (CCDC 1939356 for compound 4) and copies of NMR spectra.

Highlights

Ms. Ref. No.: TETL-D-19-01047 *Title:* Three-component couplings for the synthesis of pyrroloquinoxalinones by azomethine ylide 1,3-dipolar cycloaddition chemistry

- Multicomponent reaction of aldehydes, an amine and an alkene
- Stereoselective preparation of tetracycles
- Dipolar cycloaddition of azomethine ylides to pyrroloquinoxalinones

4