This article was downloaded by: [Universität Stuttgart] On: 17 October 2014, At: 01:10 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: <u>http://www.tandfonline.com/loi/gpss20</u>

Reaction Mechanism Studies of Solvolytic Displacement of Chloride from Phosphorus

Han Joong Koh^a, Suk Jin Kang^a & Dennis N. Kevill^b ^a Department of Science Education, Chonju National University of Education, Chonju, Korea

^b Department of Chemistry and Biochemistry, Northern Illinois University, DeKalb, Illinois, USA Published online: 21 Jun 2008.

To cite this article: Han Joong Koh , Suk Jin Kang & Dennis N. Kevill (2008) Reaction Mechanism Studies of Solvolytic Displacement of Chloride from Phosphorus, Phosphorus, Sulfur, and Silicon and the Related Elements, 183:2-3, 364-368, DOI: 10.1080/10426500701734943

To link to this article: <u>http://dx.doi.org/10.1080/10426500701734943</u>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or

indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions



Reaction Mechanism Studies of Solvolytic Displacement of Chloride from Phosphorus

Han Joong Koh,¹ Suk Jin Kang,¹ and Dennis N. Kevill²

¹Department of Science Education, Chonju National University of Education, Chonju, Korea ²Department of Chemistry and Biochemistry, Northern Illinois University, DeKalb, Illinois, USA

The extended Grunwald-Winstein equation is applied to solvolytic displacement at phosphorus. Previous studies of the solvolyses of phosphorochloridates $(RO)_2POCl, Ph_2POCl and [(CH_3)_2N]_2POCl$ are extended to cyclic phosphorochloridates, with five and six membered rings. Both kinetic and product ratio measurements are made. The equation is also applied to kinetic studies of the solvolyses of Ph_2PCl. The l and m sensitivity values are compared with previous values and with values for attack at sulfur, to give useful mechanistic information.

Keywords Chlorodiphenylphosphine; cyclic phosphorochloridates; extended Grunwald-Winstein equation; solvolysis

INTRODUCTION

The extended Grunwald-Winstein¹ equation is used to correlate the specific rates of solvolysis reactions in terms of solvent ionizing power² and solvent nucleophilicity³. In Equation (1), k and k_0 represent the specific rates of solvolysis in a given solvent and in the

$$\log(k/k_0) = lN_T + mY_x + c \tag{1}$$

standard solvent (80% ethanol), respectively; l represents the sensitivity to changes in solvent nucleophilicity (N_T); m represents the

H.J.K thanks Chonju National University of Education for the granting of a sabbatical leave during the time that a portion of this research was performed. Acknowledgement is made to the donors of the Petroleum Research Fund, administered by the American Chemical Society, for partial support of this research.

Address correspondence to Dennis N. Kevill, Department of Chemistry and Biochemistry, Northern Illinois University, Dekalb, Illinois, 60115-2862 USA. E-mail: dkevill.niu.edu

sensitivity to changes in solvent ionizing power (Y_X for a leaving group X); c represents a constant (residual) term.

The scale was initially developed for nucleophilic attack at sp³carbon but it has been successfully applied to attack at sp²-carbon⁴ and at phosphorus^{5–7} and sulfur.⁸ The solvolyses at phosphorus have included rate and product studies with dimethyl phosphorochloridate as substrate, and a direct displacement (S_N2) reaction was proposed as the pathway.⁶ In the present paper, the study is extended to the solvolyses of two cyclic phosphorochloridate esters, 2chloro-1,3,2-dioxaphospholane-2-oxide (1) and 2-chloro-5,5-dimethyl-1,3,2-dioxaphosphorinane-2-oxide (2). The study of diphenylphosphinyl chloride⁷ was also explained in terms of S_N2 attack and this has been extended to the phosphorus(III)-containing chlorodiphenylphosphine (3) (Scheme 1).



SCHEME 1

KINETIC STUDIES AND CORRELATION OF THE SPECIFIC RATES

The specific rates of solvolysis of **1** and **2** have been measured using a rapid response conductivity apparatus, in a wide range of solvents, including binary mixtures of water with 2,2,2-trifluoroethanol (TFE) and 1,1,1,3,3,3-hexafluoro-2-propanol (HFIP). The values at 25.0°C for **1** and the values at 50.0°C for **2** are determined using ethanol and methanol and their aqueous binary mixtures plus mixtures of water with acetone, TFE, and HFIP. These values are used in correlations using eqn. 1 against literature values for N_T^3 and Y_{Cl} .^{2.9} The sensitivity values together with measures of the goodness-of-fit are reported in Table I, where they are compared with literature values for solvolytic displacements of chloride from sulfur (VI) or phosphorus (V).

There have been strong indications¹⁰that nucleophilic attack at a phosphorus within a 5-membered ring shows significant pathway differences to other situations. A large^{11,12}(sometimes enormous¹¹) rate acceleration is frequently observed and, whereas inversion is normally observed, reactions of 2-chloro-4,5-dimethyl-1,3,2-dioxaphospholane-2-oxide give predominantly retention of configuration.¹² The present analyses give no support for any special effects during solvolyses, with

_						
Substrate	n ^a	T °C	l^b	m^b	R^{c}	
$(Me_2N)_2POCl^d$	31	25.0	1.20 ± 0.07	0.69 ± 0.04	0.958	
$(MeO)_2POCl^d$	18	25.0	1.27 ± 0.14	0.47 ± 0.08	0.941	
$(MeO)_2 PSCl^d$	28	25.0	1.17 ± 0.07	0.56 ± 0.03	0.966	
Ph_2POCl^d	27	25.0	1.42 ± 0.10	0.54 ± 0.07	0.956	
1	28	25.0	1.16 ± 0.06	0.60 ± 0.03	0.974	
2	27	50.0	1.03 ± 0.05	0.43 ± 0.02	0.974	
3	25	25.0	1.25 ± 0.09	0.46 ± 0.06	0.954	
p-MeOC ₆ H ₄ SO ₂ Cl ^e	37	25.0	1.10 ± 0.17	0.61 ± 0.04	0.959	
i-PrSO ₂ Cl ^e	19	25.0	1.28 ± 0.05	0.64 ± 0.03	0.988	
${ m Me_2NSO_2Cl^e}$	26	25.0	1.17 ± 0.06	0.70 ± 0.04	0.975	

TABLE I Grunwald-Winstein Correlations for Attack atPhosphorus and Sulfur

^{*a*}Number of data points; ^{*b*}from Eq. (1); ^{*c*} correlation coefficient; ^{*d*}from Ref. 7; and ^{*e*}from Ref. 8.

the l and m values for 1 being similar to those for other entries within Table I. Also, a comparison of the specific rates (Table II) shows 1 only modestly more reactive than the open-chain analog.

The phosphorus(III)- containing compound Ph₂PCl (**3**) not only shows *l* and *m* values very similar to the phosphorus(V)- containing compound Ph₂POCl⁷but the actual specific rates in a given solvent are very similar. For example, at 25.0°C, the specific rates in ethanol are 8.09×10^{-3} s⁻¹ and 7.02×10^{-3} s⁻¹ and, in 97% HFIP, they are 6.16×10^{-6} s⁻¹ and 5.91×10^{-6} s⁻¹, respectively. Studies with the replacement of phenyl by other groupings are needed to explore the generality of this

-				
Solvent	$10^4 k [(CH_3O)_2 POCl]^a$	$10^4 \mathrm{k}(1)^b$	10^4 k(2) ^b	
100%EtOH	2.96	37.8	0.13^{c}	
80%EtOH	28.5	328	0.27^{c}	
100%MeOH	11.3	48.7	0.12^c	
80%MeOH	49.6	480	_	
97%TFE	0.036	1.13	_	
50%TFE	18.6	259	_	
70%HFIP	0.71	55.4	_	

TABLE II Specific Rates of Solvolysis at 25.0°C, in a Few Typical Solvents, for Dimethyl Phosphorochloridate, 1 and 2

^{*a*}From ref. 6; ^{*b*}this Work; and ^{*c*}extrapolated using 3 or 4 values determined at higher temperatures.

Solvent ^b :	90E	70E	50E	20E	90M	70M	40M	20M	80T	70T	50T
S(2):	1.97	1.43	1.37	2.01	2.20	1.91	1.32	1.66	0.23	0.09	0.07
$S[(CH_3O)_2POCl]:$	0.36^c	0.56^{c}	0.85^c								

TABLE III Selectivity Values (S)^a in Solvolyses of 2 at 50.0°C

^{*a*}From Eq. (3); ^{*b*}E, M, T represent ethanol, methanol, and TFE, with its percentage component given (remainder water); ethanol-water and methanol-water on volume-volume basis at 25.0°C and TFE-water on weight-weight basis; ^{*c*}at 25.0°C.

observation. The acid catalysis frequently observed in substitutions at phosphorus (III)¹³ appears not to operate.

PRODUCT SELECTIVITY

The reactions of 1 were sufficiently fast for appreciable reaction to occur during mixing. For 2 infinity titers could be used to determine the product partitioning [Equation (2)]. These values can be used in conjuction with the composition of a water-alcohol mixture to



arrive at selectivity values (S) defined as in Equation 3. Values for the selectivity are

$$S = \frac{[ester in product][H_2O in solvent]}{[acid in product][ROH in solvent]}$$
(3)

compared to those previously observed for dimethyl phosphorochloridate in Table III. The values in aqueous ethanol and aqueous methanol for the open-chain compound are slightly below unity as opposed to slightly above unity for **2**. However, the variation in the energy difference between the two pathways of Equation 2 would only have to be small to account for these changes in S value. As expected, TFE does not compete effectively against water.

REFERENCES

- [1] S. Winstein, E. Grunwald, and H. W. Jones, J. Am. Chem. Soc., 73, 2700 (1951).
- [2] T.W. Bentley and G. Llewellyn, Prog. Phys. Org. Chem., 17, 121 (1990).

- [3] D. N. Kevill, In Advances in Quantitative Structure-Property Relationships, M. Charton, Ed. (JAI Press, Greenwich, CT, 1996), Vol. I, pp 81–115.
- [4] D. N. Kevill and M. J. D'Souza, J. Chem. Soc., Perkin Trans. 2, 1721 (1997).
- [5] D. N. Kevill and B. Miller, J. Org. Chem., 67, 7399 (2002).
- [6] D. N. Kevill and J. S. Carver, Org. Biomol. Chem., 2, 2040 (2004).
- [7] D. N. Kevill and H. J. Koh, J. Phys. Org. Chem., DOI:10.1002/poc.1124.
- [8] D. N. Kevill, B. -C. Park, K. -H. Park, M. J. D'Souza, L. Yaakoubd, S. L. Mlynarski, and J. B. Kyong, Org. Biomol. Chem., 4, 1580 (2006), and references therein.
- [9] (a) T. W. Bentley and G. E. Carter, J. Am. Chem. Soc., 104, 5741 (1982); (b) D.N. Kevill and M. J. D'Souza, J. Chem. Res. Synop., 174 (1993); (c) J. S. Lomas, M. J. D'Souza, and D. N. Kevill, J. Am. Chem. Soc., 117, 5891 (1995).
- [10] A. J. Kirby and S. G. Warren, *The Organic Chemistry of Phosphorus* (Elsevier, New York, 1967), Chap. 10, pp. 341–349.
- [11] P. C. Haake and F. H. Westheimer, J. Am. Chem. Soc., 83, 1102 (1961).
- [12] R. J. P. Corriu, G. F. Lanneau, and D. Leclercq, Phosphorus and Sulfur, 18, 197 (1983).
- [13] O. Dahl, Phosphorus and Sulfur, 18, 201 (1983).