


ChemComm

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: V. Dhayalan, C. Sämman and P. Knochel, *Chem. Commun.*, 2015, DOI: 10.1039/C4CC08846H.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

COMMUNICATION

Synthesis of polyfunctional secondary amines by the addition of functionalized zinc reagents to nitroso-arenes[†]

Vasudevan Dhayalan, Christoph Sämann and Paul Knochel*

Cite this: DOI: 10.1039/x0xx00000x

Received 00th Month 2014,
Accepted 00th Month 2014


DOI: 10.1039/x0xx00000x

www.rsc.org/

Addition of functionalized aryl, heteroaryl or adamantyl zinc reagents to various nitroso-arenes in the presence of magnesium salts and LiCl in THF produces after a reductive work-up with FeCl₂ and NaBH₄ in ethanol the corresponding polyfunctional secondary amines in high yields.

The preparation of arylamines is an important synthetic goal since these compounds often have useful properties for pharmaceuticals or material science applications.¹ Transition metal catalyzed aminations have been well studied,² but the use of expensive and toxic metallic catalysts reduces somewhat the utility of such synthetic methods. Another approach has been the use of electrophilic nitrogen reagents and their reactions with non-expensive and low toxic main-group organometallics.³ A few years ago, we have reported that functionalized arylmagnesium reagents add to nitroso-arenes⁴ and nitro-arenes.⁵ Although satisfactory yields were obtained, the high reactivity of the carbon-magnesium bond reduces the functional group tolerance. Furthermore, nitroso-arenes have proven to be versatile reagents for performing nitroso aldol and related reactions.⁶

Herein, we wish to report a mild synthesis of diaryl or heteroaryl(aryl)amines as well as functionalized highly sterically hindered adamantyl(aryl)amines. Thus, the treatment of an arylzinc derivative **2**, prepared either by the direct insertion of Mg in the presence of LiCl and ZnCl₂⁷ or by a I/Mg-exchange with *i*PrMgCl·LiCl⁸ followed by transmetalation with ZnCl₂, with various nitroso-arenes of type **3**⁹ affords an intermediate zincated hydroxylamine derivative **4** which after reductive work-up with













Scheme 1 Synthesis of polyfunctional secondary amines of type **5** via the addition of functionalized zinc reagents of type **2** to various nitroso compounds of type **3**.

Department of Chemistry, Ludwig-Maximilians-Universität München, Butenandstr. 5-13, 81377 Munich, Germany. Tel: +49-(0)89-2180-77679, Fax: +49-(0)89-2180-77680. E-mail: paul.knochel@cup.uni-muenchen.de

[†]Electronic Supplementary Information (ESI) available: Detailed experimental procedures and spectroscopic data for all compounds. See DOI: 10.1039/b000000x/

Table 1 Synthesis of diarylamines of type **5** via the reaction of functionalized arylzinc reagents **2a-j** with nitrosobenzene **3a**

Entry	Zn-reagent	Product, Yield (%) ^{a,b}
1	2a , PhZnCl	 5a : 85
2	 2b , FG = CF ₃	 5b : 69
3	2c , FG = CO ₂ Et ^c	5c : 76
4	2d , FG = <i>t</i> Bu	5d : 96
5	2e , FG = SCH ₃	5e : 70
6	2f , FG = OCH ₃	5f : 78
7	2g , FG = OTMS	 5g : 89 ^d
8	 2h	 5h : 72
9	 2i ^e	 5i : 64 ^f
10	 2j ^e	 5j : 75 ^f

^a General reaction conditions: arylzinc reagent (1.1 equiv), nitroso electrophile (1.0 equiv), NaBH₄ (1.0 equiv), FeCl₂ (2.0 equiv). ^b Yield of analytically pure isolated product as determined by ¹H-NMR analysis. ^c Prepared by I/Mg-exchange with *i*PrMgCl·LiCl.⁸ ^d The TMS-group was cleaved during workup and column chromatography purification. ^e The arylzinc reagents (**2i** and **2j**) were prepared from the corresponding bromides (see Supporting Information). ^f Obtained after removal of the ethylene glycol group with CF₃CO₂H in CH₂Cl₂ (see Supporting Information).

Table 2 Synthesis of polyfunctional secondary amines of type **5** by the addition of aryl and heteroaryl zinc reagents **2a-k** to various nitroso compounds **3a-g**

Ar ¹ -ZnX + Ar ² -NO		1. THF, 0 to 25 °C, 2-3 h 2. FeCl ₂ (2 equiv) NaBH ₄ (1 equiv) EtOH, 25 °C, 15 h		Ar ¹ -NH-Ar ²	Yield (%) ^{a,b}
2	3			5	
Entry	Zn-reagent (Ar ¹)	Electrophile (Ar ²)		Product	
1	2a			5k	79
2	2a			5l	77
3	2a			5m	83
4				5n	97
5				5o	90
6				5p	81
7				5q	97
8				5r	97 ^c
9				5s	63
10	2k			5t	55
11	2k			5u	60
12	2a			5v	67
13	2f			5w	96

Table 2 Continued

Entry	Zn-reagent (Ar ¹)	Electrophile (Ar ²)	Product	Yield (%) ^{a,b}
14	2c^d			83
15	2k			70

^a General reaction conditions: arylzinc reagent (1.1 equiv), nitroso electrophile (1.0 equiv), NaBH₄ (1.0 equiv), FeCl₂ (2.0 equiv). ^b Yield of analytically pure isolated product as determined by ¹H-NMR analysis. ^c The TMS-group was cleaved during the workup and column chromatography purification. ^d Prepared by I/Mg-exchange with *i*PrMgCl·LiCl.⁸

FeCl₂ and NaBH₄ in ethanol (25 °C, 15 h) produce the corresponding secondary amines of type **5** in excellent yields (Scheme 1). A range of functional groups have been tolerated in the starting arylzinc reagent as shown in Table 1.

Thus, PhZnCl (1.1 equiv) prepared by the direct insertion of Mg in the presence of LiCl and ZnCl₂ reacts with nitrosobenzene **3a** (1.0 equiv) at 25 °C within 2-3 h and produces after reductive work-up with FeCl₂ (2.0 equiv) and NaBH₄ (1.0 equiv) in ethanol (25 °C, 15 h) the corresponding diphenylamine **5a** in 85% yield (Table 1, entry 1).^{10a} The presence of both Mg salts and LiCl were found to be essential for achieving a high yield. A variety of arylzinc reagents prepared similarly were used in the addition to **3a**. Both electron withdrawing and donating groups can be attached at the aryl ring (Table 1, entries 2-8).^{10b-g} Arylzinc reagent **2c** has been prepared *via* an I/Mg-exchange,⁸ its reaction with nitrosobenzene (**3a**) furnishes the corresponding secondary amine **5c** in 76% yield (Table 1, entry 3). Although sensitive functional groups like a formyl or an acetyl group are not tolerated, the corresponding bromoacetal (**1i**) or bromoketal (**1j**) are readily converted to the zinc reagents (**2i** and **2j**) by the insertion of Mg in the presence of LiCl and ZnCl₂.⁷ The addition of nitrosobenzene (**3a**) provides after removal of the ethylene glycol protecting group (CF₃CO₂H in CH₂Cl₂ at 25 °C for 5-8 h) the secondary amines (**5i** and **5j**) in 64-75% yield (Table 1, entries 9 and 10).

Table 3 Synthesis of tertiaryalkyl(aryl)amines **7a-d** by the addition of tertiary alkylzinc reagents **6a** and **6b** to nitroso compounds **3a**, **3f** and **3h**

Entry	Zn-reagent	Electrophile	Product, Yield (%) ^{a,b}
1		3a	 7a : 50
2		3a	 7b : R = H (89)
3	6b		7c : R = NMe ₂ (71)
4	6b	3f	7d : R = OMe (56)

^a General reaction conditions: alkylzinc reagent (1.1 equiv), nitroso electrophile (1.0 equiv), NaBH₄ (1.0 equiv), FeCl₂ (2.0 equiv). ^b Yield of analytically pure isolated product as determined by ¹H-NMR analysis. ^c Prepared by transmetalation of commercially available *t*-BuMgCl with ZnCl₂.


This addition reaction can be extended to various nitroso-arenes (commercially available) or prepared according to the method of Bäckvall.¹¹ Again, electron-donating or accepting substituents are tolerated in the aryl nitroso reagents of type **3** furnishing the corresponding diarylamines **5k-r** in 77-97% yield (Table 2, entries 1-8).^{5a,10h-k} Noteworthy, a heterocyclic zinc reagent (**2k**) has also been used as well as a nitrosopyridine (**3g**)¹² leading to heteroaryl(aryl)-amines **5s-y** in 55-96% yield (Table 2, entries 9-15). Moreover, tertiary alkylzinc reagents such as *t*-BuZnCl (**6a**) and adamantylzinc chloride (**6b**)¹³ add to various nitroso-arenes under similar reaction conditions producing otherwise difficult to prepare tertiaryalkyl(aryl)amines **7a-d** in 50-89% yield (Table 3, entries 1-4).¹⁴

In summary, we have shown that aryl, heteroaryl or adamantyl zinc reagents add to various nitroso-arenes in the presence of Mg-salts and LiCl. Both Mg and Li salts are necessary to achieve high yields for the synthesis of the corresponding functionalized secondary amines. Further extensions of this work are currently underway in our laboratories.

The research leading to these results has received funding from the European Research Council under the European Community's Seventh Framework Programme (FP7/2007–2014) ERC grant agreement no. 227763. We thank the Fonds der Chemischen Industrie for financial support. We also thank Heraeus Holding GmbH (Hanau), Rockwood Lithium (Frankfurt), and BASF SE (Ludwigshafen) for the generous gift of chemicals.

Notes and references

- (a) G. Thomas, *Medicinal Chemistry*; Wiley-VCH: Weinheim, 2000; (b) S. Liu, X. Jiang, H. Ma, M. S. Liu and A. K.-Y. Jen, *Macromolecules*, 2000, **33**, 3514; (c) W.-L. Yu, J. Pei, W. Huang and A. J. Heeger, *Chem. Commun.*, 2000, 681; (d) E. Bellmann, S. E. Shaheen, S. Thayumanavan, S. Barlow, R. H. Grubbs, S. R. Marder, B. Kippelen and N. Peyghambarian, *Chem. Mater.*, 1998, **10**, 1668; (e) H. Inada, Y. Yonemoto, T. Wakimoto, K. Imai and Y. Shirota, *Mol. Cryst. Liq. Cryst.*, 1996, **280**, 331; (f) G. D'Aprano, M. Leclerc, G. Zotti and G. Schiavon, *Chem. Mater.*, 1995, **7**, 33.
- (a) M. Wolter, A. Klapars and S. L. Buchwald, *Org. Lett.*, 2001, **3**, 3803; (b) A. Klapars, J. C. Antilla, X. Huang and S. L. Buchwald, *J. Am. Chem. Soc.*, 2001, **123**, 7727; (c) C. Desmarets, R. Schneider and Y. Fort, *Tetrahedron Lett.*, 2001, **42**, 247; (d) B. H. Lipshutz and H. Ueda, *Angew. Chem., Int. Ed.*, 2000, **39**, 4492; (e) R. Shen and J. A. Porco, Jr, *Org. Lett.*, 2000, **2**, 1333; (f) A. V. Kalinin, J. F. Bower, P. Riebel and V. Snieckus, *J. Org. Chem.*, 1999, **64**, 2986; (g) B. H. Yang and S. L. Buchwald, *J. Organomet. Chem.*, 1999, **576**, 125; (h) J. P. Wolfe, S. Wagaw, J.-F. Marcoux and S. L. Buchwald, *Acc. Chem. Res.*, 1998, **31**, 805; (i) J. F. Hartwig, *Angew. Chem., Int. Ed.*, 1998, **37**, 2046; (j) M. S. Driver and J. F. Hartwig, *J. Am. Chem. Soc.*, 1996, **118**, 7217; (k) S. Wagaw and S. L. Buchwald, *J. Org. Chem.*, 1996, **61**, 7240; (l) J. P. Wolfe, S. Wagaw and S. L. Buchwald, *J. Am. Chem. Soc.*, 1996, **118**, 7215.
- (a) P. Sinha, C. C. Kofink and P. Knochel, *Org. Lett.*, 2006, **8**, 3741; (b) P. Sinha and P. Knochel, *Synlett*, 2006, 3304; (c) A. M. Berman and J. S. Johnson, *J. Org. Chem.*, 2006, **71**, 219; (d) A. M. Berman and J. S. Johnson, *J. Org. Chem.*, 2005, **70**, 364; (e) K. Narasaka and M. Kitamura, *Eur. J. Org. Chem.*, 2005, 4505; (f) A. M. Berman and J. S. Johnson, *Synlett*, 2005, 1799; (g) A. M. Berman and J. S. Johnson, *J. Am. Chem. Soc.*, 2004, **126**, 5680; (h) M. Kitamura, T. Suga, S. Chiba and K. Narasaka, *Org. Lett.*, 2004, **6**, 4619.
- (a) F. Kopp, I. Sapountzis and P. Knochel, *Synlett*, 2003, 885; (b) W. L. Waters and P. G. Marsh, *J. Org. Chem.*, 1975, **40**, 3344.
- For the reaction of nitroarenes with Grignard reagents: (a) I. Sapountzis and P. Knochel, *J. Am. Chem. Soc.*, 2002, **124**, 9390; (b) H. Gao, Q. L. Xu, M. Yousufuddin, D. H. Ess and L. Kürti, *Angew. Chem., Int. Ed.*, 2014, **53**, 2701; (c) H. Gao, D. H. Ess, M. Yousufuddin and L. Kürti, *J. Am. Chem. Soc.*, 2013, **135**, 7086; (d) L. Wylie, P. Innocenti, D. K. Wheligan and S. Hoelder, *Org. Biomol. Chem.*, 2012, **10**, 4441; (e) A. Dobbs, *J. Org. Chem.*, 2001, **66**, 638; (f) A. P. Dobbs, M. Voyle and N. Whittall, *Synlett*, 1999, 1594; (g) M. Bosco, R. Dalpozzo, G. Bartoli, G. Palmieri and M. Petrini, *J. Chem. Soc., Perkin. Trans. 2*, 1991, 657; (h) L. Barboni, G. Bartoli, E. Marcantoni, M. Petrini and R. Dalpozzo, *J. Chem. Soc. Perkin. Trans. 1*, 1990, 2133; (i) G. Bartoli, M. Petrini, E. Marcantoni, M. Bosco and R. Dalpozzo, *Tetrahedron Lett.*, 1990, **31**, 6089; (j) Y. Inagaki, R. Okazaki and N. Inamoto, *Bull. Chem. Soc. Jpn.*, 1975, **48**, 3727; (k) Y. Yost, H. R. Gutmann and C. C. Muscoplat, *J. Chem. Soc., C*, 1971, 2119; (l) G. Bartoli, G. Palmieri, M. Bosco and R. Dalpozzo, *Tetrahedron Lett.*, 1989, **30**, 2129; (m) D. Y. Curtin and J. C. Kauer, *J. Am. Chem. Soc.*, 1953, **75**, 6041.
- (a) A. Yanagisawa, S. Takeshita, Y. Izumi and K. Yoshida, *J. Am. Chem. Soc.*, 2010, **132**, 5328; (b) G.-Q. Tian, J. Yang and K. R. Perez, *Org. Lett.*, 2010, **12**, 5072; (c) M. Lu, D. Zhu, Y. Lu, X. Zeng, B. Tan, Z. Xu and G. Zhong, *J. Am. Chem. Soc.*, 2009, **131**, 4562; (d) H. Yamamoto and M. Kawasaki, *Bull. Chem. Soc. Jpn.*, 2007, **80**, 595; (e) H. Yamamoto and N. Momiyama, *Chem. Commun.*, 2005, 3514; (f) N. Momiyama and H. Yamamoto, *J. Am. Chem. Soc.*, 2005, **127**, 1080; (g) J. M. Janey, *Angew. Chem., Int. Ed.*, 2005, **44**, 4292; (h) P. Merino and T. Tejero, *Angew. Chem., Int. Ed.*, 2004, **43**, 2995; (i) A. Bøgevig, H. Sundén and A. Córdova, *Angew. Chem., Int. Ed.*, 2004, **43**, 1109; (j) Y. Hayashi, J. Yamaguchi, T. Sumiya and M. Shoji, *Angew. Chem., Int. Ed.*, 2004, **43**, 1112; (k) S. P. Brown, M. P. Brochu, C. J. Sinz and D. W. C. MacMillan, *J. Am. Chem. Soc.*, 2003, **125**, 10808; (l) G. Zhong, *Angew. Chem., Int. Ed.*, 2003, **42**, 4247; (m) N. Momiyama and H. Yamamoto, *J. Am. Chem. Soc.*, 2003, **125**, 6038; (n) N. Momiyama and H. Yamamoto, *Org. Lett.*, 2002, **4**, 3579; (o) N. Momiyama and H. Yamamoto, *Angew. Chem., Int. Ed.*, 2002, **41**, 2986.
- (a) F. M. Piller, P. Appukkuttan, A. Gavryushin, M. Helm and P. Knochel, *Angew. Chem. Int. Ed.*, 2008, **47**, 6802; (b) F. M. Piller, A. Metzger, M. A. Schade, B. A. Haag, A. Gavryushin and P. Knochel, *Chem. Eur. J.*, 2009, **15**, 7192.
- (a) A. E. Jensen, W. Dohle, I. Sapountzis, D. M. Lindsay, V. A. Vu and P. Knochel, *Synthesis*, 2002, 565; (b) I. Sapountzis, W. Dohle and P. Knochel, *Chem. Commun.*, 2001, 2068; (c) M. Rottländer, L. Boymond, L. Bérillon, A. Leprière, G. Varchi, S. Avolio, H. Laaziri, G. Quéguiner, A. Ricci, G. Cahiez and P. Knochel, *Chem. Eur. J.*, 2000, **6**, 767; (d) M. Abarbri, J. Thibonnet, L. Bérillon, F. Dehmel, M. Rottländer and P. Knochel, *J. Org. Chem.*, 2000, **65**, 4618; (e) L. Boymond, M. Rottländer, G. Cahiez and P. Knochel, *Angew. Chem. Int. Ed.*, 1998, **37**, 1701.
- (a) O. A. Blackburn, B. J. Coe and M. Helliwell, *Organometallics*, 2011, **30**, 4910; (b) B. Prievisch and K. R. Braun, *J. Org. Chem.*, 2005, **70**, 2350; (c) A. Defoin, *Synthesis*, 2004, 706; (d) S. Wang, X. Wang, L. Li and R. C. Advincula, *J. Org. Chem.*, 2004, **69**, 9073; (e) Z. Zhu and J. H. Espenson, *J. Org. Chem.*, 1995, **60**, 1326; (f) S. Sakaue, T. Tsubakino, Y. Nishiyama and Y. Ishii, *J. Org. Chem.*, 1993, **58**, 3633; (g) E. R. Møller and K. A. Jørgensen, *J. Am. Chem. Soc.*, 1993, **115**, 11814; (h) S. T. Tollari, M. Cuscuela and F. Porta, *J. Chem. Soc., Chem. Commun.*, 1993, 1510; (i) W. A. Lees and A. Burawoy, *Tetrahedron*, 1963, **19**, 419.
- (a) N. G. Gaylord, *J. Org. Chem.*, 1960, **25**, 1874; (b) Y. Yu, J. Srogl and L. S. Liebeskind, *Org. Lett.*, 2004, **6**, 2631; (c) L. Ackermann, R. Sandmann and W. Song, *Org. Lett.*, 2011, **13**, 1784; (d) D. Craig, *J. Am. Chem. Soc.*, 1935, **57**, 195; (e) H. Takeuchi, S. Hirayama, M. Mitani and K. Koyama, *J. Chem. Soc., Perkin Trans. 1*, 1988, 521; (f) M. A. Carroll and R. A. Wood, *Tetrahedron*, 2007, **63**, 11349; (g) D. Maiti and S. L. Buchwald, *J. Am. Chem. Soc.*, 2009, **131**, 17423; (h) J. Piccard and *J. Am. Chem. Soc.*, 1926, **48**, 2878; (i) J. Li, M. Cui, A. Yu and Y. Wu, *J. Organomet. Chem.*, 2007, **692**, 3732; (j) R. E. Tundel, K. W. Anderson and S. L. Buchwald, *J. Org. Chem.*, 2006, **71**, 430; (k) M. Kim and S. Chang, *Org. Lett.*, 2010, **12**, 1640.
- D. Zhao, M. Johansson and J.-E. Bäckvall, *Eur. J. Org. Chem.*, 2007, 4431.
- (a) W. Lin, A. Gupta, K. H. Kim, D. Mendel and M. J. Miller, *Org. Lett.*, 2009, **11**, 449; (b) E. C. Taylor, C.-P. Tseng and J. B. Rampal, *J. Org. Chem.*, 1982, **47**, 552.
- C. Sämman, V. Dhayalan, P. R. Schreiner and P. Knochel, *Org. Lett.*, 2014, **16**, 2418.
- (a) T. D. Quach and R. A. Batey, *Org. Lett.*, 2003, **5**, 4397; (b) T. Arnauld, D. H. R. Barton and E. Doris, *Tetrahedron*, 1997, **53**, 4137.

TOC Graphics:

Aryl, heteroaryl or adamantyl zinc reagents add in the presence of magnesium-salts and LiCl to various aryl or heteroaryl nitroso arenes producing after reductive work-up polyfunctional secondary amines.