

Article

3-Acetoxyquinuclidine as Catalyst in EDA Complex-Mediated Reactions triggered by Visible Light

Irene Bosque Martínez, and Thorsten Bach

ACS Catal., Just Accepted Manuscript • DOI: 10.1021/acscatal.9b01039 • Publication Date (Web): 23 Aug 2019

Downloaded from pubs.acs.org on August 23, 2019

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.

3-Acetoxyquinuclidine as Catalyst in EDA Complex-Mediated Reactions triggered by Visible Light

Irene Bosque^{†} and Thorsten Bach*

Department of Chemistry and Catalysis Research Center (CRC) Technische Universität München, 85747 Garching, Germany.

ABSTRACT: 3-Acetoxyquinuclidine was found to act as a catalytic electron donor species in a variety of electron donor-acceptor (EDA) complex-mediated reactions. Only substoichiometric amounts (10 to 25 mol%) were needed to trigger the desired reaction. The outcome could be tuned by selecting the nature of the formed radical to perform amino- and hydro-decarboxylation, dimerization, and cyclization reactions. Importantly, no external additives were needed in this reaction.

KEYWORDS: electron donor-acceptor complex, photochemistry, 3-acetoxyquinuclidine, homogeneous catalysis, tetrachlorophthalimide.


1. INTRODUCTION

Harnessing the power of sunlight in a productive manner has been an aim of many synthetic groups in the last decades. Different strategies such as direct excitation of molecules using UV-light, or activation via energy transfer or via electron transfer from a photocatalyst using visible light are used in order to achieve selective reactivity.¹⁻¹¹ However, UV-light can be harmful, and photocatalysts are non-innocent species that might react with the product, as they are intrinsically active molecules under visible light conditions.

Electron donor-acceptor (EDA) complexes,¹² on the other hand, emerged as an alternative to the use of photocatalysts in visible light-triggered reactions.¹²⁻¹⁶ These complexes are formed by two species, an electron donor and an electron acceptor, which show a charge transfer band only when combined. When this band occurs in the visible region of the spectra, visible light can be used to induce an electron transfer from the donor to the acceptor. Thermal activation of these complexes is also possible.^{17,18} Since only selected pairs of molecules form EDA complexes, there are

only few reports in the literature for selective reactivity, even though these complexes have been known for decades.^{12,19} In the vast majority of the examples, after electron transfer (ET) the electron donor (ED) radical cation and the electron acceptor (EA) radical anion react with each other to form the product (Scheme 1a). Elegant work from the group of Melchiorre exploited the formation of an intramolecular EDA complex via iminium formation (Scheme 1b).²⁰⁻²² However, to the best of our knowledge, there are no reports showing the use of an electron donor or acceptor as external catalytic species for visible light-mediated reactions via EDA complex formation.

Intrigued by this lack of precedent, we sought to explore new avenues of EDA complexes, in particular, the possibility of finding a molecule easy to obtain that could act as a donor of an EDA complex, but which, at the same time, could be regenerated and used in catalytic amounts in different types of reactions (Scheme 1c). In this work, we present the synthetic possibilities of using an ED catalyst in EDA complex-mediated reactions that could greatly expand the use of this strategy in synthetic chemistry.



Scheme 1. Selected literature precedents on EDA complex (top left) and reductive decarboxylative phthalimide esters chemistry (top right); Idea and advantages of this work (bottom); HE: Hantzsch ester; EY: Eosin Y.

Since phthalimide-derived esters have been largely used in borylations (Scheme 1a),^{15,23,24} in photoredox catalysis (Scheme 1d-e),²⁵⁻²⁷ or in coupling chemistry (Scheme 1f)²⁸⁻³⁰ among others, we decided to use them as initial substrates. Most of the approaches for the reductive decarboxylation of phthalimide ester substrates are

net-reductive methods that require the use of stoichiometric amounts of an external reductant (Scheme 1d-f). In this work we propose a redox-neutral process using the phthalimide moieties as acceptor partners of a catalytic donor to induce reactivity via EDA complex formation.

2. RESULTS AND DISCUSSION

2.1. Selection of 3-acetoxyquinuclidine as catalyst


Our investigations started with the synthesis of the corresponding phthalimide ester of 2-phenylbutyric acid, **1a** (Figure 1). However, no EDA complex formation was observed when mixing **1a** with a common donor such as triethylamine (Et_3N). Since the potential of the single electron reduction of phthalimide esters ($E_{1/2} = -1.24$ to -1.38 V vs. SCE²⁷) is lower than the potential of the corresponding tetrachlorophthalimide esters ($E_{1/2} = -0.70$ to -0.54 V vs. SCE, see Supporting Information) we synthesized the analogous tetrachlorophthalimide derivative. An equimolar mixture of the tetrachlorophthalimide ester of 2-phenylbutyric acid, **2a**, with Et_3N in MeCN gave a band with a maximum defined in the blue region of the spectra,

not observed for the separate components. This band was our first indication for the formation of an EDA complex. We then decided to explore amines more stable in their oxidized state than Et₃N, for instance triaryl amines such as dimethylaniline, triphenylamine and 4-methoxytriphenylamine. Unfortunately, none of them gave an observable band in the desired region. Even *N*-methylmorpholine, which has a higher stability due to the orientation of the oxygen lone pairs, showed no EDA complex formation with **2a** (See Scheme S1).


We then turned our attention to a different group of amines. In particular we focused on quinuclidine (azabicyclo[2.2.2]octane) derivatives, as we were attracted by its peculiar tethered structure which imparts unique redox properties to this trialkylamine. As opposed to non-cyclic trialkylamines, such as Et₃N, the bicyclic structure of quinuclidine avoids its degradation after oxidation, since the fixed orientation of the C-H bonds in α -position to the oxidized nitrogen center avoids α -deprotonation (Scheme 2a). For this reason, quinuclidine derivatives have been used by many research groups as reductive quenchers and hydrogen atom donors (HAT) in photo-induced processes (Scheme 2b).³¹ We hypothesized that this molecule could act not only as a

single electron donor but also as an acceptor playing the role of a SET catalyst triggered by an EDA process (Scheme 2c). To the best of our knowledge, this is the first study of this matter in the literature.


a) Difference in stability between triethylamine and quinuclidine



b) Quinuclidine as SET and HAT



c) New hypothesis:
reversible ED catalyst



Scheme 2. Unique structure of quinuclidine. PC: photocatalyst. SET: single electron transfer. HAT: hydrogen atom transfer. R: organic moiety. ED: electron donor. A: molecule acting as electron acceptor. D: electron donor catalyst quinuclidine. S: organic molecule.

2.2. Towards the aminodecarboxylation reaction

With this hypothesis in mind, 3-acetoxyquinuclidine (q-OAc) was chosen as potential EDA catalyst for its simplicity in handling compared to the non-substituted quinuclidine.

An equimolar mixture of **2a** and q-OAc gave a bright yellow solution, indicating the formation of an EDA complex that was observed in the UV-visible spectra recorded for this mixture, not observed for the separate components (Figure 1a). In order to verify the generality of the formation of an EDA complex with q-OAc in the presence of diverse tetrachlorophthalimide acid derivatives, we synthesized the corresponding ester of *N*-Boc-proline **2b** which, in the presence of q-OAc, revealed a yellow solution indicative of the formation of an EDA complex in MeCN (Figure 1b).

Based on these promising observations, we subjected the mixture of **2b** and substoichiometric amounts of q-OAc (25 mol%) in MeCN to blue LEDs irradiation ($\lambda = 455$ nm) at room temperature. A single product and complete conversion were observed after 20 h of irradiation. The characterization of the isolated material revealed the formation of the corresponding decarboxylated product **3b** in a 69% yield (Figure 1c, entry 1). No starting material was recovered and no apparent formation of

any other product was detected. Control reactions in the absence of light or q-OAc resulted in recovery of the starting material (entries 2 and 3), indicating that this is a light-mediated reaction that proceeds via EDA-complex formation. Optimization of the reaction conditions was performed. Change of solvent to 1,4-dioxane gave incomplete conversion (entry 4), but the use of dimethylformamide (DMF) notably improved the yield of this reaction, providing a 91% yield even after only 4 h of irradiation (entries 5 and 6). No conversion was observed in the absence of light but a background reaction in the absence of catalyst gave 39% yield of **3b** (entries 7 and 8) which can be attributed to the contribution of a radical cage mechanism explained by the absorption at 455 nm in DMF observed in the UV-visible spectra of **2b** (see Supporting Information for further details). The use of a higher wavelength, 470 nm, in order to minimize this background reaction gave similar results (entries 9 and 10). Further optimization of the catalyst revealed 10 mol% of q-OAc to be optimal, giving a yield of 89% of **3b** (entries 11 to 13. See Scheme S2).





Figure 1. a) UV-visible spectra of 15 mM solutions in MeCN containing **1a**, **2a** or an equimolar mixture with the corresponding amines, path length 1 cm. b) Initial observations of an EDA complex formation; picture shows a solution of **2b** (left), q-OAc (right) and a 1:1 mixture of both (center) in MeCN. c) ^a Isolated yields. ^b 12% of **2b** was recovered, ^c 6% of **2b** was recovered.

Remarkably, this reaction failed when common photocatalysts as Ru(bpy)₃Cl₂ or Eosin Y were used under the same reaction conditions (Scheme 3). For comparison, substrate **2b'** was selected since the control reaction in the absence of catalyst gave no product formation. The reduction potentials of the excited states of these two photocatalysts suggest that both should be capable of reducing the tetrachlorophthalimide

moiety. On the other hand, a second SET event needs to occur for the turnover of the photocatalyst to its ground state, which might not be the case for these catalysts. In any case, our catalyst **q-OAc** fulfils both requirements and it is capable of reducing the tetrachlorophthalimide moiety via EDA-complex formation and closing the catalytic cycle for a redox-neutral process without the need of any external additives. Furthermore, the formation of this aminodecarboxylated product would not be possible by using other EDA complex-mediated approaches due to the following reasons: (1) the use of stoichiometric amounts of a non-tethered amine, such as Et₃N, gave minimal or no product formation when substrates **2b'** (Scheme 3) or **2b** (see Scheme S6) were used, probably due to the degradation of the oxidized amine, which avoids the second SET from taking place; (2) there was no EDA complex formation observed between tetrachlorophthalimide and *N*-Boc-pyrrolidine; nor (3) between **2b** and **3b** (see Scheme S10); and (4) no concentration dependency of absorption spectrum of **2b** was observed (see Scheme S10) which excludes the possibility of intramolecular SET.




Scheme 3: Comparison of the method described in this work using q-OAc vs the use of other photocatalysts or stoichiometric EDA complex conditions in the synthesis of compound **3b'** from substrate **2b'**. Isolated yields given. Yields of recovered starting material **2b'** are given in parenthesis.


These results are in agreement with the hypothesized behavior of q-OAc in its role as reversible SET catalyst. We envisioned that the formation of the EDA complex **2b/q**-OAc and light irradiation of this complex would result in SET from the catalyst (q-OAc) to the tetrachlorophthalimide moiety. Decarboxylation to form the corresponding α -amino radical intermediate would be followed by its oxidation with q-OAc⁺, regenerating the initial q-OAc in this process, to presumably form an iminium ion that would be trapped by the previously liberated tetrachlorophthalimide anion, delivering

1
2
3 the observed product **3b** (Scheme 4a). The quantum yield of this reaction was found
4
5
6
7 to be 0.02 (See Supporting Information), suggesting that propagative pathways can
8
9
10 be ruled out of the mechanism. We believe that the close proximity of the generated
11
12
13 q-OAc^{•+} to the α -amino radical species after SET promotes a fast back SET and avoids
14
15
16 any HAT pathway from being productive. The formation of this α -amino radical
17
18
19 intermediate was proved when the reaction was run in the presence of 2 equiv. of
20
21
22
23
24 TEMPO as radical trapping agent. Starting material **2b** was recovered in 66% yield
25
26
27 and product **3b** was observed only in trace amounts. The product formed as
28
29
30 consequence of TEMPO trapping the α -amino radical ($m/z = 327$ [M+H⁺]) was
31
32
33
34
35 observed by mass-spectrometry (MS) (see Scheme S5).
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

a) Proposed mechanism for the use of q-OAc as reversible SET catalyst



b) Proposed mechanism for the use of q-OAc as SET/HAT catalyst



Scheme 4. Mechanistic proposal.

A variety of tetrachlorophthalimide esters derived from α -amino acids (**2b** to **2f**) as well as α -hydroxy acid derivatives (**2g** to **2j**) were synthesized and submitted to the optimized catalytic conditions. In order to achieve full conversion of the starting material in all cases, irradiation times were established to be 17 to 20 h. Cyclic and acyclic amines and ethers gave stable amins (**3b** – **3f**) and hemiaminals (**3g** – **3j**, through the corresponding oxonium ion) in yields up to 89% yield using only 10 mol%

of q-OAc. Olefinic double bonds in **3c**, **3e** and **3h** were tolerated as well as the NH group of products **3d** and **3f**. Interestingly, product **3d** bearing a sensitive sulfinyl group was obtained in a 76% yield. Aromatic ethers with substituents in meta- and para-positions **3i** and **3j** were obtained in 78% and 70% respectively, with special interest in **3j** due to the formation of a quaternary hemiaminal center (Scheme 5).

2.3. Cyclization reaction


Intrigued by the high efficiency of these reaction conditions, we wondered whether we could tune the outcome of this reaction by selecting the molecular structure of the starting material. As observed in examples **3c** and **3h**, products derived from the corresponding 6-*exo-trig* or 7-*exo-trig* cyclization were not observed. However, the irradiation of the tetrachlorophthalimide derivative of *O*-allyl-*N*-Boc-serine (**2k**) in the presence of an optimized 25 mol% of q-OAc (see Scheme S3 for optimization conditions) induced the 5-*exo-trig* cyclization to form the tetrahydrofuran derivative **3k** in 59% yield. The use of other photocatalysts as Eosin Y failed to give the product (See Scheme S9). Importantly, control reactions in the absence of light or q-OAc gave no product formation and led to recovery of **2k**, highlighting the essential role of the q-

OAc in the formation of the EDA complex for this reaction to proceed under blue LEDs.

Two-dimensional NMR analysis revealed the formation of a 4:1 diastomeric *cis:trans* mixture (See Supporting Information). According to the proposed mechanism, radical cyclization would be followed by the oxidation of the primary radical to the primary carbocation (from the q-OAc⁺), which is a highly energetically uphill process, and proton elimination to form a double bond. However, this behavior was not observed.

In order to better understand the mechanism we attempted to use DMF-d₇ as solvent which resulted in the formation of **3k** in trace amounts and degradation of the starting material **2k** (See Scheme S7). These results suggest that DMF has a crucial role in this cyclization, probably as a terminal hydrogen atom transfer (HAT) species (Scheme 4b), which is preceded in the literature^{32,33} (See Supporting Information for more details). This is in agreement with the absence of product formation when the reaction was performed in MeCN (see Scheme S3). Comparing bond dissociation energies (BDE) of a primary alkyl radical (BDE [C1-H in propane] = 100 kcal/mol³⁴) and DMF (BDE [H-CH₂N(Me)CHO or Me₂NC(O)-H] = 89 – 90 kcal/mol³²), HAT is feasible. Furthermore, the submission of esters **2l** and **2m** to the reaction conditions resulted in

the isolation of product **3l** in slightly lower yield (49%), and detection of product **3m** only in traces (Scheme 5).



Scheme 5. Products isolated from the photocatalytic reaction. ^a 10 mol% of q-OAc was used. ^b 25 mol% of q-OAc was used. ^c 8% of product **4k** was also isolated. ^d Isolated

as an inseparable 88:12 mixture of **3l** and **4l**. ^e Not isolated. ^f 8% of product **4o** was also isolated. ^g 14% of product **5p** was also isolated. ^h Only the *trans*-configured product was observed. ⁱ 25% of an unidentified product was also isolated. ^j 3% of product **4a** was also isolated. *Indicates the center to which the dr refers to. NTCIPhth: tetrachlorophthalimide moiety. PG: protecting group. For cyclic voltammograms (CVs) and UV-visible spectra of compounds **2** see Supporting Information.

Direct comparison of the BDEs of the bonds involved in the process could explain these results. The BDE of a C-H bond decreases with the higher substitution of the carbon center, meaning that the strength of the C-H bond decreases in the same order: $\text{RH}_2\text{C-H} > \text{R}_2\text{HC-H} > \text{R}_3\text{C-H}$.³⁴ In consequence, for the formation of product **3m** the reaction might not have a thermodynamically favorable HAT from DMF, giving no product formation (See Scheme S7). On the other hand, turnover of the catalyst must occur in a different way than indicated for the formation of amins and hemiaminals **3b** to **3j**. Using the same reasoning, we can presume that HAT is also feasible from DMF to q-OAc⁺ (BDE [q-OAc⁺-H] = 95-100 kcal/mol³⁵) (Scheme 4b). Other

1
2
3 tetrahydrofuran derivatives were synthesized, as **3n** with three consecutive
4
5
6
7 stereocenters and an excellent 9:1 dr isolated in 51% yield, and product **3o** with a
8
9
10 newly formed quaternary carbon atom obtained in 42% yield (Scheme 5).
11
12
13

14
15 Further examples using *N*-alkylated leucine, phenylalanine, or glycine derivatives
16
17
18 with unsaturation at different distances from the formed radical were tested (Scheme
19
20
21
22 5). Cyclization in order to give the corresponding pyrrolidine **3q** was observed from the
23
24
25 *N*-(3-butenyl) derivative of phenylalanine **2q**. In the case of the leucine derivative **2p**,
26
27
28 pyrrolidine **3p** was isolated in 15% yield and with an excellent dr (>9:1) together with
29
30
31 the stable enamide **4p** as major product (35%), presumably from overoxidation of the
32
33
34
35
36 formed radical, and some aminodecarboxylation product (**5p**, 14% yield, see
37
38
39 Supporting Information). Interestingly, the submission of the *N*-(4-butenyl) derivative
40
41
42 of glycine **2r** to the optimized reaction conditions resulted in the formation of products
43
44
45
46 **3r** and **4r** as the only observed products with yields of 41% and 23%, respectively.
47
48
49
50 This result is in contrast with the observed behavior of the formed radical in the case
51
52
53 of product **3h**, where products derived from the cyclization were not observed.
54
55
56
57 Oxidation of the radical formed from **2h** after SET and decarboxylation under these
58
59
60

1
2
3
4 reaction conditions might be consequently faster than cyclization, giving **3h** as single
5
6
7 product instead of products derived from 6-exo-trig or 7-endo-trig cyclization as
8
9
10 observed for **3r** and **4r** respectively. In addition, the analogue ester of **2r** with an oxygen
11
12
13 instead of a nitrogen atom at the α -position of the formed radical gave 54% of the initial
14
15
16 acid (ester **2z**, Supporting Information) with no decarboxylation observed. These
17
18
19 results suggest that both factors, the substituents (N or O) and the substitution of the
20
21
22 esters (primary or secondary), affect to the outcome of the reaction.
23
24
25
26
27

28 29 **2.4. Dimerization and hydrodecarboxylation reactions**

30
31
32

33
34 As expected, when the tetrachlorophthalimide ester of phenylacetic acid **2a** was
35
36
37 exposed to blue LEDs in the presence of 25 mol% of q-OAc in DMF, dimerization of
38
39
40 the benzylic radical was observed to form **3a** in a 67% yield with a 1:1 dr, a behavior
41
42
43 that was also observed when the diphenylacetic acid derivative **2s** was submitted to
44
45
46 the same reaction conditions. In the latter case, not only the dimerized product **3s** was
47
48
49 observed in 24%, but also alkene **4s** was isolated as orange needles in 18% yield
50
51
52
53
54 (Scheme 5).
55
56
57
58
59
60

When purely alkyl radicals were formed in the reaction, hydrodecarboxylation products **3t** to **3w** were isolated in good yields ranging 62% to 75% yield (Scheme 5) under the optimized conditions (See Scheme S4). Replacing q-OAc by Ru(bpy)₃Cl₂ or Eosin Y resulted in no product formation (See Scheme S9), highlighting the importance of q-OAc in this reaction. Interestingly, the q-OAc tolerated the presence of the pendant tetrahydrophthalimide (THPhth) group in product **3w**, which was obtained in 62% yield. A formation of an EDA complex between the q-OAc and this pendant THPhth group, stronger than between the q-OAc and the active tetrachlorophthalimide moiety, could have resulted in trapping and inactivation of the catalyst. A control reaction in the absence of q-OAc gave no product formation. Remarkably, this method proved to be very selective for hydrodecarboxylation of di-substituted acids, as mono- or tri-substituted acids were unreactive (See Scheme S8). Mechanistically, the formation of these products would occur in a similar fashion as for compounds **3k** to **3o**, from decarboxylation after SET in the EDA complex to give the corresponding secondary radical, followed by HAT presumably from DMF to deliver the final product.

3. CONCLUSIONS

In summary, we have shown the potential of q-OAc as electron donor catalyst in EDA complex-mediated reactions. Furthermore, we have demonstrated a new mode of action of q-OAc as reversible SET catalyst. It was found that q-OAc can be regenerated, in contrast to most traditional chemistry where the overall reaction consumes the DA pair. Using the tetrachlorophthalimide moiety to activate a great variety of acids and q-OAc in catalytic amounts (10 to 25 mol%) we have proved the versatility of these unique reaction conditions. By precise selection of the nature of the formed radical we have been able to achieve amino- and hydro-decarboxylation reactions as well as cyclizations and dimerizations in yields up to 89%. Control reactions have shown the necessity of light and catalytic amounts of the ED catalyst q-OAc for the reaction to be effective. Interestingly, no incorporation of the q-OAc catalyst to any product was detected or derivatives of the catalyst in any of the experiments. Further investigations on the stability and versatility of q-OAc as well as other derivatives will be performed.

ASSOCIATED CONTENT

Supporting Information. Experimental procedures, full characterization of all synthesized compounds, cyclic voltammetry experiments of the starting materials and additional experiments explained in Scheme S1 to S13 are included.

AUTHOR INFORMATION

Corresponding Author

*Email: irene.bosque@ua.es. †Present Address: Institute of Electrochemistry and Institute of Organic Synthesis, University of Alicante, Carretera San Vicente del Raspeig s/n, 03690, San Vicente del Raspeig, Alicante, Spain.

Author Contributions

I.B. performed the experiments. The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

ACKNOWLEDGMENT

I.B. gratefully thanks the Alexander von Humboldt Foundation for a postdoctoral fellowship. The Spanish Ministerio de Economía y Competitividad (No. CTQ2017-88171-P) is acknowledged. The authors would like to thank Michael Grübel for assistance in the CV measurements and Jose Carlos Gonzalez-Gomez for helpful discussions.

REFERENCES

- (1) Skubi, K. L.; Blum, T. R.; Yoon, T. P. Dual Catalysis Strategies in Photochemical Synthesis. *Chem. Rev.* **2016**, *116*, 10035–10074.
- (2) Twilton, J.; Le, C. (Chip); Zhang, P.; Shaw, M. H.; Evans, R. W.; MacMillan, D. W. C. The Merger of Transition Metal and Photocatalysis. *Nat. Rev. Chem.* **2017**, *1*, 1–18.
- (3) Zelenka, J.; Svobodová, E.; Tarábek, J.; Hoskovcová, I.; Boguschová, V.; Bailly, S.; Sikorski, M.; Roithová, J.; Cibulka, R. Combining Flavin Photocatalysis and Organocatalysis: Metal-Free Aerobic Oxidation of Unactivated Benzylic Substrates. *Org. Lett.* **2018**, *21*, 114–119.

(4) König, B.; Kümmel, S.; Svobodová, E.; Cibulka, R. Flavin Photocatalysis. *Phys. Sci. Rev.* **2018**, *3*, 20170168.

(5) Silvi, M.; Melchiorre, P. Enhancing the Potential of Enantioselective Organocatalysis with Light. *Nature* **2018**, *554*, 41–49.

(6) Brenninger, C.; Jolliffe, J. D.; Bach, T. Chromophore Activation of α , β -Unsaturated Carbonyl Compounds and Its Application to Enantioselective Photochemical Reactions. *Angew. Chemie Int. Ed.* **2018**, *57*, 14338–14349.

(7) Zou, Y.-Q.; Hörmann, F. M.; Bach, T. Iminium and Enamine Catalysis in Enantioselective Photochemical Reactions. *Chem. Soc. Rev.* **2018**, *47*, 278–290.

(8) Zhou, Q.; Zou, Y.; Lu, L.; Xiao, W. Visible-Light-Induced Organic Photochemical Reactions through Energy-Transfer Pathways. *Angew. Chemie Int. Ed.* **2018**, *58*, 1586–1604.

(9) McAtee, R. C.; McClain, E. J.; Stephenson, C. R. J. Illuminating Photoredox Catalysis. *Trends Chem.* **2019**, *1*, 111–125.

(10) Petzold, D.; Nitschke, P.; Brandl, F.; Scheidler, V.; Dick, B.; Gschwind, R. M.; König, B. Visible-Light-Mediated Liberation and In Situ Conversion of Fluorophosgene. *Chem. Eur. J.* **2019**, *25*, 361–366.

(11) Matsui, J. K.; Lang, S. B.; Heitz, D. R.; Molander, G. A. Photoredox-Mediated Routes to Radicals: The Value of Catalytic Radical Generation in Synthetic Methods Development. *ACS Catal.* **2017**, *7*, 2563–2575.

(12) Foster, R. Electron Donor-Acceptor Complexes. *J. Phys. Chem.* **1980**, *84*, 2135–2141.

(13) Lima, C. G. S.; de M. Lima, T.; Duarte, M.; Jurberg, I. D.; Paixao, M. W. Organic Synthesis Enabled by Light-Irradiation of EDA Complexes: Theoretical Background and Synthetic Applications. *ACS Catal.* **2016**, *6*, 1389–1407.

(14) Zhang, J.; Li, Y.; Xu, R.; Chen, Y. Donor–Acceptor Complex Enables Alkoxy Radical Generation for Metal-Free C(Sp³)–C(Sp³) Cleavage and Allylation/Alkenylation. *Angew. Chemie Int. Ed.* **2017**, *56*, 12619–12623.

(15) Wu, J.; He, L.; Noble, A.; Aggarwal, V. K. Photoinduced Deaminative Borylation of Alkylamines. *J. Am. Chem. Soc.* **2018**, *140*, 10700–10704.

(16) Jiang, H.; He, Y.; Cheng, Y.; Yu, S. Radical Alkynyltrifluoromethylation of Alkenes Initiated by an Electron Donor–Acceptor Complex. *Org. Lett.* **2017**, *19*, 1240–1243.

(17) Shirke, R. P.; Ramasastry, S. S. V. Organocatalytic β -Azidation of Enones Initiated by an Electron-Donor–Acceptor Complex. *Org. Lett.* **2017**, *19*, 5482–5485.

(18) Zhou, S.; Song, T.; Chen, H.; Liu, Z.; Shen, H.; Li, C. Catalytic Radical Trifluoromethylalkynylation of Unactivated Alkenes. *Org. Lett.* **2017**, *19*, 698–701.

(19) Fukuzumi, S.; Mochida, K.; Kochi, J. K. A Unified Mechanism for Thermal and Photochemical Activation of Charge-Transfer Processes with Organometals. Steric Effects in the Insertion of Tetracyanoethylene. *J. Am. Chem. Soc.* **1979**, *101*, 5961–5972.

(20) Arceo, E.; Jurberg, I. D.; Álvarez-Fernández, A.; Melchiorre, P. Photochemical Activity of a Key Donor–Acceptor Complex Can Drive Stereoselective Catalytic α -Alkylation of Aldehydes. *Nat. Chem.* **2013**, *5*, 750.

(21) Cao, Z.-Y.; Ghosh, T.; Melchiorre, P. Enantioselective Radical Conjugate Additions Driven by a Photoactive Intramolecular Iminium-Ion-Based EDA Complex. *Nat. Commun.* **2018**, *9*, 3274–3283.

(22) Bahamonde, A.; Melchiorre, P. Mechanism of the Stereoselective α -Alkylation of Aldehydes Driven by the Photochemical Activity of Enamines. *J. Am. Chem. Soc.* **2016**, *138*, 8019–8030.

(23) Candish, L.; Teders, M.; Glorius, F. Transition-Metal-Free, Visible-Light-Enabled Decarboxylative Borylation of Aryl N-Hydroxyphthalimide Esters. *J. Am. Chem. Soc.* **2017**, *139*, 7440–7443.

(24) Fawcett, A.; Pradeilles, J.; Wang, Y.; Mutsuga, T.; Myers, E. L.; Aggarwal, V. K. Photoinduced Decarboxylative Borylation of Carboxylic Acids. *Science* **2017**, *357*, 283–286.

(25) Schnermann, M. J.; Overman, L. E. A Concise Synthesis of (–)-Aplyviolene Facilitated by a Strategic Tertiary Radical Conjugate Addition. *Angew. Chemie Int. Ed.* **2012**, *51*, 9576–9580.

(26) Schwarz, J.; König, B. Metal-Free, Visible-Light-Mediated, Decarboxylative Alkylation of Biomass-Derived Compounds. *Green Chem.* **2016**, *18*, 4743–4749.

(27) Okada, K.; Okamoto, K.; Morita, N.; Okubo, K.; Oda, M. Photosensitized Decarboxylative Michael Addition through *N*-(Acyloxy) Phthalimides via an Electron-Transfer Mechanism. *J. Am. Chem. Soc.* **1991**, *113*, 9401–9402.

(28) Huihui, K. M. M.; Caputo, J. A.; Melchor, Z.; Olivares, A. M.; Spiewak, A. M.; Johnson, K. A.; DiBenedetto, T. A.; Kim, S.; Ackerman, L. K. G.; Weix, D. J.

Decarboxylative Cross-Electrophile Coupling of *N*-Hydroxyphthalimide Esters with Aryl Iodides. *J. Am. Chem. Soc.* **2016**, *138*, 5016–5019.

(29) Toriyama, F.; Cornella, J.; Wimmer, L.; Chen, T.-G.; Dixon, D. D.; Creech, G.; Baran, P. S. Redox-Active Esters in Fe-Catalyzed C–C Coupling. *J. Am. Chem. Soc.* **2016**, *138*, 11132–11135.

(30) Qin, T.; Cornella, J.; Li, C.; Malins, L. R.; Edwards, J. T.; Kawamura, S.; Maxwell, B. D.; Eastgate, M. D.; Baran, P. S. A General Alkyl-Alkyl Cross-Coupling Enabled by Redox-Active Esters and Alkylzinc Reagents. *Science* **2016**, *352*, 801–805.

(31) Shaw, M. H.; Shurtleff, V. W.; Terrett, J. A.; Cuthbertson, J. D.; MacMillan, D. W. C. Native Functionality in Triple Catalytic Cross-Coupling: Sp³ C–H Bonds as Latent Nucleophiles. *Science* **2016**, *352*, 1304–1308.

(32) Markgraf, J. H.; Chang, R.; Cort, J. R.; Durant Jr, J. L.; Finkelstein, M.; Gross, A. W.; Lavyne, M. H.; Moore, C. M.; Petersen, R. C.; Ross, S. D. Protodediazoni-ation of Aryldiazonium Fluoroborates by Dimethylformamide. *Tetrahedron* **1997**, *53*, 10009–10018.

(33) Tam, C. M.; To, C. T.; Chan, K. S. Carbon–Carbon σ -Bond Transfer Hydrogenation with DMF Catalyzed by Cobalt Porphyrins. *Organometallics* **2016**, *35*, 2174–2177.

(34) Gowenlock, B. G.; Jones, P. P.; Majer, J. R. Bond Dissociation Energies in Some Molecules Containing Alkyl Substituted CH_3 , NH_2 and OH . *Trans. Faraday Soc.* **1961**, *57*, 23–27.

(35) Liu, W.-Z.; Bordwell, F. G. Gas-Phase and Solution-Phase Homolytic Bond Dissociation Energies of $\text{H}-\text{N}^+$ Bonds in the Conjugate Acids of Nitrogen Bases. *J. Org. Chem.* **1996**, *61*, 4778–4783.

FOR TABLE OF CONTENTS ONLY

