

PII: S0040-4039(96)00992-6

Hydroxylation of Polycyclic Alkanes with Molecular Oxygen Catalyzed by N-Hydroxyphthalimide (NHPI) Combined with Transition Metal Salts

Yasutaka Ishii*, Susumu Kato, Takahiro Iwahama, and Satoshi Sakaguchi

Department of Applied Chemistry, Faculty of Engineering,

Kansai University, Suita, Osaka 564, Japan

Abstract: Adamantanes were successfully converted into the corresponding mono- and dihydroxy adamantanes with molecular oxygen in the presence of *N*-hydroxyphthalimide (NHP1) combined with cobalt salts under mild conditions. For example, exposure of adamantane under oxygen atmosphere in the presence of NHPI (10 mol%) and Co(acac)₂ (0.5 mol%) in acetic acid at 75 °C for 6 h afforded adamantan-1-ol (43 %) and adamantane-1,3-diol (40 %) along with adamantan-2-one (8 %) in 93 % conversion. Similarly, 1,3-dimethyladamantane produced 3,5-dimethyladamantane-1,3-diol (37 %). Copyright © 1996 Elsevier Science Ltd

The selective oxygenation of saturated hydrocarbons under mild conditions is one of the most challenging and promising targets in oxidation chemistry.¹ In particular, homogeneous catalytic oxidation of hydrocarbons with molecular oxygen is very important in the chemical industry.² Recently, we have developed a novel catalytic system consisting of *N*-hydroxyphthalimide (NHPI) and Co(acac)_n (n=2 or 3) that oxidizes alkanes, alkylbenzenes and alcohols with molecular oxygen without any reducing agents.^{3,4} There have been a few reports on the catalytic hydroxylation of adamantane (1a) using O₂ as the ultimate oxidant.⁵ In general, the direct hydroxylation of 1a with O₂ in the absence of any reducing agents such as aldehyde is very difficult to carry out in higher conversion and selectivity.⁶ We have achieved the first successfully catalytic hydroxylation of adamantanes, 1a and 1b, with O₂ in higher conversions and selectivities. To our knowledge, this is the best result obtained by the direct hydroxylation of adamantanes using O₂ through the catalytic process. In this paper, we describe the oxidation of polycyclic alkanes such as adamantanes with molecular oxygen catalyzed by NHPI combined with several metal salts.

The results for oxidation of **1a** under various reaction conditions are shown in Table 1. The oxidation of **1a** in the presence of a catalytic amount of NHPI (10 mol%) and Co(acac)₂(0.5 mol%) in acetic acid under oxygen



Run	Metal Salt	Time / h	(<i>c</i>	Yield / % ^{b)}		
			Conv. 7 %	2a	3a	4a
1	Co(acac) ₂	6	93	43	8	40
2	_	6	17	52	-	
3 c)	$Co(acac)_2$	6	no reaction			
4	$Co(acac)_3$	6	91	47	8	37
5	Co(OAc) ₂	6	>99	21	6	54
6	Mn(acac) ₃	6	68	56	6	31
7	VO(acac) ₂	6	95	32	7	54
8	$MoO_2(acac)_2$	6	85	54	6	31
9	Fe(acac) ₃	6	19	63	_	_
10	$Cu(OAc)_2$	6	36	64	6	8
11 ^{d)}	Co(acac) ₂	10	73	56	14	25
12	Co(acac) ₂	2	50	80	6	12
13	Co(acac) ₂	15	>99	10	5	53

 Table 1 Aerobic Oxidation of Adamantane (1a) Catalyzed by

 NHPI in the Presence of Transition Metal Salts ^{a)}

a) 1a (3 mmol) was allowed to react with dioxygen (1 atm) in the presence of NHPI (10 mol%) and transition metal salts (0.5 mol%) in acetic acid (8 mL) at 75 °C. b) Based on 1a reacted. c) In the absence of NHPI. d) Under air atmosphere.

atmosphere (1 atm) at 75 °C for 6 h (standard conditions) produced adamantan-1-ol (**2a**) (43 %), adamantane-1,3-diol (**4a**) (40 %) along with a small amount of adamantan-2-one (**3a**) (8%) (Run 1). It is worth noting that the present oxidation produces diol **4a** in relatively high yield, since **4a** is rarely produced by conventional aerobic oxidations.^{5,6} In the oxidation of **1a** with 2,6-dichloropyridine *N*-oxide catalyzed by a Ru complex, **4a** is reported to be formed in 25 % yield.⁷ Leaving out Co(acac)₂ from the catalytic system led to a significant decrease in the conversion of **1a** (Run 2). Needless to say, the oxidation did not take place in the absence of NHPI (Run 3). Oxidations of **1a** using NHPI combined with Co(acac)₃ or Co(OAc)₂ gave almost the same results as those with Co(acac)₂. Replacing Co(acac)₂ with Mn(acac)₃ resulted in a slight decrease in the conversion of **1a** to give **2a** and **4a** in good yields. Fe(acac)₃ and Cu(acac)₂ were inefficient in this oxidation. The oxidation of **1a** in air under these conditions afforded **2a** (56 %), **3a** (14 %) and **4a** (25 %) in 73 % conversion. When the reaction was prolonged for 15 h, diol **4a** was obtained in 53 % yield.

Since adamantane-1,3-diol (4a) is an interesting material as a component of diols in polymer chemistry, stepwise hydroxylation of 2a was examined. The hydroxylation of 2a under the standard conditions afforded 4a in 76 % yield along with a small amount of adamantane-1,3,5-triol (5) (18 %). 4a was prepared from 2a with ozone on silica gel in 43 %.⁸



Table 2 shows the results for the oxidation of various polycyclic alkanes by the NHPI / Co(acac), system. 1,3-Dimethyladamantane (1b) was also oxidized with dioxygen by the NHPI / Co(acac), system. As expected, 1b was more easily oxidized than 1a to form 3,5-dimethyladamantan-1-ol (2b) (47 %) and 5,7dimethyladamantane-1,3-diol (4b) (37 %). Methyl substituents of 1b did not oxidize under these reaction conditions. It is surprising that the hydroxylation of 1b is promoted even at 40 °C to form 2b in high selectivity (87 %), although the conversion was moderate (46 %).

cis-Decalin (6) was hydroxylated at 60 °C to form bicyclo[4.4.0]decan-1-ol (7) in which the tertiary C-H bond was selectively oxidized. However, the oxidation of trans-decalin (8) under the same conditions was difficult and most of 8 was recovered unchanged. The reaction of 8 at higher temperature (75 °C) yielded 7 in low selectivity. Unlike 1a, tetrahydrodicyclopentadiene (THDCP) (9) was oxidized to diketone (10)⁹ rather than hydroxyketone. Diketone 10 was formed in excellent selectivity when VO(acac), was used in place of Co(acac),. The oxidation of tetracyclo[6.2.1.0^{2,7}.1^{3,6}]dodecane (TCDD)(11) led to a complex mixture of degradation products other than diketone 12. 10

Run	Substrate	Temp. / °C	Time / h	Conv. / %	Product (Yield / %) b)
1	1b	75	6	>99	2b (47), 4b (37)
2	1b	75	15	>99	2b (23), 4b (64)
3	1b	40	24	46	2b (87), 4b (6)
4 ^{c)}	cis-decalin (6)	60	15	97	bicyclo[4.4.0]decan-1-ol (7) (56) d)
5 c)	trans-decalin (8) 75	6	49	7 (32) ^{e)}
6 ^{c)}	endo-THDCP (9) 65	15	80	bicyclo[5.2.1]decane-2,6-dione (10) (54)
7 c), i	n 9	65	6	39	10 (98)
8	TCDD (11)	85	15	69	tricyclo[6.2.1.1 ^{3,6}]dodecane-2,7-dione (12) (13)

Table 2 Aerobic Oxidation of Various Polycyclic Alkanes by NHPI / Co(acac)₂ System ^{a)}

a) Substrate was allowed to react with dioxygen (1 atm) in the presence of NHPI (10 mol%) and Co(acac)₂ (0.5 mol%) in acetic acid. b) Based on substrates reacted. c) Benzonitrile was used as a solvent. d) The ratio of *cis*-7 and *trans*-7 was 1 : 3.6. e) The ratio of *cis*-7 and *trans*-7 was 1 : 3.5. f) VO(acac)₂ was used in place of Co(acac)₂.



In conclusion, polycyclic alkanes were efficiently oxygenated with molecular oxygen using a combined catalytic system of NHPI and $Co(acac)_2$ under mild conditions. This method provides a novel, direct aerobic hydroxylation route of polycyclic alkanes.

Acknowledgment

This work was partly supported by a Grant-in-Aid (08455420) from the Ministry of Education, Science, and Culture of Japan.

REFERENCES AND NOTES

- (a) Sheldon, R. A.; Kochi, J. K, Metal-catalyzed Oxidation of Organic Compounds; Academic Press: New York, 1981. (b) Hill, C. L. Activation and Functionalization of Alkanes; Academic Press: New York, 1989. (c) Simándi, L. Catalytic Activation of Dioxygen by Metal Compounds; Kluwer Academic Publisher, 1992. (d) The Activation of Dioxygen and Homogeneous Catalytic Oxidation; Barton, D. H. R.; Martell, A. E.; Sawyer, D. T. Eds.; Plenum Press: New York, 1993. (e) Advances in Oxygenated processes; Baumstark, A. L.; Eds. vol 1, JAI Press, London 1988. (f) Reiser, O. Angew. Chem. Int. Ed. Engl. 1994, 33, 69-72. (g) Maunier, B. Chem. Rev. 1992, 92, 1411-1456 and references cited therein. (h) Jaynes, B. S.; Hill, C. L. J. Am. Chem. Soc. 1995, 117, 4704-4705. (i) Curci, R.; Dinoi, A.; Rubino, M. F. Pure Appl. Chem. 1995, 67, 811-822.
- 2. Parshall, G. W.; Ittel, S. D. *Homogeneous Catalysis*; 2nd Ed. John Wiley & Sons, Inc: New York 1992. *Ctalytic Selecte Oxidation*, Oyama, S. T.; Hightower, J. W.; ACS Symposium Series 523, ACS, Washington, DC, 1993.
- 3. Ishii, Y.; Nakayama, K.; Takeno, M.; Sakaguchi, S.; Iwahama, T.; Nishiyama, Y. J. Org. Chem. 1995, 60, 3934-3935.
- 4. Ishii, Y.; Iwahama, T.; Sakaguchi, S.; Nakayama, K.; Nishiyama, Y. J. Org. Chem., in printed.
- (a) Barton, D. H. R.; Boivin, J.; Gastiger, M.; Morzycki, J.; Hay-Motherwell, R. S.; Motherwell, W. B.; Ozbalik, N.; Schwartzentruber, K. M. J. Chem. Soc., Perkin Trans. 1, 1986, 947-955. (b) Battioni, P.; Bartori, J. F.; Leduc, P.; Fortecave, M.; Mansuy, D. J. Chem. Soc., Chem. Commun. 1987, 791-792. (c) Murahasi, S.-I.; Oda, Y.; Naota, T.; Komiya, N. J. Chem. Soc., Chem. Commun. 1993, 139-140. (d) Murahasi, S.-I.; Oda, Y.; Naota, T. J. Am. Chem. Soc. 1992, 114, 7913-7914.
- (a) Mizuno, N.; Tateishi, M.; Hirose, T.; Iwamoto, M. Chem. Lett. 1993, 2137-2140. (b) Lyons, J. E.; Ellis, P. E. Jr.; Myers, H. K. Jr. J. Catal. 1995, 155, 59-73.
- 7. Ohtake, H.; Higuchi, T.; Hirobe, M. J. Am. Chem. Soc. 1992, 114, 10660-10662.
- 8. Cohen, Z.; Keinan, E.; Mazur, Y.; Varkony, T. H. J. Org. Chem. 1975, 40, 2141-2142.
- 9. Bicyclo[5.2.1]decane-2,6-dione (10): IR 2948, 1698, 1141, 896 cm⁻¹; ¹³C NMR δ 23.6, 25.2, 30.8, 39.0, 51.5, 213.3; ¹H NMR δ 1.01-1.81 (m, 6H), 1.95-2.02 (m, 2H), 2.24-2.31 (m, 4H), 2.45-2.63 (m, 2H); MS *m/e* 166 (M⁺, 17), 138 (13), 125 (13), 97 (100).
- 10. Tetracyclo[6.2.1.1^{3.6}]dodecane-2,7-dione (12): IR 2938, 1671, 1280, 859 cm⁻¹; ¹³C NMR δ 29.8, 30.4, 52.9, 216.1; ¹H NMR δ 1.93-2.42 (m, 13H), 3.19-2.26 (m, 3H); MS m/e 192 (M⁺, 0.2) 164 (6), 151 (100), 123 (34), 95 (9).

(Received in Japan 15 April 1996; revised 20 May 1996; accepted 24 May 1996)