

Synthesis, Spectroscopic Characterization and Thermal Studies of (*E,E*)-2-Hydroxy-3-methoxybenzaldehyde Azine

RONG ZHOU^{1,2}, HUAIBO ZOU¹ and GUANGQUAN MEI^{1,*}

¹Key Laboratory of Jiangxi University for Applied Chemistry & Chemical Biology, Yichun University, Jiangxi Province, P.R. China

²College of Chemistry & Bioengineering, Yichun University, Jiangxi Province, P.R. China

*Corresponding author: Tel/Fax: +86 795 3200655; E-mail: yc_mgq@163.com

Received: 18 June 2013;

Accepted: 28 August 2013;

Published online: 15 February 2014;

AJC-14706


Hydrazone, (*E,E*)-2-hydroxy-3-methoxybenzaldehyde azine was synthesized and its spectroscopic characterization was studied via IR, ¹H NMR, MS and XRD. Its fluorescence activity and thermal stability were also analyzed.

Keywords: Hydrazone, Spectroscopic characterization, XRD, Thermal analysis.

INTRODUCTION

Hydrazone contains the azomethine group (–RC=N–) and usually formed by the condensation of a primary amine with an active carbonyl compound¹. Hydrazones are widely used as intermediates in synthesis², and in particular in hydrazone Schiff base ligands, which are among others employed in dinuclear catalysts³. In analytical chemistry, the formation of hydrazones was extensively used as ideal reagents in detection, determination and isolation of compounds containing the carbonyl group or transition metals determinations^{4,5}.

Furthermore, hydrazones demonstrate many bioactivities in the treatment of several diseases, therefore, a number of hydrazone-hydrazone derivatives have been claimed to possess interesting antibacterial⁶⁻⁷, antifungal⁸⁻⁹, anticonvulsant¹⁰, anti-inflammatory¹¹, antimalarial¹², and antituberculosis activities¹³. Hydrazones have also been used for different purposes such as herbicides, insecticides¹⁴⁻¹⁵ and plant growth regulators¹⁶. This paper contains the synthesis, characterizations and thermal stability of the hydrazone, (*E,E*)-2-hydroxy-3-methoxybenzaldehyde azine (**Scheme-I**).



Scheme-I: (*E,E*)-2-Hydroxy-3-methoxybenzaldehyde azine

EXPERIMENTAL

2-Hydroxy-3-methoxybenzaldehyde (Fluka) and hydrazine hydrochloride (Fluka) were used as received. Analytical grade ethanol was used as solvent.

Synthesis of (*E,E*)-2-hydroxy-3-methoxybenzaldehyde azine: The (*E,E*)-2-hydroxy-3-methoxybenzaldehyde azine was prepared by the condensation reaction of 2-hydroxy-3-methoxybenzaldehyde (1.52 g, 1 mmol) and hydrazine hydrochloride (0.34 g, 1 mmol) in ethanol (30 mL). The reaction mixture was refluxed and stirred at 70 °C for 24 h. The resulting clear solution was kept in air and, after several minutes, yellow crystals were formed at the bottom of the vessel. The crystals were isolated, washed three times with ethanol and dried in an infrared-ray vacuum oven at 55 °C (yield 72 %).

The IR spectra (4000-400 cm⁻¹) were recorded as KBr pellets on a TG-209/Vector-22 spectrometer. ¹H NMR spectra was elucidated using a Bruker Avance AV 400 MHz ¹H NMR spectrometer. The solvent used was DMSO-*d*₆ and the chemical shifts were given in ppm in using TMS as the internal standard. The ¹H NMR spectra were taken at 25 °C. Mass spectra was obtained on a Agilent GC-MS 5975C instrument at 70 eV. XRD was obtained on a SMART APEXII Single-Crystal Diffractometer. The fluorescence spectra was obtained on a Varian LS55 fluorescence spectrometer. The solvent used was DMSO. Thermal analysis and differential thermogravimetric (TGA-DTG) analyses were carried out in temperature range from 25 to 600 °C in nitrogen atmosphere by TGA-Q50 thermal analyzer. The experimental conditions were: aluminium crucible with 1 mg of sample, nitrogen atmosphere (nitrogen flow

50 mL/min) and heating rate of 20 °C/min. Differential scanning calorimetric (DSC) analyses were carried out in temperature range from 25 to 500 °C in nitrogen atmosphere by DSC-Q20 analyzer. The experimental conditions were: aluminium crucible with 2 mg of sample, nitrogen atmosphere (nitrogen flow 50 mL/min) and heating rate of 10 °C/min.

RESULTS AND DISCUSSION

The IR spectrum (Fig. 1) shows a broad band at 3470 cm^{-1} , which may be assigned to the -OH. The bands appeared at around 3000 cm^{-1} are assigned to the phenyl group and the strong intensity peak at 1615 cm^{-1} belongs to $\nu(\text{C}=\text{N})$, suggesting that the condensation was carried out and produced the hydrazone. The peak at 1257 cm^{-1} is the usual mode of C-O bending vibration. Bands appearing at 1577, 1460, 777, 731 cm^{-1} are the usual modes of phenyl ring vibrations.




Fig. 1. Infrared spectrum of the (*E,E*)-2-hydroxy-3-methoxybenzaldehyde azine

The ^1H NMR spectrum of the hydrazone (Fig. 2) show a singlet band at 3.83 ppm that was assigned to the proton of $-\text{OCH}_3$ group. The multiplet signals at 6.87-7.29 ppm were attributed to the protons of aromatic rings. The signal of the azomethine group was observed at 8.97 ppm, which could further prove the formation of the hydrazone and the signal of the -OH was found at 10.85 ppm.




Fig. 2. ^1H NMR spectral of the (*E,E*)-2-hydroxy-3-methoxybenzaldehyde azine

Mass Spectroscopy: The most significant m/z peaks in Fig. 3 of mass spectrum of the hydrazone are given in Table-1.

TABLE-1
MASS SPECTRUM (m/z) OF THE (*E,E*)-2-HYDROXY
-3-METHOXYBENZALDEHYDE AZINE



Origin	m/z
$\text{C}_{16}\text{H}_{16}\text{N}_2\text{O}_4$	300
$[\text{C}_{16}\text{H}_{15}\text{N}_2\text{O}_3]^+$	283
$[\text{C}_8\text{H}_8\text{NO}_2]^+$	150
$[\text{C}_7\text{H}_5\text{NO}_2]^+$	135
$[\text{C}_6\text{H}_4\text{O}_2]^+$	108
$[\text{C}_5\text{H}_3\text{O}]^+$	80
$[\text{C}_3\text{O}]^+$	52

The (*E,E*)-2-hydroxy-3-methoxybenzaldehyde azine show one peak at m/z 300 assigned to the $\text{C}_{16}\text{H}_{16}\text{N}_2\text{O}_4$ molecular ion. It also shows a series of peaks *i.e.* m/z (%): 300 (81.55), 283 (26.46), 150 (50.34), 135 (65.40), 80 (51.35) and 52 (100) corresponding to various fragments. Their signals give an idea about the formation of the hydrazone (**Scheme-II**).

X-ray structure: The X-ray structure of the (*E,E*)-2-hydroxy-3-methoxybenzaldehyde azine is shown in Fig. 4 and selected metric parameters are listed in Table-2. The whole molecule is to be almost planar. There are two half-molecules in the asymmetric unit, the mid-points of the N-N bonds lie on centers of symmetry. The molecular structure is stabilized by an intermolecular O-H...N hydrogen bond which forms a six-membered ring.

Other X-ray crystallographic data are shown below: Monoclinic, space group $\text{P2}(1)/n$, $a = 5.9986(6)$, $b = 18.3183(19)$, $c = 6.8263(7)$ Å; $\alpha = 90$, $\beta = 107.093(3)$, $\gamma = 90^\circ$; $V = 716.97(13)$ Å³; $Z = 4$; $D_{\text{calc.}} = 1.391$ mg m⁻³, $F(000) = 316$; $\mu(\text{Mo-K}\alpha) = 0.71073$; $T = 293(2)$ K.

Fluorescence spectroscopy: Fig. 5 shows the fluorescence spectra in the range 200-500 nm of the hydrazone in DMSO (1×10^{-4} mol L⁻¹). The emission spectrum indicates a peak at 275 nm due to $\pi-\pi^*$ transition, with the red shift deriving from the resonance donating effect of both -OH and -OCH₃ groups in the phenyl ring and another broad peak at 425 nm is due to $n-\pi^*$ transition.

Fig. 3. Mass spectrum of the (*E,E*)-2-hydroxy-3-methoxybenzaldehyde azineScheme-II: Fragment patterns of the (*E,E*)-2-hydroxy-3-methoxy-benzaldehyde azine




Fig. 4. Molecular structure of the (*E, E*)-2-hydroxy-3-methoxybenzaldehyde azine

TABLE-2
SELECTED BOND LENGTHS AND BOND ANGLES FOR THE COMPOUND

Bond	Bond length (Å)	Bond	Bond angle (°)
N1-N1A	1.402(3)	C7-N1-N1A	112.97(18)
N1-C7	1.281(2)	N1-C7-C6	122.22(16)
C7-C6	1.448(2)	C1-C6-C7	119.12(16)
C(1)-C(2)	1.370(3)	C5-C6-C7	121.51(16)
C(1)-C(6)	1.400(2)	C1-C6-C5	119.37(16)
C(2)-C(3)	1.393(2)	C6-C5-C4	119.41(16)
C(3)-C(4)	1.377(2)	C5-C4-C3	120.02(16)
C(4)-C(5)	1.402(2)	C4-C3-C2	120.45(17)
C(5)-C(6)	1.402(2)	C3-C2-C1	120.12(17)




Fig. 5. Emission spectrum of the (*E, E*)-2-hydroxy-3-methoxybenzaldehyde azine

TG/DTA-DSC analysis: The TG thermogram of the hydrazone (Fig. 6) shows only one stage of decomposition from 198 to 315 °C with about weight loss 100 % and the fastest rate point of the mass loss is at 308 °C in the DTG (Fig. 6).




Fig. 6. TG-DTG of the (*E, E*)-2-hydroxy-3-methoxybenzaldehyde azine

The DSC curve (Fig. 7) indicates that the melt point of the hydrazone is 185-187 °C and the maximum peak at 205 °C in DSC is observed by endothermic reaction.




Fig. 7. DSC of the (*E, E*)-2-hydroxy-3-methoxybenzaldehyde azine

Conclusion

In summary, the (*E, E*)-2-hydroxy-3-methoxybenzaldehyde azine was synthesized by the condensation reaction in a relatively good yield. Its spectroscopic characterizations were analyzed in terms of IR, ¹H NMR, MS, XRD, fluorescence spectrometry and the thermal stability was also studied.

ACKNOWLEDGEMENTS

The authors gratefully acknowledged the National Natural Science Foundation of China (21261024), Jiangxi Department of Education (GJJ11704, GJJ12603), the Natural Science Foundation of Jiangxi Province (20132BAB203002) and Key Laboratory of Jiangxi Province for Research on Active Ingredients in Natural Medicines (China) for financial support.

REFERENCES

1. M.S. Refat and A. A. Ibrahim, *Spectrochim. Acta A*, **70**, 234 (2008).
2. D.F. Taber and P.F. Guo, *J. Org. Chem.*, **73**, 9479 (2008).
3. J. Chakraborty, R.K.B. Singh, B. Samanta, C.R. Choudhury, S.K. Dey, P. Talukder, M.J. Borah and S. Mitra, *Z. Naturforsch.*, **61b**, 1209 (2006).
4. A. Irshad, K. Misbah-ul-Ain and A. Makshoof, *J. Pure Appl. Sci.*, **17**, 67 (1998).
5. S.P. Douglas and P.N. Rao, *Int. J. Eng. Sci. Technol.*, **2**, 4655 (2010).
6. N.H. Al-Sha'alan, *Molecules*, **12**, 1080 (2007).
7. S.K. Sridhar, M. Saravanan and A. Ramesh, *Eur. J. Med. Chem.*, **36**, 615 (2001).
8. V. Alptüzün, S. Parlar, H. Tasli and E. Erciyas, *Molecules*, **14**, 5203 (2009).
9. C. Mirela, I. Emilia and G. Rodica, *Rev. Roum. Chim.*, **10**, 911 (2008).
10. S.N. Pandeya, A.S. Raja and J.P. Stables, *J. Pharm. Pharmaceut. Sci.*, **3**, 266 (2002).
11. C.M. Moldovan, O. Oniga, A. Părvu, B. Tiperciuc, P. Verite, A. Pîrnu, O. Crisan, M. Bojita and R. Pop, *Eur. J. Med. Chem.*, **4**, 526 (2011).
12. S. Gemma, G. Kukreja, C. Fattorusso, M. Persico, M.P. Romano, M. Altarelli, L. Savini, G. Campiani, E. Fattorusso, N. Basilico, D. Taramelli, V. Yardley and S. Butini, *Bioorg. Med. Chem. Lett.*, **16**, 5384 (2006).
13. A. Nayyar, A. Malde, E. Coutinho and R. Jain, *Bioorg. Med. Chem.*, **14**, 7302 (2006).
14. M.K. Mao and J.E. Franz, *Synthesis*, 920 (1991).
15. M. Böger, D. Dürr, L. Gsell, R. G Hall, F. Karrer, O. Kristiansen, P. Maienfisch, A. Pascual and A. Rindlisbacher, *Pest Manage. Sci.*, **57**, 191 (2001).
16. J.B. Raoof, R. Ojani and Z. Mohammadpour, *Int. J. Electrochem. Sci.*, **5**, 177 (2010).