## JOURNAL OF THE CHEMICAL SOCIETY

## **Chemical Communications**

Number 2 1992

## A New Route to 1,3-Benzoxazepines and 1,3-Benzodiazepines *via* Intramolecular Aza-Wittig Reaction

Jyoji Kurita, Takao lwata, Shuji Yasuike and Takashi Tsuchiya\* Faculty of Pharmaceutical Sciences, Hokuriku University, Kanagawa-machi, Kanazawa 920-11, Japan

The reaction of triphenylphosphine with the *o*-acyloxy- **5** and *o*-acylamino-azidocinnamates **9**, prepared from salicylaldehyde and *o*-aminobenzaldehyde, results in ring closure to give the 1,3-benzoxazepines **7** and 1,3-benzodiazepines **12**, *via* the intramolecular aza-Wittig reaction of the iminophosphoranes **6** and **10** initially formed, respectively.

Fully unsaturated 1,3-dihetero seven-membered ring compounds (1,3-diheteroepines) have been prepared mainly by ring transformations of known heterocyclic rings. With regard to benzo compounds, 1,3-benzoxazepines<sup>1,2</sup> and 1,3-benzodiazepines<sup>3</sup> are synthesized by the photochemical rearrangement with ring expansion of isoquinoline *N*-oxides and quinoline *N*-imides, respectively. 3,1-Benzoxazepines are prepared by the photochemical rearrangement of quinoline *N*-oxides<sup>2,4</sup> or by the thermal ring expansion of 1-azidoisochromenes.<sup>5</sup> We report here a new synthetic route to 1,3-benzodiheteroepines involving no ring transformation.

Salicylaldehyde 1 was protected as the methoxymethyl (MOM) derivative 2, which was condensed with ethyl



Scheme 1 Reagents and conditions: i, NaOMe, EtOH, N<sub>3</sub>CH<sub>2</sub>CO<sub>2</sub>Et, 0-5 °C, 2 h, 60%; ii, NaOH, Me(Octyl)<sub>3</sub>N+Cl<sup>-</sup>, H<sub>2</sub>O, CH<sub>2</sub>Cl<sub>2</sub>, ClCH<sub>2</sub>OMe, room temp., 1 h, 95%; iii, acetone, 10% HCl, room temp., 8 h, 85%; iv, PhCOCl, Ac<sub>2</sub>O or ClCO<sub>2</sub>Me, pyridine, room temp., 5–8 h, 60–80%; v, PPh<sub>3</sub>, benzene, Ar, room temp., 3–4 h, 85–90%

azidoacetate to give the azidocinnamates 3. After removal of the MOM group in 3, the resulting phenolic compound 4 was treated with benzoyl chloride, acetic anhydride or methyl chloroformate giving the cinnamyl esters 5 in 30–40% yields from 1.† Treatment of 5 with triphenylphosphine in benzene



Scheme 2 Reagents and conditions: i, PPh<sub>3</sub>, benzene, Ar, room temp., 2 h, 95%; ii, xylene, reflux, 15 h, 40–50%

<sup>†</sup> Direct conversion of unprotected 1 into 4 by treatment with ethyl azidoacetate has been unsuccessful. Similar azidocinnamates obtained by the condensation of benzaldehydes with ethyl azidoacetate are known to be Z-forms (T. L. Gilchrist, C. W. Rees and J. A. R. Rodrigues, J. Chem. Soc., Chem. Commun., 1979, 672; D. M. B. Hickey, C. J. Moody and C. W. Rees, J. Chem. Soc., Chem. Commun., 1982, 1419). Satisfactory elemental analyses and spectroscopic data were obtained for all new compounds reported.

Selected data for **5a**: m.p. 95–96 °C; IR (KBr) 2128 (N<sub>3</sub>),  $v_{C=0}$  1752 and 1712 cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.16 and 4.20 (3H, t, and 2H, q, J 7 Hz, CO<sub>2</sub>Et), 7.02 (1H, s, alkenic H), 7.2–7.6 and 8.1–8.3 (7H, m, and 2H, m, Ph-H); **5b**: m.p. 66–68 °C; **5c**: m.p. 58–59 °C.

at room temperature resulted in ring closure to form the desired 1,3-benzoxazepines 7 in 85–90% yields, probably *via* the intramolecular aza-Witting reaction of the initially formed iminophosphoranes **6**, which could not be isolated, however (Scheme 1).‡ The 2-phenyl-oxazepine **7a** is stable and was isolated in 90% yield, however, the other oxazepines **7b,c** were readily hydrolysed to the ring-opened compounds **8** during isolation by chromatography and thus were purely isolated only in 30–40% yields, together with **8** (50–60%). 1,3-Benzodiheteroepines are known to readily undergo hydrolysis under mild conditions.<sup>1–3</sup>

On the other hand, in the reaction of triphenylphosphine with the o-acylaminoazidocinnamates 9 prepared from oaminobenzaldehyde via two steps, the iminophosphoranes 10 were isolated as stable crystals almost quantitatively. Heating the phosphoranes 10 in refluxing xylene resulted in aza-Wittig reaction to form the 3H-1,3-benzodiazepines 12 in 40–50% yields, presumably via the 1H-isomers 11 initially formed (Scheme 2).§ It is known that N-unsubstituted 1H-1,3-benzodiazepines are unstable and tautomerize rapidly to the

§ Selected data for 10 and 12–14: 10a: m.p. 164–166 °C; 10b: m.p. 154–156 °C; 12a: m.p. 117–118 °C; IR (KBr)  $v_{NH}$  3368 and  $v_{C=0}$  1692 cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.38 and 4.32 (3H, t, and 2H, q, J 7 Hz, CO<sub>2</sub>Et), 6.47 (1H, s, 5-H), 6.61 (1H, br, NH) and 6.7–7.8 (9H, m, Ph-H). The above spectroscopic data show that the ester carbonyl IR absorption appeared at a lower wavelength and the NH proton NMR spectral signal also appeared at a lower field. These lower shifts may be a consequence of hydrogen bonding forming a five-membered ring chelate between the NH and the ethoxycarbonyl oxygen, by analogy with *N*-unsubsituted 2-methoxycarbonyl-1*H*-1,4-benzodiazepines (H. Sashida, M. Kaname and T. Tsuchiya, *Chem. Pharm. Bull.*, 1990, 38, 2919); strongly suggesting that the diazepines 12 are the 3*H*-isomers. 13: m.p. 183–185 °C; 14: m.p. 137–138 °C; IR (KBr)  $v_{C=0}$  1740 and 1724 cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.38 and 4.30 (3H, t, and 2H, q, *J* 7 Hz, CO<sub>2</sub>Et) and 7.1–8.1 (9H, m, ring H).

3*H*-isomers.<sup>6</sup> Similarly, the tetracyclic diazepine **14** was also obtained from the phthalimido compound **13** in 60% yield.

These results indicate that the intramolecular aza-Wittig reaction to the ester carbonyl occurs at below room temperature, whereas a higher temperature is required for that to the amide carbonyl; this difference in reactivity between esters and amides is analogous to those observed in a variety of nucleophilic reactions. Although the aza-Wittig reactions have recently been utilized in the syntheses of nitrogen heterocycles, especially five- and six-membered rings,<sup>7</sup> the present result is the first example of utilization for the synthesis of seven-membered heterocyclic rings.

Received, 17th September 1991; Com. 1/04826K

## References

- C. Kaneko, S. Yamada and M. Ishikawa, *Tetrahedron Lett.*, 1966, 2145; O. Buchardt, C. Lohse, A. M. Duffield and C. Djerassi, *Tetrahedron Lett.*, 1967, 2471; O. Simonsen, C. Lohse and O. Buchardt, *Acta Chem. Scand.*, 1970, 24, 268.
- 2 F. Bellamy and J. Streith, *Heterocycles*, 1976, **4**, 1391; G. G. Spence, E. C. Taylor and O. Buchardt, *Chem. Rev.*, 1970, **70**, 231.
- 3 T. Tsuchiya, M. Enkaku, J. Kurita and H. Sawanishi, J. Chem. Soc., Chem. Commun., 1979, 534; T. Tsuchiya, M. Enkaku and S. Okajima, Chem. Commun., 1979, 534; T. Tsuchiya, M. Enkaku and S. Okajima, Chem. Pharm. Bull., 1980, 28, 2602.
- 4 C. Kaneko and S. Yamada, *Chem. Pharm. Bull.*, 1966, 14, 555; O. Buchardt, B. Jensen and I. K. Larsen, *Acta Chem. Scand.*, 1967, 21, 1841.
- 5 J.-P. Le Roux, P.-L. Desbene and J.-C. Cherton, J. Heterocycl. Chem., 1981, 18, 847.
- 6 J. Kurita, M. Enkaku and T. Tsuchiya, *Heterocycles*, 1983, 20, 2173.
- 7 For example: L. J. Leyshon and D. G. Saunders, J. Chem. Soc., Chem. Commun., 1971, 1608; S. A. Foster, L. J. Leyshon and D. G. Saunders, J. Chem. Soc., Chem. Commun., 1973, 29; Y. G. Gololobov, I. N. Zhmurova and L. F. Kasukhin, Tetrahedron, 1981, 37, 437; S. Eguchi and H. Takeuchi, J. Chem. Soc., Chem. Commun., 1989, 602; H. Takeuchi, S. Yanagida, T. Ozaki, S. Hagiwara and S. Eguchi, J. Org. Chem., 1989, 54, 431; H. Takeuchi and S. Eguchi, Tetrahedron Lett., 1989, 30, 3313; P. Molina, A. Arques and M. V. Vinader, J. Org. Chem., 1990, 55, 4724.

 $<sup>\</sup>ddagger$  Selected data for 7a: oil, IR (film)  $v_{C=O}$  1722 cm $^{-1}$ ;  $^{1}H$  NMR (CDCl<sub>3</sub>)  $\delta$  1.38 and 4.32 (3H, t, and 2H, q, J 8 Hz, CO<sub>2</sub>Et), 7.1–7.5 (7H, m, Ph-H), 7.57 (1H, s, 5-H) and 8.2–8.3 (2H, m, Ph-H); 7b: oil; 7c: oil.