

# Effect of Chelate Ring Size on the Rate of Hydride Transfer from CpRu(P–P)H (P–P = Chelating Diphosphine) to an Iminium Cation

Hairong Guan, Masanori Iimura, Matthew P. Magee, Jack R. Norton,\* and Kevin E. Janak

Department of Chemistry, Columbia University, New York, New York 10027

Received May 8, 2003

The rate constants for hydride transfer from CpRu(P–P)H (P–P = bis(diphenylphosphino)methane (dppm), bis(diphenylphosphino)ethane (dppe), bis(diphenylphosphino)benzene (dpbz), or bis(diphenylphosphino)propane (dppp)) to 1-(1-phenylethylidene)pyrrolidinium tetrafluoroborate have been measured. The bite angles of the hydride complexes CpRu(dppm)H, CpRu(dppe)H, CpRu(dpbz)H, and CpRu(dppb)H (dppb = bis(diphenylphosphino)butane) have been determined by X-ray diffraction. Hydride transfer is faster when the chelate ring of the diphosphine is smaller (i.e., CpRu(dppm)H > CpRu(dppe)H ≈ CpRu(dpbz)H > CpRu(dppp)H >> CpRu(dppb)H). Boiling CpRu(PPh<sub>3</sub>)<sub>2</sub>Cl with dpbz in benzene or toluene results in the formation of [CpRu(PPh<sub>3</sub>)(η<sup>2</sup>-dpbz)]Cl as well as CpRu(dpbz)Cl.


## Introduction

We have previously reported<sup>1</sup> the ruthenium-catalyzed hydrogenation of iminium cations (including protonated imines) by an ionic mechanism, involving heterolytic cleavage of H<sub>2</sub> and sequential transfer of H<sup>−</sup> and H<sup>+</sup> to the substrates. When chiral diphosphines are employed, C=N bonds can be hydrogenated enantioselectively with ee's up to 60%. As the literature contains few catalysts for the asymmetric hydrogenation of C=N bonds,<sup>2,3</sup> we have studied the mechanism of ours; the transfer of H<sup>−</sup> from CpRu(P–P)H to the iminium cation (Scheme 1) has proven to be turnover-limiting and enantioselectivity-determining.<sup>1</sup>

It is possible to stop the reaction after a single H<sup>−</sup> transfer if it is carried out in the absence of H<sub>2</sub> and the presence of a coordinating solvent (usually CH<sub>3</sub>CN) (eq 1). To screen phosphines for catalytic use, we have measured the rate at which their CpRu(P–P)H complexes undergo this “stoichiometric” hydride transfer. When such an experiment (eq 2) showed CpRu((S,S)-DIOP)H to be inactive at hydride transfer to 1-(1-phenylethylidene)pyrrolidinium tetrafluoroborate (**1**),<sup>1</sup> we suspected an effect of the size of the chelate ring. (The CpRu hydrides of other chelating diphosphines with one alkyl and two aryl substituents on each phosphorus transfer H<sup>−</sup> to **1**.)

The effect of chelate ring size on metal-catalyzed C–C bond forming reactions, such as hydroformylation, allylic substitution, hydrocyanation, and the amination of aryl halides, has been reviewed by van Leeuwen and

Scheme 1



co-workers.<sup>4</sup> The Brookhart group has shown that the kinetic barriers for the migratory insertion of complexes [Ph<sub>2</sub>P(CH<sub>2</sub>)<sub>n</sub>PPh<sub>2</sub>]Pd(CH<sub>3</sub>)(CO)<sup>+</sup> (*n* = 2–4) decrease with an increase in the P–Pd–P bond angle.<sup>5</sup> Vilar et al. have recently reported a similar influence of chelate ring size on the insertion of isocyanides into the Pd–C bond of (P–P)Pd(CH<sub>3</sub>)Cl complexes.<sup>6</sup>

(1) Magee, M. P.; Norton, J. R. *J. Am. Chem. Soc.* **2001**, *123*, 1778–1779.

(2) (a) Willoughby, C. A.; Buchwald, S. L. *J. Am. Chem. Soc.* **1994**, *116*, 8952–8965. (b) Broger, E. A.; Burkart, W.; Hennig, M.; Scalone, M.; Schmid, R. *Tetrahedron: Asymmetry* **1998**, *9*, 4043–4054. (c) Xiao, D.; Zhang, X. *Angew. Chem., Int. Ed.* **2001**, *40*, 3425–3428.

(3) Review on hydrogenation of imines: Blaser, H.-U.; Spindler, F. *Hydrogenation of Imine Groups*. In *Comprehensive Asymmetric Catalysis*; Jacobsen, E. N., Pfaltz, A., Yamamoto, H., Eds.; Springer-Verlag: New York, 1999; Vol. 1, pp 247–265.

(4) (a) van Leeuwen, P. W. N. M.; Kamer, P. C. J.; Reek, J. N. H.; Dierkes, P. *Chem. Rev.* **2000**, *100*, 2741–2769. (b) Kamer, P. C. J.; van Leeuwen, P. W. N. M.; Reek, J. N. H. *Acc. Chem. Res.* **2001**, *34*, 895–904.

(5) Ledford, J.; Shultz, C. S.; Gates, D. P.; White, P. S.; DeSimone, J. M.; Brookhart, M. *Organometallics* **2001**, *20*, 5266–5276.

However, little is known about the effect of chelate ring size on hydride transfer, although much work has been done on other factors influencing the rate of such transfers from transition-metal hydrides.<sup>7</sup> DuBois and co-workers have determined the equilibrium constants for several H<sup>-</sup> transfer reactions of [HM(diphosphine)<sub>2</sub>]-PF<sub>6</sub> (M = Pt, Ni) complexes and have found that a smaller chelate ring favors hydride transfer *thermodynamically*.<sup>8</sup> Knowledge of the *kinetic* effects of chelate ring size is still lacking.

In this paper we report the kinetics of hydride transfer from CpRu(P-P)H, with bidentate phosphines of different chelate ring sizes, to the iminium cation **1**. We also report the structures of these complexes and their influence on the rates of hydride transfer.

## Results and Discussion

**Preparation of CpRu(P-P)H.** The literature contains two general methods of preparing CpRu(P-P)H: (1) the synthesis of CpRu(P-P)Cl and its reduction by CH<sub>3</sub>ONa,<sup>9</sup> and (2) the displacement of carbonyl ligands from CpRu(CO)<sub>2</sub>H by diphosphines.<sup>10</sup> Method 1 is more convenient because the chlorides CpRu(P-P)Cl (**3**) are air stable; however, method 2 gives better results when the distance between the P atoms is large and bridging is a possibility. We thus prepared CpRu(dppm)H (**2a**), CpRu(dppe)H (**2b**), and CpRu(dpbz)H (**2c**) by method 1; CpRu(dppp)H (**2d**) and CpRu(dppb)H (**2e**) by method 2.

Most **3** are readily prepared by displacing PPh<sub>3</sub> from CpRu(PPh<sub>3</sub>)<sub>2</sub>Cl with the appropriate diphosphine. However, the preparation of CpRu(dpbz)Cl (**3c**) by this method in boiling benzene (eq 3) gives an interesting byproduct: a light green precipitate, **4**. (The precipitate **4** is not an intermediate in the formation of **3c**: **4** remains after 3 days in boiling benzene, or when the higher-boiling toluene is used as solvent.) The <sup>1</sup>H NMR of **4** shows, in addition to the Cp singlet (δ 4.71), 39 protons in the aromatic region; the <sup>31</sup>P{<sup>1</sup>H} NMR reveals a triplet at δ 43.39 and a doublet at δ 72.05, with *J* = 34.0 Hz. Thus **4** is [CpRu(PPh<sub>3</sub>)(η<sup>2</sup>-dpbz)]Cl, analogous to the known [CpRu(PPh<sub>3</sub>)(η<sup>2</sup>-dppm)]BF<sub>4</sub>.<sup>11,12</sup>

(6) Owen, G. R.; Vilar, R.; White, A. J. P.; Williams, D. J. *Organometallics* **2002**, *21*, 4799–4807.


(7) (a) Labinger, J. A. Nucleophilic Reactivity of Transition Metal Hydrides. In *Transition Metal Hydrides*; Dedieu, A., Ed.; VCH: New York, 1992; pp 361–379. (b) Labinger, J. A.; Komadina, K. H. *J. Organomet. Chem.* **1978**, *155*, C25–C28. (c) Kinney, R. J.; Jones, W. D.; Bergman, R. G. *J. Am. Chem. Soc.* **1978**, *100*, 7902–7915. (d) Kao, S. C.; Gaus, P. L.; Youngdahl, K.; Darensbourg, M. Y. *Organometallics* **1984**, *3*, 1601–1603. (e) Kao, S. C.; Spillett, C. T.; Ash, C.; Lusk, R.; Park, Y. K.; Darensbourg, M. Y. *Organometallics* **1985**, *4*, 83–91. (f) Gaus, P. L.; Kao, S. C.; Youngdahl, K.; Darensbourg, M. Y. *J. Am. Chem. Soc.* **1985**, *107*, 2428–2434. (g) Martin, B. D.; Warner, K. E.; Norton, J. R. *J. Am. Chem. Soc.* **1986**, *108*, 33–39. (h) Smith, K.-T.; Norton, J. R.; Tilset, M. *Organometallics* **1996**, *15*, 4515–4520. (i) Hembre, R. T.; McQueen, S. *J. Am. Chem. Soc.* **1994**, *116*, 2141–2142. (j) Hembre, R. T.; McQueen, J. S. *Angew. Chem., Int. Ed. Engl.* **1997**, *36*, 65–67. (k) Cheng, T.-Y.; Bullock, R. M. *Organometallics* **1995**, *14*, 4031–4033. (l) Cheng, T.-Y.; Brunshwig, B. S.; Bullock, R. M. *J. Am. Chem. Soc.* **1998**, *120*, 13121–13137. (m) Cheng, T.-Y.; Bullock, R. M. *Organometallics* **2002**, *21*, 2325–2331. (n) Sarker, N.; Bruno, J. W. *J. Am. Chem. Soc.* **1999**, *121*, 2174–2180.

(8) Berning, D. E.; Noll, B. C.; DuBois, D. L. *J. Am. Chem. Soc.* **1999**, *121*, 11432–11447.

(9) Bruce, M. I.; Humphrey, M. G.; Swincer, A. G.; Wallis, R. C. *Aust. J. Chem.* **1984**, *37*, 1747–1755.


(10) White, C.; Cesarotti, E. *J. Organomet. Chem.* **1985**, *287*, 123–125.

(11) Bruce, M. I.; Humphrey, M. G.; Patrick, J. M.; White, A. H. *Aust. J. Chem.* **1983**, *36*, 2065–2072.



**Figure 1.** Structure of [CpRu(PPh<sub>3</sub>)(η<sup>2</sup>-dpbz)]Cl (**4**) (20% probability level). Selected bond lengths (Å) and angles (deg): Ru–P1 2.3478(6), Ru–P2 2.2787(7), Ru–P3 2.3136(7), P1–Ru–P2 97.26(2), P1–Ru–P3 97.58(2), P3–Ru–P2 82.87(2).

and [CpRu(PPh<sub>3</sub>)(η<sup>2</sup>-dppe)]BF<sub>4</sub><sup>12</sup> (prepared by treating CpRu(dppm)Cl or CpRu(dppe)Cl with PPh<sub>3</sub> and NH<sub>4</sub>-BF<sub>4</sub> in CH<sub>3</sub>OH).



A dimeric or oligomeric structure for **4** in solution cannot be ruled out, although we are not aware of any case where dpbz has served as a bridging ligand. We have been able to grow single crystals of **4**, by slow diffusion of hexanes into a concentrated CH<sub>2</sub>Cl<sub>2</sub> solution. X-ray diffraction shows (Figure 1)<sup>13</sup> a monomeric structure in the solid state.

The ruthenium in **4** sits in a very crowded environment, with both the PPh<sub>3</sub> and the dpbz coordinated. The dissociation of the Cl<sup>-</sup> in eq 3 parallels its dissociation in polar solvents, such as CH<sub>3</sub>OH, C<sub>2</sub>H<sub>5</sub>OH, and DMSO. Treichel and co-workers have measured the rates of halide ion solvolysis of CpRuL<sub>2</sub>Cl (L = PMe<sub>3</sub>, PPhMe<sub>2</sub>, PPh<sub>2</sub>Me, PPh<sub>2</sub>(OMe), P(OEt)<sub>3</sub>, PMe(OMe)<sub>2</sub>; L<sub>2</sub> = dppe) in DMSO and CH<sub>3</sub>CN.<sup>14</sup> (They found that a less electron-donating phosphine retards Cl<sup>-</sup> dissociation, while silver, ammonium, or sodium salts facilitate it.) However, in nonpolar solvents such as benzene, toluene, or decalin, phosphine dissociation from CpRu(P-P)Cl

(12) Ashby, G. S.; Bruce, M. I.; Tomkins, I. B.; Wallis, R. C. *Aust. J. Chem.* **1979**, *32*, 1003–1016.


(13) The solvent molecule is severely disordered, and a suitable model cannot be constructed for the solvate. The program SQUEEZE (see van der Sluis, P.; Spek, A. L. *Acta Crystallogr.* **1990**, *A46*, 194–201) was used to refine the solvent molecule as a diffuse contribution to overall scattering without specific atom positions; however, the empirical formula, density, and absorption coefficient reflect the full formula.

(14) (a) Treichel, P. M.; Komar, D. A.; Vincenti, P. J. *Inorg. Chim. Acta* **1984**, *88*, 151–152. (b) Treichel, P. M.; Vincenti, P. J. *Inorg. Chem.* **1985**, *24*, 228–230.


is generally more facile,<sup>15</sup> so the Cl<sup>−</sup> dissociation in eq 3 is surprising.

Most of the hydrides **2** proved to be stable in halogenated solvents, but **2d**, regardless of the batch or the synthetic methods used to prepare it, gave CpRu(dppp)-Cl within minutes in CD<sub>2</sub>Cl<sub>2</sub>.<sup>16,17</sup> This type of halogenation is known to be a radical process<sup>18</sup> and potentially retarded by radical inhibitors, so we examined the stability of **2d** in the presence of several such inhibitors. After 1 day with a suspension of hydroquinone (only slightly soluble) in CD<sub>2</sub>Cl<sub>2</sub>, about 85% of the **2d** remained, but 4-*tert*-butylcatechol, which has better solubility in halogenated solvents, reacted immediately with **2d** (to give complex products that were not identified). However, TEMPO (2,2,6,6-tetramethyl-1-piperidinyloxy) proved unreactive toward **2d** and efficient at preventing its reaction with CD<sub>2</sub>Cl<sub>2</sub>: no significant decomposition of **2d** in CD<sub>2</sub>Cl<sub>2</sub> solution was observed after 30 h in the presence of 1% (by weight) TEMPO.

The fact that TEMPO does not itself abstract H<sup>•</sup> from **2d** warrants comment. TEMPO abstracts H<sup>•</sup> from group 14 organometallic hydrides (e.g., Bu<sub>3</sub>SnH) at 80 °C,<sup>19</sup> forming the hydroxylamine **5** (eq 4). As the O–H bond dissociation energy (BDE) of **5** in benzene is 69 kcal/mol<sup>20</sup> and the Ru–H BDE value for **2d** can be estimated at 65 cal/mol,<sup>21</sup> H<sup>•</sup> abstraction from **2d** by TEMPO should be thermodynamically favorable. The reason it does not occur at an appreciable rate must be kinetic: the cyclopentadienyl ring and dppp ligand in **2d** must make the hydride ligand inaccessible to a bulky radical like TEMPO.



**Kinetics of Hydride Transfer from the Hydride Complexes **2** to the Iminium Cation **1**.** When H<sup>−</sup> transfer to **1** from any of the hydride complexes **2a–d** is carried out in the presence of CH<sub>3</sub>CN, the cation **6** results (eq 5). Monitoring the disappearance of the Cp signal of the hydride **2b** by <sup>1</sup>H NMR in the presence of excess **1** (10 equiv) gave a pseudo-first-order rate constant *k*<sub>obs</sub>. Variation of [**1**] from 0.10 to 0.26 M (Table S-1, Figure 2) showed that *k*<sub>obs</sub> for **2b** was linear with [**1**], implying that the hydride transfer reaction in eq 5




**Figure 2.** Plot of *k*<sub>obs</sub> vs [**1**] for hydride transfer from **2b** to the iminium cation **1** at 300 K with MeCN present (●). (The value of *k*<sub>obs</sub> without MeCN present is also displayed in this figure (□).)

**Table 1. Rates of Hydride Transfer from CpRu(P–P)H to the Iminium Cation **1** at 300 K in CD<sub>2</sub>Cl<sub>2</sub>**

complex	chelating ligand	bite angle P–Ru–P (deg)	rate constant <i>k</i> (M <sup>−1</sup> s <sup>−1</sup> )
<b>2a</b>	dppm	72.01(3)	6.4(3) × 10 <sup>−1a</sup>
<b>2b</b>	dppe	84.497(16)	2.0(1) × 10 <sup>−2</sup>
<b>2c</b>	dpbz	84.22(4)	2.1(2) × 10 <sup>−2</sup>
<b>2d</b>	dppp	91.9(1) <sup>b</sup>	2.8(2) × 10 <sup>−3</sup>
<b>2e</b>	dppb	93.96(3)	no reaction (<4 × 10 <sup>−6</sup> )

<sup>a</sup> Average of two kinetic experiments with different concentrations of **1** and **2a**. <sup>b</sup> See ref 31.

is second-order overall (eq 6). The rate constant *k* for **2b** was calculated from the slope of Figure 2 and inserted in Table 1.



$$-\frac{d[\mathbf{2}]}{dt} = k_{\text{obs}}[\mathbf{2}] = k[\mathbf{1}][\mathbf{2}] \quad (6)$$

Without CH<sub>3</sub>CN *k*<sub>obs</sub> does not change appreciably (Figure 2), although the amine **7** becomes coordinated instead of the CH<sub>3</sub>CN.<sup>22</sup> The rate-determining step in reaction 5 is clearly H<sup>−</sup> transfer, with CH<sub>3</sub>CN serving as an efficient trap. Sixteen-electron half-sandwich ruthenium diphosphine complexes, with and without agostic interactions, have been isolated and structurally characterized. They are particularly stable when a Cp<sup>\*</sup> ring and bulky phosphine ligands such as (tPr)<sub>2</sub>PCH<sub>2</sub>–CH<sub>2</sub>P(tPr)<sub>2</sub> or two PMe(tPr)<sub>2</sub> are both present,<sup>23</sup> but can also be made when there is a less bulky diphosphine.<sup>24</sup> (Sixteen-electron complexes with less bulky diphosphines are more reactive.)

The *k*<sub>obs</sub> value for **2c** was obtained by monitoring the appearance of the coordinated CH<sub>3</sub>CN signal of **6c**,

(22) Iimura, M.; Magee, M. P.; Norton, J. R. Unpublished work.

(23) Jiménez-Tenorio, M.; Mereiter, K.; Puerta, M. C.; Valerga, P. *J. Am. Chem. Soc.* **2000**, *122*, 11230–11231.

(15) Davies, S. G.; McNally, J. P.; Smallridge, A. J. *Adv. Organomet. Chem.* **1990**, *30*, 1–76, and references therein.

(16) The fact that the dppp hydride **2d** is more reactive toward CD<sub>2</sub>Cl<sub>2</sub> than **2a** and **2b** is probably due to the more negative potential of **2d**,<sup>17</sup> which enables it to initiate a radical chain more easily by electron transfer to the solvent.

(17) Jia, G.; Morris, R. H. *J. Am. Chem. Soc.* **1991**, *113*, 875–883.


(18) (a) Shackleton, T. A.; Mackie, S. C.; Fergusson, S. B.; Johnson, L. J.; Baird, M. C. *Organometallics* **1990**, *9*, 2248–2253. (b) Brown, T. L. In *Organometallic Radical Processes*; Trogler, W. C., Ed.; Journal of Organometallic Chemistry Library 22; Elsevier: New York, 1990; pp 67–107, and references therein.

(19) Lucarini, M.; Marchesi, E.; Pedulli, G. F.; Chatgililoglu, C. *J. Org. Chem.* **1998**, *63*, 1687–1693.

(20) Mahoney, L. R.; Mendenhall, G. D.; Ingold, K. U. *J. Am. Chem. Soc.* **1973**, *95*, 8610–8614.

(21) The BDE of CpRu(CO)<sub>2</sub>H has been reported as 65 kcal/mol, and phosphine substitution usually has a negligible effect on M–H energies. See: (a) Tilset, M.; Parker, V. D. *J. Am. Chem. Soc.* **1989**, *111*, 6711–6717; correction, *J. Am. Chem. Soc.* **1990**, *112*, 2843. (b) Kristjánssdóttir, S. S.; Norton, J. R. Acidity of Hydrido Transition Metal Complexes in Solution. In *Transition Metal Hydrides*; Dedieu, A., Ed.; VCH: New York, 1992; pp 309–359.





**Figure 3.** Plot of  $\ln\{([RuH]_0/[RuH]_t) + 1\}$  vs time  $t$  for hydride transfer from **2a** to the iminium cation **1** at 300 K in  $CD_2Cl_2$  with MeCN present. Initially  $[2a] = 7.5 \times 10^{-3}$  M,  $[1] = 1.5 \times 10^{-2}$  M.

because the Cp of **6c** has a  $^1H$  NMR chemical shift ( $\delta$  4.635) very close to that of the Cp of **2c** ( $\delta$  4.639). Variation of  $[1]$  from 0.10 to 0.25 M (Table S-2, Figure S-2) also showed a linear relationship between  $k_{obs}$  and  $[1]$ , confirming that the hydride transfer is a second-order reaction. The rate constant  $k$  for **2c** has been calculated from the slope of Figure S-2 and is shown in Table 1.

Measuring the kinetics of  $H^-$  transfer to **1** from **2d** required the addition of TEMPO to suppress H/Cl exchange between **2d** and  $CD_2Cl_2$ . (A control experiment with the stable hydride **2b** showed that the presence of TEMPO had no effect on the rate of hydride transfer.<sup>25</sup>) The kinetics behavior of **2d** was similar to that of **2b** (Table S-3, Figure S-3). The rate constant  $k$  for **2d** was calculated from the slope of Figure S-3 and is shown in Table 1.

The transfer of hydride to **1** from **2a** was so fast that it was difficult to measure  $k_{obs}$  under pseudo-first-order conditions (the reaction of **2a** with **1** was complete within minutes). The rate constant was measured using a 1:2 ratio of  $[2a]$  to  $[1]$ . Under these conditions, a second-order reaction between **2a** and **1** will behave according to eq 7,<sup>26</sup> where  $[RuH]_t$  is the concentration of **2a** at time  $t$ ,  $[RuH]_0$  is the initial concentration of **2a**, and  $k$  is the second-order rate constant.


$$\ln\{1 + (\Delta_0[RuH]_0/[2a]_0[RuH]_t)\} = \ln([1]_0/[2a]_0) + k\Delta_0 t \quad (7)$$

$$\Delta_0 = [1]_0 - [2a]_0$$

When  $[1]_0 = 2[2a]_0$ ,  $\Delta_0 = [2a]_0$ , eq 7 can be rewritten as eq 8.

$$\ln\{1 + ([RuH]_0/[RuH]_t)\} = \ln 2 + k[2a]_0 t \quad (8)$$

Experimental data confirm a linear relationship between  $\ln\{1 + [RuH]_0/[RuH]_t\}$  and  $t$  (Figure 3). Fur-



**Figure 4.** Molecular structure of  $CpRu(dppm)H$  (**2a**) (20% probability level). Selected bond lengths (Å) and angle (deg): Ru-P1 2.2435(9), Ru-P2 2.2399(9), P1-Ru-P2 72.01(3).

thermore, the  $y$ -intercept ( $0.744 \pm 0.072$ ) is consistent with  $\ln 2$  in eq 8, again confirming that  $H^-$  transfer from **2** to **1** is a second-order reaction.<sup>27</sup> The rate constant  $k$  for **2a** has been derived from the slope ( $k[2a]_0$ ) and is shown in Table 1.

**Structures of Hydride Complexes  $CpRu(P-P)H$  (**2**).** The literature contains surprisingly little crystallographic information about half-sandwich ruthenium phosphine hydrides. Structures had been reported for the bisphosphine complexes  $CpRu(PPh_3)_2H$ <sup>28</sup> and  $CpRu(PMe_3)_2H$ <sup>29</sup> and for the chelating diphosphine complexes  $CpRu(dppf)H$  ( $dppf$  = bis(diphenylphosphino)ferrocene)<sup>30</sup> and **2d**.<sup>31</sup>

Crystals of **2a** were obtained during a study of the mechanism of ionic hydrogenation.<sup>32</sup> Crystals of **2b**, **2c**, and **2e** were grown by adding  $CH_3OH$  to a saturated benzene solution of the hydrides. Thermal ellipsoid plots of the solid-state structures of **2a**, **2b**, **2c**, and **2e** are shown in Figures 4–7. (Figure 6 shows only one of the two independent molecules of **2c** in a unit cell.<sup>33</sup>)

The coordination spheres of Ru in all these hydride complexes adopt pseudo-octahedral geometries. (All the hydride ligands were located except the one in **2e**.) The Ru–P distances of these hydrides (from 2.21 Å in **2c** to 2.25 Å in **2e**) are similar to those found in the structures previously reported. The only significant differences are in the P–Ru–P angles, which vary from 72.01(3)° in **2a** to 93.96(3)° in **2e**. Relevant bite angles are shown in Table 1.

**Relationship between Chelate Bite Angle and Hydride Transfer Rate.** As Table 1 shows,  $k$  decreases

(27) Two experiments with different initial concentrations gave the same second-order rate constants within experimental error:  $k_2 = 6.5(2) \times 10^{-1} s^{-1} M^{-1}$  when  $[2a] = 7.5 \times 10^{-3} M$ ,  $[1] = 1.5 \times 10^{-2} M$ ;  $k_2 = 6.3(3) \times 10^{-1} s^{-1} M^{-1}$  when  $[2a] = 1.0 \times 10^{-2} M$ ,  $[1] = 2.0 \times 10^{-2} M$ .

(28) Smith, K. T.; Rømming, C.; Tilset, M. *J. Am. Chem. Soc.* **1993**, *115*, 8681–8689.

(29) (a) Lemke, F. R.; Brammer, L. *Organometallics* **1995**, *14*, 3980–3987. (b) Brammer, L.; Klooster, W. T.; Lemke, F. R. *Organometallics* **1996**, *15*, 1721–1727.

(30) Bruce, M. I.; Butler, I. R.; Cullen, W. R.; Koutsantonis, G. A.; Snow, M. R.; Tiekinck, E. R. T. *Aust. J. Chem.* **1988**, *41*, 963–969.

(31) Litster, S. A.; Redhouse, A. D.; Simpson, S. J. *Acta Crystallogr.* **1992**, *C48*, 1661–1663.

(32) Single crystals of **2a** for structure analysis were grown from a saturated solution of **2a**, a slight excess of **1** (1.5 equiv), and an excess of free amine **7** (ca. 7 equiv) at  $-35^\circ C$ .<sup>22</sup>

(33) The aromatic ring in the backbone of the other molecule is disordered.

(24) Aneetha, H.; Jiménez-Tenorio, M.; Puerta, M. C.; Valerga, P.; Sapunov, V. N.; Schmid, R.; Kirchner, K.; Mereiter, K. *Organometallics* **2002**, *21*, 5334–5346. Extensive earlier theoretical work on such systems is cited therein.

(25)  $k_{obs} = 3.6(1) \times 10^{-3} s^{-1}$  when  $[2b] = 1.5 \times 10^{-2} M$ ,  $[1] = 1.5 \times 10^{-1} M$ ;  $k_{obs} = 3.5(2) \times 10^{-2} s^{-1}$  when  $[2b] = 1.5 \times 10^{-2} M$ ,  $[1] = 1.5 \times 10^{-1} M$ , with 1% (by weight) TEMPO.

(26) Espenson, J. H. *Chemical Kinetics and Reaction Mechanisms*, 2nd ed.; McGraw-Hill: New York, 1995; p 25, eq 2–34.



Table 2. Summary of Crystallographic Data

	4·C <sub>6</sub> H <sub>14</sub>	2a·CH <sub>2</sub> Cl <sub>2</sub>	2b·C <sub>6</sub> H <sub>6</sub>	2c	2e(–H)
empirical formula	C <sub>59</sub> H <sub>58</sub> ClP <sub>3</sub> Ru	C <sub>31</sub> H <sub>30</sub> Cl <sub>2</sub> P <sub>2</sub> Ru	C <sub>37</sub> H <sub>36</sub> P <sub>2</sub> Ru	C <sub>35</sub> H <sub>30</sub> P <sub>2</sub> Ru	C <sub>33</sub> H <sub>33</sub> P <sub>2</sub> Ru
fw	996.48	636.46	643.67	613.60	592.60
temp, K	293(2)	293(2)	243(2)	293(2)	293(2)
cryst syst	monoclinic	monoclinic	monoclinic	monoclinic	monoclinic
space group	<i>P</i> 2 <sub>1</sub> / <i>c</i>	<i>P</i> 2 <sub>1</sub> / <i>c</i>	<i>P</i> 2 <sub>1</sub> / <i>c</i>	<i>P</i> 2 <sub>1</sub> / <i>c</i>	<i>P</i> 2 <sub>1</sub> / <i>c</i>
<i>a</i> , Å	9.7432(6)	9.7901(5)	11.7095(6)	11.0831(7)	10.6447(5)
<i>b</i> , Å	23.3215(16)	17.0869(10)	18.9662(9)	27.3950(17)	19.8601(9)
<i>c</i> , Å	20.5948(14)	17.2365(10)	14.3963(7)	18.6843(11)	13.3136(6)
α, deg	90	90	90	90	90
β, deg	94.0850(10)	95.2370(10)	107.3910(10)	92.2170(10)	98.7860(10)
γ, deg	90	90	90	90	90
<i>V</i> , Å <sup>3</sup>	4667.8(5)	2871.3(3)	3051.0(3)	5668.7(6)	2781.5(2)
<i>Z</i>	4	4	4	8	4
<i>d</i> <sub>calc</sub> , g/cm <sup>3</sup>	1.418	1.472	1.401	1.438	1.415
λ(Mo Kα), Å	0.71073	0.71073	0.71073	0.71073	0.71073
μ, mm <sup>–1</sup>	0.537	0.862	0.643	0.689	0.699
no. of data collected	31 301	19 076	20 068	33 512	18 736
no. of unique data	10 391	6469	6851	12 872	6319
no. of data/restraints/params	10391/0/524	6469/0/330	6851/0/366	12 872/0/694	6319/0/326
GOP on <i>F</i> <sup>2</sup>	1.028	1.028	1.029	1.011	1.030
R1, wR2 ( <i>I</i> > 2σ( <i>I</i> ))	0.0366, 0.0724	0.0449, 0.1176	0.0250, 0.0633	0.0531, 0.1347	0.0358, 0.0937
R1, wR2 (all data)	0.0667, 0.0791	0.0694, 0.1285	0.0291, 0.0652	0.0671, 0.1437	0.0548, 0.1017

**CpRu(η<sup>2</sup>-dpbz)Cl (3c) and [CpRu(PPh<sub>3</sub>)(η<sup>2</sup>-dpbz)]Cl (4).** Under a nitrogen atmosphere 300 mL of benzene was added to a mixture of CpRu(PPh<sub>3</sub>)<sub>2</sub>Cl (1.162 g, 1.6 mmol) and dpbz (734 mg, 1.64 mmol), giving an orange-red solution. While the solution was boiled for 24 h, a light green precipitate formed, which was removed by filtration after the mixture was cooled to room temperature. The volume of the orange filtrate was reduced to 40 mL, then hexanes were added to cause precipitation. Solvent was removed by cannula, and the orange solid was washed with hexanes three times, then dried under vacuum to give 591 mg of the orange powder **3c** (57% yield). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 4.62 (s, Cp, 5H), 6.92–6.96 (m, Ar, 4H), 7.09–7.13 (m, Ar, 4H), 7.22–7.27 (m, Ar, 2H), 7.36–7.42 (m, Ar, 8H), 7.67–7.69 (m, Ar, 2H), 7.90–7.94 (m, Ar, 4H). <sup>31</sup>P NMR (121.1 MHz, CDCl<sub>3</sub>): δ 75.21 (s). Its LRMS (FAB<sup>+</sup>) showed a parent ion peak at *m/e* 648.1 (<sup>102</sup>Ru), with the appropriate isotopic distribution. Anal. Calcd for C<sub>35</sub>H<sub>29</sub>P<sub>2</sub>RuCl: C, 64.87; H, 4.51; Cl, 5.47. Found: C, 64.75; H, 4.70; Cl, 5.47. The light green solid after first filtration was washed with benzene twice and hexanes three times, then dried under vacuum to afford 400 mg of **4** as a light green powder (27% yield). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 4.71 (s, Cp, 5H), 6.33–6.38 (m, Ar, 4H), 6.40–6.50 (m, Ar, 2H), 6.80–6.90 (m, Ar, 2H), 6.95–7.05 (m, Ar, 12H), 7.07–7.12 (m, Ar, 4H), 7.19–7.26 (m, Ar, 7H), 7.30–7.34 (m, Ar, 4H), 7.66–7.67 (m, Ar, 4H). <sup>31</sup>P NMR (121.1 MHz, CDCl<sub>3</sub>): δ 43.39 (t, *J*<sub>P–P</sub> = 34.0 Hz), 72.05 (d, *J*<sub>P–P</sub> = 34.0 Hz). Its LRMS (FAB<sup>+</sup>) showed *m/e* 875.1 for [CpRu(PPh<sub>3</sub>)(η<sup>2</sup>-dpbz)]<sup>+</sup> (<sup>102</sup>Ru) and *m/e* 613.1 for [CpRu(η<sup>2</sup>-dpbz)]<sup>+</sup> (<sup>102</sup>Ru) with the appropriate isotopic distribution. Anal. Calcd for C<sub>53</sub>H<sub>44</sub>P<sub>3</sub>RuCl: C, 69.92; H, 4.87; Cl, 3.89. Found: C, 70.10; H, 4.97; Cl, 3.60.

**CpRu(η<sup>2</sup>-dpbz)H (2c).** Sodium (110 mg, 4.78 mmol) was added to a suspension of 575 mg of CpRu(dp bz)Cl (**3c**) (0.66 mmol) in 30 mL of CH<sub>3</sub>OH. Boiling the resulting mixture for 2 h gave a yellow suspension, from which (after cooling to room temperature) the solvent was removed by cannula. The solid was washed with CH<sub>3</sub>OH three times, then dried under vacuum to give 242 mg of yellow powder **2c** (48% yield). <sup>1</sup>H NMR (400 MHz, CD<sub>2</sub>Cl<sub>2</sub>): δ –13.36 (t, RuH, *J*<sub>P–H</sub> = 33.2 Hz, 1H), 4.63 (s, Cp, 5H), 7.16–7.27 (m, Ar, 18H), 7.32–7.34 (m, Ar, 2H), 7.52–7.56 (m, Ar, 4H). <sup>31</sup>P NMR (CD<sub>2</sub>Cl<sub>2</sub>): δ 90.26 (s). Its LRMS (FAB<sup>+</sup>) showed a parent ion peak at *m/e* 613.6 (<sup>102</sup>Ru), with the appropriate isotopic distribution. Anal. Calcd for C<sub>35</sub>H<sub>30</sub>P<sub>2</sub>Ru: C, 68.51; H, 4.93. Found: C, 67.82; H, 4.77.

**NMR Measurements of Rate Constants for Hydride Transfer from the Hydride Complexes 2a–d to the Iminium Cation 1.** In a typical experiment, **2b** (11.3 mg, 20 μmol) and **1** (52.2 mg, 200 μmol) were placed in a vial in an inert atmosphere box. CD<sub>2</sub>Cl<sub>2</sub> (1.0 mL) containing CH<sub>3</sub>CN (4.2 μL, 80 μmol) was added by a gastight syringe, and the solution was immediately transferred to a J. Young NMR tube. The first <sup>1</sup>H NMR spectrum was recorded within 5 min, and single-pulse spectra were taken every 120 s for three to five half-lives. The height of the Cp resonance of the hydride (δ 4.66) was compared to that of residual CDHCl<sub>2</sub> (δ 5.32). Zero-filling was used to ensure adequate digital resolution. The probe temperature (300 K) was calibrated using anhydrous methanol.<sup>38</sup>

For **2a** the procedure was similar except that 2 equiv of **1** was used. For **2d**, 1% (by weight) TEMPO was dissolved in CD<sub>2</sub>Cl<sub>2</sub> before CD<sub>2</sub>Cl<sub>2</sub> was added to the starting materials. For **2c** hexamethylcyclotrisiloxane was used as internal standard, and the reaction was monitored by observing the growth of the signal of coordinated CH<sub>3</sub>CN (δ 0.99) in the product [CpRu(dp bz)(CH<sub>3</sub>CN)]BF<sub>4</sub> (**6c**). For **2e** <5% reaction was observed after 24 h with excess **1** (0.15 M), implying *k*<sub>obs</sub> < 6 × 10<sup>–7</sup> s<sup>–1</sup> and *k* < 4 × 10<sup>–6</sup> M<sup>–1</sup> s<sup>–1</sup>.

**X-ray Structure Determinations.** Crystal data collection and refinement parameters are summarized in Table 2. Data were collected on a Bruker P4 diffractometer equipped with a SMART CCD detector. The structures were solved using direct methods and standard difference map techniques and refined by full-matrix least-squares procedures using SHELXTL. Hydrogen atoms on carbon were included in calculated positions.

**Acknowledgment.** This research was supported by NSF grant CHE-0211310. We thank Dr. D. L. DuBois and Dr. R. M. Bullock for helpful discussions, and Prof. G. Parkin for assistance with the X-ray structure determinations.

**Supporting Information Available:** Complete details of the crystallographic study (PDF and CIF), kinetic data, and plots of these data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OM0303404

(37) Leonard, N. J.; Paukstelis, J. V. *J. Org. Chem.* **1963**, *28*, 3021–3024.

(38) (a) van Geet, A. L. *Anal. Chem.* **1970**, *42*, 679–680. (b) Raidford, D. S.; Fisk, C. L.; Becker, E. D. *Anal. Chem.* **1979**, *51*, 2050–2051.