

Synthesis of 3,4-Dihydro-2*H*-pyrroles from γ,δ -Unsaturated Ketone *O*-Acetyloximes


Masayuki Yoshida, Mitsuru Kitamura, and Koichi Narasaka*

Department of Chemistry, Graduate School of Science, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033

(Received October 29, 2001; CL-011072)

3,4-Dihydro-2*H*-pyrroles were synthesized from γ,δ -unsaturated ketone *O*-acetyloximes by treatment with acetic acid and 1,4-cyclohexadiene in the presence of a catalytic amount of 1,5-naphthalenediol. During the cyclization, *syn-anti* isomerization of the *O*-acetyloximes easily occurred and so, both stereoisomers of oximes could be employed.

Recently, we reported unusual intramolecular SN2-type substitution on sp² nitrogen atom of oximes.¹ For example, *O*-methylsulfonyloximes having an active methine group are converted to cyclic imines by treatment with 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU).^{1e,f} Due to a characteristic nature of SN2 reaction, only *anti*-isomers² of oximes are converted to cyclic imines. This stereospecificity seems to be a serious drawback in applying this reaction for organic synthesis.



It was expected that both of the stereoisomers of oxime derivatives could be employed in the cyclization if *syn-anti* isomerization occurs under the reaction conditions. Although oximes themselves easily isomerize under acidic conditions, isomerization of *O*-substituted oximes hardly occurs under mild conditions.³ We investigated oxime derivatives that isomerize under mild conditions, and *O*-acetyloximes were found to be isomerized by treatment with acetic acid. The isomerization proceeded slowly, but it was accelerated in the presence of a catalytic amount of 1,4-hydroquinone or 1,5-naphthalenediol. That is, isomerization of (*E*)-4-phenyl-2-butanone *O*-acetyloxime attained an equilibrium (*E* : *Z* = 3 : 1) by treatment with 5 molar amounts of acetic acid in toluene-*d*₈ at 80 °C after 3 h, while it took only 1.5 h in the presence of a 5% molar amount of 1,4-hydroquinone.

Based on these findings, we examined the intramolecular cyclization of γ,δ -unsaturated ketone *O*-acetyloxime **1a** (*syn* : *anti* = 1 : 1), expecting the formation of 2-acetoxy-methyl-5-phenethyl-3,4-dihydro-2*H*-pyrrole (**2a**) by intramolecular nucleophilic attack of the olefinic moiety on sp² nitrogen of the *O*-acetyloxime (Table 1).


By treating **1a** with 2 molar amounts of acetic acid and a catalytic amount of 1,4-hydroquinone or 1,5-naphthalenediol in refluxing 1,4-dioxane, the expected cyclization product **2a** was

obtained in about 20% yield with 45% yield of 2-methyl-5-phenethyl-3,4-dihydro-2*H*-pyrrole (**3a**) (Table 1, Entries 1 and 2). It is noteworthy that the addition of 1,4-cyclohexadiene increased the total yield of **2a** and **3a** to 85–86% as shown in Entries 4 and 5. Without the phenol derivatives, the reaction proceeded very slowly and the products yield was low (Entries 3 and 6). Thus, 1,4-hydroquinone and 1,5-naphthalenediol have not only a significant effect on the isomerization of **1a** but also on the acceleration of the cyclization reaction.

Table 1. Cyclization reaction of *O*-acetyloxime **1a**


Entry	Additive 1 (5 mol%)	Additive 2 (1000 mol%)	Time / h	Yield / %	
				2a	3a
1	1,4-hydroquinone	none	6	22	44
2	1,5-naphthalenediol	none	6	23	43
3	none	none	24	19	20
4	1,4-hydroquinone	1,4-cyclohexadiene	6	32	53
5	1,5-naphthalenediol	1,4-cyclohexadiene	6	34	52
6	none	1,4-cyclohexadiene	24	12	29

The formation of the unexpected product **3a** could not be explained by the SN2-type mechanism, but was considered to be generated from radical intermediate.^{4,5} In fact, by the reaction in toluene-*d*₈ without 1,4-cyclohexadiene, the cyclic imine **3a** was obtained in 17% yield containing 65% deuterium in the methyl group along with 26% yield of **2a**.



A plausible mechanism of the formation of **2a** and **3a** is depicted in Scheme 1. Imine **2a** is caused by the nucleophilic substitution reaction. The olefinic moiety of **1a** intramolecularly attacks on the nitrogen of *O*-acetyloxime **1a**, and imine **2a** is generated by trapping the cationic intermediate **A** with acetic acid. For the formation of **3a**, one-electron transfer occurs from 1,5-naphthalenediol to **1a**.^{5c,f,g,7} The resulting anion radical then cyclizes to generate alkyl radical **B**,^{5g} which captures hydrogen from 1,4-cyclohexadiene. The catalyst, 1,5-naphthalenediol, is regenerated from hydroxynaphthalenoxyl radical by hydrogen abstraction from 1,4-cyclohexadiene.

The reaction of various *O*-acetyloximes was examined and the results are shown in Table 2. By the reaction of *O*-



Scheme 1.


Table 2. Preparation of imines **2**, **3** from *O*-acetyloxime **1**^a

Entry	R ¹	R ²	R ³	Time / h	Yield / %
1	PhCH ₂ CH ₂ ^b	H	H	6	2a 34 3a 52
2	PhCH ₂ CH ₂ ^b	Me	Me	6	2b 72 3b 5
3	PhCH ₂ CH ₂ ^c	H	CN	12	2c 0 3c 69
4	PhCH ₂ CH ₂ ^b	H	CO ₂ Et	12	2d 0 3d 72
5	Ph ^d	H	H	8	2e 0 3e 75
6	Ph(CH ₂) ₃ O ₂ C ^{d,e}	H	H	6	2f 0 3f 61

^aCyclization was carried out under the following conditions; 1,5-naphthalenediol (5 mol%), acetic acid (2 molar amounts), 1,4-cyclohexadiene (10 molar amounts). ^b*syn* : *anti* = 1 : 1. ^c*syn* : *anti* = 1 : 2. ^dOnly *syn* isomers were used. ^e*O*-Pivaloyloxime was used instead of *O*-acetyloxime.

acetyloxime having an electron-rich olefinic moiety, acetoxy imine **2b** was yielded as a major product (Entry 2). In contrast, only radical cyclization products **3c** and **3d** were obtained from oximes having electron deficient olefinic moieties (Entries 3 and 4). Radical cyclization product **3e** was obtained exclusively from *O*-acetyloxime of phenyl ketone (Entry 5). Presumably, the steric repulsion between phenyl and acetoxy group give rise to preferential formation of the *syn*-isomer, which prevents the nucleophilic substitution with the alkenyl group. *O*-Pivaloyloxime⁶ of α -keto ester also cyclized to imino ester **3f** in 61% yield (Entry 6).

Thus the cyclization of *O*-acetyl ketoximes proceeded successfully, whereas *O*-acetyl aldoxime **4** did not cyclize but gave nitrile **5** by the Beckmann fragmentation.



In conclusion, a synthetic method of 3,4-dihydro-2*H*-

pyrroles from γ,δ -unsaturated ketone *O*-acetyloximes was developed with the use of acetic acid and 1,4-cyclohexadiene in the presence of a catalytic amount of 1,5-naphthalenediol. In this reaction, both stereoisomers of oximes can be employed, because *O*-acetyloximes are readily isomerized under the reaction conditions.

This research was supported by a Grant-in-Aid for Scientific Research on Priority Areas (A) "Exploitation of Multi-Element Cyclic Molecules" from the Ministry of Education, Culture, Sports, Science and Technology, Japan.

We dedicate this article to Professor Teruaki Mukaiyama for his 75th birthday.

References and Notes

- a) H. Kusama, Y. Yamashita, and K. Narasaka, *Chem. Lett.*, **1995**, 5. b) H. Kusama, K. Uchiyama, Y. Yamashita, and K. Narasaka, *Chem. Lett.*, **1995**, 715. c) H. Kusama, Y. Yamashita, K. Uchiyama, and K. Narasaka, *Bull. Chem. Soc. Jpn.*, **70**, 965 (1997). d) S. Mori, K. Uchiyama, Y. Hayashi, K. Narasaka, and E. Nakamura, *Chem. Lett.*, **1998**, 111. e) K. Uchiyama, M. Yoshida, Y. Hayashi, and K. Narasaka, *Chem. Lett.*, **1998**, 607. f) M. Yoshida, K. Uchiyama, and K. Narasaka, *Heterocycles*, **52**, 681 (2000).
- In this manuscript, *syn* isomers mean the oximes having leaving group on oxime nitrogen and nucleophile or olefin in the same side of oxime carbon-nitrogen double bond and *anti* isomers mean the other.
- a) J. E. Johnson, N. M. Silk, E. A. Nalley, and M. Arfan, *J. Org. Chem.*, **46**, 546 (1981). b) J. E. Johnson, N. M. Silk, and M. Arfan, *J. Org. Chem.*, **47**, 1958 (1982). c) Y. Kawamura, R. Takayama, M. Nishiuchi, and M. Tsukayama, *Tetrahedron Lett.*, **41**, 8101 (2000).
- For recent review of alkylideneaminy radicals. A. G. Fallis and I. M. Brinza, *Tetrahedron*, **53**, 17543 (1997).
- a) J. Boivin, E. Fouquet, A.-M. Schiano, and S. Z. Zard, *Tetrahedron*, **50**, 1769 (1994). b) J. Boivin, A.-C. Callier-Dublanchet, B. Quiclet-Sire, A.-M. Schiano, and S. Z. Zard, *Tetrahedron*, **51**, 6517 (1995). c) K. Uchiyama, Y. Hayashi, and K. Narasaka, *Chem. Lett.*, **1998**, 1261. d) J. Boivin, A.-M. Schiano, S. Z. Zard, and H. Zhang, *Tetrahedron Lett.*, **40**, 4531 (1999). e) X. Lin, D. Stien, and S. M. Weinreb, *Org. Lett.*, **1**, 637 (1999). f) K. Uchiyama, Y. Hayashi, and K. Narasaka, *Tetrahedron*, **55**, 8915 (1999). g) T. Mikami and K. Narasaka, *Chem. Lett.*, **2000**, 338. h) A. J. McCarroll and J. C. Walton, *J. Chem. Soc., Perkin Trans. 2*, **2000**, 1868.
- Because 3-phenylpropyl 2-oxo-hexenoic acid *O*-acetyloxime was easily hydrolyzed, *O*-pivaloyloxime **1f** was used in stead of the *O*-acetyloxime.
- M. Horspool, G. Hynd, and U. Ixkes, *Tetrahedron Lett.*, **40**, 8295 (1999).
- O*-acetyloxime was prepared in 70–80% yield from ketones^{5f} by treatment with hydroxylamine hydrochloride and pyridine in ethanol, and then triethylamine and acetic anhydride in dichloromethane.