


Direct Cyclopropanation of 1-Alkynylphosphonates by $\text{Cp}_2\text{ZrCl}_2/2\text{EtMgBr}/2\text{AlCl}_3$ To Afford Cyclopropylmethylphosphonates

Abed Al Aziz Quntar and Morris Srebnik*

Department of Medicinal Chemistry and Natural Products, School of Pharmacy,
Hebrew University in Jerusalem, Jerusalem 99120, Israel

msrebn@md.huji.ac.il

Received October 16, 2005



The reagent system $\text{Cp}_2\text{ZrCl}_2/2\text{EtMgBr}/2\text{AlCl}_3$ converts 1-alkynylphosphonates into cyclopropylmethylphosphonates **3** in good isolated yields. Ethers, chlorides, and other cyclopropyl groups are compatible with the reaction conditions. Deuterium labeling is consistent with the formation of stable cyclopropylmethylbimetallic phosphonates by ring contraction of the corresponding aluminacyclopentenylphosphonate. Temperature is crucial; apparently, the cyclopropylmethylbimetallic phosphonates are in equilibrium with the aluminacyclopentenylphosphonates. Low temperature favors the former. We surmise that the negative charges of the intermediate are stabilized by the phosphonate group. Thus, diphenylacetylene and 3-hexyne failed to give cyclopropyl products under the same reaction conditions.

Introduction

The proximity of cyclopropyl and phosphonate groups on the same molecule may endow it with pharmacological properties. For instance, cyclopropylphosphonates can act as *N*-methyl-D-aspartate (NMDA) receptor antagonists,¹ are selective anti-HBV agents,² are insecticides³ and cytostatic agents,⁴ possess antiproliferation properties,⁵ are virostatics,⁶ antidiabetics,⁷ and antitumor agents,⁸ and display antiviral activity.⁹ The methods for synthesizing cyclopropylphosphonates depend on the relative placement of the cyclopropyl and phosphonate groups.¹⁰

Zirconium reagents have been used to prepare cyclopropanes from alkynes.¹¹ Dzhemilev reported that symmetrical alkynes reacted with excess AlEt_3 in the catalytic presence of Cp_2ZrCl_2

to afford aluminacyclopentenones that converted to 1,1-disubstituted cyclopropanes after workup with alkylsulfonates.¹² Szymoniak has used homoallylic ethers and $\text{Cp}_2\text{ZrCl}_2/2\text{n-BuLi}$ to produce substituted cyclopropanes,¹³ and Taguchi has shown that γ,γ -dialkoxyallylic zirconium species react with aldehydes under neutral workup conditions to afford *gem*-dialkoxy cyclopropyl alcohols.¹⁴ Negishi reacted aluminacyclopentenones, obtained from the corresponding alkynes and excess of AlEt_3 in the presence of 10 mol % Cp_2ZrCl_2 , with $\text{BrCH}_2\text{OCH}_3$ to give vinyl cyclopropanes.¹⁵ Negishi also reported that cyclopropanation of enynes could be effected with the reagent system $\text{Et}_2\text{Zn}/0.1\text{XTi}(i\text{-PrO})_3/0.2\text{EtMgBr}/\text{BrCH}_2\text{OCH}_3$ to give fused vinylcyclopropanes;¹⁶ however, simple alkynes did not undergo cyclopropanation. Kulinkovich-type¹⁷ conversions of esters or acyl chlorides to cyclopropanols using zirconocene reagents have been reported.¹⁸

Vinylphosphonates are highly valued products and reagents.¹⁹ Over the past few years we have been investigating the conversion of 1-alkynylphosphonate to highly substituted vinylphosphonates with group IV reagents.²⁰ We have shown that the intermediate metallocycles are very sensitive to the reaction conditions. Various substituted vinylphosphonates can be obtained from the same electrophile by changing both the quantities of reagents required to generate the reactive group IV species and the metal additives.²¹ In this paper we report our results on the reaction of 1-alkynylphosphonates with $\text{Cp}_2\text{ZrCl}_2/2\text{EtMgBr}$ in the presence of 2AlCl_3 . Although our initial aim was to obtain substituted vinylphosphonates, cyclopropylmethylphosphonates

(1) Dappen, M. S.; Pellicciari, J. R.; Natalini, B.; Monahan, J. B.; Chiorri, C.; Cordis, A. A. *J. Med. Chem.* **1991**, *34*, 161–168.

(2) Choi, J.-R.; Cho, D.-G.; Roh, K. Y.; Hwang, J.-T.; Ahn, S.; Jang, H. S.; Cho, W.-Y.; Kim, K. W.; Cho, Y.-G.; Kim, J.; Kim, Y. Z. *J. Med. Chem.* **2004**, *47*, 2864–2869.

(3) Reid, J. R.; Marmor, R. S. *J. Org. Chem.* **1978**, *43*, 999–1001.

(4) Zidek, Z.; Potmesil, P.; Holy, A. *Toxicol. Appl. Pharmacol.* **2003**, *192*, 246–253.

(5) Hatse, S.; Naesens, L.; De Clercq, E.; Balzarini, J. *Biochem. Pharmacol.* **1999**, *58*, 311–323.

(6) Zidek, Z.; Potmesil, P.; Kmoniekova, E.; Holy, A. *Eur. J. Pharmacol.* **2003**, *475*, 149–59.

(7) Van Poelje, P. D.; Erion, M. D.; Fujiwara, T. *PCT Int. Appl.*, 2002.

(8) Naesens, L.; Hatse, S.; Segers, C.; Verbeken, E.; De Clercq, E.; Waer, M.; Balzarini, J. *Oncol. Res.* **1999**, *11*, 195–203.


(9) Hockova, D.; Holy, A.; Masojidkova, M.; Andrei, G.; Snoeck, R. T.; De Clercq, E. *J. Med. Chem.* **2003**, *46*, 5064–5073.

were formed instead. While $\text{Cp}_2\text{ZrCl}_2/2\text{EtMgBr}$ is a known reagent,²² $\text{Cp}_2\text{ZrCl}_2/2\text{EtMgBr}/2\text{AlCl}_3$ has been recently reported.²³


(10) Cyclopropanation of unsaturated phosphonates by copper-catalyzed decomposition of ethyl diazoacetate: (a) Seyferth, D.; Marmor, R.; Hilbert, P. *J. Org. Chem.* **1971**, *36*, 1379–1386. (b) Tsuji, T.; Nishida, S. In *The Chemistry of the Cyclopropyl Group*; Rappoport, Z., Ed.; John Wiley and Sons Ltd.: Chichester, 1987; Chapter 7, pp 330–333. (c) Paul-Roth, C.; De Montigny, F.; Rethore, G.; Simonneaux, G.; Gulea, M.; Masson, S. *J. Mol. Catal. A: Chem.* **2003**, *201*, 79–91. (d) Simonneaux, G.; De Montigny, F.; Paul-Roth, C.; Gulea, M.; Masson, S. *Tetrahedron Lett.* **2002**, *43*, 3685–3687. (e) Lewis, R. T.; Motherwell, W. B. *Tetrahedron Lett.* **1988**, *29*, 5033–5036. (f) Lewis, R. T.; Motherwell, W. B.; Shipman, M.; Slawin, A. M. Z.; Williams, D. J. *Tetrahedron* **1995**, *51*, 3289–3302. Arbuzov reaction of 1-methyl-2,2-dichlorocyclopropylcarbonyl chlorides with P(OR)_3 : Mitrasov, Yu. N.; Simakova, E. A.; Antonova, I. I.; Krasnova, A. A.; Alekseeva, O. O.; Skvortsov, V. G. *Russ. J. Gen. Chem.* **2004**, *74*, 637–638 (CAN 141:243627 AN 2004:485032). Rearrangement of epoxyphosphonates: Kuznetsova, N. G.; Amori, L.; Pellicciari, R. *Ufim. Russ. Bashkirkii Khim. Zh.* **2002**, *9*, 40–41 (CAN 139:149676 AN 2003:120994). Amination/phosphonation of alkylcyclopropanone acetals to give the corresponding aminophosphonates: Tesson, N.; Dorigneux, B.; Fadel. *Tetrahedron: Asymmetry* **2002**, *13*, 2267–2276. Reaction of cobalt(0) or magnesium on α -(halomethyl)phosphonates and c -PrCOOR to give c -PrCOCH₂P(O)(OEt)₂: Orsini, F.; Di Teodoro, E.; Ferrari, M. *Synthesis* **2002**, 1683–1688. Michael induced ring closure (MIRC) of trialkyl phosphites with the corresponding α -bromoalkylidene cyanoacetates and malonates: Stevens, C. V.; Van Heecker, G.; Barbero, C.; Patora, K.; De Kimpe, N.; Verhe, R. *Synlett* **2002**, 1089–1092. 1,3-Dipolar cycloaddition of diazomethane to 1,1-difluoroalkylphosphonates followed by photolysis: Yokomatsu, T.; Suemune, K.; Murano, T.; Shibuya, S. *Heterocycles* **2002**, *56*, 273–282. 1,3-Dipolar addition of diazomethane to 1-acetoxyphosphonates followed by thermolysis: Chen, R.; Liu, Z.; Chen, L. *Gaodeng Xuexiao Huaxue Xuebao* **1988**, *9*, 295–297 (CAN 110:75647 AN 1989:75647). Reaction of phosphorane or arsonane with diisopropyl (*E*)-(3,3,3-trifluoroprop-1-en-1-yl)phosphonate to give trifluoromethylated cyclopropylphosphonates: Shen, Y.; Qi, M. *J. Chem. Res. Synop.* **1996**, 328–329. Reaction of cyclopropanes with Et_2PCl followed by H_2S and Arbuzov reaction with $(\text{PrO})_3\text{P}$: Kormachev, V. V.; Bolesov, I. G.; Ishmuratov, A. S. *Zh. Obshch. Khim.* **1987**, *57*, 704–7055 (CAN 108:21972 AN 1988:21972). Lithiation of $\text{ClCH}_2\text{P(O)(OEt)}_2$ and reaction with RR_1CO to give epoxyalkylphosphonates: Coutrot, P.; Savignac, P. *Synthesis* **1978**, 34–36. Intramolecular cyclization of 2-allylbenzoylphosphonates with trimethyl phosphite via carbene intermediates: Griffiths, D. V.; Harris, J. E.; Karim, K.; Whitehead, B. J. *ARKIVOC* **2000**, *1*, <http://www.arkat.org/arkat/journal/Issue3/onweb17/gj17.htm>. Double olefination of 1,1-bis phosphonates to give 4-alkyl-1,3-butadienylphosphonates: Shen, Y.; Li, P.; Ni, J.; Sun, J. *J. Org. Chem.* **1998**, *63*, 9396–9398. Peterson olefination of the reaction of aldehydes and ketones to α -lithiated- α -fluoro- α -trimethylsilylmethylphosphonate: Waschbusch, R.; Carran, J.; Savignac, P. *Tetrahedron* **1996**, *52*, 14199–14216. Reaction of $\text{RR}_1\text{C:CH}_2$ with $(\text{MeO})_2\text{P(O)H}$ to give $\text{RR}_1\text{CHCH}_2\text{P(O)(OMe)}_2$: Kormachev, V. V.; Mitrasov, Y. N.; Anisimova, E. A.; Kolyamshin, O. A. *Zh. Obshch. Khim.* **1992**, *62*, 1428–29 (CAN 118:102083 AN 1993:102083). Bis-alkylation of α -functional methylphosphonates with dibromoalkanes in the presence of base: Nasser, J.; About-Jaudet, E.; Collignon, N. *Phosphorus, Sulfur Silicon Relat. Elem.* **1990**, *54*, 171–179. Addition of organolithium reagents to di-Et 1-(trimethylsilyl)vinylphosphonate followed by quenching with acid chlorides and alkyl isocyanates: Hong, S.; Chang, K.; Ku, B.; Oh, D. Y. *Tetrahedron Lett.* **1989**, *30*, 3307–3308. Various transformations of existing cyclopropylphosphonates: (g) from 1-hydroxyiminophosphonates and their reduction with NaBH_4 in the presence of transition metal compounds to give 1-aminophosphonates: Demir, A. S.; Tanyeli, C.; Sesenoglu, O.; Demic, S.; Evin, O. O. *Tetrahedron Lett.* **1996**, *37*, 407–10. (h) by formylation of 1,1-dialkylphosphonates: Teulade, M. P.; Savignac, P. *Synth. Commun.* **1987**, *17*, 125–136. (i) by reductive amination of β -ketophosphonates: Varlet, J. M.; Collignon, N.; Savignac, P. *Tetrahedron* **1981**, *37*, 3713–3721. (j) (*E*)-Vinylphosphonates by indium-promoted borohydride reduction of alkynylphosphonates: Wang, C.; Pan, Y.; Yang, D. *China J. Organomet. Chem.* **2005**, *690*, 1705–1709 (CAN 142:482099 AN 2005:280585). (k) Phosphaisocoumarins by Cu(I)-catalyzed intramolecular cyclization of *o*-ethynylphenylphosphonic acid monoethyl esters: Peng, A.-Y.; Ding, Y.-X. *J. Am. Chem. Soc.* **2003**, *125*, 15006–15007. (l) Cyclopropane nucleotide analogues by reduction of cyclopropylalkanones possessing a difluoromethylenephosphonate group at the ring: Yokomatsu, T.; Yamagishi, T.; Suemune, K.; Abe, H.; Kihara, T.; Soeda, S.; Shimeno, H.; Shibuya, S. *Tetrahedron* **2000**, *56*, 7099–7108. (m) Addition of cyclopropylphosphonates to aldehydes: Hirao, T.; Nakamura, T.; Hagihara, M.; Agawa, T. *J. Org. Chem.* **1985**, *50*, 5860–5862.

Results and Discussion

The zirconacyclopentenylphosphonate **2b** was obtained in >97% (based on GCMS and NMR analysis) from $\text{Cp}_2\text{ZrCl}_2/2\text{EtMgBr}$. Intermediates **2** are thermally stable at room temperature and smoothly insert various electrophiles.²⁴ Surprisingly, during our present effort to insert other functional groups to **2b**, we found that addition of 2 equiv of AlMe_3 or AlEt_3 led to the exclusive formation of cyclopropylmethyl phosphate **3b** in 65% yield (based on ^{31}P NMR). Optimization of the reaction conditions by using AlCl_3 rather than AlR_3 improved the yield of **3b** to 95% (eq 1).



The temperature is crucial for the course of the reaction. We found that AlCl_3 must be added to **2b** at -30 to -40 °C, after which the reaction is maintained at the same temperature for 1 h and is quenched with dilute HCl . Adding AlCl_3 at a higher temperature or allowing the temperature to rise above -30 °C resulted in the formation of predominantly ethylated vinylphosphonates (eq 2). Also, 2 equiv of AlCl_3 is essential for optimum



yield; using less than 2 equiv of AlCl_3 substantially decreased the yield of **3b**. The products **3** were extracted, separated on

- (11) For a review of cyclopropylmethylmetal reagents, see: *Houben-Weyl, Methods of Organic Chemistry*; de Meijere, A., Ed.; Thieme: Stuttgart, 1997; Vol. E17, p 2686 and references therein.
- (12) (a) Dzhemilev, U. M.; Ibragimov, A. G.; Khafizova, L. O.; Ramazanov, I. R.; Yalalova, D. F.; Tolstikov, G. A. *J. Organomet. Chem.* **2001**, *636*, 76–81. (b) A Review: Dzhemilev, U. M.; Ibragimov, A. G. *Russ. Chem. Bull.* **1998**, *47*, 786–794.
- (13) Gandon, V.; Laroche, C.; Szymoniak, J. *Tetrahedron Lett.* **2003**, *44*, 4827–4829.
- (14) (a) Ito, H.; Kuroi, H.; Ding, H.; Taguchi, T. *J. Am. Chem. Soc.* **1998**, *120*, 6623–6624. (b) Ito, H.; Sato, A.; Kusanagi, T.; Taguchi, T. *Tetrahedron Lett.* **1999**, *40*, 3397–3398.
- (15) Negishi, E.; Montchamp, J. L.; Anastasia, L.; Elizarov, A.; Choueiry, D.-le. *Tetrahedron Lett.* **1998**, *39*, 2503–2506.
- (16) Montchamp, J.-L.; Negishi, E. *J. Am. Chem. Soc.* **1998**, *120*, 5345–5346.
- (17) Kulinkovich, O. G.; de Meijere, A. *Chem. Rev.* **2000**, *100*, 2789–2834.
- (18) (a) Fujita, K.; Yorimitsu, H.; Shinokubo, H.; Oshima, K. *J. Am. Chem. Soc.* **2004**, *126*, 6776–6783. (b) Gandon, V.; Bertus, P.; Szymoniak, J. *Eur. J. Org. Chem.* **2000**, 3713–3719.
- (19) Dembitsky, V. M.; Aziz Quntar, A. A.; Haj-Yehia, A.; Srebnik, M. *Mini Rev. Org. Chem.* **2005**, *2*, 343–357.
- (20) Aziz Quntar, A. A.; Srebnik, M. *J. Organomet. Chem.* **2005**, *690*, 2504–2514.
- (21) Aziz Quntar, A. A.; Srebnik, M. *Chem. Commun.* **2003**, 58–59.
- (22) (a) Negishi, E.; Takahashi, T. *Bull. Chem. Soc. Jpn.* **1998**, *71*, 755. (b) Negishi, E.; Takahashi, T. *Acc. Chem. Res.* **1994**, *27*, 124–130.
- (23) Xi, Li, Z. *Top. Organomet. Chem.* **2004**, *10*, 27–56.
- (24) Ben-Valid, S.; Aziz Quntar, A. A.; Srebnik, M. *J. Org. Chem.* **2005**, *70*, 3554–3559.

TABLE 1. Formation of Cyclopropylmethylphosphonates, 3

Entry	3	Conversion ^a /Yield ^b
3a		95/75
3b		95/81
3c		95/72
3d		95/72
3e		95/73
3f		95/68
3g		80/57
3h		95/71
3i		95/68


^a Determined by ³¹P NMR. ^b Isolated yield after chromatography.

silica gel chromatography, and analyzed by NMR spectroscopy, GCMS, and elemental analysis. For instance, in the ¹H NMR spectrum of **3b**, the two multiplets in the region 0.3–0.5 ppm, and the ¹³C peaks in ¹³C NMR spectrum in the region ~12 ppm are indicative of a cyclopropyl structure. The doublet in the region 1.43–1.77 ppm (²J_{PH} = 14.7–18.0 Hz) corresponds to the methylene hydrogens α to phosphorus. Furthermore, ³¹P NMR of **3b** in the region 25.9–31.9 ppm is consistent with phosphorus attached to sp³ carbon rather than to a sp² or sp carbon. The reaction is sensitive to the R group of the 1-alkynylphosphonate. When R was phenyl, no cyclopropane product was observed, but rather the respective ethylated vinyl phosphonate was obtained (eq 2).


The ester moiety affects the rate of the reaction. The diphenyl phosphonate required overnight stirring to achieve an 85% yield of the zirconacyclopentenediphenyl phosphonate (**3f** intermediate), apparently due to steric effect. In addition, the present cyclopropanation procedure seems to be restricted to aluminum reagents. Other metals (Ni, Cu, Zn, Pt, Ce, Pd etc) were unsuccessful. Various substituents are tolerated by the reaction conditions, such as ethers (Table 1, **3d**) or chlorides (Table 1, **3g**). Another cyclopropyl group may also be incorporated (Table 1, **3h**). The conversion, based on starting 1-alkynylphosphonate is essentially complete (³¹P NMR). Isolated yields after silica gel chromatography are good. These novel cyclopropylmethylphosphonates are thermally and air stable compounds and are soluble in most solvents. Results are summarized in Table 1.

Mechanistic Insights. Both Negishi¹⁵ and Dzhemilev¹² have reported cyclopropanation reactions of alkynes with group IV metals. Negishi has investigated the mechanism of formation and subsequent reactions of aluminacyclopentenes derived from simple alkynes and 3AlEt₃/Cp₂ZrCl₂ 10 mol %.²⁵ XCH₂OCH₃ is required to convert aluminacyclopentenes to vinylcyclopropanes. Dzhemilev, using an almost identical reagent system as

SCHEME 1



that of Negishi, 3AlEt₃/Cp₂ZrCl₂ 10 mol %, reported that workup with dialkylsulfonates of aluminacyclopentenes derived from simple alkynes gave *gem*-dialkylcyclopropanes.¹² In the absence of this workup procedure, cyclopropanation did not occur. Workup with tosylates afforded diethylated alkenes. Clearly, the system, 3AlEt₃/Cp₂ZrCl₂ 10 mol % is capable of reacting differently by slight modification of the reaction conditions. Using the Negishi/Dzhemilev protocol with 1-alkynylphosphonates, no cyclopropanation products were observed and only the starting material was recovered. Apparently, the phosphonate oxygens coordinate with the AlEt₃ and Cp₂ZrCl₂. Indeed when hex-1-ynylphosphonate was stirred with 1 equiv of AlCl₃ in CDCl₃, an upfield shift in the ³¹P NMR was observed from ca. δ -4 to ca. δ -8 ppm, which is indicative of Lewis acid complexation with the phosphonate oxygens. Thus, it is necessary to form the much more reactive zirconacyclopentene, which subsequently can be transmetalated with AlCl₃ to afford the aluminacyclopentenes. Even with preformed zirconacyclopentenes **2**, 2 equiv of AlCl₃ are required. One equivalent reacts with **2** to form the aluminacyclopentenylphosphonate, and we conjecture that the other equivalent remains coordinated to the phosphonate oxygens. The aluminacyclopentenylphosphonate apparently is in equilibrium with the ring-contracted cyclopropylmethylbimetallic phosphonate. Low temperature favors the latter. Deuteriolysis of the reaction mixture at -30 °C gave a 2 [2H] species, which is consistent with formation of a cyclopropylmethylbimetallic phosphonate prior to workup (Scheme 1). Hydrolysis of the same reaction mixture at 25 °C led to the ethylated product. That low temperature favors the cyclopropylmethylbimetallic species must be due to the phosphonate group's ability to stabilize the adjacent negative charges. To verify this observation, diphenyl acetylene and 3-hexyne were reacted under the same conditions as for the 1-alkynylphosphonates (eq 3). No cyclopropanation occurred under our



(25) Negishi, E.; Kondakov, D. Y.; Choueiry, D.; Kasai, K.; Takahishi, T. *J. Am. Chem. Soc.* **1996**, *118*, 9577–9588.

standard conditions. Only the ethylated products could be detected.

These results support the notion that the phosphonate group must stabilize the negative charge of the intermediate cyclopropylmethylbimetallic phosphonates, whereas alkyl and phenyl groups do not. Our method is unique in that it gives cyclopropylmethyl compounds directly without requiring additional alkylating reagents.

Conclusion

We have described a new utility for the reagent system $\text{Cp}_2\text{-ZrCl}_2/2\text{EtMgBr}/2\text{AlCl}_3$ that can transform 1-alkynylphosphonates to cyclopropylmethylphosphonates in good isolated yields. Temperature is crucial to the success of the reaction. We have found that -30 to -40 °C is optimal for generating **3**. Working up the same reaction mixture at 25 °C results only in the formation of ethylated vinylphosphonates. Apparently there is equilibrium between the aluminacyclopentenylphosphonates and the cyclopropylbimetallic phosphonates. Low temperature favors the latter. This must be due to the presence of the phosphonate group, which can stabilize the adjacent negative charges of the cyclopropylbimetallic phosphonates. Though diphenylacetylene and 3-hexyne failed to give cyclopropylmethyl derivatives under the same reaction conditions, it was successful for 1-alkynylphosphonates. The reaction is compatible with ethers, chlorides, and other cyclopropyl groups. We conjecture that other EWGs (electron-withdrawing groups) stable under the conditions of the reaction should also provide the corresponding cyclopropylmethyl derivatives.

Experimental Section

General Procedure for the Synthesis of 3b. To 0.306 g (1.05 mmol) of zirconocene dichloride dissolved in 7 mL of dry THF at

-78 °C was added 1.05 mL of 2 M EtMgBr (2.1 mmol) dropwise in a 25 mL round-bottom flask. After 5 min of stirring at -78 °C, 1 mmol (0.22 g) of diethyl hex-1-ynylphosphonate was added. The reaction was gradually warmed to 25 °C and stirred for 2 h. The reaction was cooled to -30 °C and maintained at -30 – -40 °C while 2 equiv of AlCl_3 was added. The reaction was stirred for 1 h at -30 °C and was worked up with dilute HCl. The product was extracted with diethyl ether (2×15 mL), separated on silica gel column (80% petroleum ether/20% ethyl acetate), and analyzed by GCMS, elemental analysis, and NMR spectroscopy.

Diethyl (1-Butylcyclopropyl)methylphosphonate (3b). ^1H NMR (300 MHz): δ 0.35 (m, 2H), 0.42 (m, 2H), 0.86 (t, 3H, $J_{\text{HH}} = 6.6$ Hz), 1.30 (dt, 6H, $J_{\text{HH}} = 6.9$, $^4J_{\text{PH}} = 0.3$ Hz), 1.30–1.45 (overlap, 6H), 1.75 (d, 2H, $^2J_{\text{PH}} = 18.0$ Hz), 4.00–4.18 (m, 4H). ^{31}P NMR (121.4 MHz): δ 31.89. ^{13}C NMR (75.5 MHz): δ 12.7, 12.8, 14.1, 15.0 (d, $^2J_{\text{PC}} = 4.3$ Hz), 16.4 (d, $^3J_{\text{PC}} = 6.0$ Hz), 22.8, 28.7, 32.3 (d, $^1J_{\text{PC}} = 140.1$ Hz), 36.7, 61.3 (d, $^2J_{\text{PC}} = 6.6$ Hz). MS m/z : 248 (28.4), 233 (16.7), 219 (66.7), 205 (25.5), 191 (29.5), 163 (49.8), 149 (50.0), 138 (44.1), 125 (35.3), 111 (100), 97 (43.1), 81 (82.4), 67 (47.1), 57 (43.1), 41 (71.6), 29 (48.0). Anal. Calcd for $\text{C}_{12}\text{H}_{25}\text{O}_3\text{P}$: C, 58.05; H, 10.15; P, 12.47. Found: C, 58.24; H, 10.21; P, 12.52.

Acknowledgment. We thank the Israel Science Foundation (grant 138/01) for supporting this work A.A.Q. thanks the Israel Ministry of Science for providing financial support.

Supporting Information Available: General experimental procedures; ^1H , ^{13}C , and ^{31}P spectra; GCMS spectra; and elemental analysis for compounds **3**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JO052157L