

SYNTHESIS OF PHENAZINE 5,10-DIOXIDES FROM BENZOFUROXAN CATALYZED BY MOLECULAR SIEVES

Tohru Takabatake, Tomoyuki Miyazawa, Mahiro Kojo,
and Minoru Hasegawa*

College of Pharmacy, Nihon University, 7-7-1 Narashinodai,
Funabashi-shi, Chiba 274-8555, Japan

Abstract - The synthesis of substituted phenazines was carried out. Phenazine 5,10-dioxides (**3**) were obtained from benzofuroxan (**1**) with dihydroxybenzene derivatives (**2**) catalyzed by molecular sieves at room temperature.

INTRODUCTION

Benzofuroxan (**1**) has been shown to have numerous pharmacological and industrial applications.^{1a-c} Reactions of benzofuroxan with phenolic compounds in basic medium provide the corresponding phenazine 5,10-dioxide derivatives.² Certain phenazine derivatives are known to have antibacterial activity and two of which, iodinin and myxin, are microbial metabolites.^{3, 4} As a part of benzofurazan chemistry, reactions of benzofuroxans with active methylene compounds lead to the corresponding quinoxaline 1,4-dioxides catalyzed by silica gel⁵ or molecular sieves^{6, 7} and the antibacterial activity of quinoxaline 1,4-dioxides has been reported.⁸ Pyrido[2,3-*b*]pyrazine 1,4-dioxides and pyrido[2,3-*b*]pyrazine 1-oxides have been obtained from pyrido[2,3-*c*]furoxan catalyzed

by silica gel, alumina, or molecular sieves⁹ and the antibacterial activity of pyrido[2,3-*b*]pyrazine 1,4-dioxides and pyrido[2,3-*b*]pyrazine 1-oxides has been reported.¹⁰ In this study, phenazine 5,10-dioxide derivatives were obtained from benzofuroxan and dihydroxybenzene derivatives catalyzed by molecular sieves or silica gel or alumina at room temperature.

RESULTS AND DISCUSSION


Phenazine 5,10-dioxide derivatives were synthesized as follows. A solution of compound (**1**) and 1,4-dihydroxybenzene derivatives (**2**) in methanol was evaporated in the presence of molecular sieves 4A, aluminum oxide, or silica gel. Both starting materials were adsorbed on the molecular sieves or other catalysts followed by standing at room temperature, 40 °C or 90 °C. The reaction mixture was chromatographed on silica gel to give the corresponding phenazine 5,10-dioxide derivatives (**3**).

Table 1

Catalysts	3a ¹¹ (%)	1 (% , recovery)	Reaction condition	
Molecular sieves 4A	87	0	rt, 2 h	
Aluminum oxide, basic	37	0	90 °C, 1 h	
	56	8	40 °C, 2 h	
	acidic	21	7	40 °C, 2 h
	neutral	29	0	40 °C, 2 h
Silica gel C-200	16	39	90 °C, 2 h	
None	0	97	rt, 2 h	

Various catalysts and temperature in the reaction of compound (1) with 1,4-dihydroxybenzene (2a) were examined (Table 1). In using catalysts, reaction efficacy varied considerably with the catalyst used. Phenazine 5,10-dioxide derivatives were obtained in good yield using molecular sieves 4A as catalyst at room temperature. The most suitable catalyst was molecular sieves 4A.

Table 2




Compound No.	Yield (%) of Phenazine
2a	87 (3a)
4	0 (3a, 6)
	1.1* (3a)
5	0 (6)
	0* (6)

* 90 °C, 1 h

to good yield (See Table 4). Phenazines (**3a~3f**), the assignments were made based on the following; When $^1\text{H-NMR}$ signals of phenazines were compared, $^1\text{H-NMR}$ spectrum of compound (**3d**) showed two doublet signals (δ 7.45 and 8.91 ppm) and the coupling constant was 10.27 Hz. These signals may possible consist of protons 3 and 4. Phenazines including the electron-donating group have two singlet signals consisting of protons 1 and 4.

Table 4

R_2 = electron-withdrawing group



Compound No.	R_2	Yield (%)	1 (Recovery)
3d	CHO	52	29
3e	COCH ₃	26	44
3f	COCH ₂ CH ₃	34	47
3g	COOH	0	90

The ethyl 2-hydroxyphenazine-1-yl ketone 5,10-dioxide (**3f**) has interesting $^1\text{H-NMR}$ spectrum which showed two quartets at δ 2.63 and 2.99 ppm with each 1H integration for protons at the methylene of **3f** at 25 °C. The methylene group may thus rotate slowly enough to show different chemical shifts of two kinds of protons. The temperature-

dependent ^1H -NMR spectra for compound (**3f**) are shown in Figure 1. The two quartet signals broadened slightly at 80 °C and then changed to one signal (δ 2.82 ppm) at 140 °C.





Figure 1

Scheme 1 shows the positions of protons at the methylene of **3f** to be close to oxygen at *N*-oxide, and consequently oxygen affects the rotation of the two protons at methylene. The oxygen may possibly function as a barrier against conformation of methylene and so methylene group rotation is slow enough that different chemical shifts are apparent. Molecular sieves 4A or other catalysts would thus appear to affect dehydration in the reaction mechanism and the reaction proceed not in but on molecular sieves in that the



3f


Scheme 1

pore diameter of molecular sieves 4A is approximately 0.4 nm. Windows to the cages are too small to allow the organic molecules access to cages.

The enol form of carbonyl compounds was previously shown necessary for the formation of quinoxalines or pyridopyrazines, whose yields depends on enol content in 1,3-diketones.^{7,9} Molecular sieves or other catalysts may serve to enhance the stability of the enol form of carbonyl compound and the dehydration capacity of catalysts may determine significantly the possibility of synthesis.

Molecular sieves may have similar effect as electron-withdrawing groups on dihydroxybenzenes (see Scheme 2). Electron-withdrawing groups and molecular sieves surface force dihydroxybenzenes to couple at the ortho position followed by cyclization and elimination of water to yield phenazines. The different attacking positions in the reaction on molecular sieves may lead to different structures of phenazines as products. In the case of reaction with 2,5-dihydroxybenzoic acid, no product was obtained, possibly owing to steric hindrance.

The present method is simple and efficient for the preparation of phenazine 5,10-dioxide derivatives under mild conditions.



Scheme 2

EXPERIMENTAL

Melting points were determined with a Yanagimoto micromelting point apparatus and are uncorrected. The IR spectra were recorded on a JASCO IR-810 spectrophotometer. The ¹H-NMR spectra were recorded on a JNM-GSX 400 FT NMR System with TMS as the internal standard. The MS spectra were recorded on a Hitachi M-2000 and JEOL JMS-GCmate spectrometers with an electron beam energy of 70 eV. Microanalysis was

performed at the microanalytical laboratory of the Center for Instrumental Analysis in College of Science & Technology, Nihon University.

General Procedure (Table 1).

To a solution of **1** (34.5 mg, 0.25 mmol) and dihydroxybenzene (**2a**) (30.3 mg, 0.275 mmol) in methanol (5 mL) was added silica gel [Wako Pure Chemical Industries, Wakogel C-200, 1 g] or aluminium oxide [Merck, Aluminium oxide 90 active basic for column chromatography (Art. 1076), Aluminium oxide 90 active neutral for column chromatography (Art. 1077), Aluminium oxide 90 active acidic for column chromatography (Art. 1078), 1 g] or molecular sieves [4A powder, Union Showa, 1 g]. The mixture was evaporated in an evaporator at 30 °C. The adsorbent containing the adsorbed reagents was allowed to stand for 2 h at rt and then introduced onto the silica gel column (Wako Pure Chemical Industries, Wakogel C-200). The product 2-Hydroxyphenazine 5,10-dioxide (**3a**) was eluted with dichloromethane/methanol (95:5) and purified by preparative TLC (Merck, Silica gel plate 60 F₂₅₄ Art. 5717) with dichloromethane/methanol (95:5). Yield 49.8 mg (87 %).

2-Hydroxyphenazine 5,10-dioxide (**3a**)

Deep red powder, mp 243-245 °C(decomp) (lit.,¹¹ mp 234 °C(decomp)). IR (KBr) cm⁻¹: 1615, 1595, 1353, 1238, 1078, 837, 762. ¹H-NMR (DMSO-d₆) δ : 7.49 (dd, 1H, *J*_{1,3} = 2.2 Hz, *J*_{3,4} = 9.5 Hz, C3-H), 7.77 (d, 1H, *J*_{1,3} = 2.2 Hz, C1-H), 7.84-7.93 (m, 2H, C7-H, C8-H), 8.48 (d, 1H, *J*_{3,4} = 9.5 Hz, C4-H), 8.52-8.56 (m, 2H, C6-H, C9-H), 11.33 (br s, 1H, OH, deuterium oxide-exchangeable). EI-MS *m/z*: 228.0502 (Calcd for C₁₂H₈N₂O₃: 228.0534). **3a** was characterized by comparison of IR spectra with an authentic sample.¹¹

2-Hydroxy-3-methylphenazine 5,10-dioxide (**3b**)

Deep red powder, mp 223-225 °C. IR (KBr) cm⁻¹: 1620, 1457, 1346, 1254, 1076, 830, 760. ¹H-NMR (DMSO-d₆) δ : 2.41 (s, 3H, CH₃), 7.78 (s, 1H, C1-H), 7.78-7.90 (m, 2H, C7-H, C8-H), 8.33 (s, 1H, C4-H), 8.46-8.55 (m, 2H, C6-H, C9-H), 11.28 (br s, 1H, OH, deuterium oxide-exchangeable). EI-MS *m/z*: 242.0672 (Calcd for C₁₃H₁₀N₂O₃ :

242.0690). *Anal.* Calcd for C₁₃H₁₀N₂O₃: C, 64.46; H, 4.16; N, 11.56. Found: C, 64.46; H, 4.19; N, 11.57.

2-Hydroxy-3-methoxyphenazine 5,10-dioxide (**3c**)

Deep red powder, mp 248-250 °C. IR (KBr) cm⁻¹: 1620, 1524, 1470, 1347, 1223, 1185, 1082, 852, 763. ¹H-NMR (DMSO-d₆) δ : 4.06 (s, 3H, OCH₃), 7.80 (s, 1H, C1-H), 7.81 (s, 1H, C4-H), 7.84-7.89 (m, 2H, C7-H, C8-H), 8.52-8.57 (m, 2H, C6-H, C9-H), 11.47 (br s, 1H, OH, deuterium oxide-exchangeable). EI-MS m/z: 258.0639 (Calcd for C₁₃H₁₀N₂O₄ : 258.0639). *Anal.* Calcd for C₁₃H₁₀N₂O₄: C, 60.47; H, 3.90; N, 10.85. Found: C, 60.42; H, 4.01; N, 10.80.

2-Hydroxy-1-phenazinecarboxaldehyde 5,10-dioxide (**3d**)

Orange powder, mp 221-224 °C. IR (KBr) cm⁻¹: 1630, 1583, 1472, 1355, 1290, 1170, 1104, 832, 776. ¹H-NMR (DMSO-d₆) δ : 7.45 (d, 1H, *J*_{3,4} = 10.2 Hz, C3-H), 7.85-7.96 (m, 2H, C7-H, C8-H), 8.68-8.77 (m, 2H, C6-H, C9-H), 8.91 (d, 1H, *J*_{3,4} = 10.2 Hz, C4-H), 11.99 (s, 1H, CHO), 15.23 (s, 1H, OH, deuterium oxide-exchangeable). EI-MS m/z: 256.0489 (Calcd for C₁₃H₈N₂O₄ : 256.0483). *Anal.* Calcd for C₁₃H₈N₂O₄: C, 60.94; H, 3.15; N, 10.93. Found: C, 60.85; H, 3.29; N, 10.73.

2-Hydroxyphenazine-1-yl methyl ketone 5,10-dioxide (**3e**)

Deep red powder, mp 182-183 °C (mp. 180-181 °C). ¹¹ IR (KBr) cm⁻¹: 1706, 1555, 1415, 1337, 1278, 1173, 816, 768. ¹H-NMR (DMSO-d₆) δ : 2.56 (s, 3H, CH₃), 7.61 (d, 1H, *J*_{3,4} = 9.5 Hz, C3-H), 7.83-7.88 (m, 2H, C7-H, C8-H), 8.40 (d, 1H, *J*_{3,4} = 9.5 Hz, C4-H), 8.45-8.51 (m, 2H, C6-H, C9-H), 11.45 (s, 1H, OH, deuterium oxide-exchangeable). EI-MS m/z: 270.0686 (Calcd for C₁₄H₁₀N₂O₄ : 270.0640). **3e** was characterized by comparison of IR spectra with an authentic sample. ¹¹

Ethyl 2-hydroxyphenazine-1-yl ketone 5,10-dioxide (**3f**)

Deep red powder, mp 180-182 °C. IR (KBr) cm^{-1} : 1675, 1590, 1555, 1420, 1348, 1285, 1180, 1113, 831, 763. $^1\text{H-NMR}$ (DMSO- d_6) δ : 1.17 (t, 3H, $J = 6.8$ Hz, $\text{CH}_3\text{-CH}_2$), 2.63 (q, 1H, $J = 6.8$ Hz, $\text{CH}_3\text{-CHaHb}$), 2.99 (q, 1H, $J = 6.8$ Hz, $\text{CH}_3\text{-CHaHb}$), 7.56 (d, 1H, $J_{3,4} = 9.7$ Hz, C3-H), 7.87 (m, 1H, C7-H, C8-H), 8.40 (m, 1H, C9-H), 8.50 (d, 1H, $J_{3,4} = 9.7$ Hz, C4-H), 8.52 (m, 1H, C6-H, C9-H), 11.36 (s, 1H, OH, deuterium oxide-exchangeable). EI-MS m/z : 284.0804 (Calcd for $\text{C}_{15}\text{H}_{12}\text{N}_2\text{O}_4$: 284.0796). *Anal.* Calcd for $\text{C}_{15}\text{H}_{12}\text{N}_2\text{O}_4$: C, 63.38; H, 4.25 ; N, 9.85. Found: C, 63.22; H, 4.38; N, 9.86.

REFERENCES AND NOTES

1. For comprehensive reviews see: [a] A. J. Boulton and P. B. Ghosh, 'Advances in Heterocyclic Chemistry: Benzofuroxans.' Vol. 10, ed. by A. R. Katritzky and A. J. Boulton, Academic Press, Inc., London, 1969, pp. 1-41. [b] A. Gasco and A. J. Boulton, 'Advances in Heterocyclic Chemistry: Furoxans and Benzofuroxans.' Vol. 29, ed. by A. R. Katritzky and A. J. Boulton, Academic Press, Inc., London, 1981, pp. 251-340. [c] P. Ghosh, B. Ternai, and M. Whitehouse, 'Medicinal Research Reviews: Benzofurazans and Benzofuroxans.' Vol. 1, John Wiley & Sons, Inc. 1981, pp. 159-187.
2. M. J. Abu El-Haj, B. W. Dominy, J. D. Johnston, M. J. Haddadin, and C. H. Issidorides, *J. Org. Chem.*, 1972, **37**, 589.
3. G. R. Clemo and H. McIlwain, *J. Chem. Soc.*, **1938**, 479.
4. [a] A. E. Peterson, D. C. Gillespie, and F. D. Cook, *Can. J. Microbiol.*, 1966, **12**, 221. [b] M. Weigele and L. Leimgruber, *Tetrahedron Lett.*, **1967**, 715.
5. M. Hasegawa and T. Takabatake, *Synthesis*, **1985**, 938.
6. T. Takabatake and M. Hasegawa, *J. Heterocycl. Chem.*, 1987, **24**, 529.
7. T. Takabatake, Y. Hasegawa, and M. Hasegawa, *J. Heterocycl. Chem.*, 1993, **30**, 1477.

8. T. Takabatake, Y. Takabatake, T. Miyazawa, and M. Hasegawa, *Yakugaku Zasshi*, 1996, **116**, 491.
9. T. Takabatake, T. Miyazawa, and M. Hasegawa, *Heterocycles*, 1997, **45**, 107.
10. T. Miyazawa, T. Takabatake, and M. Hasegawa, *Yakugaku Zasshi*, 1997, **117**, 126.
11. M. L. Edwards, R. E. Bambury, and H. K. Kim, *J. Heterocycl. Chem.*, 1976, **13**, 653.