

Structure-activity relationships and kinetic studies of peptidic antagonists of CBX chromodomains

Jacob I Stuckey, Catherine Simpson, Jacqueline L. Norris-Drouin, Stephanie H Cholensky, Junghyun Lee, Ryan Pasca, Nancy Cheng, Bradley M Dickson, Kenneth H. Pearce, Stephen V. Frye, and Lindsey I. James

J. Med. Chem., **Just Accepted Manuscript** • DOI: 10.1021/acs.jmedchem.6b00801 • Publication Date (Web): 29 Aug 2016

Downloaded from <http://pubs.acs.org> on September 3, 2016

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.

Structure-activity relationships and kinetic studies of peptidic antagonists of CBX chromodomains

Jacob I. Stuckey, Catherine Simpson, Jacqueline L. Norris-Drouin, Stephanie H. Cholensky, Junghyun Lee, Ryan Pasca, Nancy Cheng, Bradley M. Dickson†, Kenneth H. Pearce, Stephen V. Frye, Lindsey I. James**

Center for Integrative Chemical Biology and Drug Discovery, Division of Chemical Biology and Medicinal Chemistry, UNC Eshelman School of Pharmacy, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina 27599, USA.


ABSTRACT. To better understand the contribution of methyl-lysine (Kme) binding proteins to various disease states, we recently developed and reported the discovery of **1** (UNC3866), a chemical probe that targets two families of Kme binding proteins, CBX and CDY chromodomains, with selectivity for CBX4 and -7. The discovery of **1** was enabled in part by the use of molecular dynamics simulations performed with CBX7 and its endogenous substrate. Herein, we describe the design, synthesis and structure-activity relationship studies that led to the development of **1** and provide support for our model of CBX7-ligand recognition by examining the binding kinetics of our antagonists with CBX7 as determined by surface-plasmon resonance.

Introduction

Controlled regulation of gene expression is essential for the proper development and function of multicellular organisms. Cell-type specific gene regulation is achieved by expression of specific transcription factors and through a complex language of chemical modifications to both DNA and histone proteins that are interpreted through the binding of various proteins and protein complexes.¹ These ‘reader’ complexes then modulate the transcriptional outcome by controlling the accessibility of the underlying gene to transcription factors.^{2,3}

One key chemical modification that regulates gene expression is the posttranslational methylation of histone lysine residues.² The lysine ϵ -nitrogen can be mono-, di- or tri-methylated (Kme1, Kme2 or Kme3, respectively). Methyl-lysine (Kme) reader proteins recognize Kme in a manner that is specific to the methylation state of the lysine and often to the sequence surrounding the modified lysine.^{3,4} Kme readers bind methylated-lysine through an aromatic cage that engages the lysine side chain through cation- π and van der Waals’ interactions. The size and shape of the aromatic cage allows the Kme reader to discriminate between different methylation states, while the surrounding protein residues dictate sequence selectivity.³ Previous studies have characterized the ability of various Kme1 and Kme2 reader proteins to accommodate non-natural methyl-lysine analogs in their aromatic cages;⁵⁻⁷ however, little is known about the preference of Kme3 reader proteins for different Kme mimetics. Initial efforts toward the discovery of Kme3 reader antagonists were focused on the development of peptidic inhibitors wherein the key Kme3 residue was maintained and potency was achieved through the variation of surrounding residues.^{8,9}

We recently reported the development of **1** (UNC3866), a peptide-based chemical probe that contains an unnatural diethyl-lysine in lieu of Kme3 and selectively targets two families of Kme3 reader proteins (**Figure 1**).¹⁰ These two families of proteins belong to the chromodomain superfamily of Kme readers that are essential for proper genomic regulation in numerous organisms, spanning fungi, plants and animals.¹¹ Compound **1** targets the Polycomb (Pc) CBX family of chromodomains⁴ and the lesser explored CDY family of chromodomains.¹² In mammals, the Pc family of chromodomains consists of five proteins, CBX2, -4, -6, -7 and -8. These proteins compete with each other for incorporation into Polycomb Repressive Complex 1 (PRC1) where they regulate numerous cellular processes including differentiation, growth and proliferation.¹³⁻¹⁷



Kme Reader Domain	1 (UNC3866)	Interaction with PRC1	UNC3866 ITC K_d (μM)
CBX2			1.8 ± 0.21
CBX4			0.094 ± 0.017
CBX6			0.610 ± 0.0078
CBX7			0.097 ± 0.0024
CBX8			1.2 ± 0.021
CDY1			6.3 ± 0.92
CDYL1b			0.91 ± 0.076
CDYL2			0.85 ± 0.076

Figure 1. Compound 1 and its chromodomain targets. (Top) Structure of **1**, a cell-active peptidic antagonist of CBX and CDY chromodomains. (Bottom) Domain maps of human CBX and CDY chromodomains as annotated in Uniprot. The K_d of compound **1** as determined by ITC for each of the chromodomains is listed.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
Compound **1** is most potent for the chromodomains of CBX4 and -7 ($K_d \sim 100$ nM). It is 6- to 63-fold selective for these chromodomains over the other CBX and CDY chromodomains.¹⁰ Interestingly, our selectivity studies revealed that **1** does not significantly interact ($K_d > 100$ μ M) with the HP1 family of chromodomains (CBX1, -3 and -5),¹⁰ which share a high degree of structural similarity to the Pc family of chromodomains but are not associated with PRC1.⁴ Compound **1** was also highly selective versus >250 other molecular targets including chromatin regulatory proteins and a general pharmacology panel. Our previous studies characterized the effects of **1** in PC3 cells, an advanced stage metastatic prostate cancer model previously shown to have its growth regulated by CBX7.¹⁸ Consistent with CBX7 genetic manipulation, compound **1** is able to inhibit PC3 cell proliferation by inducing a senescent-like state.^{14, 18} These studies suggest that Pc CBX antagonists may one day serve as a novel class of therapeutically useful agents.

33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Herein we report the structure-activity relationship (SAR) studies leading to the development of compound **1**. We explored the effects of various modifications (**Figure 2a**) on the potency and selectivity versus three representative members of the Pc CBX chromodomains (CBX4, -7 and -8), with CBX4 and -7 being the most potent targets of **1**, and CBX8 representing the remaining Pc CBX chromodomains (CBX2, -6, and -8), all of which display somewhat weaker affinity for **1**.¹⁰ Compounds were also tested against one member of the CDY family of chromodomains, CDYL2, and a representative of the HP1 family of chromodomains, CBX5, in order to better understand the molecular determinants for selectivity between these three structurally related families of chromodomains. Additionally, kinetic studies with CBX7 provided key insights into the induced-fit recognition mechanism of these ligands by CBX7.




Figure 2. (a) X-ray crystal structure of compound **1** bound to CBX7 (PDB 5EPJ). Compound **1** (yellow) binds in an extended groove along the surface of the chromodomain of CBX7 (light blue). The interaction involves multiple hydrogen bonds (black dashes) between CBX7 and the backbone amides of **1**. The high affinity of this interaction is driven by hydrophobic contacts between the *t*-butyl group of **1** and the side chains of Leu48, Asp49 and Arg51. The different regions of compound **1** explored in this study are indicated. (b) Structure of compound **2**. The biotin affinity tag was added to the C-terminus of **1** which is solvent exposed, and **2** retains high affinity for CBX7.

Results and Discussion

We first developed an AlphaScreen^{®19} assay for each of the Pc CBX and CDY chromodomains using His-tagged recombinant protein and **2** (UNC4195, **Figure 2b**), a biotinylated analog of **1**, as the bait ligand.¹⁰ It should be noted that our previously reported AlphaScreen[®] studies with compound **1** were performed utilizing a similar assay with a biotinylated H3K9me3 bait ligand and CBX7. The improved potency of **2** for our target chromodomains over the H3K9me3 peptide allows us to reduce the amount of all reagents used

1
2
3 in the assay and also reduce the tight binding limit. For CBX5, which does not appreciably bind
4 to **2**, we again utilized a biotinylated version of the endogenous substrate, H3K9me3⁴ (see
5
6
7
8 **Supplementary Table 1** for specific assay concentrations).
9

10
11 Compound **1** is based on a peptide fragment from the methyltransferase SETDB1 previously
12 found to bind more tightly to Pc CBX chromodomains than H3K27me3⁴. The initial SETDB1
13 sequence identified was a 10-residue peptide containing an A-L-Kme3-S motif. Therefore, our
14 initial efforts sought to determine the minimal sequence from this peptide necessary for Pc CBX
15 antagonism. Using solution-phase peptide synthesis and an N-terminal benzoyl cap to aid in
16 purification (**Supplementary Information, Synthetic Schemes 1 and 2**), we generated a series
17 of peptides based on the ALKme3S motif and found that, minimally, a 5 residue peptide
18 (compound **5**) is required for antagonism of the CBX chromodomains in our panel (**Table 1**,
19 compounds **3-7**). Compound **5** displayed modest affinity for CBX4 and -7, weak affinity for
20 CBX8 and CBX5, and submicromolar potency for CDYL2. In contrast to the Pc CBX
21 chromodomains, the chromodomain of CDYL2 maintains weak affinity for the shorter peptides,
22 compounds **3**, **4** and **7**. Extending the peptide length to six residues resulted in compound **6**
23 which displays somewhat reduced affinity for each panel member relative to **5**, suggesting that
24 there are subtle differences in the way these different chromodomains recognize the N-terminus
25 of these molecules. Peptide **7** binding affinities were consistent with previous observations that
26 binding of CBX chromodomains requires the presence of a serine C-terminal to the methyl-
27 lysine.^{4, 8}
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51
52 Concomitantly with our studies to investigate the minimal length requirements for
53 chromodomain engagement, we had observed robust binding of compound **6** with CBX7 in
54 isothermal titration calorimetry (ITC) studies ($K_d = 10.3 \pm 0.9 \mu\text{M}$, **Supplementary Figure 1**).
55
56
57
58
59
60

1
2
3 Therefore, we sought to use **6** as a scaffold for further SAR exploration in parallel. *N*_α-
4 methylation has previously been shown to improve the membrane permeability of peptides²⁰ so
5 we explored the effect of methylating select backbone amides of **6** on chromodomain binding
6 (peptides **8-10**). Methylation of the alanine amide nitrogen (**8**) resulted in loss of binding to all
7 chromodomains tested. Methylation of the phenylalanine amide nitrogen (**9**) selectively bound to
8 CDYL2 while methylation of the glycine amide in compound **10** had little effect on binding to
9 any of our panel members when compared to the corresponding unmethylated analog (**6**). These
10 results combined with prior structural studies^{4, 21} suggest that the alanine N-H of these peptides
11 makes a critical hydrogen bond that is necessary for binding to chromodomains. The amide N-H
12 of the Phe residue of these and related peptides does not appear to be positioned to form any
13 hydrogen bonds when bound to Pc chromodomains,^{10, 21} suggesting that this modification may
14 instead introduce a steric clash. Finally, when combined with prior structural studies,^{4, 10, 21} these
15 results indicate that the benzamide cap of both **6** and **10** is solvent-exposed when bound to
16 CBX4, -7 or CDYL2 and the Gly amide is likely not making any significant protein contacts.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35

36
37 In parallel with our studies in **Table 1**, we focused on finding a suitable replacement for the
38 Kme3 of **6** in an effort to increase passive membrane permeability and enhance stability toward
39 demethylase enzymes²² (**Table 2**). Consistent with previous studies, the non-methylated lysine
40 analog of **6**, compound **11**, displayed no antagonism of any of the chromodomains in the panel.⁴
41 We performed a series of reductive aminations on **11** (**Supplementary Information, Synthetic**
42 **Scheme 3**) to explore the effect of replacing the quaternary amine with bulky secondary amines
43 (compounds **12-16**) and in general, found that the CBX chromodomains have limited affinity for
44 these secondary amines, while the CDYL2 chromodomain is able to bind these ligands with
45 modest affinity. For example, simple isopropylation of the lysine (**12**) showed no affinity for
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 CBX chromodomains but single-digit micromolar affinity for CDYL2. Incorporation of a
4 cyclopentyl or cyclohexyl (compounds **13** and **14**) did not recover CBX activity. Interestingly,
5
6 fusion of the 5- and 6-membered rings in norbornyl-lysine (**15**) showed weak antagonism of
7
8 CBX7 and equipotent antagonism of CDYL2 when compared to cyclopentyl alone (**13**).
9
10 Adamantyl substitution (**16**) resulted in no measurable CBX7 activity and CDYL2 activity
11
12 comparable to the cyclohexyl compound (**14**), indicating that increasing the size of the fused ring
13
14 system beyond norbornyl (**15**) is not beneficial to chromodomain binding. Taken together, this
15
16 data suggests that CDYL2 is more accommodating to various secondary amine Kme3
17
18 replacements than the CBX chromodomains.
19
20
21
22
23
24

25
26 We next focused on the effects of tertiary substitutions at the ϵ -nitrogen of the lysine residue.
27
28 Unsurprisingly,^{4, 12} we found that dimethylation (**17**) reduced the activity for both CBX7 and
29
30 CDYL2 about 3-fold relative to the analogous Kme3 compound (**6**). However, incorporation of a
31
32 slightly larger diethyl-lysine as in compound **18** resulted in similar affinities for all
33
34 chromodomains as compound **6**, albeit with slighted improved affinity for CBX8.
35
36 Encouragingly, this supported the notion that a quaternary amine was not likely to be required
37
38 for potent CBX chromodomain binding. To generate cyclic tertiary amines as in compounds **19-**
39
40 **21**, we reacted **11** with a series of dibromoalkanes (**Supplementary Information, Synthetic**
41
42 **Scheme 3**). Cyclization of the diethyl-lysine to generate the pyrrolidinyl derivative, **19**, resulted
43
44 in a reduction in potency across the panel, while increasing the ring size to a piperidinyl group
45
46 (**20**) or a 7-membered azepane (**21**) also did not appear to be beneficial. Incorporation of a
47
48 bulkier dicyclobutyl-lysine (**22**) also did not result in any significant enhancements in potency.
49
50
51
52
53
54

55 We were intrigued by the ability of the norbornyl functionality to weakly interact with CBX7
56
57 and more potently with CDYL2. It is possible that binding of this functionality to the aromatic
58
59
60

1
2
3 cage could sterically hinder any cation- π interactions between the cage residues and the ϵ -
4 nitrogen, relying less on the presence of a basic nitrogen for binding. We had simultaneously
5 prepared an acetylated version of **15** (compound **23**) and although potency was reduced
6 approximately 7-fold for CDYL2, the presence of a basic nitrogen does not appear to be an
7 absolute requirement for CDYL2 binding. We further interrogated the nature of the cation- π
8 interaction at this position by synthesizing the more strongly basic tetramethylguanidine
9 derivative by reacting **11** with TBTU in the absence of a carboxylic acid to give compound **24**,
10 and additionally prepared dibasic compounds **25** and **26** via reductive amination
11 (**Supplementary Information, Synthetic Scheme 3**). All three of these compounds showed no
12 activity for the CBX chromodomains and weakened affinity for CDYL2. Taken together, the
13 SAR data resulting from modification of the lysine indicates that the presence of a quaternary
14 amine is not required for engagement of our chromodomain panel; however, only the diethyl-
15 lysine containing compound (**18**) demonstrates a binding profile comparable to that of **6**.
16 Substantially improving potency for Pc CBX chromodomains through modification of the lysine
17 amine presents a significant challenge, but may be a more viable strategy for CDY
18 chromodomain ligand optimization.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41

42 Our previously described *in silico* studies¹⁰ on the interaction of CBX7 and an H3K9me3
43 peptide provided insight into the mechanism of induced-fit recognition of Kme3 peptides by
44 CBX7. These studies suggested that the chromodomain of CBX7 first recognizes the N-terminal
45 cap residue at the (-4) position from the methyl-lysine, allowing the peptide to engage the
46 chromodomain and leading CBX7 to close around the histone and engage the Kme3 with its
47 newly formed aromatic cage.¹⁰ This induced-fit binding mechanism supports peptidomimetics as
48 a likely choice for CBX7 inhibitors and the absence of a preformed aromatic cage makes the
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 discovery of traditional small molecule inhibitors a significant challenge. Previous studies from
4 the Zhou lab have reported weak, non-peptidic small molecule CBX7 ligands; however, SAR
5 studies around these molecules struggled to produce significant improvements in potency.^{21, 23}
6
7
8
9

10
11 Our *in silico* studies led us to hypothesize that the potency of our peptidic antagonists could be
12 improved through modification of the N-terminus. Diethyl-lysine was chosen from our studies in
13 **Table 2** as an optimal Kme3 replacement and hence was incorporated in future antagonists while
14 we varied the N-terminus (**Table 3**, compounds **1** and **27-41**; **Supplementary Information**,
15 **Synthetic Schemes 1 and 4**). Our studies in **Table 1** indicated that the glycine residue of **6** was
16 dispensable, which led us to prepare compound **27**, which is the diethyl-lysine analog of
17 compound **5**. This compound overall showed similar or improved activities toward each of the
18 chromodomains bound by **5**, further confirming that diethyl-lysine is a suitable Kme3
19 replacement in the context of this shorter peptide. Interestingly, **27** demonstrates measurable
20 affinity for CBX5 in contrast to compound **18**, providing initial evidence that the aromatic cage
21 of CBX5 can also accommodate non-natural lysine analogs.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36

37
38 We next began exploring substituents on the N-terminal aromatic ring of **27** by placing a
39 methyl substituent at the *ortho*-, *meta*- and *para*-positions. Methylation at the *ortho*-position (**28**)
40 of the ring dramatically reduced potency for all chromodomains except for CDYL2, causing a
41 20-fold decrease in binding for CBX7 relative to compound **27**. Affinity for all chromodomains
42 except CBX5 is recovered to levels comparable to **27** when the methyl group is shifted to the
43 *meta*-position (**29**). Installation of *para*-Me (**30**) results in further improvements in binding to
44 CBX7 and CBX4, with submicromolar affinities in each case. Interestingly, CBX8 did not
45 display the same preference as CBX4 and -7 for **30** over **29**, while methylation had little to no
46 effect on CDYL2 binding.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Having determined that, in general, alkylation is preferred at the *para*-position by Pc
4 chromodomains, we next explored the effect of increasing steric bulk at this position
5 (compounds **31-32** and **1**). Overall, increasing lipophilicity had opposing effects on the affinity
6 for Polycomb vs HP1 chromodomains. Introduction of a *para*-ethyl or *para*-isopropyl substituent
7 as in compounds **31** and **32**, respectively, increased the affinity ~2-5 fold for CBX4, -7, and -8,
8 as well as CDYL2, relative to **30**, while the affinity for CBX5 was largely unchanged or slightly
9 weakened. *t*-Butyl substitution (**1**) further improved potency for CBX4 and -7 while having no
10 significant effect on the affinity for CBX8 and causing a loss of detectable CBX5 affinity. Taken
11 together, these data indicate that potency and selectivity can be tuned by systematically adjusting
12 both the position of alkylation and degree of lipophilicity on the benzoyl cap at the N-terminus.
13 These studies enabled us to dramatically improve the potency of our compounds for CBX4 and -
14 7, leading to compounds with IC₅₀ values of less than 100 nM. These results also provided strong
15 support for our molecular dynamics studies, which suggested that initial recognition of histones
16 occurs through hydrophobic contacts formed between CBX7 and the histone side chain at this
17 position.¹⁰

18
19
20 We also evaluated the effect of incorporating electron-withdrawing groups at the *para*-position
21 of the N-terminal phenyl cap with compounds **33** and **34**. A methyl ester at the *para*-position
22 (**33**) resulted in an approximately 15-fold loss in potency for both CBX4 and -7 relative to
23 compound **1**, but only an approximately 3-fold loss in potency for CBX8, further suggesting that
24 CBX4 and -7 depend more strongly on the presence of bulky alkyl substituents at this position.
25 The strongly electron-withdrawing trifluoromethyl substituent of **34** also weakened binding to
26 the Pc chromodomains, although to a lesser extent than compound **33**. CDYL2 affinity is largely
27 unaffected by replacement of *para*-alkyl substituents with electron-withdrawing groups,
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 consistent with the generally less stringent binding preferences of CDYL2. Inclusion of an
4 electron-donating group (**35**) also did not further improve binding to Pc CBX chromodomains,
5
6 indicating that alkyl substituents at this position are preferred. In contrast, these results suggest
7
8 that CBX5 may prefer less sterically demanding functional groups at the N-terminus.
9
10

11
12
13 We next investigated the importance of the spacing between the N-terminal hydrophobic group
14 and the rest of the peptide. As can be seen with compound **36**, addition of a single methylene
15 group relative to compound **1** resulted in an approximately 100-fold loss of potency for CBX4
16 and -7 and almost a 20-fold loss for CBX8, while CDYL2 was largely unable to distinguish
17 between the *t*-butyl phenyl and *t*-butyl benzyl moieties. Additionally, larger bicyclic aromatic
18 capping groups with a methylene spacer (**37** and **38**) were also less well tolerated by CBX4, -7
19 and -8. Consistent with prior results, the increased steric demand of these functionalities was
20 detrimental to CBX5 activity.
21
22
23
24
25
26
27
28
29
30
31

32
33 Our studies in **Table 2** indicated that the tolerance of Pc CBX chromodomains to unnatural
34 Kme analogs is fairly limited, yet we were interested in exploring a small number of lysine
35 modifications, as well as the native Kme3 residue, in the context of our more potent peptide
36 scaffold. We generated the Kme3 analog of **1**, compound **39**, and two asymmetric tertiary amine
37 derivatives, compounds **40** and **41**. The isopropyl group of **40** and **41** was hypothesized to make
38 additional contacts with the aromatic cage of the Pc CBX proteins and to more completely fill
39 the cage based on the crystal structures of **1** bound to these chromodomains (**Figure 3a**).
40 Mirroring prior results, the Kme3 containing compound (**39**) was about equipotent to **1**, while the
41 bulkier lysine mimetics of compounds **40** and **41** also did not appear to pick up any additional
42 interactions to further improve potency.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 3. Compound 1 bound to CBX7 (PDB 5EPJ). (a) Interaction of the CBX7 aromatic cage with diethyl-lysine. (b) Binding pocket of CBX7 occupied by the Ala side chain of 1. (c) Position of Phe of 1 when bound to CBX7.

Previous studies from the Hof group found that the residue at the (-2) position from the modified lysine can have profound effects on the selectivity and potency of peptide antagonists for Pc CBX chromodomains.⁹ We explored this in the context of our antagonists by testing the effect of increasing the size of the alkyl side chain at this position (**Table 4**, compounds **42** and **43**; **Supplementary Information, Synthetic Scheme 4**). We found that replacing the methyl side chain of alanine with an ethyl or isopropyl group reduced affinity for CBX4, -7, and CDYL2 while slightly increasing the affinity for CBX8. This was not surprising as previously solved co-crystal structures with **1** revealed that the pocket occupied by this alanine side chain is larger in CBX8 than in CBX4 and -7 owing to substitution of Val13 in CBX4/7 with Ala13 in CBX8 (**Figure 3b**).¹⁰ The increased potency and modest selectivity of **43** for CBX8 may make it a useful starting point for the development of more potent and selective CBX8 compounds.

Modifications to the C-terminus of compound **1** (compounds **44-46**; **Supplementary Information, Synthetic Schemes 4 and 5**), were reasonably well tolerated. While the methyl ester of **1** is the preferred functionality at this position, replacement with the corresponding acid (**44**) or dimethyl amide (**45**), or removal of the C-terminal carbonyl entirely (**46**), resulted in a

1
2
3 mere 2- to 4-fold reduction in potency for Pc CBX chromodomains. Interestingly, compound **44**
4
5 resulted in a slightly larger reduction in potency for CDYL2 (5-fold), suggesting that additional
6
7 exploration of the C-terminus of these molecules may be useful for further tuning selectivity.
8
9

10
11 The observed tolerance to modifications at the C-terminus previously led us to generate a
12
13 biotinylated analog of **1** (compound **2**). This tool compound showed minimal loss in potency for
14
15 CBX7, and importantly was used to evaluate the selectivity of compound **1** more broadly and
16
17 served as a key chemiprecipitation reagent to demonstrate cellular target engagement.¹⁰ The co-
18
19 crystal structures of the Pc CBX proteins with compound **1** revealed that the phenylalanine of **1**
20
21 is largely solvent exposed (**Figure 3c**); therefore, we hypothesized that this was an additional
22
23 position we could use for tagging to access additional tool compounds. An amine functional
24
25 group was installed on the phenylalanine side chain, and as expected from the crystallographic
26
27 information, compound **47** displayed similar affinities to **1** for each of the chromodomains in our
28
29 panel. Further, because the boc group of **48** does not interfere with binding, we reasoned that this
30
31 would be a suitable location to place larger chemical moieties. To this end, we generated a
32
33 biotinylated derivative of **47** (**biotin-47**; **Supplementary Information, Synthetic Scheme 6**) via
34
35 modification of the free amine. Compound **2** was previously shown to selectively
36
37 chemiprecipitate CBX4, -7 and -8, but not CBX2 or -6 from PC3 cell lysates, despite having
38
39 demonstrated affinity for these proteins.¹⁰ Similarly, we found that **biotin-47** was unable to pull-
40
41 down CBX2 and -6, while it also lost the ability to pull-down CBX7, indicating that pull-down
42
43 of these endogenous proteins from cell lysates is highly sensitive to the positioning of the
44
45 chemical tag (**Supplementary Figure 2**). It is possible that biotinylation of the ligand produces a
46
47 steric clash in the context of specific intact PRC1 complexes that can't be anticipated from CBX-
48
49 only co-crystal structures.
50
51
52
53
54
55
56
57
58
59
60

1
2
3 While modifications to the lysine side chain did not improve the *in vitro* potency of our
4 compounds, we sought to compare the cellular activity of compound **39** (containing Kme3) to
5 compound **1** since we initially hypothesized that replacement of the Kme3 would improve
6 passive membrane permeability. We first measured the intracellular concentration of **39**.
7
8 Treatment at a 30 μM extracellular dose of **1** for 24 hours results in an intracellular concentration
9 of $1.4 \pm 0.3 \mu\text{M}$ in PC3 cells.¹⁰ Under the same conditions, we observed an intracellular
10 concentration of $0.61 \pm 0.2 \mu\text{M}$ of compound **39**. It was surprising that incorporation of
11 trimethyl-lysine resulted in only an approximately 2-fold decrease in the permeability of the
12 compound, and we were intrigued to investigate whether **39** was functionally available for
13 chromodomain binding in cells since it is known that highly charged compounds may
14 accumulate in endosomes and lysosomes.^{24,25}
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29

30 Previously, we have shown that **1** inhibits PC3 cell proliferation through antagonism of
31 polycomb CBX chromodomains. In order to assess the effect of Kme3 substitution on functional
32 activity, we compared the anti-proliferative effects of **1** and **39** to our negative control analog of
33 **1**, compound **49** (UNC4219). Compound **49** contains a single backbone methylation which
34 ablates its ability to interact with chromodomains.¹⁰ Despite having nearly identical *in vitro*
35 potencies for CBX chromodomains as **1** and only a 2-fold reduction in permeability, compound
36 **39** had no effect on PC3 cell proliferation, analogous to the negative control (**Figure 4**),
37 supporting our hypothesis that Kme3 replacement is required for cellular activity. We suspect
38 that the lack of biological activity observed upon treatment with compound **39** may be a result of
39 endosomal and lysosomal trapping of the molecule.^{24,25}
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 4. Effect of compounds 1, 39, and 49 on PC3 cell proliferation. Cells were treated for 72 hours with the indicated compounds (30 μ M) or DMSO, fixed and DAPI stained. Cell counts were numerated using high content microscopy and normalized to DMSO control ($n \geq 4$).

As discussed previously, our proposed binding model involves an induced-fit mode of recognition in which the N-terminus of these ligands makes the first specific contact with CBX7 and that initial hydrophobic interaction nucleates a series of backbone hydrogen bonds, allowing CBX7 to close and the aromatic cage to form to generate the stably bound state. While we were able to apply this model to develop more potent chromodomain antagonists, largely through modification of the N-terminal aromatic capping group, we were motivated to further interrogate the validity of the model. We therefore turned to surface-plasmon resonance (SPR) studies with CBX7 and a subset of our compounds that differ at the N-terminus in order to determine if the binding kinetics could provide any evidence to further support this induced-fit hypothesis.

When first comparing the SPR data for compound **18** (Figure 5 and Supplementary Figure 3), which contains the larger benzoyl glycine N-terminus, to the *ortho*-Me phenyl derivative (**28**), both of which have similar IC_{50} 's for CBX7, we observed that the k_{on} for compound **28** was slightly faster but that the k_{off} increased in a proportional manner, resulting in comparable affinities. When the methyl group of compound **28** is shifted to the *meta*-position (compound **29**), the higher potency observed for this compound is largely driven by a significant increase in k_{on} . Moving the methyl substituent to the *para*-position (compound **30**) did not substantially

1
2
3 change the k_{on} relative to compound **29** but did slightly improve k_{off} , resulting in the 3-fold
4
5 improvement in CBX7 potency for **30** over **29** observed in AlphaScreen®. Inclusion of a *para-t*-
6
7 butyl moiety in **1** further improves both k_{on} and k_{off} , resulting in one of our most potent CBX7
8
9 inhibitors to date.
10
11

12
13
14 We were intrigued by the fact that the majority of the potency improvements resulting from
15
16 modification of the N-terminus were driven primarily by enhanced association rates instead of
17
18 decreased dissociation rates.²⁶ Based on molecular dynamics simulations, we believe that N-
19
20 terminal portion of these molecules forms the first stable interaction with CBX7, which is
21
22 consistent with modifications at this position primarily affecting the association kinetics. More
23
24 specifically, the improvement in association kinetics may result from a reduction in the
25
26 conformational flexibility at the N-terminus, thereby increasing the likelihood of a productive
27
28 binding collision between CBX7 and the antagonist. SPR results with CBX7 and compound **36**,
29
30 which is approximately 100-fold less potent than **1** for CBX7, demonstrate that insertion of a
31
32 flexible methylene linker primarily reduced k_{on} while k_{off} was minimally affected. Finally,
33
34 analysis of the k_{on} and k_{off} rates of these select compounds with respect to their K_d 's (**Figure 5c**)
35
36 revealed that the binding is better correlated with k_{on} as opposed to k_{off} , providing even stronger
37
38 support for this hypothesis.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 5. (a) Comparison of compound-CBX7 association rates. (b) Comparison of compound-CBX7 dissociation rates. Negative log values of the dissociation rates (1/s) are shown. Data are reported as the mean \pm the standard error of the mean ($n \geq 5$). (c) CBX7 binding affinities of compounds with N-terminal variations are better correlated with the association rates as compared to the dissociation rates. The log transformations of k_{on} (1/M/s), k_{off} (1/s) and K_d (nM) are plotted.

Conclusions


The studies presented herein build on previous work investigating structure-activity relationships of peptide-based antagonists for Pc CBX chromodomains.⁸⁻¹⁰ In particular, we performed the first systematic effort to replace Kme3 with non-natural lysine derivatives and found that tertiary amine Kme3 mimetics are not only tolerated, but are able to maintain similar affinities for chromodomains. Replacement of Kme3 was also found to be critical for cellular activity of these ligands.

A thorough examination of the N-terminus of our inhibitors has also begun to establish the requirements for optimizing potency and selectivity for the Pc, HP1 and CDY families of chromodomains. While CDYL2 appears to be much more tolerant of various modifications at

1
2
3 this position in our studies, CBX chromodomains show a strong preference for alkyl substituents
4 located at the *para*-position of an N-terminal benzoyl cap. Pc chromodomains tend to prefer
5 larger alkyl substituents at this position while the CBX5 chromodomain interacts more favorably
6 with smaller substituents or an unmodified benzoyl group at this position. Further studies will
7 help to elaborate on the requirements for CBX selectivity at this position, and we are currently
8 pursuing alternative approaches toward the discovery and optimization of selective
9 chromodomain ligands via an affinity-based, one-bead one-compound, target class screening
10 strategy.²⁷
11
12
13
14
15
16
17
18
19
20
21
22

23 Our SPR studies have provided strong support for the hypothesized induced-fit binding
24 mechanism for these antagonists with CBX7. These studies have revealed what is, to our
25 knowledge, the first example of a protein-ligand system with an exploitable SAR to control
26 potency via optimization of association kinetics. Further, these studies have demonstrated that
27 the residence time of compound **1** on CBX7 is quite short (~100 seconds). There are an
28 increasing number of examples where the cellular efficacy of drugs and probes are better
29 correlated with the residence time than the binding constant.²⁶ Therefore, we are currently
30 working to apply our knowledge of the binding pathway to increase the residence time of these
31 ligands in order to assess whether this will improve their cellular and *in vivo* efficacy.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60


Table 1. Determination of peptide length requirements for CBX antagonism of ALKme3S peptides.









Compound	R ₁	R ₂	R ₃	IC ₅₀ (μM) ^a				
				Polycomb			CDY	HP1
				CBX4	CBX7	CBX8	CDYL2	CBX5 (HP1α)
3	Bz	-CONMe	-H	>60	>60	>60	54 ± 16	>60
4	Bz	-CO-Ser-OMe	-H	>60	>60	>60	16 ± 1.9	>60
5	Bz-Phe	-CO-Ser-OMe	-H	6.9 ± 1.3	2.6 ± 0.39	33 ± 7.1	0.57 ± 0.042	49 ± 4.1
6	Bz-Gly-Phe	-CO-Ser-OMe	-H	58 ± 5.3	18 ± 4.5	>60	1.0 ± 0.23	>60
7	Bz-Phe	-H	-H	>60	>60	>60	37 ± 4.2	>60
8	Bz-Gly-Phe	-CO-Ser-OMe	-Me	>60	>60	>60	>60	>60
9	Bz-Gly-N(me)Phe	-CO-Ser-OMe	-H	>60	>60	>60	15 ± 6.4	>60
10	Bz-N(me)Gly-Phe	-CO-Ser-OMe	-H	57 ± 6.0	26 ± 8.3	>60	0.47 ± 0.14	>60

^aIC₅₀ values are reported as the mean ± the standard deviation of at least two biological replicates each performed in duplicate.

Table 2. Exploration of secondary and tertiary amine Kme3 replacements.




Compound	R ₁	IC ₅₀ (μM) ^a				
		Polycomb			CDY	HP1
		CBX4	CBX7	CBX8	CDYL2	CBX5
11		>60	>60	>60	>60	>60
12		>60	>60	>60	2.3 ± 0.48	>60
13		>60	>60	>60	3.0 ± 0.28	>60
14		>60	>60	>60	17 ± 4.3	>60
15		>60	51 ± 10	>60	5.0 ± 0.48	>60
16		>60	>60	>60	14 ± 2.0	>60
17		>60	51 ± 10	>60	2.8 ± 0.41	>60
18		46 ± 19	16 ± 3.7	47 ± 19	0.67 ± 0.071	>60
19		>60	53 ± 5.6	>60	1.1 ± 0.25	>60
20		>60	41 ± 6.6	>60	0.54 ± 0.039	>60















21		>60	28 ± 3.6	31 ± 7.8	0.59 ± 0.080	>60
22		>60	32 ± 8.9	34 ± 27	1.8 ± 0.56	>60
23		>60	>60	>60	38 ± 13	>60
24		>60	>60	>60	2.5 ± 0.48	>60
25		>60	>60	>60	8.6 ± 0.62	>60
26		>60	>60	>60	24 ± 9.7	>60

^aIC₅₀ values are reported as the mean \pm the standard deviation of at least two biological replicates each performed in duplicate.

Table 3. Effect of N-terminal variations and select lysine modifications.




Compound	R ₁	R ₂	IC ₅₀ (μM) ^a				
			Polycomb			CDY	HP1
			CBX4	CBX7	CBX8	CDYL2	CBX5
27			3.0 ± 0.82	1.2 ± 0.69	4.8 ± 1.4	0.34 ± 0.021	6.9 ± 0.52
28			58 ± 5.4	25 ± 2.5	>60	0.56 ± 0.054	>60
29			2.7 ± 1.0	0.74 ± 0.11	5.1 ± 1.4	0.34 ± 0.12	>60
30			0.77 ± 0.030	0.22 ± 0.049	4.8 ± 0.39	0.28 ± 0.16	21 ± 4.3
31			0.19 ± 0.074	0.081 ± 0.020	1.5 ± 0.19	0.10 ± 0.025	14 ± 2.4
32			0.16 ± 0.021	0.061 ± 0.016	1.9 ± 0.46	0.10 ± 0.026	38 ± 6.3
1			0.083 ± 0.028	0.037 ± 0.017	1.3 ± 0.48	0.14 ± 0.043	>60
33			1.2 ± 0.45	0.57 ± 0.087	4.8 ± 0.87	0.11 ± 0.026	26 ± 2.7
34			0.48 ± 0.11	0.23 ± 0.041	3.0 ± 0.23	0.13 ± 0.019	33 ± 8.6

35			0.75 ± 0.11	0.25 ± 0.026	3.3 ± 0.44	0.26 ± 0.018	12 ± 2.3
36			9.7 ± 4.4	4.4 ± 0.67	23 ± 2.3	0.20 ± 0.025	>60
37			3.4 ± 0.66	1.7 ± 0.61	15 ± 2.3	0.11 ± 0.031	>60
38			5.2 ± 0.72	1.1 ± 0.23	24 ± 4.2	0.084 ± 0.028	>60
39			0.075 ± 0.021	0.035 ± 0.0064	1.5 ± 0.52	0.11 ± 0.034	>60
40			0.075 ± 0.014	0.035 ± 0.0084	1.0 ± 0.21	0.13 ± 0.010	>60
41			0.062 ± 0.016	0.032 ± 0.0078	1.3 ± 0.34	0.11 ± 0.022	>60

^aIC₅₀ values are reported as the mean \pm the standard deviation of at least two biological replicates each performed in duplicate.

Table 4. Exploration of modifications to the phenylalanine, alanine, and C-terminus of compound 1.



Compound	R ₁	R ₂	R ₃	IC ₅₀ (μM) ^a				
				Polycomb			CDY	HP1
				CBX4	CBX7	CBX8	CDYL2	CBX5
42	-H	-Et	COOMe	0.64 ± 0.19	0.24 ± 0.017	0.98 ± 0.22	0.50 ± 0.071	>60
43	-H	-iPr	COOMe	7.9 ± 1.0	3.1 ± 0.87	0.70 ± 0.15	0.75 ± 0.026	>60
44	-H	-Me	COOH	0.24 ± 0.027	0.13 ± 0.021	2.0 ± 0.92	0.74 ± 0.12	>60
45	-H	-Me	CON(Me) ₂	0.14 ± 0.038	0.047 ± 0.012	2.1 ± 0.50	0.36 ± 0.055	>60
46	-H	-Me	-H	0.20 ± 0.031	0.075 ± 0.012	3.5 ± 0.68	0.41 ± 0.061	>60
47	-CH ₂ NH ₂	-Me	COOMe	0.17 ± 0.046	0.063 ± 0.028	2.4 ± 0.61	0.075 ± 0.019	53 ± 4.7
48	-CH ₂ NHBoc	-Me	COOMe	0.098 ± 0.011	0.056 ± 0.016	1.7 ± 0.32	0.16 ± 0.012	44 ± 11

^aIC₅₀ values are reported as the mean ± the standard deviation of at least two biological replicates each performed in duplicate.

EXPERIMENTAL SECTION

Expression constructs

The chromodomains of CBX4 (residues 8-65 of NP_003646), CBX5 (residues 18-75 of NP_036429) and CBX8 (residues 8-61 of NP_065700) were expressed with N-terminal His-tags in pET28 expression vectors. The chromodomain of CBX7 (residues 8-62 of NP_783640) and

1
2
3 CDYL2 (residues 1-75 of NP_689555) were expressed with a C-terminal His-tag in a pET30
4
5 expression vector.
6
7

8 9 **AlphaScreen® procedure and conditions**

10
11 The AlphaScreen® assay was generally performed as previously described.¹⁹ In brief, compound
12
13 plates (1 μ L at 10 mM highest concentration; 3-fold, 10-point dilutions in DMSO) were diluted
14
15 in 1X assay buffer (20 mM TRIS pH 7.5, 75 mM NaCl, 2 mM DTT and 0.05% Tween-20) to 1
16
17 mM using a Multimek robotic pipettor (Nanoscreen) and 1 μ L was spotted into the wells of 384-
18
19 well low-volume Proxiplates (PerkinElmer). To these plates 9 μ L of protein-peptide mix in 1X
20
21 assay buffer was added by Multidrop (Thermo) to bring the final compound concentration to 100
22
23 μ M and incubated for 30 min at room temperature. Next, 2 μ L of a 1:1 mixture of streptavidin-
24
25 conjugate donor and nickel-chelate or α -GST acceptor beads (45 μ g/mL in 1X assay buffer) were
26
27 added and the plates were allowed to incubate for an additional 30 min in the dark at room
28
29 temperature. After incubation, the plates were read on an EnVision multi-label reader equipped
30
31 with an HTS AlphaScreen laser (Perkin Elmer). Dose-response curves were fit using a 4-
32
33 parameter or 3-parameter fixed top binding equation.
34
35
36
37
38
39

40 41 **Compound permeability studies**

42
43 PC3 cells were cultured until reaching ~80% confluency in a T75 cell culture flask. The media
44
45 was exchanged with 10 mL fresh media (0.3% DMSO) containing 30 μ M compound. Cells were
46
47 incubated with compound for 24 hours. After 24 hours, cells were washed with PBS, trypsinized,
48
49 and wash again with PBS. The resulting cell pellets (3) were flash frozen with liquid nitrogen
50
51 and submitted to Drumetix Laboratories (Greensboro, NC) for analysis by LC-MS/MS.
52
53
54

55 56 **Pull-down studies**

57
58
59
60

1
2
3 Cells were cultured on a T175 tissue culture flask until reaching ~80-90% confluency. Following
4 trypsinization, the pellet was washed twice with PBS and lysed with 500 μ L of lysis buffer.
5
6 When applicable, vehicle control or 100 μ M of compound **1** was added to an aliquot of 1000 μ g
7
8 of total protein and diluted to 500 μ L with 20 mM Tris (pH 8)/150 mM NaCl/0.1% Tween-20
9
10 (TBST). Pull-down reagents were bound to Streptavidin M-280 Dynabeads® by rotating 30 μ L
11
12 of beads with a 20-fold excess of pull-down reagent for 30 minutes at room temperature in.
13
14 Unbound pull-down reagent was then removed and the beads were washed 3X with 200 μ L of
15
16 TBST. The lysate was then transferred to an Eppendorf tube containing 30 μ L of Streptavidin M-
17
18 280 Dynabeads® that had been pre-bound to biotin-46. The mixture was rotated for 2 hr at 4°C.
19
20 The depleted lysate was removed and the beads were washed 3X with 300-500 μ L of TBST. The
21
22 beads were re-suspended with 30 μ L of 1X Laemmli sample buffer and heated at 95°C for 3
23
24 minutes. Fifteen microliters of the sample was then loaded into a gel and analyzed via western
25
26 blotting using the appropriate primary antibody and detected using a LI-COR Odyssey®
27
28 instrument and the appropriate fluorescent secondary antibodies (diluted 1:10,000 in PBST).
29
30 One percent of input was used for western blotting of all proteins.
31
32
33
34
35
36
37
38
39

40 **Cell Proliferation Experiments**

41 PC3 cells were seeded at 400 cells/well into 96-well plates. Cells were allowed to adhere
42
43 overnight. The media (DMEM supplemented with 10 % FBS) was then exchanged with fresh
44
45 media containing DMSO, or 30 μ M of the indicated compound. On day three, the cells were
46
47 fixed with ice-cold methanol for 30 sec. and rehydrated with PBS. Nuclei of the cells were
48
49 stained with DAPI (0.05 μ g/ml) and numerated using High Content Microscopy (Array Scan™
50
51 High Content Analysis, Thermo Fisher #NX10002L).
52
53
54
55
56

57 **Surface Plasmon Resonance Experiments**

58
59
60

1
2
3 SPR experiments were performed on a ProteOn™ XPR36 Interaction Array System. C-terminally
4 His-tagged CBX7 was immobilized using a ProteOn™ HTE sensor chip. Compounds were
5
6 flowed at a rate of 50-100 $\mu\text{L}/\text{minute}$ with a contact time of 150 seconds. Dissociation steps were
7
8 run for 300 seconds at a flow rate of 100 $\mu\text{L}/\text{minute}$. Double referencing subtraction was done
9
10 with buffer and protein blank channels to account for nonspecific binding to the sensor chip.
11
12 Sample Sensorgrams are shown in **Supplementary Figure 3**.
13
14
15
16
17
18
19

20 ASSOCIATED CONTENT

21 22 23 **Supporting Information.**

24
25
26 Descriptions of protein expression and purification, ITC experiments, cell culture and lysis
27
28 conditions, general chemistry procedures, compound characterization data, and additional figures
29
30 illustrating ITC results, pull-down results and SPR sensorgrams. This material is available free of
31
32 charge via the Internet at <http://pubs.acs.org>.”
33
34
35
36

37 Author Contributions

38
39 Compound synthesis was performed by JIS, JL, and RP. AlphaScreen® assays were developed
40
41 by CS, JIS and KHP, screening was performed by CS. JND and SHC expressed and purified all
42
43 protein constructs. NC and JIS performed cell proliferation and compound permeability studies.
44
45 JL performed pull-down experiments. JS and BDM designed all compounds. JIS performed all
46
47 SPR studies. JS, KHP, SVF and LIJ planned and discussed all experiments. The manuscript was
48
49 written through contributions of all authors. All authors have given approval to the final version
50
51 of the manuscript.
52
53
54
55

56 57 Funding Sources

1
2
3 The research described here was supported by the National Institute of General Medical
4 Sciences, US National Institutes of Health (NIH, grant R01GM100919), the Carolina Partnership
5 and the University Cancer Research Fund, University of North Carolina at Chapel Hill.
6
7
8
9
10

11 12 13 ACKNOWLEDGMENT

14
15 We would like to acknowledge Ashutosh Tripathy (UNC Chapel Hill) for his assistance with the
16 SPR experiments in this work. We thank Noah Sciaky (UNC Chapel Hill) for his help with the
17 high content microscopy experiments. We would also like to acknowledge Mark Bedford (MD
18 Anderson) for the CDYL2 chromodomain constructs and Cheryl Arrowsmith (SGC Toronto) for
19 the CBX chromodomain constructs.
20
21
22
23
24
25
26
27

28 29 ABBREVIATIONS

30 CBX2, chromobox homolog protein 2; CBX4, chromobox homolog protein 4; CBX5,
31 chromobox homolog protein 5; CBX6, chromobox homolog protein 6; CBX7, chromobox
32 homolog protein 7; CBX8, chromobox homolog protein 8; HP1, Heterochromatin protein 1;
33 Kme, methyl-lysine; PRC1, Polycomb repressive complex 1; SPR, surface plasmon resonance;
34 Pc, Polycomb; CDYL2, chromodomain protein, Y-like 2; DMSO, dimethyl sulfoxide.
35
36
37
38
39
40
41
42

43 44 AUTHOR INFORMATION

45 46 Corresponding Authors

47
48 E-mail: ingerman@email.unc.edu (t: 919-962-4870) and svfrye@email.unc.edu (t: 919-843-
49 5486)
50
51
52
53

54 55 Present Addresses

56
57 †Center for Epigenetics, Van Andel Research Institute, Grand Rapids, Michigan 49503, USA
58
59
60

REFERENCES

1. Strahl, B. D.; Allis, C. D. The language of covalent histone modifications. *Nature* **2000**, *403*, 41-45.
2. Arrowsmith, C. H.; Bountra, C.; Fish, P. V.; Lee, K.; Schapira, M. Epigenetic protein families: a new frontier for drug discovery. *Nat. Rev. Drug Discovery* **2012**, *11*, 384-400.
3. Yun, M.; Wu, J.; Workman, J. L.; Li, B. Readers of histone modifications. *Cell Res.* **2011**, *21*, 564-578.
4. Kaustov, L.; Ouyang, H.; Amaya, M.; Lemak, A.; Nady, N.; Duan, S.; Wasney, G. A.; Li, Z.; Vedadi, M.; Schapira, M.; Min, J.; Arrowsmith, C. H. Recognition and specificity determinants of the human cbx chromodomains. *J. Biol. Chem.* **2011**, *286*, 521-529.
5. James, L. I.; Korboukh, V. K.; Krichevsky, L.; Baughman, B. M.; Herold, J. M.; Norris, J. L.; Jin, J.; Kireev, D. B.; Janzen, W. P.; Arrowsmith, C. H.; Frye, S. V. Small-molecule ligands of methyl-lysine binding proteins: optimization of selectivity for L3MBTL3. *J. Med. Chem.* **2013**, *56*, 7358-7371.
6. Herold, J. M.; James, L. I.; Korboukh, V. K.; Gao, C.; Coil, K. E.; Bua, D. J.; Norris, J. L.; Kireev, D. B.; Brown, P. J.; Jin, J.; Janzen, W. P.; Gozani, O.; Frye, S. V. Structure-activity relationships of methyl-lysine reader antagonists. *Med. Chem. Commun.* **2012**, *3*, 45-51.
7. Perfetti, M. T.; Baughman, B. M.; Dickson, B. M.; Mu, Y.; Cui, G.; Mader, P.; Dong, A.; Norris, J. L.; Rothbart, S. B.; Strahl, B. D.; Brown, P. J.; Janzen, W. P.; Arrowsmith, C. H.; Mer, G.; McBride, K. M.; James, L. I.; Frye, S. V. Identification of a fragment-like small molecule ligand for the methyl-lysine binding protein, 53BP1. *ACS Chem. Biol.* **2015**, *10*, 1072-1081.
8. Simhadri, C.; Daze, K. D.; Douglas, S. F.; Quon, T. T.; Dev, A.; Gignac, M. C.; Peng, F.; Heller, M.; Boulanger, M. J.; Wulff, J. E.; Hof, F. Chromodomain antagonists that target the

1
2
3 polycomb-group methyllysine reader protein chromobox homolog 7 (CBX7). *J. Med. Chem.*
4
5 **2014**, *57*, 2874-2883.

6
7
8 9. Milosevich, N.; Gignac, M. C.; McFarlane, J.; Simhadri, C.; Horvath, S.; Daze, K. D.;
9
10 Croft, C. S.; Dheri, A.; Quon, T. T.; Douglas, S. F.; Wulff, J. E.; Paci, I.; Hof, F. Selective
11
12 inhibition of CBX6: a methyllysine reader protein in the polycomb family. *ACS Med. Chem.*
13
14 *Lett.* **2016**, *7*, 139-144.

15
16
17 10. Stuckey, J. I.; Dickson, B. M.; Cheng, N.; Liu, Y.; Norris, J. L.; Cholensky, S. H.;
18
19 Tempel, W.; Qin, S.; Huber, K. G.; Sagum, C.; Black, K.; Li, F.; Huang, X. P.; Roth, B. L.;
20
21 Baughman, B. M.; Senisterra, G.; Pattenden, S. G.; Vedadi, M.; Brown, P. J.; Bedford, M. T.;
22
23 Min, J.; Arrowsmith, C. H.; James, L. I.; Frye, S. V. A cellular chemical probe targeting the
24
25 chromodomains of Polycomb repressive complex 1. *Nat. Chem. Biol.* **2016**, *12*, 180-187.

26
27
28 11. Eissenberg, J. C. Structural biology of the chromodomain: form and function. *Gene* **2012**,
29
30 *496*, 69-78.

31
32
33 12. Fischle, W.; Franz, H.; Jacobs, S. A.; Allis, C. D.; Khorasanizadeh, S. Specificity of the
34
35 chromodomain Y chromosome family of chromodomains for lysine-methylated ARK(S/T)
36
37 motifs. *J. Biol. Chem.* **2008**, *283*, 19626-19635.

38
39
40 13. Vandamme, J.; Volkel, P.; Rosnoblet, C.; Le Faou, P.; Angrand, P. O. Interaction
41
42 proteomics analysis of polycomb proteins defines distinct PRC1 complexes in mammalian cells.
43
44 *Mol. Cell Proteomics* **2011**, *10*, M110.002642-1-23.

45
46
47 14. Gil, J.; Bernard, D.; Martinez, D.; Beach, D. Polycomb CBX7 has a unifying role in
48
49 cellular lifespan. *Nat. Cell Biol.* **2004**, *6*, 67-72.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
15. Ren, X.; Vincenz, C.; Kerppola, T. K. Changes in the distributions and dynamics of polycomb repressive complexes during embryonic stem cell differentiation. *Mol. Cell Biol.* **2008**, *28*, 2884-2895.
16. O'Loghlen, A.; Munoz-Cabello, A. M.; Gaspar-Maia, A.; Wu, H. A.; Banito, A.; Kunowska, N.; Racek, T.; Pemberton, H. N.; Beolchi, P.; Lavial, F.; Masui, O.; Vermeulen, M.; Carroll, T.; Graumann, J.; Heard, E.; Dillon, N.; Azuara, V.; Snijders, A. P.; Peters, G.; Bernstein, E.; Gil, J. MicroRNA regulation of Cbx7 mediates a switch of Polycomb orthologs during ESC differentiation. *Cell Stem Cell* **2012**, *10*, 33-46.
17. Klauke, K.; Radulovic, V.; Broekhuis, M.; Weersing, E.; Zwart, E.; Olthof, S.; Ritsema, M.; Bruggeman, S.; Wu, X.; Helin, K.; Bystrykh, L.; de Haan, G. Polycomb Cbx family members mediate the balance between haematopoietic stem cell self-renewal and differentiation. *Nat. Cell Biol.* **2013**, *15*, 353-362.
18. Bernard, D.; Martinez-Leal, J. F.; Rizzo, S.; Martinez, D.; Hudson, D.; Visakorpi, T.; Peters, G.; Carnero, A.; Beach, D.; Gil, J. CBX7 controls the growth of normal and tumor-derived prostate cells by repressing the Ink4a/Arf locus. *Oncogene* **2005**, *24*, 5543-5551.
19. Wigle, T. J.; Herold, J. M.; Senisterra, G. A.; Vedadi, M.; Kireev, D. B.; Arrowsmith, C. H.; Frye, S. V.; Janzen, W. P. Screening for inhibitors of low-affinity epigenetic peptide-protein interactions: an AlphaScreen-based assay for antagonists of methyl-lysine binding proteins. *J. Biomol. Screening* **2010**, *15*, 62-71.
20. Burton, P. S.; Conradi, R. A.; Ho, N. F. H.; Hilgers, A. R.; Borchardt, R. T. How structural features influence the biomembrane permeability of peptides. *J. Pharm. Sci.* **1996**, *85*, 1336-1340.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
21. Ren, C.; Morohashi, K.; Plotnikov, A. N.; Jakoncic, J.; Smith, S. G.; Li, J.; Zeng, L.; Rodriguez, Y.; Stojanoff, V.; Walsh, M.; Zhou, M. M. Small-molecule modulators of methyl-lysine binding for the CBX7 chromodomain. *Chem. Biol.* **2015**, *22*, 161-168.
22. Hopkinson, R. J.; Walport, L. J.; Munzel, M.; Rose, N. R.; Smart, T. J.; Kawamura, A.; Claridge, T. D.; Schofield, C. J. Is JmjC oxygenase catalysis limited to demethylation? *Angew Chem. Int. Ed. Engl.* **2013**, *52*, 7709-7713.
23. Ren, C.; Smith, S. G.; Yap, K.; Li, S.; Li, J.; Mezei, M.; Rodriguez, Y.; Vincek, A.; Aguilo, F.; Walsh, M. J.; Zhou, M. M. Structure-guided discovery of selective antagonists for the chromodomain of polycomb repressive protein CBX7. *ACS Med. Chem. Lett.* **2016**, *7*, 601-605.
24. Goldman, S. D.; Funk, R. S.; Rajewski, R. A.; Krise, J. P. Mechanisms of amine accumulation in, and egress from, lysosomes. *Bioanalysis* **2009**, *1*, 1445-1459.
25. Van Dyke, R. W.; Faber, E. D.; Meijer, D. K. Sequestration of organic cations by acidified hepatic endocytic vesicles and implications for biliary excretion. *J. Pharmacol. Exp. Ther.* **1992**, *261*, 1-11.
26. Copeland, R. A. The drug-target residence time model: a 10-year retrospective. *Nat. Rev. Drug Discovery* **2016**, *15*, 87-95.
27. Barnash, K. D.; Lamb, K. N.; Stuckey, J. I.; Norris, J. L.; Cholensky, S. H.; Kireev, D. B.; Frye, S. V.; James, L. I. Chromodomain ligand optimization via target-class directed combinatorial repurposing. *ACS Chem. Biol.* **2016**, DOI: 10.1021/acscchembio.6b00415.

TABLE OF CONTENTS GRAPHIC

