

# Metabolism of 1'- and 4-Hydroxymidazolam by Glucuronide Conjugation Is Largely Mediated by UDP-Glucuronosyltransferases 1A4, 2B4, and 2B7

Kyung-Ah Seo, Soo Kyung Bae, Young-Kil Choi, Chang Soo Choi, Kwang-Hyeon Liu, and Jae-Gook Shin

Department of Pharmacology and Pharmacogenomics Research Center, Inje University College of Medicine, Busan, Korea (K.-A.S., S.K.B., K.-H.L., J.-G.S.); and Departments of Clinical Pharmacology (S.K.B., J.-G.S.) and Surgery (Y.-K.C., C.S.C.), Inje University Busan Paik Hospital, Busan, Korea

Received July 5, 2010; accepted August 16, 2010

## ABSTRACT:

Midazolam undergoes oxidative hydroxylation by CYP3A to its metabolites, which are excreted mainly as glucuronidated conjugates into the urine. In this study, we examined the glucuronidation of hydroxymidazolam in human liver microsomes (HLMs) and characterized the UDP-glucuronosyltransferases (UGTs) involved in 1'- and 4-hydroxymidazolam glucuronidation. Among the 12 UGT isoforms tested, the *O*- and *N*-glucuronidation of 1'-hydroxymidazolam was mediated by UGT2B4/2B7 and 1A4, respectively. In contrast, the glucuronidation of 4-hydroxymidazolam was mediated by UGT1A4. Consistent with these observations, the UGT1A4 inhibitor hecogenin and the

UGT2B7 substrate diclofenac potentially inhibited the *N*- and *O*-glucuronidation of 1'-hydroxymidazolam in HLMs, respectively. A correlation analysis of UGT enzymatic activity and the formation rate of glucuronide metabolites from 1'- and 4-hydroxymidazolam in 25 HLMs showed that hydroxymidazolam glucuronidation is correlated with UGT1A4-mediated lamotrigine glucuronidation and UGT2B7-mediated diclofenac glucuronidation activity. Taken together, these findings indicate that UGT1A4, 2B4, and 2B7 are major isoforms responsible for glucuronide conjugate formation from 1'- and 4-hydroxymidazolam, which are the two major oxidative metabolites of midazolam.

## Introduction

Glucuronidation is an important detoxification and elimination pathway for endogenous and exogenous compounds, and it accounts for ~35% of all phase II drug metabolism (Evans and Relling, 1999). UDP-glucuronosyltransferases (UGTs) mediate glucuronidation. In humans, the UGT superfamily is composed of two families (UGT1 and UGT2) and three subfamilies (UGT1A, 2A, and 2B) (Mackenzie et al., 2005). The lone UGT1A locus on chromosome 2q37 encodes nine functional proteins: UGT1A1, 1A3, 1A4, 1A5, 1A6, 1A7, 1A8, 1A9, and 1A10. The nomenclature of UGT1A is based on the relative position of the first variable exon with respect to four common exons in the unspliced mRNA. The UGT2A subfamily has multiple loci located on chromosome 4q13 and comprises the following functional proteins: UGT2A1, 2B4, 2B7, 2B10, 2B11, 2B15, 2B17, and 2B28 (Guillemette, 2003).

This study was supported by the Korean Health 21 R&D Project, Ministry of Health, Welfare, and Family Affairs [Grant A030001]; and by the Ministry of Education, Science, and Engineering [Korean Science and Engineering Foundation Grant R13-2007-023-00000-0].

Article, publication date, and citation information can be found at <http://dmd.aspetjournals.org>.

doi:10.1124/dmd.110.035295.

Individual UGTs exhibit distinct, but overlapping, substrate selectivities and differ in their regulation of expression, genetic polymorphisms, and other factors known to influence the activity of drug-metabolizing enzymes in humans (Miners et al., 2002, 2004). Despite recognition of the importance of glucuronidation as a clearance and detoxification mechanism for drugs, environmental chemicals, and endogenous compounds, relatively few selective substrates and inhibitors useful for the phenotyping of glucuronidated compounds have been identified (Court, 2005).

Midazolam is a potent benzodiazepine with sedative activities. It is metabolized to 1'- and 4-hydroxymidazolam by CYP3A4 and CYP3A5 (Kronbach et al., 1989) and is further metabolized to glucuronide conjugates by UGT (Fig. 1). Glucuronide conjugates of 1'-hydroxymidazolam account for 60 to 70% of the midazolam dose (Heizmann and Ziegler, 1981). Zhu et al. (2008) recently identified an *N*-linked glucuronide conjugate of 1'-hydroxymidazolam by incubation with human liver microsomes (HLMs). Although the *O*- and *N*-glucuronidation of 1'-hydroxymidazolam is catalyzed by UGT2B4/2B7 and 1A4, respectively (Zhu et al., 2008), the specific hepatic UGT isoforms involved in 4-hydroxymidazolam glucuronidation have not been identified.

The objective of this study was to identify and kinetically characterize and compare the UGT isoforms responsible for 1'- and 4-hydroxymidazolam glucuronidation in vitro. Our data will

**ABBREVIATIONS:** UGT, UDP-glucuronosyltransferase; HLM, human liver microsome; UDPGA, uridine diphosphoglucuronic acid; LC, liquid chromatography; MS/MS, tandem mass spectrometry.




FIG. 1. Proposed pathway of midazolam metabolism in HLMs.

allow a better understanding of the factors affecting midazolam pharmacokinetics and drug interactions.

### Materials and Methods

**Chemicals and Reagents.** Midazolam, 1'-hydroxymidazolam, and 4-hydroxymidazolam were purchased from Ultrafine Chemical Co. (Manchester, UK). Alamethicin (from *Trichoderma viride*), uridine diphosphoglucuronic acid (UDPGA), saccharic acid-1, 4-lactone, diclofenac, hecogenin, and lamotrigine were obtained from Sigma-Aldrich (St. Louis, MO). The solvents were of high-performance liquid chromatography grade (Thermo Fisher Scientific, Waltham, MA); all other chemicals were of the highest quality available. Recombinant human UGT isoforms (UGT1A1, 1A3, 1A4, 1A6, 1A7, 1A8, 1A9, 1A10, 2B4, 2B7, 2B15, and 2B17) and pooled and 10 single-donor HLMs were purchased from BD Gentest (Woburn, MA). The vendor supplied information regarding the protein concentration. Fifteen single-donor HLMs were obtained from the Inje Pharmacogenomics Research Center Tissue Bank (Busan, Korea).

**In Vitro Metabolism of Hydroxymidazolam by HLMs.** For the microsomal phase II metabolism of hydroxymidazolam, 0.25 mg of pooled HLMs (H161; BD Gentest), 1'- or 4-hydroxymidazolam (100  $\mu$ M), and 1.5 mg/ml alamethicin were reconstituted in 0.5 M Tris-HCl buffer (pH 7.5) and preincubated on ice for 15 min. The reaction was initiated by addition of UDPGA (5 mM) and run for 60 min at 37°C in a shaking water bath. In this study, a 60-min incubation time was selected because the reaction showed linearity until 60 min from the preliminary experiments. The reaction was terminated by the addition of 100  $\mu$ l of acetonitrile on ice and then centrifuged at 10,000g for 5 min at 4°C. Aliquots of the supernatant were analyzed by liquid chromatography/tandem mass spectrometry (LC-MS/MS) for the identification of glucuronide conjugate metabolites of 1'- and 4-hydroxymidazolam.

**Analysis of the Glucuronide Conjugates of Hydroxymidazolam by LC-MS/MS.** For the identification of the glucuronide metabolites of hy-

droxymidazolam, an API 3000 LC-MS/MS system (Applied Biosystems, Foster City, CA), coupled with an 1100 series high-performance liquid chromatography system (Agilent Technologies, Santa Clara, CA) was used. Separation was performed on a Polar-RP column (2 mm i.d.  $\times$  250 mm, 4- $\mu$ m particle size; Phenomenex, Torrance, CA) with a mobile phase consisting of water (A) and acetonitrile (B) containing 0.1% formic acid. The analytes were eluted at a flow rate of 0.2 ml/min using a linear gradient. Solvent B started at 30%, then increased as follows: 30% (1 min), 50% (6–8 min), and 30% (9 min). The TurboIonSpray interface was operated in the positive ion mode at 5500 V and 500°C. The operating conditions were as follows: nebulizing gas flow, 7 psi; curtain gas flow, 12 psi; and collision energy, 35 eV. Quadrupoles Q1 and Q3 were set on unit resolution. Multiple-reaction-monitoring mode using specific precursor/product ion transition was used for quantification. Detection of the ions was performed by monitoring the transitions of  $m/z$  518  $\rightarrow$  324 and 518  $\rightarrow$  342 for the hydroxymidazolam glucuronides and 339  $\rightarrow$  163 for 7-hydroxycoumarin glucuronide (internal standard). Peak areas for all compounds were automatically integrated using Analyst software (version 1.2; Applied Biosystems).

**Glucuronidation of Hydroxymidazolam in HLMs or Recombinant UGT Isoforms.** The optimal conditions for microsomal incubation were determined in the linear range for the formation of glucuronide metabolites from hydroxymidazolam. In all experiments, 1'- and 4-hydroxymidazolam were dissolved and serially diluted with methanol to the required concentrations. The final concentration of organic solvent did not exceed 1% in the final incubation mixtures. Typical incubation mixtures (total volume, 125  $\mu$ l), containing either 0.25 mg of HLMs or recombinant UGT (0.1 mg), 1'- or 4-hydroxymidazolam (1–200  $\mu$ M), and 1.5 mg/ml alamethicin, were reconstituted in 0.5 M Tris-HCl buffer (pH 7.5) and preincubated on ice for 15 min. The reaction was initiated by adding UDPGA (5 mM) and run for 60 min at 37°C in a shaking water bath. The reaction was terminated by the addition of 100  $\mu$ l of acetonitrile on ice, and the mixture was centrifuged

at 10,000g for 5 min at 4°C. Aliquots of the supernatant were used for LC-MS/MS.

**Chemical Inhibition Studies with HLMs.** The inhibitory effects of the known UGT1A4 inhibitor hecogenin (final concentration, 10  $\mu$ M) or the UGT2B7 substrate diclofenac (50  $\mu$ M) on the formation of 1'- and 4-hydroxymidazolam glucuronides were evaluated to determine the UGT isoforms involved in this metabolic pathway. Hecogenin was dissolved in dimethyl sulfoxide and diclofenac was dissolved in methanol. The final concentration of the organic solvents in the reaction mixture was <1% (v/v). The formation ratios of 1'- and 4-hydroxymidazolam glucuronides from 1'- and 4-hydroxymidazolam were determined from reaction mixtures incubated in the presence or absence of hecogenin or diclofenac. Except for the addition of hecogenin or diclofenac, all other incubation conditions were similar to those described previously.

**Correlation Experiments.** For the correlation analysis, the comparative metabolic rates of hydroxymidazolam in 25 different HLMs were investigated by the incubation of 1'- and 4-hydroxymidazolam (50  $\mu$ M) with 0.25 mg/ml microsomal protein for 60 min. The activity of each UGT isoform was determined by LC-MS/MS as described previously for lamotrigine (Rowland et al., 2006) and diclofenac glucuronidation (King et al., 2001). The correlation coefficients between the formation rates of glucuronide metabolites of 1'- and 4-hydroxymidazolam and the activity of each UGT isoform in the various HLMs were calculated by Spearman's correlation coefficient analysis (SAS version 8.01; SAS Institute Inc., Cary, NC).  $p < 0.05$  was considered to be statistically significant.

**Data Analysis.** All results are expressed as the mean  $\pm$  S.D. of estimates obtained from three different HLMs in triplicate experiments. Because of the absence of authentic standards for the hydroxymidazolam glucuronides, the conjugates were semiquantified using a hydroxymidazolam calibration curve. The apparent kinetic parameters for the biotransformation of 1'- and 4-hydroxymidazolam ( $K_m$  and  $V_{max}$ ) were determined by fitting an one-enzyme Michaelis-Menten equation or a Hill equation ( $V = V_{max} \cdot [S]^n / (K_m + [S]^n)$ ). The calculated parameters include the maximum rate of formation ( $V_{max}$ ), the Michaelis-Menten constant (apparent  $K_m$ ), the intrinsic clearance ( $CL_{int} = V_{max}/\text{apparent } K_m$ ), and Hill coefficient ( $n$ ). All calculations were performed using WinNonlin software (Pharsight, Mountain View, CA). The percent inhibition was calculated from the ratio of the amount of metabolite formed with and without the specific inhibitor.

## Results

**Identification of Glucuronide Metabolites of Hydroxymidazolam in HLMs.** After the incubation of 1'-hydroxymidazolam with HLMs in the presence of UDPGA, two metabolites were profiled. These metabolites were confirmed as glucuronide conjugates by full-scan analysis, which showed the addition of 176 Da (glucuronic acid) to the parent ( $m/z$  518). These two metabolites were tentatively identified as 1'-hydroxymidazolam *O*- and *N*-glucuronide based on the MS/MS fragmentation pattern. A fragment ion at  $m/z$  324 (loss of glucuronic acid and a water molecule) was the base peak observed with 1'-hydroxymidazolam *O*-glucuronide, whereas that at  $m/z$  342 (loss of glucuronic acid) was the major fragment ion for 1'-hydroxymidazolam *N*-glucuronide (Fig. 2). These results are consistent with those from previous reports (Zhu et al., 2008). When 4-hydroxymidazolam was incubated with the HLMs, two glucuronide conjugates ( $m/z$  518) were detected. On the basis of the structures of the 1'-hydroxymidazolam *O*- and *N*-glucuronides, which were reported by Zhu et al. (2008), it is reasonable to assume that glucuronic acid was conjugated to 4-hydroxymidazolam at the same N and OH positions. These two metabolites were designated as 4-hydroxymidazolam glucuronide I and II because we could not distinguish 4-hydroxymidazolam *O*-glucuronide from the *N*-glucuronide.

**Identification of the UGT Isoforms Involved in the Metabolism of Hydroxymidazolams.** Two concentrations of 1'- and 4-hydroxymidazolam (20 and 100  $\mu$ M) were incubated with a panel of recombinant UGT isoforms. The formation rates of hydroxymidazo-




FIG. 2. Multiple-reaction-monitoring (MRM) chromatogram of glucuronide metabolites obtained by LC-MS/MS using HLM incubates of 1'- (A) and 4-hydroxymidazolam (B) in the presence of UDPGA.

lam glucuronides after the incubation of 1'- or 4-hydroxymidazolam with the UGT isoforms are shown in Fig. 3. UGT1A4 and 2B4/2B7 metabolized 1'-hydroxymidazolam to 1'-hydroxymidazolam *N*- and *O*-glucuronide, respectively, more efficiently than any other UGT (Fig. 3, A and B), whereas UGT1A4 metabolized 4-hydroxymidazolam to its glucuronide metabolites with little contribution from UGT1A3, 1A6, and 2B7 (Fig. 3, C and D). Hydroxymidazolam glucuronide formation from 1'- and 4-hydroxymidazolam (50  $\mu$ M) was also studied using a UGT isoform inhibitor or substrate (Fig. 4). 1'-Hydroxymidazolam *N*-glucuronidation was inhibited by the UGT1A4 inhibitor hecogenin ( $IC_{50} = 1.5 \mu$ M), whereas 1'-hydroxymidazolam *O*-glucuronidation was inhibited by the UGT2B7 substrate diclofenac ( $IC_{50} = 2.0 \mu$ M). The formation of 4-hydroxymidazolam glucuronide I and II was also inhibited by hecogenin ( $IC_{50} = 0.8 \mu$ M) and diclofenac ( $IC_{50} = 46.6 \mu$ M), respectively (Table 1). The formation rates of 1'-hydroxymidazolam *N*-glucuronide and 4-hydroxymidazolam glucuronide I were significantly correlated with the activity of UGT1A4, whereas that of 1'-hydroxymidazolam *O*-glucuronide was correlated with the activity of UGT2B7 and 1A4 (Fig. 5).

**Enzyme Kinetic Analysis.** Kinetic analyses of 1'- and 4-hydroxymidazolam metabolite formation were performed using three different HLM preparations. Under the experimental conditions used, the metabolism of 1'- and 4-hydroxymidazolam by the HLMs was best fitted by a Hill equation (Fig. 6). The fitting of the data to the Michaelis-Menten two-enzyme model did not significantly improve the regression, compared with fitting of the data to a one-enzyme model. The kinetic parameters estimated from the three HLMs are shown in Table 2. The sums of the formation ( $CL_{int}$ ) of both metabolites were 346.2 and 90.3  $\mu$ l/(min  $\cdot$  mg protein) for 1'- and 4-hydroxymidazolam, respectively (Table 2).

Next, we examined the enzyme kinetic parameters for the formation of glucuronide conjugates from each hydroxymidazolam




FIG. 3. Representative plots of the formation of 1'-hydroxymidazolam *N*-glucuronide (A) and *O*-glucuronide (B) from 1'-hydroxymidazolam, from 4-hydroxymidazolam glucuronides I (C) and II (D) and from 4-hydroxymidazolam by recombinant human UGT isoforms. IS, internal standard.

(0–200 μM) upon incubation with recombinant human UGT1A4, 2B4, and 2B7 (Fig. 7). Under the experimental conditions used, the metabolism of 1'- and 4-hydroxymidazolam by these UGT isoforms was also best described by a Hill equation. The UGT1A4-

mediated *N*-glucuronidation of 1'-hydroxymidazolam showed a lower  $K_m$  and higher  $V_{max}$  than did UGT2B4- and 2B7-catalyzed *O*-glucuronidation, resulting in a higher  $CL_{int}$  [29.5 versus 6.8 μl/(min · mg protein)] (Table 3). The in vitro intrinsic clearance for




FIG. 4. Effects of the UGT isoform-selective inhibitor hecogenin (10 μM, UGT1A4) or the substrate diclofenac (50 μM, UGT2B7) on the formation of glucuronide metabolites from 1'-hydroxymidazolam (A) and 4-hydroxymidazolam (B) by HLMs. Each bar represents the mean of duplicate determinations.

TABLE 1

<i>IC<sub>50</sub> values for the inhibition of UGT-mediated hydroxymidazolam glucuronidation by the UGT isoform-selective inhibitor or substrate in pooled HLMs</i>					
UGT Isoform	Inhibitor	IC <sub>50</sub> Values <sup>a</sup>			
		1'-Hydroxymidazolam <i>N</i> -glucuronide	1'-Hydroxymidazolam <i>O</i> -glucuronide	4-Hydroxymidazolam Glucuronide I	4-Hydroxymidazolam Glucuronide II
<i>μM</i>					
1A4	Hecogenin	1.5	>5	0.8	>5
2B7	Diclofenac	802.8	2.0	941.1	46.6

<sup>a</sup> Averages of duplicate determinations.






FIG. 5. Simple regression analysis of lamotrigine or diclofenac glucuronidation and the rates of 1'-hydroxymidazolam *O*-glucuronidation (A) and *N*-glucuronidation (B) and 4-hydroxymidazolam glucuronide I formation (C) in 25 different HLMs.

glucuronide conjugate formation from 1'-hydroxymidazolam by UGT1A4 was similar to that for 4-hydroxymidazolam by UGT1A4 [29.5 versus 26.1  $\mu\text{L}/(\text{min} \cdot \text{mg protein})$ ].

### Discussion

The glucuronidation of midazolam and its oxidized metabolites is an important part of midazolam metabolism in humans. It has been reported that the amount of hydroxymidazolam *O*-glucuronide excreted in the urine is quite variable (Heizmann and Ziegler, 1981). This variability could be one factor responsible for the large interindividual differences seen in midazolam metabolism. In the present study, the glucuronidation of 1'- and 4-hydroxymidazolam in HLMs and the recombinant UGT isoforms mediating this metabolism was thoroughly evaluated.

In this study, we reconfirmed that 1'-hydroxymidazolam is metabolized to *N*- and *O*-glucuronide in the presence of UDPGA in HLMs and that UGT1A4 and 2B4/2B7 are the major enzymes responsible for 1'-hydroxymidazolam *N*- and *O*-glucuronidation, respectively. Recombinant human UGT2B4/2B7 and 1A4 metabolized 1'-hydroxymidazolam to 1'-hydroxymidazolam *O*- and *N*-glucuronide, respectively, whereas the other UGT isoforms did not (Fig. 3). The formation rates of 1'-hydroxymidazolam *N*-glucuronide were strongly inhibited by hecog-

nin (10  $\mu\text{M}$ , ~90%), a potent UGT1A4-selective inhibitor (Fig. 4) (Uchaipichat et al., 2006), whereas those of 1'-hydroxymidazolam *O*-glucuronide were inhibited by diclofenac (50  $\mu\text{M}$ , ~95%), a selective UGT2B7 substrate (King et al., 2001). We found a significant correlation between 1'-hydroxymidazolam *N*-glucuronidation in HLMs ( $n = 25$ ) and the glucuronidation activity of the UGT1A4 substrate lamotrigine (Rowland et al., 2006), whereas 1'-hydroxymidazolam *O*-glucuronidation was correlated with glucuronidation activity mediated by the UGT2B7 substrates diclofenac (King et al., 2001) and lamotrigine. The significant correlation observed between the activity of UGT1A4 and 1'-hydroxymidazolam *O*-glucuronidation in the panel of HLMs tested may not directly involve UGT1A4. Because our inhibition and recombinant experiments do not support a significant role for UGT1A4 in 1'-hydroxymidazolam *O*-glucuronidation, the observed correlation may be derived from the involvement of UGT2B7 in lamotrigine glucuronidation (Rowland et al., 2006). These results are compatible with those from a previous report by Zhu et al. (2008).

In addition, we demonstrated for the first time that 4-hydroxymidazolam is transformed to two glucuronide metabolites (I and II). Formation of the major metabolite, 4-hydroxymidazolam glucuronide I, was catalyzed by UGT1A4 but not by the other UGT isoforms




FIG. 6. Representative Michaelis-Menten (left panels) and Eadie-Hofstee (right panels) plots of the glucuronide conjugation of 1'- (A and B) and 4-hydroxymidazolam (C and D). HLMs were incubated with 0 to 200  $\mu$ M 1'-hydroxymidazolam or 4-hydroxymidazolam at 37°C for 60 min in the presence of UDPGA (5 mM). The data points represent the averages obtained from three different HLMs.

(Fig. 3C). The other isoforms, including UGT1A3, 1A6, 2B4, and 2B7, contributed to the formation of the minor metabolite 4-hydroxymidazolam glucuronide II (Fig. 3D). We also found that the formation rates of 4-hydroxymidazolam glucuronide I were strongly inhibited by hecogenin (10  $\mu$ M, ~95%) (Fig. 4). In addition, diclofenac, a well known UGT2B7 substrate, inhibited 4-hydroxymidazolam glucuronide II formation (50  $\mu$ M, ~50%). Our data significantly correlated 4-hydroxymidazolam glucuronide I formation in HLMs with UGT1A4-mediated lamotrigine *N*-glucuronidation (Fig. 5). Taken together, these results suggest that 4-hydroxymidazolam glucuronide I is the major glucuronide metabolite of 4-hydroxymidazolam and was formed by UGT1A4.

The glucuronidation of 1'- and 4-hydroxymidazolam in HLMs was characterized by Hill kinetics, suggesting the involvement of a single enzyme or more than one enzyme with similar affinities (Fig. 6). Eadie-Hofstee plots of 1'-hydroxymidazolam glucuronide formation




FIG. 7. Representative Michaelis-Menten plots for glucuronide conjugate formation from 1'-hydroxymidazolam (A and B) and 4-hydroxymidazolam (C) by recombinant UGT isoforms 1A4, 2B4, and 2B7. The recombinant UGT isoforms (0.25 mg/ml) were incubated with 0 to 100  $\mu$ M 1'- or 4-hydroxymidazolam at 37°C for 60 min in the presence of UDPGA (5 mM). Each data point represents the average of triplicate incubations.

showed convex relationships (Fig. 6B), indicating positive cooperativity ( $n > 1$ ), whereas 4-hydroxymidazolam glucuronide I and II formation showed convex ( $n = 1.2$ ) and concave ( $n = 0.5$ ) relationships (Fig. 6D), respectively. The intrinsic clearance for 1'-hydroxymidazolam *O*-glucuronidation was roughly 3-fold higher than that for *N*-glucuronidation (Table 2). A similar kinetic profile was reported by Zhu et al. (2008), although the difference in intrinsic clearance for metabolite formation was higher than observed in this study. This may be accounted for by differences between the assay conditions in their study (100 mM phosphate buffer, 2 mM  $MgCl_2$ , 0.5 mg/ml HLMs, and 2 mM UDPGA) and in the present study. A different kinetic profile for nicotine *N*-glucuronidation has also been shown to be dependent on the incubation conditions (Ghosheh et al., 2001; Nakajima et al., 2002). In the current study, 4-hydroxymidazolam glucuronide I formation was a major metabolic pathway for 4-hydroxymidazolam glucuronidation [the  $CL_{int}$  values for 4-hydroxymidazolam glucuronide I and II formation were 87.0 and 3.3  $\mu$ L/(min  $\cdot$  mg protein), respectively]. In contrast to 4-hydroxymidazolam, 1'-hydroxymidazolam showed high enzyme affinity (low  $K_m$  values), resulting in relatively high metabolic clearance (Table 2).

We observed large interindividual variability in 1'-hydroxymidazolam *O*-glucuronidation (roughly 9-fold), *N*-glucuronidation (~20 fold), and 4-hydroxymidazolam glucuronide I formation (approximately 13-fold) in the HLMs ( $n = 25$ ). Large interindividual differences in the percentage of glucuronide conjugates excreted in the urine may be due to interindividual differences in the formation of these metabolites (i.e., differences in the catalytic properties of the UGT isoforms, rather than in the process of excretion) (Nakajima et al., 2002). There are genetic polymorphisms in certain UGT isoforms, including UGT1A1, 1A4, 1A6, 2B4, 2B7, and 2B15 (Tukey and Strassburg, 2000). Polymorphisms in the UGT1A4, 2B4, and 2B7

TABLE 2

Mean enzyme kinetic parameters for glucuronide conjugate formation from 1'- and 4-hydroxymidazolam in HLMs

The values are the means of estimates from three different HLM preparations.

	Kinetic Parameter			
	$K_m$	$V_{max}$	Hill Coefficient ( $n$ )	$CL_{int}$ ( $V_{max}/K_m$ )
	$\mu$ M	pmol/(min $\cdot$ mg protein)		$\mu$ L/(min $\cdot$ mg protein)
1'-Hydroxymidazolam				
<i>N</i> -Glucuronide	9.1	790.0	1.6	86.8
<i>O</i> -Glucuronide	12.7	3285.8	1.5	259.4
4-Hydroxymidazolam				
Glucuronide I	41.5	3609.9	1.2	87.0
Glucuronide II	976.9	3210.5	0.5	3.3

TABLE 3

Mean enzyme kinetic parameters for glucuronide conjugate formation from 1'- and 4-hydroxymidazolam by recombinant human UGT isoforms

Each data set is the average of triplicate incubations.

UGT Isoform	Metabolite	Kinetic Parameters			
		$K_m$	$V_{max}$	Hill Coefficient (n)	$CL_{int} (V_{max}/K_m)$
	$\mu M$	$pmol/(min \cdot mg \text{ protein})$		$\mu l/(min \cdot mg \text{ protein})$	
1'-Hydroxymidazolam					
1A4	N-Glucuronide	15.9	468.3	1.31	29.5
2B4	O-Glucuronide	29.8	127.0	1.00	4.3
2B7		35.5	90.1	0.83	2.5
4-Hydroxymidazolam					
1A4	Glucuronide I	40.8	1064.2	1.23	26.1

isoforms might cause the interindividual differences seen in 1'- and 4-hydroxymidazolam glucuronidation in humans.

In conclusion, we thoroughly characterized the glucuronidation of 1'- and 4-hydroxymidazolam in HLMs. Using UGT isoform-specific inhibitors, correlation analyses, and recombinant UGT isoforms, we found that UGT1A4 and 2B4/2B7 were responsible for the formation of 1'-hydroxymidazolam N- and O-glucuronide, respectively, from 1'-hydroxymidazolam. In addition, the glucuronidation of 4-hydroxymidazolam was mainly metabolized by UGT1A4. The available data suggest that UGT enzymes exhibit overlapping substrate selectivity (Burchell et al., 1995; King et al., 2000). Therefore, identifying the substrate of each UGT isoform could be useful in UGT phenotyping. For this purpose, 4-hydroxymidazolam could be used in vitro as a probe substrate for UGT1A4 as has been suggested for 1'-hydroxymidazolam (Zhu et al., 2008).

### References

- Burchell B, Brierley CH, and Rance D (1995) Specificity of human UDP-glucuronosyltransferases and xenobiotic glucuronidation. *Life Sci* **57**:1819–1831.
- Court MH (2005) Isoform-selective probe substrates for *in vitro* studies of human UDP-glucuronosyltransferases. *Methods Enzymol* **400**:104–116.
- Evans WE and Relling MV (1999) Pharmacogenomics: translating functional genomics into rational therapeutics. *Science* **286**:487–491.
- Ghosheh O, Vashishtha SC, and Hawes EM (2001) Formation of the quaternary ammonium-linked glucuronide of nicotine in human liver microsomes: identification and stereoselectivity in the kinetics. *Drug Metab Dispos* **29**:1525–1528.
- Guillemette C (2003) Pharmacogenomics of human UDP-glucuronosyltransferase enzymes. *Pharmacogenomics J* **3**:136–158.
- Heizmann P and Ziegler WH (1981) Excretion and metabolism of  $^{14}C$ -midazolam in humans following oral dosing. *Arzneimittelforschung* **31**:2220–2223.
- King C, Finley B, and Franklin R (2000) The glucuronidation of morphine by dog liver microsomes: identification of morphine-6-O-glucuronide. *Drug Metab Dispos* **28**:661–663.
- King C, Tang W, Ngui J, Tephly T, and Braun M (2001) Characterization of rat and human UDP-glucuronosyltransferases responsible for the *in vitro* glucuronidation of diclofenac. *Toxicol Sci* **61**:49–53.
- Kronbach T, Mathys D, Umeno M, Gonzalez FJ, and Meyer UA (1989) Oxidation of midazolam and triazolam by human liver cytochrome P450III<sub>A4</sub>. *Mol Pharmacol* **36**:89–96.
- Mackenzie PI, Bock KW, Burchell B, Guillemette C, Ikushiro S, Iyanagi T, Miners JO, Owens IS, and Nebert DW (2005) Nomenclature update for the mammalian UDP glycosyltransferase (UGT) gene superfamily. *Pharmacogenet Genomics* **15**:677–685.
- Miners JO, McKinnon RA, and Mackenzie PI (2002) Genetic polymorphisms of UDP-glucuronosyltransferases and their functional significance. *Toxicology* **181–182**:453–456.
- Miners JO, Smith PA, Sorich MJ, McKinnon RA, and Mackenzie PI (2004) Predicting human drug glucuronidation parameters: application of *in vitro* and *in silico* modeling approaches. *Annu Rev Pharmacol Toxicol* **44**:1–25.
- Nakajima M, Tanaka E, Kwon JT, and Yokoi T (2002) Characterization of nicotine and cotinine N-glucuronidations in human liver microsomes. *Drug Metab Dispos* **30**:1484–1490.
- Rowland A, Elliot DJ, Williams JA, Mackenzie PI, Dickinson RG, and Miners JO (2006) *In vitro* characterization of lamotrigine N<sub>2</sub>-glucuronidation and the lamotrigine-valproic acid interaction. *Drug Metab Dispos* **34**:1055–1062.
- Tukey RH and Strassburg CP (2000) Human UDP-glucuronosyltransferases: metabolism, expression, and disease. *Annu Rev Pharmacol Toxicol* **40**:581–616.
- Uchaipichat V, Mackenzie PI, Elliot DJ, and Miners JO (2006) Selectivity of substrate (trifluoperazine) and inhibitor (amitriptyline, androsterone, canrenoic acid, hecogenin, phenylbutazone, quinidine, quinine, and sulfapyrazole) “probes” for human UDP-glucuronosyltransferases. *Drug Metab Dispos* **34**:449–456.
- Zhu B, Bush D, Doss GA, Vincent S, Franklin RB, and Xu S (2008) Characterization of 1'-hydroxymidazolam glucuronidation in human liver microsomes. *Drug Metab Dispos* **36**:331–338.

**Address correspondence to:** Dr. J.-G. Shin, Department of Pharmacology and Pharmacogenomics Research Center, Inje University College of Medicine, #633-165 Gaegum-Dong, Busanjin-Gu, Busan 614-735, Korea. E-mail: phshinjg@inje.ac.kr