



A Journal of the Gesellschaft Deutscher Chemiker

Angewandte Chemie

GDCh

International Edition

www.angewandte.org

Accepted Article

Title: A General and Highly Selective Palladium-Catalyzed Hydroamidation of 1,3-Diynes

Authors: Jiawang Liu, Carolin Schneider, Ji Yang, Zhihong Wei, Haijun Jiao, Robert Franke, Ralf Jackstell, and Matthias Beller

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: *Angew. Chem. Int. Ed.* 10.1002/anie.202010768

Link to VoR: <https://doi.org/10.1002/anie.202010768>

RESEARCH ARTICLE

A General and Highly Selective Palladium-Catalyzed Hydroamidation of 1,3-Diynes

Jiawang Liu,^[a] Carolin Schneider,^[a] Ji Yang,^[a] Zhihong Wei,^[a,b] Haijun Jiao,^[a,*] Robert Franke,^[c] Ralf Jackstell,^[a] and Matthias Beller^[a,*]

[a] Dr. J. Liu,[†] C. Schneider,[†] Dr. J. Yang, Dr. Z. Wei, Dr. H. Jiao, Dr. R. Jackstell, Prof. Dr. M. Beller
Leibniz-Institut für Katalyse e.V., Albert-Einstein-Str. 29a, Rostock, 18059, Germany
E-mail: Haijun.Jiao@catalysis.de; matthias.beller@catalysis.de

[b] Dr. Z. Wei
Institute of Molecular Science, Key Laboratory of Materials for Energy Conversion and Storage of Shanxi Province, Shanxi University, Taiyuan 030006, P. R. China

[c] Prof. Dr. R. Franke
Evonik Performance Materials GmbH, Paul-Baumann-Str. 1, 45772 Marl, Germany and Lehrstuhl für Theoretische Chemie, Ruhr-Universität Bochum, 44780 Bochum, Germany

[†] These authors contributed equally to this work.

Supporting information for this article is given via a link at the end of the document.

Abstract: A chemo-, regio- and stereoselective mono-hydroamidation of (un)symmetrical 1,3-diynes is described. Key for the success of this novel transformation is the utilization of an advanced palladium catalyst system with the specific ligand **L6** (Neolephos). The synthetic value of this general approach to synthetically useful α -alkynyl- α , β -unsaturated amides is showcased by diversification of several structurally complex molecules and marketed drugs. Control experiments and density functional theory (M06L-SMD) computations also suggest the crucial role of the substrate in controlling the regioselectivity of unsymmetrical 1,3-diynes.


Introduction

Transition metal catalyzed carbonylation reactions belong to the important examples of industrially applied homogeneous catalytic reactions. Since their discovery, they became a powerful tool for the synthesis of numerous value-added carbonyl containing compounds.^[1] Thus, carbonylation reactions are frequently applied in organic synthesis due to the versatility of the carbonyl groups and the possibility to easily expand carbon chains.^[2] Within this class of reactions, hydroamidations, also called aminocarbonylations, represent a straightforward route for the conversion of olefins or alkynes, carbon monoxide and amines into the corresponding amides. Compared to the well-explored aminocarbonylation of alkenes using catalytic systems based on cobalt,^[3a, 3b] nickel,^[3c] iron,^[3d] ruthenium,^[3e] rhodium^[3f] and mainly palladium,^[3g-3i] the related reactions of alkynes leading to α , β -unsaturated amides have received much less attention.


α , β -Unsaturated amides are valuable intermediates and building blocks in organic synthesis and are used as functional molecules in material science.^[4] In addition, they are occurring in natural products and bio-active pharmaceuticals.^[5] Traditionally, they are prepared by nucleophilic condensation of carboxylic acid derivatives with amines in the presence of stoichiometric amounts of activating or coupling reagents.^[6] Obviously, such processes result in significant waste formation. In this context, the aminocarbonylation of alkynes can be a more atom-economic and benign methodology. However, aminocarbonylations of alkynes (and olefins) are intrinsically challenging due to the problems associated with the acidity of the active metal hydride catalysts

and the basicity of the amine reagent. Moreover, alkynes are prone to side reactions such as hydroamination and oligomerization/polymerization. Nevertheless, by selecting appropriate catalysts and optimization of reaction conditions notable progress was achieved in the past decades.^[7]


(a) Aminocarbonylation of terminal alkynes



(b) Aminocarbonylation of unsymmetrical internal alkynes



(c) Unprecedented regioselective aminocarbonylation of unsymmetrical 1,3-diynes (this work)



Scheme 1. Selected catalytic hydroamidations of alkynes.

So far, almost all these reactions are restricted to terminal alkynes, which can be easily controlled to afford branched ^[7c-n,13] or linear^[7h,7m,7n-q] α , β -unsaturated amides (Scheme 1, a). In fact, only few aminocarbonylations of "simple" unsymmetrical internal alkynes were reported,^[7o,7s] leading to unsatisfactory regioselectivity (Scheme 1, b). Clearly, carbonylations of unsymmetrical internal alkynes are more demanding because of their low reactivity and the difficulty in achieving high regioselectivity. Notable exceptions include the recent works on hydroformylation and alkoxy carbonylation reported by the groups of Breit,^[8a] You,^[8b] and Ma,^[9] respectively. In these studies, regioselectivity could be controlled by hydrogen bonding

RESEARCH ARTICLE

interactions between substrates and catalysts, steric hinderance and/or chelation.

Based on our continuous interest in carbonylation reactions, recently we started to investigate the selective carbonylation of unsymmetrical 1,3-diyne, which can be conveniently generated via alkyne coupling processes^[10] and provide synthetically highly useful intermediates. Obviously, the selective hydroamidation of unsymmetrical 1,3-diyne compared to internal monoalkynes implies further challenges: 1) due to their higher reactivity competitive hydroamination also becomes potentially easier;^[11] 2) the two unbiased alkyne moieties of unsymmetrical diynes create more complex selectivity problems, including chemo-, stereo-, and regioselectivity.^[12] Thus, it is not surprising that to the best of our knowledge no such process has been reported, yet. In fact, only in 2018 Huang^[13] and co-workers reported the first aminocarbonylations using symmetrical 1,3-diyne. The need of a comparably high palladium catalyst loading (5 mol%), and relative harsh conditions (140 °C) offers also potential for improvements.


Herein, we present the first examples of regioselective Pd-catalyzed hydroamidation of unsymmetrical 1,3-diyne. Notably, the mild conditions can also be applied in various symmetrical substrates and a wide range of α -alkynyl- α , β -unsaturated amides are obtained in general in good yields with excellent selectivities (Scheme 1, c).

In the past three years, inspired by the seminal work of Drent and co-workers,^[14] we prepared several novel bidentate pyridyl-substituted phosphine ligands.^[15] Some of these ligands showed significantly improved performance for Pd-catalyzed alkoxy carbonylations of olefins or alkynes. Mechanistic studies revealed that the nitrogen atoms of the pyridyl groups act as proton shuttle to accelerate the nucleophilic attack on the intermediate Pd acyl complex and thereby increasing the rate of the overall transformation.^[16] We envisioned that these ligands might also improve the reactivity in Pd-catalyzed aminocarbonylations. Hence, in our initial studies we compared the influence of these and related ligands for the hydroamidation of 1-(5-(trimethylsilyl)penta-2,4-diyne-1-yl)pyrrolidine-2,5-dione (**1a**) with aniline at room temperature.

Results and Discussion

As shown in Scheme 2, applying 1,1'-ferrocenediyl-bis(*tert*-butyl(pyridin-2-yl)phosphine) **L1** a highly selective monocarbonylation (regioselectivity: >99/1) occurred leading to product **3aa**. Unfortunately, the reactivity of this system is too low (15% yield). In the presence of other pyridyl-substituted ligands such as 1,3-bis(*tert*-butyl(pyridin-2-yl)phosphanyl)-propane **L2**, 1,4-bis(*tert*-butyl(pyridin-2-yl)phosphanyl)butane **L3**, 1,2-bis((*tert*-butyl(pyridin-2-yl)phosphanyl)methyl)benzene **L4** and 2,2'-((2,7-di-*tert*-butyl-9,9-dimethyl-9H-xanthene-4,5-di-yl)bis-(*tert*-butylphosphanediy))dipyridine **L5**, under these mild conditions no conversion is observed, although most of these ligands have been proved to be active in Pd-catalyzed alkoxy carbonylations. Surprisingly, using 2,2'-bis(*tert*-butyl(pyridin-2-yl)phosphanyl)-1,1'-binaphthalene **L6** (Neolephos), the desired product **3aa** was afforded in 84% yield and with excellent regioselectivity (>99/1). In addition, the corresponding acid, which is generated via hydroxycarbonylation, was isolated as a side product in 16%. It should be noted that although ligand **L6** showed high

chemoselectivity for mono-alkoxy carbonylation of 1,3-diyne, it was not the most efficient ligand in Pd-catalyzed alkoxy carbonylation of dienes^[15d] or diyne^[15e]. Commercially available ligands **L7-L16**, including mono and bidentate phosphine ligands, which are commonly used in other carbonylations,^[17] were examined, too; however, in all cases no product was detected under the standard conditions. Comparison of **L6** with **L1-L5** demonstrated that the backbones of ligand, which may lead to different bite angles in Pd complexes, are also crucial for this transformation. To improve the benchmark reaction further, we evaluated the influence of critical reaction parameters in the presence of **L6** (for more details, see Table S7-S6 in SI). It is worth pointing out that there was no reaction without acid or in the presence of weak acid (HOAc or PhCO₂H, Table S2), indicating the importance of the strong acid for the generation of catalytically active Pd hydride species (Figure 1). As a result, the formation of the acid by-product could be prevented using triflic acid instead of PTSA·H₂O.




Scheme 2. Pd-catalyzed aminocarbonylation of unsymmetrical 1,3-diyne **1a**: Influence of phosphine ligands. Reaction conditions: **1a** (0.25 mmol), Pd(TFA)₂ (1.0 mol%), ligand (4.0 mol% for **L1-L12**, and 8.0 mol% for **L13-L16**), PTSA·H₂O (16.0 mol%), PhNH₂ (**2a**, 0.25 mmol), CO (40 atm), 23 °C and toluene (1.0 mL), 20 h. The ratio of **3aa/4aa** and yield were determined by GC and GC-MS analysis.

With optimized reaction conditions established, we set up to explore the scope of different unsymmetrical 1,3-diyne using **2a** as the nucleophile. As shown in Table 1, the catalytic system can be conveniently applied to various unsymmetrical 1,3-diyne derivatized from propargylamine or alcohol. In general, good to high yields (50-94%) were obtained with substrates based on succinimide, glutarimide or phthalimide derivatives (**3aa-3na**). While the corresponding free propargylic alcohol derivative gave no clean conversion, the acetylated substrates **1o-1q** reacted smoothly and gave the corresponding amides in 50-76% yields. Notably, in all these cases excellent regioselectivity (>99/1) was observed and exclusive formation of the *E*-regioisomers took

RESEARCH ARTICLE


place. In addition, the reaction of 1,3-diyne bearing two different aromatic substituents also proceeded well to afford the corresponding product in 86% yield with low regioselectivity (60/40) (Scheme S1). Apart from unsymmetrical 1,3-diyne, a variety of symmetrical substrates provided the corresponding α -alkynyl- α , β -unsaturated amides under very mild conditions, too (Table 2). In fact, without further optimization products **6aa-6ma** were obtained in 68-96% with excellent stereoselectivity. More specifically, aromatic 1,3-diyne **5a-5f** with either electron-donating (OMe, Me, ^tBu) or electron-withdrawing (F, CF₃) substituents on the phenyl ring provided **6aa-6fa** in high yields (70-96%) and excellent selectivity. Substituents in the *ortho*-position of the phenyl ring have no significant influence on both reactivity and selectivity of this reaction; thus, **6ia-6ja** were produced in 78-92% yield. In addition, the thiofuryl-substituted substrate was well tolerated by the catalyst to afford **6ma**. Next, the reactivity of aliphatic 1,3-diyne was investigated. Gratifyingly, also in these cases the palladium-catalyzed hydroamidation proceeded selectively, affording carbonylative products **6na-6wa** in 60-96% yields. It should be noted that the obtained products may undergo isomerization processes in the presence of palladium hydride species; but no side-products were observed in all cases. Furthermore, functional groups such as cyano, ester and trimethylsilyl survived in products **6ra** and **6ta-6wa**.

Table 1. Pd-catalyzed aminocarbonylation of unsymmetrical 1,3-diyne




Reaction conditions: **1** (0.25 mmol), Pd(acac)₂ (1.0 mol%), **L6** (2.0 mol%) HOTf (8.0 mol%), aniline (**2a**, 0.25 mmol), CO (40 atm), 40 °C in toluene (1.0 mL), 20 h. Yields of isolated products are shown.

Table 2. Pd-catalyzed aminocarbonylation of symmetrical 1,3-diyne^a



[a] Reaction conditions: **5** (0.25 mmol), Pd(acac)₂ (1.0 mol%), **L6** (2.0 mol%) HOTf (8.0 mol%), aniline (**2a**, 0.25 mmol), CO (40 atm), 40 °C in toluene (1.0 mL), 20 h. Yields of isolated products are shown. [b] 60 °C. [c] PTSA-H₂O (16.0 mol%) was used instead of HOTf.

Table 3. Pd-catalyzed aminocarbonylation of 1,3-diyne **1a** or **5a** with different arylamines^a




[a] Reaction conditions: **1a** or **5a** (0.25 mmol), Pd(acac)₂ (1.0 mol%), **L6** (2.0 mol%) HOTf (8.0 mol%), arylamines (**2**, 0.25 mmol), CO (40 atm), 40 °C in toluene (1.0 mL), 20 h. Yields of isolated products are shown. [b] 60 °C. [c] PTSA-H₂O (16.0 mol%) was used instead of HOTf.

Next, we evaluated the scope of this novel hydroamidation process with respect to the reactivity of amines using unsymmetrical (**1a**) and symmetrical (**5a**) 1,3-diyne as a standard coupling partner, respectively (Table 3). A variety of arylamines with electron-neutral, electron-deficient, and electron-rich

RESEARCH ARTICLE

substituents led to the corresponding carbonylative products in good yields (51-90%) and again with excellent regioselectivities (>99/1). Apparently, the substituents on the arylamines have no real impact on the catalysis. Specifically, the reactions of arylamines bearing Me, F, Cl, Ac, CN, CO₂Et, MeO and MeS substituents proceeded smoothly, providing the corresponding products **3ab-3ae**, **3ah-3ak** in good isolated yields.^[19] Notably, bromine- and even iodine-substituted arylamines, which are known to be sensitive to palladium catalysis, also worked well and afforded the corresponding products (**3af-3ag**). The position of substituents on the phenyl ring has no influence on the reaction outcome. Hence, arylamines **2n-2p** afforded **3an-3ap** in 81-89% yields. Interestingly, the 2-aminophenol (**2q**) and 2-aminobenzamide (**2r**), which contain two nucleophilic positions, also reacted selectively to give the amides **3aq** and **3ar** in 84% and 90% yield, respectively. Furthermore, the heterocycle-substituted amine **2s** proved to be a viable coupling partner and gave **3as** in 73% yield. While secondary aromatic amines (**2t** and **2u**) gave the desired products in good yields (**3at** and **3au**), in case of benzyl amine or *n*-butylamine no product was detected under these conditions. It should be noted that almost all these tested amines have similar behaviour in the hydroamidation of symmetrical 1,3-diyne **5a**, providing products **6ab-6aq** and **6at-6au** in 45-94% yield. Unfortunately, there was no conversion at all when aliphatic amines such as benzylic amine or *n*-pentylamine was used instead of aniline (Scheme S2), and this is probably due to the higher basicity of aliphatic amines than aniline.^[20]

Table 4. Pd-catalyzed aminocarbonylation of 1,3-diyne containing biologically or pharmaceutically relevant skeletons^a




[a] Reaction conditions: diynes (0.25 mmol), arylamines (0.25 mmol), Pd(acac)₂ (1.0 mol%), **L6** (2.0 mol%), HOTf (8.0 mol%), CO (40 atm), 40 °C in toluene (1.0 mL), 20 h. Yields of isolated products are shown and in all cases the regioselectivities are >99/1.

In general, our methodology *vide supra* allows to functionalize aromatic amines to the corresponding enyne amides in a straightforward manner under very mild conditions. We thought that this unusual transformation, which also increases the molecular complexity, can be of interest for many life science applications including the diversification of natural products and synthetic drugs. Notably, the resulting 1,3-enyne motif can be additionally modified in several ways.^[18] In this respect, the hydroamidation of several structurally more complex molecules

was studied to prove functional group tolerance and compatibility with bio-relevant derivatives (Table 4). We were particularly delighted to find that this Pd-catalyzed regioselective hydroamidation of 1,3-diyne progressed well with substrates **7a** and **7b** derivatized from uracil, one of the nucleobases of RNA, and furnished products **7aa** and **7ba** in high yields and excellent regioselectivity. Compound **7c**, containing dehydrocholic acid, which is a drug for stimulation of biliary lipid secretion, could also be applied to afford the amide product **7ca** in decent yield. Moreover, nonsteroidal anti-inflammatory drugs such as ibuprofen, indomethacin and isoxepac are viable components for 1,3-diyne substrates **7d-7f** in this transformation. Finally, the products **7at-7ct** were prepared from the corresponding secondary aniline.

To understand the observed regioselectivity and to gain more mechanistic insights, several control experiments with unsymmetrical 1,3-diyne were conducted. As shown in Scheme 3, the different substituents on the diyne unit control the regioselectivity of the carbonylation reaction to a significant extent. In general, the following selectivity order is observed: TMS ≤ prim. alkyl < aryl << CH₂OAc ≈ CH₂NR₂. Hence, testing **8** with our catalyst under standard conditions, aminocarbonylation proceeded preferentially at the phenyl-substituted alkyne group (regioselectivity: 70/30) (Eq. 1). However, applying propargylamine/alcohol derivatives, hydroamidations regioselectively took place at the triple bond substituted with these functional groups. As an example, the protected propargyl alcohol derivative **9** was synthesized and gave the amide product **9b** in 56% yield with >99/1 regioselectivity (Eq. 2), which demonstrates the crucial role of the oxygen atom for determining the regioselectivity.



Scheme 3. Pd-catalyzed hydroamidation of 1,3-diyne: Control experiments. Standard condition: diynes (0.25 mmol), Pd(acac)₂ (1.0 mol%), **L6** (2.0 mol%), HOTf (8.0 mol%), aniline (0.25 mmol), CO (40 atm), 40 °C in toluene (1.0 mL), 20 h. Yields of isolated products are shown and the ratio of regioisomers were determined by GC and GC-MS analysis as well as ¹H NMR analysis. [a] 70 °C.

Interestingly, the steric nature of the substituents on the alkyne also plays a decisive role. Hence, the regioselectivity is increased to 98/2 when trimethyl(phenylbuta-1,3-diyne-1-yl)silane **10** was used (Eq. 3) instead of **8**. In order to figure out, whether the

RESEARCH ARTICLE

carbonyl group plays a decisive role in the control of regioselectivity, the unsymmetrical 1,3-diyne **11** without nitrogen atom but with carbonyl group was prepared and tested. Here, product **11a** was obtained with a completely different regioselectivity (Eq. 4), which disproved the involvement of the carbonyl group in the determination of the regioselectivity.

To understand the experimentally observed regioselectivity further, density functional theory computations using different methods in gas phase as well as in solution with and without dispersion corrections were carried (see *Supporting Information* for more details). If not otherwise mentioned, we used the M06L/TZVP computed Gibbs free energies under the consideration of solvation effect (toluene) on the basis of the B3PW91/TZVP gas phase optimized geometries for discussion and other results are given in the Supporting Information for comparison. In our study we chose substrates **9** and **10** using the complex of palladium and ligand **L6** as active catalyst resulting in totally opposite regioselectivity. These results are further established with those of substrate **1a**. Due to the chirality of the ligand and the possible diastereomers of the formed complex, we used the C_2 -symmetrical (*R,R*)-ligand for our computation with the nitrogen atoms of the pyridyl groups in the Pd complex pointing towards the Pd center. This enables a hemilabile coordination of the ligand; and such complexation modes have been found in the molecular structures of similar complexes.^[16a]

Since the regioselectivity of metal-catalyzed carbonylation reactions of non-symmetrical C=C or C≡C bonds comes from the M-H insertion resulting in the formation of alkyl or alkenyl complex, we computed this elementary step to elucidate the origin of the regioselectivity. For the protonated active catalyst (Figure 1), there are two stable isomers, and the one with proton on the Pd center (**LPd-H**) is more stable than the one with proton on the N atom of one 2-pyridyl ring (**LPd-NH**) by 2.4 kcal/mol. Such small energy difference reveals the possibility of their mutual exchange upon the change of the coordination sphere. It is noted that **LPd-NH** is not stable and has been optimized directly to **LPd-H** at the MN15/TZVP level. In both complexes, one 2-pyridyl nitrogen is coordinated to the Pd center. With the coordination of substrate **9**, it shows that complexes with N-H functionality are more stable than those with Pd-H group; and the most stable complex has the C≡C coordination terminated with CH₂OTBS substituent, and that with Ph substituent is less stable by 3.7 kcal/mol. For the coordination of substrate **10**, the same energetic patterns have been found and the most stable complex has the C≡C coordination terminated with Ph substituent and that with TMS substituent is less stable by 2.7 kcal/mol. This energetic order reveals the increasing steric effect of CH₂OTBS, Ph and TMS substitutions.

Based on these complexes, we computed this elementary step and the simplified potential energy surface for the regioselective formation of the alkenyl complexes is shown in Figure 2. For substrate **9**, two transition states for the N-H insertion into the C≡C bond have been located, which differs from the traditional Pd-H insertion. The transition state with the sterically less hindered CH₂OTBS group at the C≡C bond (**LPd-NH-9-OTBSCH₂/C₂Ph-TS**) is more stable than that one with Ph group (**LPd-NH-9-Ph/C₂CH₂OTBS-TS**) at the C≡C bond by 3.0 kcal/mol, and this reveals a kinetic preference.

Following these transition states, the corresponding intermediates with C-H agostic interaction have been located (**LPd-9-**




Figure 1. Relative energy of the active catalysts as well as corresponding alkyne complexes of substrates **9** and **10**

OTBSCH₂-C-H/C₂Ph, -7.5 kcal/mol and **LPd-9-Ph-C-H/C₂CH₂OTBS**, -4.7 kcal/mol). It is interesting to note that the presence of the hemilabile 2-pyridyl ligand results in the more stable alkenyl intermediates (**LPd-9-OTBSCH₂-CH/C₂Ph** and **LPd-9-R₁-CH/C₂R₂**); and the former is also more stable than the latter by 2.5 kcal/mol, which indicates a thermodynamic performance. These energy differences rationalize the observed regioselectivity of substrate **9**, which is favored both kinetically and thermodynamically. For substrate **10**, different results have been found (Figure 2). For example, the transition state with sterically less hindered Ph group at the coordinated C-C bond is higher in energy than that with the sterically more hindered TMS group by 2.1 kcal/mol (**LPd-NH-10-Ph/C₂TMS-TS** vs. **LPd-NH-10-TMS/C₂Ph-TS**). Such reversed energetic difference is indeed understandable, since the C atom bonded to Ph is less negatively charged than that bonded to TMS (-0.241 vs. -0.725) in the transition states due to the difference in electronegativity of carbon and silicon atoms. Consequently, the former will have weaker attractive interaction than the latter. Next, it is not possible to locate the C-H agostic intermediate following **LPd-NH-10-Ph/C₂TMS-TS**, and all attempts resulted in the direct formation of the corresponding alkenyl complex **LPd-10-PhCH/C₂TMS**. In contrast, the corresponding C-H agostic intermediate (**LPd-10-**

RESEARCH ARTICLE




Figure 2. Potential energy surface for the regioselective formation of alkenyl complexes of **9** (top) and **10** (bottom).

TMS-C-H/C₂Ph) following **LPd-NH-10-TMS/C₂Ph-TS** was located (-5.0 kcal/mol). Subsequently, the more stable alkenyl intermediate (**LPd-10-TMSCH/C₂Ph**) was located. On the potential energy surface, the alkenyl intermediate **LPd-10-PhCH/C₂TMS** is more stable than **LPd-10-TMSCH/C₂Ph** (-17.8 vs. -12.5 kcal/mol) due to the enhanced steric difference between Ph and TMS, although the formation of the former is less favored kinetically than the latter formation. Consequently, the observed regioselectivity of substrate **10** should be governed

thermodynamically by considering the reversibility between (**LPd-NH-10-TMS/C₂Ph-TS**) and (**LPd-10-TMS-C-H/C₂Ph**).

A similar potential energy surface for substrate **1a** has been found (Figure S3). For the CH₂R terminated C≡C bond, the reaction has an insertion free energy barrier of 9.9 kcal/mol and goes directly to the corresponding alkenyl intermediate with exergonic reaction free energy of 20.4 kcal/mol. For the TMS terminated C≡C bond, the reaction has an insertion free energy barrier of 9.6 kcal/mol followed by an C-H agostic intermediate (-4.6 kcal/mol) and the

RESEARCH ARTICLE

reaction goes to the corresponding alkenyl intermediate with exergonic reaction free energy of 13.5 kcal/mol. This indicates once again the thermodynamic origin of the observed regioselectivity.

To analyze the steric effect of different substituents, we computed the relative energies of the products (Figure 3). For example, **9a** is more stable than **9b** by 1.7 kcal/mol, while **10a** is more stable than **10b** by 10.6 kcal/mol. Furthermore, **3aa** is more stable than **4aa** by 10.0 kcal/mol (Figure S3). To confirm this thermodynamic trend, we computed the relative energy of isomers with H substitution as reference and the substitution of CH₂OTBS, Ph and SiMe₃ is less favored by 1.7, 2.4 and 11.1 kcal/mol, respectively, indicating the increasing steric interaction. This energetic order is in line with the observed regioselectivity.




Figure 3. Relative energy (kcal/mol) of the regioisomers as well as the simplified reference molecules.

Conclusion

In summary, we have developed a general and convenient Pd-catalyzed hydroamidation of (un)symmetrical 1,3-diyne. For the first time differently substituted 1,3-diyne undergo highly chemo-, regio-, and stereoselective transformation to the corresponding α,β -unsaturated amides. This novel catalytic transformation was enabled by the “built-in-base” ligand (**L6**, Neolephos) under mild conditions and provided a general approach to a variety of interesting functionalized synthetic building blocks in good to high yields. The utility of the catalytic system is showcased in versatile modifications of several structurally complex molecules and marketed drugs. Mechanistic studies and M06L-SMD density functional theory computations revealed the key role of the intrinsic substituents of substrates and the ligand in determining the selectivity. We believe these findings will provide new impetus for other selectivity-controlled carbonylation reactions using unsymmetrical substrates.

Acknowledgements

This work is supported by Evonik Performance Materials GmbH, the BMBF (Bundesministerium für Bildung und Forschung), and the State of Mecklenburg-Vorpommern. We thank the analytical team of LIKAT for their kind support. J. Y. thanks the Chinese Scholarship Council (CSC) for financial support.


Keywords: catalysis • regioselectivity • hydroamidation • P ligand • amides

- [1] a) P. W. N. M. van Leeuwen, C. Claver, *Rhodium Catalyzed Hydroformylation*, Vol. 22, Springer, Netherlands, **2002**; b) *Catalytic Carbonylation Reactions* (Ed.: M. Beller), Springer, Berlin Heidelberg, **2006**; c) B. Breit, in *Metal Catalyzed Reductive C@C Bond Formation: A Departure from Preformed Organometallic Reagents* (Ed.: M. J. Krische), Springer, Berlin, Heidelberg, **2007**, pp. 139-172; d) *Modern Carbonylation Methods*, Wiley-VCH, Weinheim, **2008**; e) *Transition Metals for Organic Synthesis: Building Blocks and Fine Chemicals* (Eds.: M. Beller, C. Bolm), Wiley-VCH, Weinheim, **2008**; f) R. Franke, D. Selent, A. Bçrner, *Chem. Rev.* **2012**, *112*, 5675-5732.
- [2] a) X.-F. Wu, H. Neumann, M. Beller, *Chem. Soc. Rev.* **2011**, *40*, 4986-5009; b) X.-F. Wu, X. Fang, L. Wu, R. Jackstell, H. Neumann, M. Beller, *Acc. Chem. Res.* **2014**, *47*, 1041-1053; c) B. Sam, B. Breit, M. J. Krische, *Angew. Chem. Int. Ed.* **2015**, *54*, 3267-3274; d) S. D. Friis, A. T. Lindhardt, T. Skrydstrup, *Acc. Chem. Res.* **2016**, *49*, 594-605; e) J.-B. Peng, F.-P. Wu, X.-F. Wu, *Chem. Rev.* **2019**, *119*, 2090-2127.
- [3] a) P. Pino, P. Paleari, *Gazz. Chim. Ital.* **1951**, *81*, 64; b) S. I. Lee, S. U. Son, Y. K. Chung, *Chem. Commun.* **2002**, 1310-1311; c) W. Reppe, H. Main, *Chem. Abstr.* **1953**, *47*, 5428; d) A. Striegler, J. Weber, *J. Prakt. Chem.* **1965**, *29*, 281-295; e) Y. Tsuji, T. Ohsumi, T. Kondo, Y. Watanabe, *J. Organomet. Chem.* **1986**, *309*, 333-344; f) K. Dong, X. Fang, R. Jackstell, G. Laurenczy, Y. Li, M. Beller, *J. Am. Chem. Soc.* **2015**, *137*, 6053-6058; g) C. Jim8nez-Rodríguez, A. A. NflicÇez-Magro, T. Seidensticker, G. R. Eastham, M. R. L. Furst, D. J. Cole-Hamilton, *Catal. Sci. Technol.* **2014**, *4*, 2332-2339; h) H. Liu, N. Yan, P. J. Dyson, *Chem. Commun.* **2014**, *50*, 7848-7851; i) X. Fang, R. Jackstell, M. Beller, *Angew. Chem. Int. Ed.* **2013**, *52*, 14089-14093; j) G. Zhang, B. Gao, H. Huang, *Angew. Chem. Int. Ed.* **2015**, *54*, 7657-7661; k) T. Xu, F. Sha, H. Alper, *J. Am. Chem. Soc.* **2016**, *138*, 6629-6635; l) J. Liu, H. Li, A. Spannenberg, R. Franke, R. Jackstell, M. Beller, *Angew. Chem. Int. Ed.* **2016**, *55*, 13544-13548.
- [4] a) T. K. Hyster, T. Rovis, *Chem. Sci.* **2011**, *2*, 1606-1610; b) J.-H. Fan, W.-T. Wei, M.-B. Zhou, R.-J. Song, J.-H. Li, *Angew. Chem., Int. Ed.* **2014**, *53*, 6650-6654; c) X. Mu, T. Wu, H.-Y. Wang, Y.-L. Guo, G. Liu, *J. Am. Chem. Soc.* **2012**, *134*, 878-881; d) *The Amide Linkage: Selected Structural Aspects in Chemistry, Biochemistry and Materials Science*; A. Greenberg, C. M. Breneman, J. F. Liebman, Eds.; Wiley-VCH: New York, **2000**; e) M. J. Caulfield, G. G. Qiao, D. H. Solomon, *Chem. Rev.* **2002**, *102*, 3067-3084.
- [5] a) C. Marrano, P. de Macedo, J. W. Keillor, *Bioorg. Med. Chem.* **2001**, *9*, 1923-1928; b) C.-C. Hung, W.-J. Tsai, L.-M. Yang Kuo, Y.-H. Kuo, *Bioorg. Med. Chem.* **2005**, *13*, 1791-1797; c) S. Chaudhury, T. R. Welch, B. S. J. Blagg, *ChemMedChem* **2006**, *1*, 1331-1340.
- [6] a) E. Valeur, M. Bradley, *Chem. Soc. Rev.* **2009**, *38*, 606-631; b) V. J. Pattabiraman, J. W. Bode, *Nature*, **2011**, *480*, 471-479; c) J. M. Concellón, J.A. Pérez-Andrés, H. Rodríguez-Solla, *Angew. Chem., Int. Ed.* **2000**, *39*, 2773-2775; d) S. Kim, C. J. Lim, *Angew. Chem., Int. Ed.* **2004**, *43*, 5378-5380; e) M. K. Hadden, B. S. J. Blagg, *J. Org. Chem.* **2009**, *74*, 46974704.
- [7] For reviews, see: a) A. Brennführer, H. Neumann, M. Beller, *ChemCatChem*, **2009**, *1*, 28-41; b) S. Quintero-Duque, K. M. Dybala, I. Fleischer, *Tetrahedron Lett.* **2015**, *56*, 2634-2650; For selected examples, see: c) S. Torii, H. Okumoto, M. Sadakane, L. H. Xu, *Chem. Lett.* **1991**, *20*, 1673-1676; d) E. Drent, P. Arnoldy, P. H. M. Budzelaar, *J. Organomet. Chem.* **1993**, *455*, 247-253; e) B. El Ali, A. M. ElGhanam, M. Fettouhi, J. Tijani, *Tetrahedron Lett.* **2000**, *41*, 5761-5764; f) B. El Ali, J. Tijani, A. M. El-Ghanam, *Appl. Organomet. Chem.* **2002**, *16*, 369-376; g) B. El Ali, J. Tijani, A. M. El-Ghanam, *J. Mol. Catal. A: Chem.* **2002**, *187*, 17-33; h) B. El Ali, J. Tijani, *Appl. Organomet. Chem.* **2003**, *17*, 921-931; i) Y. Li, H. Alper, Z. Yu, *Org. Lett.* **2006**, *8*, 5199-5201; j) S.-M. Lu, H. Alper, *J. Am. Chem. Soc.* **2008**, *130*, 6451-6455; k) R. Suleiman, J. Tijani, B. El Ali, *Appl. Organomet. Chem.* **2010**, *24*, 38-46; l) H. Liu, G. P. S. Lau, P. J. Dyson, *J. Org. Chem.* **2015**, *80*, 386-391; m) F. Sha, H. Alper, *ACS Catal.* **2017**, *7*, 2220-2229; n) D.-L. Wang, W.-D. Guo, L. Liu, Q. Zhou, W.-Y. Liang, Y. Lu, Y. Liu, *Catal. Sci. Technol.*, **2019**, *9*, 1334-1337; o) K. M. Driller, S. Prateptongkum, R. Jackstell, M. Beller, *Angew. Chem. Int. Ed.* **2011**, *50*, 537-541; p) M. Pizzetti, A. Russo, E. Petricci, *Chem. Eur. J.* **2011**, *17*, 4523-4528; q) Z. Huang, Y. Dong, Y. Li, M.

RESEARCH ARTICLE

- Makha, Y. Li, *ChemCatChem*, **2019**, *11*, 5236-5240; r) T. Sugihara, Y. Okada, M. Yamaguchi, M Nishizawa, *Synlett*, **1999**, *6*, 768-770; s) J. H. Park, S. Y. Kim, S. M. Kim, Y. K. Chung, *Org. Lett.* **2007**, *9*, 2465-2468.
- [8] a) W. Fang, B. Breit, *Angew. Chem. Int. Ed.*, **2018**, *57*, 14817-14821; b) G. Tan, Y. Wu, Y. Shi, J. You, *Angew. Chem. Int. Ed.* **2019**, *58*, 7440-7444. For other regioselective hydroformylation of internal alkynes, see: c) B. G. Van den Hoven, H. Alper, *J. Org. Chem.* **1999**, *64*, 9640-9646; d) B. G. Van den Hoven, H. Alper, *J. Org. Chem.* **1999**, *64*, 3964-3968; e) J. R. Johnson, G. D. Cuny, S. L. Buchwald, *Angew. Chem. Int. Ed. Engl.* **1995**, *34*, 1760-1761.
- [9] C. Huang, H. Qian, W. Zhang, S. Ma, *Chem. Sci.*, **2019**, *10*, 5505-5512.
- [10] a) C. Glaser, *Ber. Dtsch. Chem. Ges.* **1869**, *2*, 422-424; b) C. Glaser, *Ann. Chem. Pharm.* **1870**, *154*, 137-171; c) A. S. Hay, *J. Org. Chem.* **1960**, *25*, 1275-1276; d) A. Lei, M. Srivastava, X. Zhang, *J. Org. Chem.* **2002**, *67*, 1969-1971; e) Y. Nishihara, K. Ikegashira, K. Hirabayashi, J. -i. Ando, M. Mori, T. Hiyama, *J. Org. Chem.* **2000**, *65*, 1780-1787; f) W. Yin, C. He, M. Chen, H. Zhang, A. Lei, *Org. Lett.*, **2009**, *11*, 709-712.
- [11] a) T. Shimada, Y. Yamamoto, *J. Am. Chem. Soc.*, **2002**, *124*, 12670-12671; b) D. H. Camacho, S. Saito, Y. Yamamoto, *Tetrahedron Lett.*, **2002**, 1085-1088.
- [12] a) Z. Yan, X.-A. Yuan, Y. Zhao, C. Zhu, J. Xie, *Angew. Chem. Int. Ed.* **2018**, *57*, 12906-12910; b) J. Liu, J. Yang, W. Baumann, R. Jackstell, M. Beller, *Angew. Chem. Int. Ed.* **2019**, *58*, 10683-10687; c) S. Cembellin, T. Dalton, T. Pinkert, F. Schäfers, F. Glorius, *ACS Catal.* **2020**, *10*, 197-202; d) H. L. Sang, C. Wu, G. G. D. Phua, S. Ge, *ACS Catal.*, **2019**, *9*, 10109-10114.
- [13] X. Ji, B. Gao, X. Zhou, Z. Liu, H. Huang, *J. Org. Chem.*, **2018**, *83*, 10134-10141.
- [14] a) E. Drent, P. Arnoldy, H. M. Budzelaar, *J. Organomet. Chem.*, **1993**, *455*, 247-253; b) E. Drent, P. Arnoldy, P. H. M. Budzelaar, *J. Organomet. Chem.* **1994**, *475*, 57-63.
- [15] Several ligands were developed for different reactions, **L4**: a) K. Dong, X. Fang, S. Gülak, R. Franke, A. Spannenberg, H. Neumann, R. Jackstell, M. Beller, *Nat. Commun.* **2017**, *8*, 14117; **L1**: b) K. Dong, R. Sang, X. Fang, R. Franke, A. Spannenberg, H. Neumann, R. Jackstell, M. Beller, *Angew. Chem. Int. Ed.* **2017**, *56*, 5267-5271; **L2** and **L3**: c) J. Liu, K. Dong, R. Franke, H. Neumann, R. Jackstell, M. Beller, *J. Am. Chem. Soc.* **2018**, *140*, 10282-10288; HeMaRaphos: d) J. Yang, J. Liu, H. Neumann, R. Franke, R. Jackstell, M. Beller, *Science*, **2019**, *366*, 1514-1517; **L6**: e) J. Liu, J. Yang, C. Schneider, R. Frank, R. Jackstell, M. Beller, *Angew. Chem. Int. Ed.* **2020**, *59*, 9032-9040.
- [16] a) K. Dong, R. Sang, Z. Wei, J. Liu, R. Dühren, A. Spannenberg, H. Jiao, H. Neumann, R. Jackstell, R. Franke, M. Beller, *Chem. Sci.* **2018**, *9*, 2510-2516; b) L. Crawford, D. J. Cole-Hamilton, E. Drent, M. Bühl, *Chem.-Eur. J.*, **2014**, *20*, 13923-13926; c) L. Crawford, D. J. Cole-Hamilton, M. Bühl, *Organometallics*, **2015**, *34*, 438-449.
- [17] a) G. Vasapollo, A. Scarpa, G. Mele, L. Ronzini, B. El Ali, *Appl. Organomet. Chem.* **2000**, *14*, 739-743; b) C. J. Rodriguez, D. F. Foster, G. R. Eastham, D. J. Cole-Hamilton, *Chem. Commun.*, **2004**, 1720-1721; c) T. Xu, H. Alper, *J. Am. Chem. Soc.* **2014**, *136*, 16970-16973; d) P. Roesle, L. Caporaso, M. Schnitte, V. Goldbach, L. Cavallo, S. Mecking, *J. Am. Chem. Soc.*, **2014**, *136*, 16871-16881; e) P. W. van Leeuwen, P. C. Kamer, *Catal. Sci. Technol.* **2018**, *8*, 26-113; f) J. Y. Wang, A. E. Strom, J. F. Hartwig, *J. Am. Chem. Soc.* **2018**, *140*, 7979-7993; g) S. J. Byrne, A. J. Fletcher, P. Hebeisen, M. C. Willis, *Org. Biomol. Chem.* **2010**, *8*, 758-760; h) J. K. Stille, P. K. Wong, *J. Org. Chem.* **1975**, *40*, 532-534; i) S. Klaus, H. Neumann, A. Zapf, D. Strübing, S. Hübner, J. Almena, T. Riermeier, P. Groß, M. Sarich, W.-R. Krahnert, K. Rossen, M. Beller, *Angew. Chem. Int. Ed.* **2006**, *45*, 154-158.
- [18] a) D. Kang, J. Kim, S. Oh, P. H. Lee, *Org. Lett.* **2012**, *14*, 5636-5639; b) H. Qian, X. Yu, J. Zhang, J. Sun, *J. Am. Chem. Soc.* **2013**, *135*, 18020-18023; c) Q. Yao, Y. Liao, L. Lin, X. Lin, J. Ji, X. Liu, X. Feng, *Angew. Chem., Int. Ed.* **2016**, *55*, 1859-1863; d) P. H. Poulsen, Y. Li, V. H. Lauridsen, D. K. B. Jørgensen, T. A. Palazzo, M. Meazza, K. A. Jørgensen, *Angew. Chem., Int. Ed.* **2018**, *57*, 10661-10665; e) Z.-G. Ma, J.-L. Wei, J.-B. Lin, G.-J. Wang, J. Zhou, K. Chen, C.-A. Fan, S.-Y. Zhang, *Org. Lett.* **2019**, *21*, 2468-2472; see also reference 15e and the references therein.
- [19] CCDC 1998320 (**3ah**) and 2001174 (**3ai**) contain the supplementary crystallographic data for this paper. These data are provided free of charge by the joint Cambridge Crystallographic Data Centre and Fachinformationszentrum Karlsruhe Access Structures service www.ccdc.cam.ac.uk/structures.
- [20] P. L. H. Edward, G. L. Sharon, *J. Phys. Chem. Ref. Data*, **1998**, *27*, 413-686.

RESEARCH ARTICLE



Select one of the twins: The advanced Pd catalyst system with **L6** (Neolephos) as ligand enables highly selective hydroamidation of unbiased (un)symmetrical 1,3-diyne. In this way, a wide range of synthetically useful α -alkynyl- α,β -unsaturated amides are afforded in good to high yields with excellent chemo-, regio-, and stereoselectivities.