

Visible-Light-Driven Stereoselective Annulation of Alkyl Anilines and Dibenzoylthylenes via Electron Donor–Acceptor Complexes

August Runemark, Savannah C. Zacharias, and Henrik Sundén*



Cite This: *J. Org. Chem.* 2021, 86, 1901–1910



Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information

ABSTRACT: A catalyst-free, stereoselective visible-light-driven annulation reaction between alkenes and *N,N*-substituted dialkyl anilines for the synthesis of substituted tetrahydroquinolines is presented. The reaction is driven by the photoexcitation of an electron donor–acceptor (EDA) complex, and the resulting products are obtained in good to high yields with complete diastereoselectivity. Mechanistic rationale and photochemical characterization of the EDA-complex are provided.



INTRODUCTION

Electron donor–acceptor (EDA) complexes have in recent years gained considerable attention in the field of organic chemistry due to their photochemical properties which can be used to mediate a number of advanced chemical transformations.^{1–5} An EDA complex is the result of a weak association between an electron-rich donor and an electron-deficient acceptor and is characterized by a charge-transfer band in the absorption spectrum.⁴ Due to the bathochromic shift of the charge-transfer band, with respect to the donor and acceptor, visible light can often be employed to induce a single electron transfer from the donor to the acceptor. The resulting radical pair can be used in synthetic chemistry,^{1–4,6–12} including enantioselective alkylations,^{13–17} aromatic alkylations,¹⁶ and biaryl couplings.¹⁸


Recently, we reported an EDA-complex-driven protocol for the UV-light-induced generation of α -aminoalkyl radicals and their addition to maleimides to form fused tetrahydroquinolines (THQs) (Scheme 1a).¹⁹

The THQ is a highly desirable structural target in synthesis, as it can be found in a variety of biologically active compounds.^{20,21} Examples include molecules with antiviral,^{22–24} antibiotic,^{25,26} and cytotoxic^{27,28} activity. Thus, a diverse set of methodologies for the synthesis of the THQ-scaffold has been developed, and among the most versatile strategies are the aza-Diels–Alder, the Grieco, and the Povarov reactions.^{22–24}


Additionally, several methods for the visible-light-driven construction of substituted THQ have been reported (Scheme 1b,c).^{29–34} However, the use of photocatalysts is typically required, and substrates are generally limited to cyclic alkenes (Scheme 1a,c) or alkenes that cannot undergo photoisomerization (Scheme 1b).^{35–37} To the best of our knowledge, no diastereoselective annulations between tertiary amines and alkenes that can undergo photoisomerization have been reported under photochemical conditions. Clearly, the possibility of alkene *E/Z*-isomerization under photochemical

Scheme 1. Light-Induced Construction of Tetrahydroquinolines

Previous work:



This work:



conditions poses a significant synthetic challenge for selectivity in the desired cyclization. But if successful, it could lead to the diastereoselective functionalization of THQ in the 3 and 4 positions (Scheme 1d). Inspired by these opportunities, we

Received: November 25, 2020

Published: January 5, 2021



envisioned that the more challenging acyclic internal alkenes could function as acceptors in novel EDA complexes with aromatic amines for the synthesis of substituted THQ (Scheme 1d) and that the cumbersome photoisomerization could be kept to a minimum with a light source operating in the visible region of the light spectrum.

Herein, we present a catalyst-free EDA-mediated diastereoselective synthesis of substituted THQ from tertiary amines and 1,2-dibenzoyl ethylene (DBE).

RESULTS AND DISCUSSION

To initiate our study, the interaction between 4'-N,N-trimethylaniline (**1a**) and 1,2-DBE (**2a**) was investigated (Figure 1). UV-vis measurements of **1a**, **2a**, and a mixture of




Figure 1. (A) Photos of **1a**, **2a**, and **1a** + **2a** in acetonitrile. (B) UV-vis absorption spectra of **1a** (0.1 M), **2a** (0.1 M), and **1a** + **2a** in acetonitrile.

the two compounds were performed. Upon mixing, a color change from light yellow to orange was observed (Figure 1A), and a new absorption band appeared in the visible region (Figure 1B), indicating an EDA interaction.

Next, the impact of visible light irradiation of a mixture of **1a** and **2a** was studied. Compact fluorescent light bulbs (CFLs) were used as the irradiation source (SI, Figure S2 for emission spectrum). The desired cyclized product **3a** was formed in 25% yield using acetonitrile as solvent (Table 1, entry 1). 1,4-Dioxane proved to be the best solvent for the reaction, providing the desired THQ in 65% yield (Table 1, entry 8). Other solvents, both polar and nonpolar, showed inferior results, correlating with previously published findings.^{19,38} The impact of irradiation time on the formation of **3a** was investigated using gas chromatography, and a reaction time of 4 h was determined to be optimal (SI, Figure S5). Significant degradation of the desired product was observed after prolonged irradiation. A large stoichiometric excess of the amine was proven to be important, as using 4 molar equiv lowered the yield significantly (Table 1, entry 9). This can be rationalized as a result of increased concentration of the photoactive EDA complex with higher loadings of the donor ($K_{\text{EDA}} = 0.42 \text{ M}^{-1}$, calculated using the Benesi-Hildebrand method, Figure S3). The role of the oxidant, oxygen, was then evaluated. When the reaction mixture was irradiated under inert atmosphere, the reaction was suppressed (Table 1, entry 10). Irradiating under an atmosphere of oxygen gas on the other hand resulted in a low yield and a sluggish reaction (Table 1, entry 11). Changing the oxidant to persulfate resulted in decreased yield (Table 1, entry 13). Addition of acetic acid increased the yield of **3a** to 73% (Table 1, entry 16), a result that correlates with the literature.³⁹ Alternative


Table 1. Screening of Reaction Conditions^a

entry	solvent	light source	yield of 3a (%) ^b	d.r. ^b
1	acetonitrile	CFL	25	>25:1
2	methanol	CFL	14	>25:1
3	tetrahydrofuran	CFL	29	>25:1
4	ethyl acetate	CFL	17	>25:1
5	dichloromethane	CFL	40	>25:1
6	1,2-dichloroethane	CFL	41	>25:1
7	1,2-dimethoxyethane	CFL	32	>25:1
8	1,4-dioxane	CFL	65	>25:1
9	1,4-dioxane	CFL	30 ^c	>25:1
10	1,4-dioxane	CFL	0 ^d	—
11	1,4-dioxane	CFL	12 ^e	>25:1
12	1,4-dioxane	CFL	37 ^f	>25:1
13	1,4-dioxane/water (2:1)	CFL	47 ^f	>25:1
14	1,4-dioxane	blue LED	23	>25:1
15	1,4-dioxane	CFL	68 ^g	>25:1
16	1,4-dioxane	CFL	73 ^h	>25:1
17	1,4-dioxane	UV-CFL	30	>25:1
18	1,4-dioxane	—	0 ⁱ	—
19	1,4-dioxane	blue LED	60 ^{h,j}	>25:1

^aReaction conditions: **1a** (0.7 mmol) and **2a** (0.1 mmol) in 3 mL of solvent irradiated with two 15 W compact fluorescent lamps in room temperature for 4 h. ^bDetermined by NMR with 1,2,4,5-tetramethylbenzene as internal standard. ^c4 equiv of amine. ^dUnder Ar. ^eUnder O₂. ^fK₂S₂O₈ (2 equiv) used as an additive. ^gAcetic acid (30 equiv) used as additive. ^hAcetic acid (80 equiv) used as additive. ⁱReaction performed in absence of light. ^jReaction time of 12 h.

light sources, such as blue LED (450 nm) or UV-CFL (365 nm) performed worse than the regular CFL (Table 1, compare entries 14 and 17 with 8). However, the use of blue LED under the optimized reaction conditions (1,4-dioxane as solvent and acetic acid as additive) and longer reaction time provide the desired product in the yield of 60% (Table 1, entry 19). The exclusion of light resulted in complete suppression of the formation of the desired product, confirming the need for an excitation source (Table 1, entry 18).

With our optimized reaction conditions in hand, the generality of the reaction was examined (Scheme 2). Excellent diastereoselectivity toward the *anti*-diastereomer (confirmed by X-ray analysis, entry 3q) was obtained for all substrates, and the *syn*-isomer of **3** was never observed. Symmetric DBEs with simple aliphatic substituents provided the corresponding THQs in moderate yields in combination with 4'-N,N-trimethylaniline (Scheme 2, 3b–e). A clear effect of the electronic properties of substituents in the *p*-position on the 1,2-DBE can be observed as the yield decreases when more electron-donating substituents are introduced: *p*-H 73%, *p*-Me 63%, *i*-Bu 52%, and *tert*-Bu 45% (Scheme 2, 3a–3d). The introduction of two methyl groups in *m*- and *p*-position likewise decreased the yield (Scheme 2, 3e) as well as *p*-OMe (Scheme 2, 3f). Mildly σ -withdrawing groups such as *p*-Br or *p*-Cl resulted in no change or a slight increase in the yield (Scheme 2, 3h–i). When an unsymmetrical dibenzoyl ethylene was used as the substrate, the product was obtained as an inseparable mixture of the two regioisomers (Scheme 2, 3j).


Scheme 2. Substrate Scope of Anilines and Dibenzoylthylenes^{a,b}

^aReaction conditions: Substituted 1,2-DBE (0.25 mmol) and aniline (1.75 mmol) in 1,4-dioxane (6 mL) and acetic acid (1 mL) was irradiated with CFL lamp for 4 h. ^bn.d. = not determined.

Next, the effect of substituents on the aniline reaction partner was evaluated. Electron-donating groups proved to be well tolerated (Scheme 2, **3n**) compared to σ -withdrawing groups such as *p*-F or *p*-Br (Scheme 2, **3p–q**). Introduction of a methoxy group in *m*-position resulted in a significantly decreased the yield (Scheme 2, **3o**). A significant steric effect was observed when a methyl group was introduced in the *o*-position on the aniline, as the reaction was completely

suppressed (Scheme 2, **3i**). This result could be explained by a significantly weaker EDA complex formed between the reaction partners, something also indicated by the lack of color change upon mixing. The steric clash between the *o*-Me and the *N*-Me groups leads to a less planar and less conjugated aromatic amine weakening the interaction with the alkene. Introduction of two methyl groups in the less sterically demanding *m*-positions did not lead to a similar decrease in yield (Scheme 2, **3m**). We were also interested in investigating the regioselectivity of the reaction regarding the aniline reactant. Changing one of the *N*-methyl substituents to aliphatic in benzylic groups resulted in a complete selectivity toward reaction of the *N*-methyl group (Scheme 2, **3s–u**). When *N,N*-diethylaniline was used, no reaction took place (Scheme 2, **3w**). These results highlight the selectivity of the reaction toward *N*-methyl substituted anilines. However, subjecting monosubstituted *N*-methyl aniline to the reaction conditions resulted in complete suppression of reactivity (Scheme 2, **3v**). The impact of geometrical isomerism of the DBE substrate was then examined. It is reported that *E*-DBE undergoes isomerization to the *Z*-isomer under visible light irradiation.⁴⁰ In order to evaluate the impact of this background reaction, *Z*-DBE was subjected to the reaction conditions to furnish the desired product **3a**, albeit in a lower yield of 55% (Scheme 3). Complete diastereoselectivity toward


Scheme 3. Impact of Stereochemistry of the Substrate Dibenzoylthylene



the *anti*-diastereomer was observed, suggesting that the cyclization reaction is not of concerted character. Notably, other acyclic activated olefins were not tolerated in the reaction such as benzylideneacetophenone, benzylidenemalonitrile, cinnamaldehyde, nitrostyrene, diethylfumarate, or diethylmaleate. These olefins did not provide the corresponding desired annulation products under the developed conditions.


To further probe the impact of light on the reaction, the illumination was cycled on/off, resulting in suppression of product formation in the absence of light (SI, Figure S6). This affirms the fact that the reaction requires constant illumination. In addition, the quantum yield of the reaction was determined to be 4.5 (SI, Section 8), which suggests that a radical chain process is involved in the mechanism. The role of oxygen was also investigated. The observation that the reaction is suppressed in the absence of oxygen (Scheme 1, entry 10) and the evidence for presence of hydrogen peroxide in the reaction mixture after irradiation (SI, Figure S4), demonstrates the role of ambient oxygen as an external oxidant. Based on these findings and support in the literature,^{19,41–43} we propose the following mechanism (Scheme 4): The reaction commences with the formation of an EDA complex between reactants **1a** and **2a**. Upon irradiation with light, an electron transfer occurs, a key step which is supported by calculation of the Gibbs free energy change for the photoinduced electron transfer (SI Section 1.8). Subsequent proton-transfer results in the formation of the α -amino alkyl radical **A** and enol radical **B**. The enol radical **B** reacts with oxygen, regenerating **2a** and

Scheme 4. Proposed Reaction Mechanism



forming a hydroperoxyl radical and thus starting the self-propagating radical chain mechanism. In the next step, **2a** reacts with α -amino alkyl radical **A** to form **C**, and after cyclization, the radical intermediate **D** is formed. In the final step, **D** is oxidized to the desired product **3a** via two possible alternative pathways. The cyclohexadienyl radical intermediate **D** is oxidized by molecular oxygen^{41,42} yielding **3a** and a second hydroperoxyl radical which in turn can propagate a radical chain reaction by oxidizing **1a** to **A**.^{43,44} Alternatively, **D** can be intercepted by a hydroperoxyl radical yielding **3a** and hydrogen peroxide as a byproduct.


In order to improve the step economy of the reaction, we were interested in the possibility of running the annulation as a multicomponent version with **2a** formed *in situ*. The one-pot version would circumvent a purification step and give potential access to a broader scope of the target compounds (**3**) while using simpler starting materials. As it turns out, the reaction between **1a**, phosphonium ylide **4**, and phenylglyoxal **5** works well, and **3a** could be obtained in 40% yield (Scheme 5), which corresponds to a yield per bond formed of 80%.

Scheme 5. Multicomponent Synthesis of **3a**

To investigate the synthetic usefulness of the 3,4-dibenzoyl-THQ structure, compound **3a** was treated with ammonium acetate in acetic acid⁴⁵ to yield fused 3H-pyrrole derivative **6** (Scheme 6). Subjecting **3a** to acid in acetic anhydride afforded fused furan derivative **7**.⁴⁶ These results highlight the possibility of efficient construction of fused heterocycles from the 3,4-dibenzoyl-THQ.

In summary, a protocol for the visible-light-mediated synthesis of substituted THQs has been developed. The reaction requires no photocatalyst and relies on photo-excitation of an EDA complex formed between *N*-alkyl-*N*-

Scheme 6. Synthetic Applications



^a NH_4OAc , AcOH, 120 °C, 5 h. ^b Ac_2O , HCl, 80 °C, 18 h.

methylaniline and a 1,2-dibenzoyl ethylene, where atmospheric oxygen functions as the terminal oxidant. A broad substrate scope is presented, demonstrating the tolerance for common functional groups in the reaction. The resulting 3,4-dibenzoyl-THQ structure is further derivatized and proven to be a useful building block for the construction of fused heterocycles. Synthetic applications of other EDA complexes are currently being explored by our research group.

EXPERIMENTAL SECTION

General Information. All reagents and solvents were purchased from Sigma-Aldrich and Alfa Aesar and used without any further purification unless specified. Purification of products was performed by an automated column chromatography Biotage Isolera Spektra One with Biotage SNAP-10 g KP-silica column together with a 1 g sample cartridge using petroleum ether (40–60 °C)/ethyl acetate as the solvent mixture unless otherwise noted. ¹H (400 MHz) and ¹³C (101 MHz) NMR spectra were acquired on an Agilent NMR machine at 25 °C. The chemical shifts for ¹H and ¹³C NMR spectra are reported in parts per million (ppm) relative to the residual peak from solvent CDCl_3 as the internal standard: ¹H NMR at δ 7.26 ppm and ¹³C NMR at δ 77.0 ppm for CDCl_3 . All coupling constants (*J*) are reported in hertz (Hz) and multiplicities are indicated by s (singlet), d (doublet), dd (doublet of doublet), td (triplet of doublet), ddd (doublet of doublets of doublets), triplet (t), dt (doublet of triplet) and m (multiplet). Fourier-transform infrared (FT-IR) spectra were recorded on a PerkinElmer series FT-IR spectrometer and are reported in wavenumber (cm^{-1}). High-resolution mass spectrometry measurements (HRMS) were performed by CMSI service at Chalmers University of Technology, Gothenburg using an Agilent 6520 equipped with an electrospray interface operated in the positive ionization mode with quadrupole time-of-flight mass analyzer. UV–vis absorption spectra were recorded on a Cary 4000 UV–vis

spectrophotometer, using 1×1 cm quartz cuvettes. All light promoted reactions were carried out in Biotage microwave vials (10–20 mL) under irradiation with two 20 W white CFL bulbs (Osram, 1200 lm) at a distance of 5 cm. Gas chromatographic studies were carried out using an Agilent 7820A gas chromatograph with an Agilent HP-5 19091J-413 column, and detection was accomplished using a flame ionization detector. Crystals of **3q** for single-crystal X-ray diffraction were grown using the layering technique from **3q** in dichloromethane and fresh hexane. The colorless yellow prism-shaped crystals appeared after 4 days. A Bruker D8 VENTURE Kappa Duo with a PHOTON III detector was used for the data collection. Collections were carried out at low temperature [120(2) K] using a Cryostream SAINT (version 7.60a; Bruker AXS, 2016) software to perform data reduction and unit cell refinement. The atomic coordinates were located using direct methods employed by SHELXS.⁴⁷ The successive refinements, once the atoms were placed in their postulated positions, were made using SHELXL.⁴⁸ X-Seed⁴⁹ was used for the data refinement. All non-hydrogen atoms were then refined anisotropically. The hydrogen atoms were placed in idealized positions in a riding model, after location on a Fourier difference map. An isotropic refinement was used for all hydrogen atoms, and temperature factors of 1.2 or 1.5 times that of the parent atoms were assigned.

Synthesis of Starting Materials. Amines **1i–k** were synthesized following a reported method.⁵⁰ To a solution of *N*-methylaniline (1.07 g, 10 mmol, 1 equiv) and potassium carbonate (4.8 g, 35 mmol, 3.5 equiv) in acetonitrile (9 mL) was added appropriate alkyl bromide (12 mmol, 1.2 equiv). The mixture was refluxed for 72 h and then cooled to room temperature. Solvent was removed under reduced pressure, and the residue was partitioned between water and dichloromethane. The aqueous phase was extracted three times with dichloromethane, and the combined organic phases were dried over anhydrous sulfate and concentrated under reduced pressure to yield an oily residue that was purified using column chromatography (SiO_2 , petroleum spirits/ethyl acetate, 9:1).

***N*-Ethyl-*N*-methylaniline (1i).** Purified using column chromatography (SiO_2 , petroleum spirits/ethyl acetate, 9:1), yellow oil, 900 mg (67%). Spectroscopic data in accordance with the literature:⁵¹ ^1H NMR (400 MHz, chloroform-*d*) δ 7.24 (m, 2H), 6.76–6.66 (m, 3H), 3.41 (qd, $J = 7.1, 1.6$ Hz, 2H), 2.91 (s, 3H), 1.16–1.10 (m, 3H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform-*d*) δ 149.1, 129.2, 116.0, 112.4, 46.8, 37.4, 11.2 ppm.

***N*-Butyl-*N*-methylaniline (1j).** Purified using column chromatography (SiO_2 , petroleum spirits/ethyl acetate, 9:1), yellow oil 1.45 g (89%). Spectroscopic data in accordance with the literature:⁵² ^1H NMR (400 MHz, chloroform-*d*) δ 7.27–7.21 (m, 2H), 6.75–6.65 (m, 3H), 3.36–3.29 (m, 2H), 2.93 (s, 3H), 1.63–1.51 (m, 2H), 1.43–1.29 (m, 2H), 1.01–0.92 (t, 3H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform-*d*) δ 149.5, 129.3, 115.9, 112.2, 52.7, 38.4, 29.0, 20.5, 14.1 ppm.

***N*-Benzyl-*N*-methylaniline (1k).** Purified using column chromatography (SiO_2 , petroleum spirits/ethyl acetate, 9:1), yellow oil, 1.2 g (60%). Spectroscopic data in accordance with the literature:⁵² ^1H NMR (400 MHz, chloroform-*d*) δ 7.35–7.22 (m, 6H), 7.19 (m, 2H), 6.63–6.56 (m, 2H), 4.51 (s, 2H), 3.01 (s, 3H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform-*d*) δ 148.7, 138.5, 131.9, 128.8, 127.2, 126.7, 114.0, 108.5, 56.7, 38.9 ppm.

1,2-Dibenzoyl ethylenes **2b–j** were synthesized according to a reported method.⁴⁰ The appropriate acetophenone (2 mmol, 1 equiv), iodine (4 mmol, 2 equiv), and copper(II)bromide (0.4 mmol, 0.2 equiv) were heated in DMF (2 mL) at 80 °C for 18–24 h. The reaction mixture was then cooled to room temperature, and excess iodine was quenched by addition of sodium thiosulfate solution. The aqueous mixture was extracted three times with ethyl acetate, and the combined organic phases were dried over sodium sulfate. After removal of the solvent, the solid residue was recrystallized from boiling heptane/ethyl acetate to yield the desired substituted Z-1,2-DBEs **2b–j**.

(*E*)-1,4-Di-*p*-tolylbut-2-ene-1,4-dione (2b). Yellow solid, 100 mg (38%), spectroscopic data in accordance with the literature:⁴⁰ ^1H

NMR (400 MHz, chloroform-*d*) δ 8.01–7.94 (m, 6H), 7.35–7.30 (m, 4H), 2.44 (s, 6H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform-*d*) δ 189.5, 145.1, 135.1, 134.6, 129.7, 129.2, 21.9 ppm.

(*E*)-1,4-Bis(4-isobutylphenyl)but-2-ene-1,4-dione (2c). Yellow solid, 193 mg (55%), spectroscopic data in accordance with the literature:⁴⁰ ^1H NMR (400 MHz, chloroform-*d*) δ 8.04–7.96 (m, 6H), 7.30 (d, $J = 8.2$ Hz, 4H), 2.57 (d, $J = 7.2$ Hz, 4H), 1.93 (dt, $J = 13.6, 6.8$ Hz, 2H), 0.92 (d, $J = 6.6$ Hz, 12H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform-*d*) δ 189.6, 148.8, 135.1, 134.9, 129.8, 129.1, 45.6, 30.3, 22.5 ppm.

(*E*)-1,4-Bis(4-*tert*-butylphenyl)but-2-ene-1,4-dione (2d). Yellow solid, 180 mg (52%), spectroscopic data in accordance with the literature:⁴⁰ ^1H NMR (400 MHz, chloroform-*d*) δ 8.05–7.99 (m, 6H), 7.58–7.52 (m, 4H), 1.36 (s, 18H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform-*d*) δ 189.6, 158.0, 135.1, 134.6, 129.1, 126.0, 35.4, 31.2 ppm.

(*E*)-1,4-Bis(3,4-dimethylphenyl)but-2-ene-1,4-dione (2e). Yellow solid, 200 mg (68%); mp 139–141 °C; ^1H NMR (400 MHz, chloroform-*d*) δ 8.00 (d, $J = 2.4$ Hz, 2H), 7.88–7.78 (m, 4H), 7.31–7.24 (m, 2H), 2.35 (s, 12H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform-*d*) δ 189.6, 143.7, 137.4, 134.9 (2C), 130.1, 123.0, 126.7, 20.2, 19.8 ppm; ATR-FTIR $\nu_{\text{max}} = 1645, 1601, 1409, 1299, 961, 752, 723$ cm^{-1} ; HRMS (ESI) m/z calcd $\text{C}_{20}\text{H}_{21}\text{O}_2$ $[\text{M} + \text{H}]^+$ 293.1536, found 293.1550.

(*E*)-1,4-Bis(4-methoxyphenyl)but-2-ene-1,4-dione (2f). Yellow solid, 148 mg (50%), spectroscopic data in accordance with the literature:⁴⁰ ^1H NMR (400 MHz, chloroform-*d*) δ 8.12–8.04 (m, 4H), 8.02 (s, 2H), 7.04–6.95 (m, 4H), 3.90 (s, 6H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform-*d*) δ 188.3, 164.3, 134.8, 131.5, 130.3, 114.3, 55.7 ppm.

(*E*)-1,4-Bis(4-(trifluoromethyl)phenyl)but-2-ene-1,4-dione (2g). Yellow solid, 86 mg (24%), spectroscopic data in accordance with the literature:⁴⁰ ^1H NMR (400 MHz, chloroform-*d*) δ 8.20–8.14 (m, 4H), 8.02 (s, 2H), 7.85–7.79 (m, 4H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform-*d*) δ 188.7, 139.4 (t, C–F, $^4J_{\text{C–F}} = 1.3$ Hz), 135.35 (q, C–F, $^2J_{\text{C–F}} = 33$ Hz), 135.27, 129.3, 126.2 (q, C–F, $^3J_{\text{C–F}} = 3.7$ Hz), 123.6 (q, C–F, $^1J_{\text{C–F}} = 272.8$ Hz) ppm.

(*E*)-1,4-Bis(4-bromophenyl)but-2-ene-1,4-dione (2h). Yellow solid, 275 mg (70%); mp 191–192 °C; ^1H NMR (400 MHz, chloroform-*d*) δ 7.97 (s, 2H), 7.91 (m, 4H), 7.72–7.65 (m, 4H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform-*d*) δ 188.7, 135.6, 135.0, 132.5, 130.5, 129.6 ppm; HRMS (ESI) m/z calcd $\text{C}_{16}\text{H}_{11}\text{Br}_2\text{O}_2$ $[\text{M} + \text{H}]^+$ 392.9120, found 392.9124.

(*E*)-1,4-Bis(4-chlorophenyl)but-2-ene-1,4-dione (2i). Yellow solid, 100 mg (33%), spectroscopic data in accordance with the literature:⁴⁰ ^1H NMR (400 MHz, chloroform-*d*) δ 8.02–7.98 (m, 4H), 7.97 (s, 2H), 7.56–7.44 (m, 4H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform-*d*) δ 188.4, 140.7, 135.2, 135.0, 130.4, 129.4 ppm.

(*E*)-1-(4-Methoxyphenyl)-4-phenylbut-2-ene-1,4-dione (2j). Isolated using column chromatography (SiO_2 , 5% ethyl acetate in petroleum ether), yellow solid, 50 mg (19%), spectroscopic data in accordance with the literature:⁴⁰ ^1H NMR (400 MHz, chloroform-*d*) δ 8.12–8.03 (m, 4H), 8.00 (dd, $J = 2.1, 0.5$ Hz, 2H), 7.66–7.59 (m, 1H), 7.56–7.48 (m, 2H), 7.03–6.94 (m, 2H), 3.89 (s, 3H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform-*d*) δ 190.06, 188.05, 164.34, 137.08, 135.41, 134.52, 133.90, 131.46, 130.11, 128.98, 114.26, 55.70 ppm.

General Procedure for the Oxidative Annulation Reaction.

To a 10 mL microwave vial were added substituted 1,2-DBE 2 (0.25 mmol), appropriate aniline derivative **1** (1.75 mmol), 1,4-dioxane (6 mL), and glacial acetic acid (1 mL). The mixture was irradiated using two 20 W white light CLF lamps, with a distance from the vial of 5 cm, for 4 h. To allow sufficient atmospheric oxygen into the reaction mixture, the vial was kept open during the course of the reaction. Solvent was then removed under reduced pressure, and the residue was purified using column chromatography (SiO_2 , petroleum spirits/ethyl acetate) to yield the THQ **3**.

(1,6-Dimethyl-1,2,3,4-tetrahydroquinoline-3,4-diyl)bis-(phenylmethanone) (3a). Purified using column chromatography (SiO_2 , 5% ethyl acetate in hexane), yellow solid, 67 mg, (73%); mp

123–124 °C; ^1H NMR (400 MHz, chloroform-*d*) δ 8.11 (dd, *J* = 8.4, 1.4 Hz, 2H), 7.99–7.94 (m, 2H), 7.64–7.55 (m, 2H), 7.54–7.44 (m, 4H), 7.01–6.95 (m, 1H), 6.70–6.59 (m, 2H), 5.39 (dt, *J* = 9.2, 0.9 Hz, 1H), 4.49 (td, *J* = 9.6, 4.8 Hz, 1H), 3.49 (dd, *J* = 11.3, 4.9 Hz, 1H), 3.30 (dd, *J* = 11.3, 9.8 Hz, 1H), 2.90 (s, 3H), 2.11 (s, 3H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform-*d*) δ 202.5, 200.5, 144.0, 138.0, 136.1, 133.53, 133.48, 129.1 (2C), 129.0 (2C), 128.9 (2C), 128.65 (2C), 128.6, 128.2, 126.9, 122.5, 112.2, 53.2, 46.2, 45.5, 39.5, 20.5 ppm; ATR-FTIR ν_{max} = 1687, 1669, 1511, 1213, 967, 774, 690 cm^{-1} ; HRMS (ESI) *m/z* calcd $\text{C}_{25}\text{H}_{24}\text{NO}_2$ [*M* + *H*] $^+$ 370.1807, found 370.1825.

(1,6-Dimethyl-1,2,3,4-tetrahydroquinoline-3,4-diyl)bis(*p*-tolyl-methanone) (3b). Purified using column chromatography (SiO_2 , 5% ethyl acetate in hexane), yellow solid, 63 mg (63%); mp 150–153 °C; ^1H NMR (400 MHz, chloroform-*d*) δ 8.03–7.96 (m, 2H), 7.88–7.81 (m, 2H), 7.32–7.21 (m, 4H), 6.95 (ddd, *J* = 8.3, 1.4, 0.8 Hz, 1H), 6.67–6.58 (m, 2H), 5.34 (d, *J* = 9.3 Hz, 1H), 4.44 (td, *J* = 9.7, 4.8 Hz, 1H), 3.45 (dd, *J* = 11.2, 4.8 Hz, 1H), 3.27 (dd, *J* = 11.3, 9.9 Hz, 1H), 2.89 (s, 3H), 2.43 (s, 3H), 2.40 (s, 3H), 2.09 (s, 3H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform-*d*) δ 202.1, 200.1, 144.44, 144.35, 144.1, 135.6, 133.6, 129.7 (2C), 129.6 (2C), 129.3 (2C), 128.9 (2C), 128.6, 128.5, 126.8, 122.9, 112.1, 53.4, 46.0, 45.36, 39.60, 21.9, 21.8, 20.5 ppm; ATR-FTIR ν_{max} = 1665, 1604, 1515, 1271, 1182, 828, 789 cm^{-1} ; HRMS (ESI) *m/z* calcd $\text{C}_{27}\text{H}_{28}\text{NO}_2$ [*M* + *H*] $^+$ 398.2120, found 398.2132.

(1,6-Dimethyl-1,2,3,4-tetrahydroquinoline-3,4-diyl)bis((4-isobutylphenyl)methanone) (3c). Purified using column chromatography (SiO_2 , 5% ethyl acetate in hexane), yellow solid, 61.9 mg (52%); mp 113–114 °C; ^1H NMR (400 MHz, chloroform-*d*) δ 8.00 (d, *J* = 8.3 Hz, 2H), 7.87 (d, *J* = 8.3 Hz, 2H), 7.29–7.19 (m, 4H), 6.95 (ddt, *J* = 8.3, 2.1, 0.8 Hz, 1H), 6.66–6.60 (m, 2H), 5.35 (d, *J* = 9.4 Hz, 1H), 4.46 (td, *J* = 9.7, 4.8 Hz, 1H), 3.47 (dd, *J* = 11.2, 4.8 Hz, 1H), 3.29 (dd, *J* = 11.2, 9.9 Hz, 1H), 2.90 (s, 3H), 2.54 (dd, *J* = 11.1, 7.2 Hz, 4H), 2.10 (s, 3H), 1.99–1.82 (m, 2H), 0.96–0.87 (m, 12H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform-*d*) δ 202.0, 200.1, 148.0, 147.9, 143.9, 135.8, 133.8, 129.5 (2C), 129.4 (2C), 128.9 (2C), 128.6 (2C), 128.50, 128.3, 126.6, 122.8, 112.0, 53.3, 46.1, 45.5, 45.4, 45.2, 39.5, 30.1, 22.4, 22.4, 22.3, 20.4 ppm; ATR-FTIR ν_{max} = 1668, 1603, 1514, 1270, 1181, 860, 799 cm^{-1} ; HRMS (ESI) *m/z* calcd $\text{C}_{33}\text{H}_{40}\text{NO}_2$ [*M* + *H*] $^+$ 482.3079, found 482.3070.

(1,6-Dimethyl-1,2,3,4-tetrahydroquinoline-3,4-diyl)bis((4-*tert*-butylphenyl)methanone) (3d). Purified using column chromatography (SiO_2 , 5% ethyl acetate in hexane), yellow solid, 54.5 mg (45%); mp 179–181 °C; ^1H NMR (400 MHz, chloroform-*d*) δ 8.02 (d, *J* = 8.6 Hz, 2H), 7.89 (d, *J* = 8.6 Hz, 2H), 7.47 (dd, *J* = 15.8, 8.6 Hz, 3H), 6.99–6.92 (m, 1H), 6.67–6.59 (m, 2H), 5.35 (d, *J* = 9.4 Hz, 1H), 4.46 (td, *J* = 9.7, 4.8 Hz, 1H), 3.47 (dd, *J* = 11.3, 4.8 Hz, 1H), 3.28 (dd, *J* = 11.3, 9.9 Hz, 1H), 2.90 (s, 3H), 2.10 (s, 3H), 1.36 (s, 9H), 1.33 (s, 9H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform-*d*) δ 202.0, 199.9, 157.2, 157.0, 143.9, 135.4, 133.4, 128.9 (2C), 128.6 (2C), 128.5, 128.3, 126.7, 125.8 (2C), 125.7 (2C), 122.8, 111.9, 53.2, 45.9, 45.2, 39.5, 35.1, 35.1, 31.1, 31.0, 20.4 ppm; ATR-FTIR ν_{max} = 1668, 1604, 1514, 1270, 1188, 803 cm^{-1} ; HRMS (ESI) *m/z* calcd $\text{C}_{33}\text{H}_{40}\text{NO}_2$ [*M* + *H*] $^+$ 482.3059, found 482.3069.

(1,6-Dimethyl-1,2,3,4-tetrahydroquinoline-3,4-diyl)bis((3,4-dimethylphenyl)methanone) (3e). Purified using column chromatography (SiO_2 , 5% ethyl acetate in hexane), yellow solid, 43.4 mg (46%); mp 129–130 °C; ^1H NMR (400 MHz, chloroform-*d*) δ 7.85 (dd, *J* = 10.2, 2.4 Hz, 2H), 7.70 (dd, *J* = 10.2, 2.5 Hz, 2H), 7.25–7.18 (m, 2H), 6.95 (dd, *J* = 8.3, 2.2 Hz, 1H), 6.67–6.59 (m, 2H), 5.34 (d, *J* = 9.4 Hz, 1H), 4.45 (td, *J* = 9.7, 4.7 Hz, 1H), 3.49–3.41 (m, 1H), 3.28 (dd, *J* = 11.2, 10.0 Hz, 1H), 2.90 (d, *J* = 1.1 Hz, 3H), 2.34 (s, 3H), 2.32 (s, 3H), 2.31 (s, 3H), 2.29 (s, 3H), 2.09 (s, 3H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform-*d*) δ 202.4, 200.4, 144.0, 143.2, 143.1, 137.23, 137.20, 136.1, 134.1, 130.2, 130.14, 130.08, 129.9, 128.6, 128.5, 127.0, 126.8, 126.5, 123.1, 112.0, 53.5, 46.0, 45.3, 39.6, 20.5, 20.22, 20.18, 20.0, 19.9 ppm; ATR-FTIR ν_{max} = 1662, 1603, 1518, 1267, 1208, 794 cm^{-1} ; HRMS (ESI) *m/z* calcd $\text{C}_{29}\text{H}_{32}\text{NO}_2$ [*M* + *H*] $^+$ 426.2433, found 426.2436.

(1,6-Dimethyl-1,2,3,4-tetrahydroquinoline-3,4-diyl)bis((4-methoxyphenyl)methanone) (3f). Purified using column chromatography (SiO_2 , 5% ethyl acetate in hexane), off-white solid, 53.3 mg (50%); mp 139–140 °C; ^1H NMR (400 MHz, chloroform-*d*) δ 8.08 (d, *J* = 9.0 Hz, 2H), 7.95 (d, *J* = 8.9 Hz, 2H), 7.08–6.86 (m, 5H), 6.70–6.56 (m, 2H), 5.32 (dt, *J* = 9.7, 1.0 Hz, 1H), 4.44 (td, *J* = 10.0, 4.8 Hz, 1H), 3.88 (s, 3H), 3.86 (s, 3H), 3.44 (dd, *J* = 11.2, 4.8 Hz, 1H), 3.28 (dd, *J* = 11.2, 10.2 Hz, 1H), 2.90 (s, 3H), 2.09 (s, 3H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform-*d*) δ 200.9, 199.0, 163.9, 163.8, 144.0, 131.5 (2C), 131.2, 131.0 (2C), 129.1, 128.5, 128.4, 126.8, 123.1, 114.1 (2C), 114.0 (2C), 112.2, 55.59, 55.58, 53.6, 46.0, 45.0, 39.6, 20.5 ppm; ATR-FTIR ν_{max} = 1661, 1598, 1572, 1509, 1316, 1255, 1170, 845 cm^{-1} ; HRMS (ESI) *m/z* calcd $\text{C}_{27}\text{H}_{28}\text{NO}_4$ [*M* + *H*] $^+$ 430.2018, found 430.2024.

(1,6-Dimethyl-1,2,3,4-tetrahydroquinoline-3,4-diyl)bis((4-(trifluoromethyl)phenyl)methanone) (3g). Purified using column chromatography (SiO_2 , 5% ethyl acetate in hexane), thick yellow-orange oil, 35 mg (28%); ^1H NMR (400 MHz, chloroform-*d*) δ 8.25–8.13 (m, 2H), 8.08–8.01 (m, 2H), 7.83–7.71 (m, 4H), 6.99 (dq, *J* = 8.2, 0.9 Hz, 1H), 6.66 (d, *J* = 8.4 Hz, 1H), 6.54–6.48 (m, 1H), 5.32 (d, *J* = 9.5 Hz, 1H), 4.46 (td, *J* = 9.7, 5.1 Hz, 1H), 3.55–3.44 (m, 1H), 3.29 (dd, *J* = 11.2, 10.1 Hz, 1H), 2.90 (d, *J* = 0.7 Hz, 3H), 2.10 (s, 3H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform-*d*) δ 201.6, 199.8, 143.8, 140.5, 138.7, 134.8 (q, C–F, $^2J_{\text{C–F}}$ = 33 Hz), 134.7 (q, C–F, $^2J_{\text{C–F}}$ = 33 Hz), 129.4 (2C), 129.0 (2C), 128.2, 127.3, 126.4–125.8 (m, C–F, 4C), 123.7 (q, C–F, $^1J_{\text{C–F}}$ = 272.6 Hz), 123.6 (q, C–F, $^1J_{\text{C–F}}$ = 272.6 Hz), 121.9, 121.2, 112.4, 52.9, 46.6, 46.2, 39.5, 20.46 ppm; ^{19}F NMR (470 MHz, chloroform-*d*) δ –63.16 (s, 3F), –63.23 (s, 3F) ppm; ATR-FTIR ν_{max} = 1683, 1512, 1322, 1170, 1128, 1067 cm^{-1} ; HRMS (ESI) *m/z* calcd $\text{C}_{27}\text{H}_{22}\text{F}_6\text{NO}_2$ [*M* + *H*] $^+$ 506.1555, found 506.1548.

(1,6-Dimethyl-1,2,3,4-tetrahydroquinoline-3,4-diyl)bis((4-bromophenyl)methanone) (3h). Purified using column chromatography (SiO_2 , 5% ethyl acetate in hexane), yellow solid, 113.4 mg (84%); mp 128–130 °C; ^1H NMR (400 MHz, chloroform-*d*) δ 8.00–7.88 (m, 2H), 7.85–7.77 (m, 2H), 7.71–7.56 (m, 4H), 6.97 (ddt, *J* = 8.3, 2.2, 0.8 Hz, 1H), 6.65 (d, *J* = 8.3 Hz, 1H), 6.52 (dt, *J* = 2.0, 0.9 Hz, 1H), 5.30–5.16 (m, 1H), 4.39 (td, *J* = 9.9, 5.0 Hz, 1H), 3.44 (dd, *J* = 11.3, 5.0 Hz, 1H), 3.26 (dd, *J* = 11.3, 10.1 Hz, 1H), 2.90 (s, 3H), 2.09 (s, 3H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform-*d*) δ 201.4, 199.6, 143.8, 136.67, 134.7, 132.34 (2C), 132.28 (2C), 130.6 (2C), 130.2 (2C), 128.99, 128.98, 128.8, 128.3, 127.1, 122.3, 112.3, 53.2, 46.3, 45.6, 39.5, 20.5 ppm; ATR-FTIR ν_{max} = 2856, 1670, 1654, 1582, 1507, 1067, 1005, 816 cm^{-1} ; HRMS (ESI) *m/z* calcd $\text{C}_{25}\text{H}_{22}\text{Br}_2\text{NO}_2$ [*M* + *H*] $^+$ 526.0017, found 526.0025.

(1,6-Dimethyl-1,2,3,4-tetrahydroquinoline-3,4-diyl)bis((4-chlorophenyl)methanone) (3i). Purified using column chromatography (SiO_2 , 5% ethyl acetate in hexane), yellow solid, 83.1 mg (72%); mp 149–150 °C; ^1H NMR (400 MHz, chloroform-*d*) δ 8.03 (d, *J* = 8.6 Hz, 2H), 7.89 (d, *J* = 8.7 Hz, 2H), 7.46 (dd, *J* = 16.1, 8.6 Hz, 4H), 6.97 (ddt, *J* = 8.3, 2.1, 0.8 Hz, 1H), 6.65 (d, *J* = 8.4 Hz, 1H), 6.54 (dt, *J* = 2.0, 0.9 Hz, 1H), 5.28 (dd, *J* = 9.6, 1.1 Hz, 1H), 4.41 (td, *J* = 9.9, 5.0 Hz, 1H), 3.45 (dd, *J* = 11.3, 5.0 Hz, 1H), 3.27 (dd, *J* = 11.2, 10.1 Hz, 1H), 2.90 (s, 3H), 2.10 (s, 3H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform-*d*) δ 201.2, 199.4, 143.8, 140.2, 140.2, 136.3, 134.3, 130.5 (2C), 130.1 (2C), 129.34 (2C), 129.28 (2C), 128.8, 128.3, 127.0, 122.3, 112.2, 53.2, 46.3, 45.6, 39.5, 20.5 ppm; ATR-FTIR ν_{max} = 2859, 1673, 1587, 1512, 1399, 1204, 1090, 818 cm^{-1} ; HRMS (ESI) *m/z* calcd $\text{C}_{25}\text{H}_{22}\text{Cl}_2\text{NO}_2$ [*M* + *H*] $^+$ 438.1028, found 438.1028.

(3-Benzoyl-1,6-dimethyl-1,2,3,4-tetrahydroquinolin-4-yl)((4-methoxyphenyl)methanone) (3ja) and (4-benzoyl-1,6-dimethyl-1,2,3,4-tetrahydroquinolin-3-yl)((4-methoxyphenyl)methanone) (3jb). Purified using column chromatography (SiO_2 , 1% ethyl acetate in hexane), thick yellow oil, 66 mg (64%); ^1H NMR (400 MHz, chloroform-*d*) δ 8.11–8.06 (m, overlapping 3ja and 3jb, 4H), 7.98–7.92 (m, overlapping 3ja and 3jb, 4H), 7.62–7.53 (m, overlapping 3ja and 3jb, 2H), 7.52–7.42 (m, overlapping 3ja and 3jb, 4H), 7.00–6.90 (m, overlapping 3ja and 3jb, 6H), 6.67–6.57 (m, overlapping 3ja and 3jb, 4H), 5.38–5.31 (m, overlapping 3ja and 3jb, 2H), 4.52–4.40 (m,

overlapping 3ja and 3ja, 2H), 3.89 (s, OMe 3ja, 3H), 3.86 (s, OMe 3jb, 3H), 3.50–3.42 (m, overlapping 3ja and 3jb, 2H), 3.34–3.25 (m, overlapping 3ja and 3jb, 2H), 2.90 (s, NMe, 3H), 2.89 (s, NMe, 3H), 2.10 (s, ArMe 3ja, 3H), 2.09 (s, ArMe 3jb, 3H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform-*d*) δ 202.7, 200.8, 200.7, 198.8, 163.93, 163.89, 144.0, 138.1, 136.1, 133.5, 133.4, 131.5 (2C), 131.12, 131.05 (2C), 130.5, 129.1 (2C), 128.92 (2C), 128.85 (2C), 128.7 (2C), 128.6, 128.52, 128.49 (2C), 128.2, 126.81, 126.76, 122.8, 114.1, 114.0, 112.11, 112.09, 55.6 (2C), 53.5, 53.3, 46.3, 45.8, 45.4, 45.1, 39.57, 39.55, 20.49, 20.47 ppm; ATR-FTIR ν_{max} = 1660, 1598, 1572, 1509, 1317, 1257, 1170 cm^{-1} ; HRMS (ESI) m/z calcd $\text{C}_{26}\text{H}_{26}\text{NO}_3$ $[\text{M} + \text{H}]^+$ 400.1911, found 400.1913.

(1-Methyl-1,2,3,4-tetrahydroquinoline-3,4-diyl)bis-(phenylmethanone) (3k). Purified using column chromatography (SiO_2 , 5% ethyl acetate in hexane), yellow oil, 44.3 mg (59%); mp 104–105 $^\circ\text{C}$; ^1H NMR (400 MHz, chloroform-*d*) δ 8.09 (d, J = 8.4 Hz, 2H), 8.00–7.94 (m, 2H), 7.59 (dtd, J = 8.5, 7.3, 1.3 Hz, 2H), 7.53–7.43 (m, 4H), 7.14 (tt, J = 7.2, 0.9 Hz, 1H), 6.78 (dd, J = 7.6, 1.3 Hz, 1H), 6.71 (dt, J = 8.4, 1.2 Hz, 1H), 6.60–6.54 (m, 1H), 5.38 (dd, J = 9.4, 1.2 Hz, 1H), 4.55–4.46 (m, 1H), 3.56–3.46 (m, 1H), 3.35 (ddd, J = 11.2, 10.1, 1.1 Hz, 1H), 2.93 (d, J = 1.1 Hz, 3H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform-*d*) δ 202.1, 200.5, 145.8, 137.9, 136.0, 133.5, 133.4, 129.0 (2C), 128.84 (2C), 128.77 (2C), 128.6 (2C), 127.9, 127.8, 122.1, 117.3, 111.7, 52.9, 46.1, 45.0, 39.2 ppm; ATR-FTIR ν_{max} = 1669, 1598, 1578, 1506, 1448, 1228, 1202, 753, 703 cm^{-1} ; HRMS (ESI) m/z calcd $\text{C}_{24}\text{H}_{22}\text{NO}_2$ $[\text{M} + \text{H}]^+$ 356.1651, found 356.1661.

(1,5,7-Trimethyl-1,2,3,4-tetrahydroquinoline-3,4-diyl)bis-(phenylmethanone) (3m). Purified using column chromatography (SiO_2 , 5% ethyl acetate in hexane), white solid, 39.9 mg (42%); mp 198–199 $^\circ\text{C}$; ^1H NMR (400 MHz, chloroform-*d*) δ 8.10–8.02 (m, 2H), 7.78 (dd, J = 7.4, 1.2 Hz, 2H), 7.63–7.52 (m, 2H), 7.52–7.40 (m, 4H), 6.52–6.38 (m, 2H), 5.46–5.33 (m, 1H), 3.83 (q, J = 1.5 Hz, 1H), 3.46 (dd, J = 11.9, 3.4 Hz, 1H), 3.41–3.31 (m, 1H), 2.71 (s, 3H), 2.26 (s, 3H), 2.06 (s, 3H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform-*d*) δ 201.8, 198.9, 146.9, 137.0, 136.3, 136.1, 135.6, 133.5, 133.0, 129.1 (2C), 128.8 (2C), 128.6 (2C), 128.3 (2C), 121.6, 116.0, 111.0, 50.4, 44.3, 43.2, 40.1, 21.8, 20.6 ppm; ATR-FTIR ν_{max} = 1680, 1577, 1447, 1331, 1268, 1214, 970, 702 cm^{-1} ; HRMS (ESI) m/z calcd $\text{C}_{26}\text{H}_{26}\text{NO}_2$ $[\text{M} + \text{H}]^+$ 384.1964, found 384.1991.

(6-Methoxy-1-methyl-1,2,3,4-tetrahydroquinoline-3,4-diyl)bis-(phenylmethanone) (3n). Purified using column chromatography (SiO_2 , 5% ethyl acetate in hexane), yellow solid, 58.6 mg (61%); mp 129–130 $^\circ\text{C}$; ^1H NMR (400 MHz, chloroform-*d*) δ 8.14–8.04 (m, 2H), 7.96 (dt, J = 7.0, 1.5 Hz, 2H), 7.58 (dddd, J = 9.0, 7.0, 5.2, 1.4 Hz, 2H), 7.53–7.39 (m, 4H), 6.81–6.61 (m, 2H), 6.43 (dd, J = 2.9, 1.3 Hz, 1H), 5.41 (d, J = 9.2 Hz, 1H), 4.51 (tdd, J = 9.5, 4.9, 1.3 Hz, 1H), 3.57 (s, 3H), 3.46 (ddd, J = 11.2, 5.0, 1.3 Hz, 1H), 3.29–3.19 (m, 1H), 2.87 (s, 3H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform-*d*) δ 201.9, 200.4, 152.0, 140.9, 138.0, 136.1, 133.57, 133.56, 129.1 (2C), 129.0 (2C), 128.9 (2C), 128.7 (2C), 123.9, 114.6, 113.3, 113.2, 55.7, 53.5, 46.2, 45.4, 40.0 ppm; ATR-FTIR ν_{max} = 1663, 1594, 1579, 1510, 1447, 1209, 803, 689 cm^{-1} ; HRMS (ESI) m/z calcd $\text{C}_{25}\text{H}_{24}\text{NO}_3$ $[\text{M} + \text{H}]^+$ 386.1756, found 386.1753.

(7-Methoxy-1-methyl-1,2,3,4-tetrahydroquinoline-3,4-diyl)bis-(phenylmethanone) (3o). Purified using column chromatography (SiO_2 , 5% ethyl acetate in hexane), yellow solid, 26.9 mg (28%); mp 152–153 $^\circ\text{C}$; ^1H NMR (400 MHz, chloroform-*d*) δ 8.07–8.00 (m, 2H), 7.85–7.77 (m, 2H), 7.57–7.49 (m, 2H), 7.46–7.38 (m, 4H), 7.13 (td, J = 8.3, 0.6 Hz, 1H), 6.42 (dd, J = 8.4, 0.8 Hz, 1H), 6.30 (dd, J = 8.2, 0.9 Hz, 1H), 5.46 (d, J = 5.5 Hz, 1H), 4.00 (ddd, J = 6.5, 5.5, 3.4 Hz, 1H), 3.50 (s, 3H), 3.44 (ddd, J = 11.7, 3.4, 0.6 Hz, 1H), 3.36–3.28 (m, 1H), 2.83 (s, 3H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform-*d*) δ 203.3, 199.6, 157.0, 148.0, 137.0, 136.1, 133.2, 132.7, 128.8 (2C), 128.57 (2C), 128.55 (2C), 128.4 (2C), 128.2, 110.5, 105.8, 100.6, 55.1, 51.6, 44.6, 41.5, 40.0 ppm; ATR-FTIR ν_{max} = 1677, 1579, 1447, 1211, 967, 700 cm^{-1} ; HRMS (ESI) m/z calcd $\text{C}_{25}\text{H}_{24}\text{NO}_3$ $[\text{M} + \text{H}]^+$ 386.1756, found 386.1751.

(6-Fluoro-1-methyl-1,2,3,4-tetrahydroquinoline-3,4-diyl)bis-(phenylmethanone) (3p). Purified using column chromatography

(SiO_2 , 5% ethyl acetate in hexane), yellow solid, 49.9 mg (51%); mp 132–133 $^\circ\text{C}$; ^1H NMR (400 MHz, chloroform-*d*) δ 8.13–8.07 (m, 2H), 8.00–7.90 (m, 2H), 7.65–7.55 (m, 2H), 7.54–7.43 (m, 4H), 6.85 (dddd, J = 9.0, 8.1, 2.9, 0.8 Hz, 1H), 6.63 (dd, J = 9.0, 4.8 Hz, 1H), 6.54 (ddd, J = 9.4, 2.9, 1.1 Hz, 1H), 5.38 (dd, J = 8.9, 1.0 Hz, 1H), 4.46 (td, J = 9.2, 4.9 Hz, 1H), 3.50 (dd, J = 11.4, 4.9 Hz, 1H), 3.29 (dd, J = 11.3, 9.5 Hz, 1H), 2.88 (s, 3H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform-*d*) δ 201.5, 200.1, 155.5 (d, C–F, $^1J_{\text{C–F}}$ = 236.2 Hz), 142.8 (d, C–F, $^4J_{\text{C–F}}$ = 1.9 Hz), 137.5, 135.9, 133.8, 133.7, 129.10 (2C), 129.06 (2C), 128.9 (2C), 128.6 (2C), 123.4 (d, C–F, $^3J_{\text{C–F}}$ = 6.9 Hz), 114.82 (d, C–F, $^2J_{\text{C–F}}$ = 23.3 Hz), 114.31 (d, C–F, $^2J_{\text{C–F}}$ = 21.8 Hz), 112.8 (d, C–F, $^3J_{\text{C–F}}$ = 7.7 Hz), 53.00, 45.86, 45.07, 39.77 ppm; ^{19}F NMR (470 MHz, chloroform-*d*) δ –127.15 to –127.23 (m) ppm; ATR-FTIR ν_{max} = 1670, 1591, 1577, 1505, 1448, 1211, 906, 729, 685 cm^{-1} ; HRMS (ESI) m/z calcd $\text{C}_{24}\text{H}_{21}\text{FNO}_2$ $[\text{M} + \text{H}]^+$ 374.1556, found 374.1572.

(6-Bromo-1-methyl-1,2,3,4-tetrahydroquinoline-3,4-diyl)bis-(phenylmethanone) (3q). Purified using column chromatography (SiO_2 , 5% ethyl acetate in hexane), off-white solid, 26 mg (24%); mp 192–194 $^\circ\text{C}$; ^1H NMR (400 MHz, chloroform-*d*) δ 8.10–8.03 (m, 2H), 7.97–7.88 (m, 2H), 7.67–7.55 (m, 2H), 7.54–7.42 (m, 4H), 7.22 (ddt, J = 8.7, 1.6, 0.7 Hz, 1H), 6.88–6.84 (m, 1H), 6.55 (dd, J = 8.9, 1.2 Hz, 1H), 5.36–5.30 (m, 1H), 4.44–4.33 (m, 1H), 3.53 (ddd, J = 11.5, 4.7, 1.3 Hz, 1H), 3.34 (ddd, J = 11.8, 9.3, 1.2 Hz, 1H), 2.88 (d, J = 1.3 Hz, 3H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform-*d*) δ 201.6, 200.0, 145.1, 137.5, 135.9, 133.9, 133.7, 130.8, 130.6, 129.2 (2C), 129.1 (2C), 129.0 (2C), 128.7 (2C), 123.8, 113.3, 109.3, 52.5, 45.8, 44.8, 39.4 ppm; ATR-FTIR ν_{max} = 1686, 1668, 1591, 1499, 1447, 1209, 699, 691 cm^{-1} ; HRMS (ESI) m/z calcd $\text{C}_{24}\text{H}_{21}\text{BrNO}_2$ $[\text{M} + \text{H}]^+$ 434.0756, found 434.0752.

(1-Methyl-1,2,3,4-tetrahydrobenzo[h]quinoline-3,4-diyl)bis-(phenylmethanone) (3r). Purified using column chromatography (SiO_2 , 5% ethyl acetate in hexane), white solid, 19.2 mg (18%); mp 197–200 $^\circ\text{C}$; ^1H NMR (400 MHz, chloroform-*d*) δ 8.26–8.15 (m, 2H), 8.04–7.99 (m, 2H), 7.77–7.72 (m, 1H), 7.67–7.42 (m, 9H), 7.38 (d, J = 8.6 Hz, 1H), 6.99 (d, J = 8.5 Hz, 1H), 5.85 (d, J = 9.8 Hz, 1H), 4.82 (ddd, J = 11.3, 9.8, 3.3 Hz, 1H), 3.66 (dd, J = 13.5, 3.3 Hz, 1H), 3.39–3.28 (m, 1H), 3.28 (s, 3H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform-*d*) δ 203.0, 200.4, 138.3, 136.1, 133.7, 133.7, 133.6, 129.4 (2C), 129.09 (2C), 129.05 (2C), 128.99, 128.68, 128.6 (2C), 128.5, 126.0, 125.9, 124.0 (2C), 123.96, 123.89, 54.7, 46.0, 44.7, 40.3 ppm; ATR-FTIR ν_{max} = 1667, 1596, 1446, 1266, 1216, 1001, 986, 797, 783 cm^{-1} ; HRMS (ESI) m/z calcd $\text{C}_{28}\text{H}_{24}\text{NO}_2$ $[\text{M} + \text{H}]^+$ 406.1807, found 406.1826.

(1-Ethyl-1,2,3,4-tetrahydroquinoline-3,4-diyl)bis-(phenylmethanone) (3s). Purified using column chromatography (SiO_2 , 1% ethyl acetate in hexane), yellow oil, 42.4 mg (46%); ^1H NMR (400 MHz, chloroform-*d*) δ 8.13–8.06 (m, 2H), 8.03–7.91 (m, 2H), 7.64–7.55 (m, 2H), 7.54–7.42 (m, 4H), 7.11 (t, J = 7.8 Hz, 1H), 6.82–6.69 (m, 2H), 6.53 (dd, J = 8.3, 6.7 Hz, 1H), 5.38 (d, J = 9.4 Hz, 1H), 4.59–4.40 (m, 1H), 3.64–3.18 (m, 4H), 1.16 (t, J = 6.88 Hz, 3H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform-*d*) δ 202.3, 200.9, 144.3, 138.2, 136.2, 133.6, 133.5, 129.1 (2C), 129.0 (2C), 128.9 (2C), 128.7 (2C), 128.4, 128.1, 121.6, 116.7, 111.6, 50.3, 46.6, 45.7, 44.6, 11.1 ppm; ATR-FTIR ν_{max} = 1675, 1597, 1495, 1449, 1346, 743, 691 cm^{-1} ; HRMS (ESI) m/z calcd $\text{C}_{25}\text{H}_{24}\text{NO}_2$ $[\text{M} + \text{H}]^+$ 370.1807, found 370.1817.

(1-Butyl-1,2,3,4-tetrahydroquinoline-3,4-diyl)bis-(phenylmethanone) (3t). Purified using column chromatography (SiO_2 , 1% ethyl acetate in hexane), yellow oil, 57.4 mg (60%); ^1H NMR (400 MHz, chloroform-*d*) δ 8.12–8.07 (m, 2H), 8.00–7.93 (m, 2H), 7.64–7.53 (m, 2H), 7.54–7.42 (m, 4H), 7.10 (dddd, J = 8.2, 7.3, 1.6, 0.8 Hz, 1H), 6.76 (dt, J = 7.6, 1.4 Hz, 1H), 6.69 (dd, J = 8.4, 1.1 Hz, 1H), 6.51 (td, J = 7.4, 1.2 Hz, 1H), 5.37 (d, J = 9.5 Hz, 1H), 4.47 (ddd, J = 10.1, 9.4, 4.2 Hz, 1H), 3.57 (dd, J = 11.6, 4.3 Hz, 1H), 3.47–3.31 (m, 2H), 3.28–3.15 (m, 1H), 1.64–1.52 (m, 2H), 1.42–1.30 (m, 2H), 0.94 (t, J = 7.3 Hz, 3H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform-*d*) δ 202.3, 200.8, 144.5, 138.2, 136.1, 133.63, 133.56, 129.1 (2C), 129.0 (2C), 128.9 (2C), 128.7 (2C), 128.4, 128.1, 121.5, 116.8, 111.8, 51.7, 51.3, 46.6, 44.5, 28.5, 20.5, 14.1 ppm;

ATR-FTIR ν_{\max} = 1678, 1599, 1502, 1457, 1449, 1218, 744 cm^{-1} ; HRMS (ESI) m/z calcd $\text{C}_{27}\text{H}_{28}\text{NO}_2$ $[\text{M} + \text{H}]^+$ 398.2120, found 398.2110.

(1-Benzyl-1,2,3,4-tetrahydroquinoline-3,4-diyl)bis-(phenylmethanone) (**3u**). Purified using column chromatography (SiO_2 , 1% ethyl acetate in hexane), yellow oil, 26.1 mg (25%); ^1H NMR (400 MHz, chloroform- d) δ 8.16–8.10 (m, 2H), 7.89–7.82 (m, 2H), 7.65–7.59 (m, 1H), 7.57–7.50 (m, 3H), 7.43–7.37 (m, 2H), 7.37–7.26 (m, 5H), 7.11–7.01 (m, 1H), 6.83 (d, J = 7.4 Hz, 1H), 6.74 (d, J = 8.3 Hz, 1H), 6.59 (t, J = 7.5 Hz, 1H), 5.48 (d, J = 9.2 Hz, 1H), 4.64–4.47 (m, 3H), 3.67 (dd, J = 11.8, 4.3 Hz, 1H), 3.52–3.41 (m, 1H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform- d) δ 202.3, 200.5, 145.0, 138.2, 138.0, 135.9, 133.6, 129.1 (2C), 129.0 (2C), 128.89 (2C), 128.87 (2C), 128.6 (2C), 128.4, 128.2, 127.9, 127.0 (2C), 125.9, 121.4, 117.4, 112.3, 55.4, 51.2, 46.4, 44.7 ppm; ATR-FTIR ν_{\max} = 1677, 1599, 1497, 1449, 1220, 749, 696 cm^{-1} ; HRMS (ESI) m/z calcd $\text{C}_{30}\text{H}_{26}\text{NO}_2$ $[\text{M} + \text{H}]^+$ 432.1958, found 432.1971.

Multicomponent Synthesis of 3a. For the multicomponent reaction, Wittig reagent **4** (74.4 mg, 0.2 mmol, 1 equiv), phenylglyoxal (35.5 mg, 0.26 mmol, 1.4 equiv), and amine **1a** (164 mg, 1.2 mmol, 6.2 equiv) were mixed in 1,4-dioxane (6 mL), and the reaction mixture was irradiated according to the general procedure for 18 h. Solvent was then removed under reduced pressure, and the product was isolated using flash chromatography (silica gel, 5% ethyl acetate in petroleum ethers) to yield the desired product **3a** (29 mg, 40%).

Synthesis of 3a on 1 mmol Scale. To a round-bottom flask were added 1,2-DBE **2a** (245.4 mg, 1 mmol), 4',*N*-trimethylaniline **1a** (947 mg, 7 mmol), 1,4-dioxane (25 mL), and glacial acetic acid (4 mL). The mixture was irradiated using two 20 W white light CLF lamps, with a distance from the flask of 5 cm, for 18 h. To allow sufficient atmospheric oxygen into the reaction mixture, the flask was kept open during the course of the reaction. Solvent was then removed under reduced pressure, and the residue was purified using column chromatography (SiO_2 , 5% ethyl acetate in petroleum spirits) to yield the THQ **3a** (204.1 mg, 53%).

Condensation Reaction of 3a with Ammonium Acetate to Pyrrole Derivative 6. Following a modified published procedure,⁴⁵ diketo compound **3a** (106.3 mg, 0.29 mmol, 1 equiv) was dissolved in glacial acetic acid (1.8 mL). Ammonium acetate (189 mg, 2.4 mmol, 8.2 equiv) was then added, and the mixture was heated to 120 °C using a heating block in a sealed vial for 4 h. The resulting deep red solution was concentrated in vacuo, and the product was isolated using column chromatography (SiO_2 , 5% methanol in dichloromethane) as dark red crystals, 43 mg (43%); mp 223–225 °C; ^1H NMR (400 MHz, chloroform- d) δ 8.50 (s, 1H), 8.09 (dd, J = 2.1, 1.0 Hz, 1H), 7.81 (ddd, J = 8.2, 7.2, 1.3 Hz, 4H), 7.47 (t, J = 7.5 Hz, 2H), 7.43–7.29 (m, 4H), 7.22–7.14 (m, 2H), 3.99 (s, 3H), 2.30 (s, 3H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform- d) δ 145.1, 142.6, 137.5, 137.0, 136.6, 134.9, 130.7, 129.7 (2C), 128.8 (2C), 128.4 (2C), 128.0 (2C), 127.6, 127.4, 127.3, 124.2, 124.0, 117.4, 116.3, 115.4, 43.4, 21.5 ppm; ATR-FTIR ν_{\max} = 1591, 1454, 1356, 1225, 807, 765, 694 cm^{-1} ; HRMS (ESI) m/z calcd $\text{C}_{25}\text{H}_{21}\text{N}_2$ $[\text{M} + \text{H}]^+$ 349.1704, found 349.1717.

Synthesis of Furan Derivative 7. Following a modified published procedure,⁴⁶ diketo compound **3a** (40.7 mg, 0.11 mmol) was dissolved in acetic anhydride (1.0 mL). Concentrated hydrochloric acid (0.3 mL) was then added at 0 °C. The mixture was heated to 80 °C using a heating block in a sealed vial for 18 h. The resulting orange solution was concentrated in vacuo, and the product was isolated using column chromatography (SiO_2 , 2% ethyl acetate in hexane) as an air sensitive yellow-orange oil, 21.8 mg (56%); ^1H NMR (400 MHz, chloroform- d) δ 7.88–7.81 (m, 2H), 7.61–7.55 (m, 2H), 7.51 (d, J = 2.1 Hz, 1H), 7.49–7.40 (m, 4H), 7.40–7.33 (m, 1H), 7.30 (ddt, J = 7.5, 5.8, 1.0 Hz, 1H), 7.05–6.98 (m, 1H), 6.74 (d, J = 8.3 Hz, 1H), 4.30 (s, 2H), 2.97 (s, 3H), 2.19 (s, 3H) ppm; $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, chloroform- d) δ 146.4, 145.1, 144.0, 131.9, 130.9, 129.0 (2C), 128.9 (2C), 128.7, 128.3, 127.9, 127.4 (2C), 127.3, 125.19 (2C), 125.15, 119.24, 119.16, 117.5, 113.2, 49.4, 39.5, 20.6 ppm; ATR-FTIR ν_{\max} = 2924, 1598, 1493, 1448, 1242,

1126 cm^{-1} ; HRMS (ESI) m/z calcd $\text{C}_{25}\text{H}_{22}\text{NO}$ $[\text{M} + \text{H}]^+$ 352.1701, found 352.1697.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.joc.0c02819>.

Photophysical measurements, synthesis procedures, NMR characterization data, and X-ray crystallographic data for **3q** (CCDC 2008622) (PDF)

Accession Codes

CCDC 2008622 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

■ AUTHOR INFORMATION

Corresponding Author

Henrik Sundén – Department of Chemistry and Chemical Engineering, Chalmers University of Technology, 41296 Gothenburg, Sweden; Chemistry and Molecular Biology, University of Gothenburg, 41296 Gothenburg, Sweden; orcid.org/0000-0001-6202-7557; Email: henrik.sunden@chem.gu.se

Authors

August Runemark – Department of Chemistry and Chemical Engineering, Chalmers University of Technology, 41296 Gothenburg, Sweden; orcid.org/0000-0002-3650-1881
Savannah C. Zacharias – Chemistry and Molecular Biology, University of Gothenburg, 41296 Gothenburg, Sweden; orcid.org/0000-0003-0895-9120

Complete contact information is available at: <https://pubs.acs.org/doi/10.1021/acs.joc.0c02819>

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This work was supported by grants from the Swedish research council FORMAS (2019-00699) and from Olle Engkvists Stiftelse. The X-ray crystallographic data used in this work was collected by Ivana Cisařová, Department of Inorganic Chemistry, Faculty of Science, Charles University, Hlavova 2030/8, 128 43 Prague 2, Czech Republic.

■ REFERENCES

- Yuan, Y.; Majumder, S.; Yang, M.; Guo, S. Recent Advances in Catalyst-Free Photochemical Reactions via Electron-Donor-Acceptor (EDA) Complex Process. *Tetrahedron Lett.* **2020**, *61* (8), 151506.
- Wei, Y.; Zhou, Q.-Q.; Tan, F.; Lu, L.-Q.; Xiao, W.-J. Visible-Light-Driven Organic Photochemical Reactions in the Absence of External Photocatalysts. *Synthesis* **2019**, *51* (16), 3021–3054.
- Crisenza, G. E. M.; Mazzarella, D.; Melchiorre, P. Synthetic Methods Driven by the Photoactivity of Electron Donor-Acceptor Complexes. *J. Am. Chem. Soc.* **2020**, *142* (12), 5461–5476.
- Lima, C. G. S.; de M. Lima, T.; Duarte, M.; Jurberg, I. D.; Paixão, M. W. Organic Synthesis Enabled by Light-Irradiation of EDA Complexes: Theoretical Background and Synthetic Applications. *ACS Catal.* **2016**, *6* (3), 1389–1407.

- (5) Postigo, A. Electron Donor-Acceptor Complexes in Perfluoroalkylation Reactions. *Eur. J. Org. Chem.* **2018**, 2018 (46), 6391–6404.
- (6) Guo, W.; Tao, K.; Xie, Z.; Cai, L.; Zhao, M.; Tan, W.; Liu, G.; Mei, W.; Deng, L.; Fan, X.; Zheng, L. Photodriven Photocatalyst/Metal-Free Direct C-C/C-N Bond Formation: Synthesis of Indoles via EDA Complexes. *J. Org. Chem.* **2019**, 84 (21), 14168–14178.
- (7) Li, Z.; Ma, P.; Tan, Y.; Liu, Y.; Gao, M.; Zhang, Y.; Yang, B.; Huang, X.; Gao, Y.; Zhang, J. Photocatalyst- And Transition-Metal-Free α -Alkylation of: N -Aryl Tetrahydroisoquinolines Mediated by Visible Light. *Green Chem.* **2020**, 22 (3), 646–650.
- (8) Yang, X.; Zhu, Y.; Xie, Z.; Li, Y.; Zhang, Y. Visible-Light-Induced Charge Transfer Enables Csp³-H Functionalization of Glycine Derivatives: Access to 1,3-Oxazolidines. *Org. Lett.* **2020**, 22 (4), 1638–1643.
- (9) Saritha, R.; Annes, S. B.; Saravanan, S.; Ramesh, S. Carbazole Based Electron Donor Acceptor (EDA) Catalysis for the Synthesis of Biaryl and Aryl-Heteroaryl Compounds. *Org. Biomol. Chem.* **2020**, 18 (13), 2510–2515.
- (10) Chen, L.; Liang, J.; Chen, Z.; Chen, J.; Yan, M.; Zhang, X. A Convenient Synthesis of Sulfones via Light Promoted Coupling of Sodium Sulfinates and Aryl Halides. *Adv. Synth. Catal.* **2018**, 361 (5), 956–960.
- (11) Morack, T.; Mück-Lichtenfeld, C.; Gilmour, R. Bioinspired Radical Stetter Reaction: Radical Umpolung Enabled by Ion-Pair Photocatalysis. *Angew. Chem., Int. Ed.* **2019**, 58 (4), 1208–1212.
- (12) Bartolomeu, A. de A.; Silva, R. C.; Brocksom, T. J.; Noël, T.; de Oliveira, K. T. Photoarylation of Pyridines Using Aryldiazonium Salts and Visible Light: An EDA Approach. *J. Org. Chem.* **2019**, 84 (16), 10459–10471.
- (13) Arceo, E.; Bahamonde, A.; Bergonzini, G.; Melchiorre, P. Enantioselective Direct α -Alkylation of Cyclic Ketones by Means of Photo-Organocatalysis. *Chem. Sci.* **2014**, 5 (6), 2438–2442.
- (14) Cao, Z.-Y.; Ghosh, T.; Melchiorre, P. Enantioselective Radical Conjugate Additions Driven by a Photoactive Intramolecular Iminium-Ion-Based EDA Complex. *Nat. Commun.* **2018**, 9 (1), 3274.
- (15) Woźniak, Ł.; Murphy, J. J.; Melchiorre, P. Photo-Organocatalytic Enantioselective Perfluoroalkylation of β -Ketoesters. *J. Am. Chem. Soc.* **2015**, 137 (17), 5678–5681.
- (16) Kandukuri, S. R.; Bahamonde, A.; Chatterjee, I.; Jurberg, I. D.; Escudero-Adán, E. C.; Melchiorre, P. X-Ray Characterization of an Electron Donor-Acceptor Complex That Drives the Photochemical Alkylation of Indoles. *Angew. Chem., Int. Ed.* **2015**, 54 (5), 1485–1489.
- (17) Arceo, E.; Jurberg, I. D.; Álvarez-Fernández, A.; Melchiorre, P. Photochemical Activity of a Key Donor-Acceptor Complex Can Drive Stereoselective Catalytic α -Alkylation of Aldehydes. *Nat. Chem.* **2013**, 5 (9), 750–756.
- (18) Tobisu, M.; Furukawa, T.; Chatani, N. Visible Light-Mediated Direct Arylation of Arenes and Heteroarenes Using Diaryliodonium Salts in the Presence and Absence of a Photocatalyst. *Chem. Lett.* **2013**, 42 (10), 1203–1205.
- (19) Hsu, C.-W.; Sundén, H. α -Aminoalkyl Radical Addition to Maleimides via Electron Donor-Acceptor Complexes. *Org. Lett.* **2018**, 20 (7), 2051–2054.
- (20) Sridharan, V.; Suryavanshi, P. A.; Menéndez, J. C. Advances in the Chemistry of Tetrahydroquinolines. *Chem. Rev.* **2011**, 111 (11), 7157–7259.
- (21) Muthukrishnan, I.; Sridharan, V.; Menéndez, J. C. Progress in the Chemistry of Tetrahydroquinolines. *Chem. Rev.* **2019**, 119 (8), S057–S191.
- (22) Su, D.-S.; Lim, J. J.; Tinney, E.; Wan, B.-L.; Young, M. B.; Anderson, K. D.; Rudd, D.; Munshi, V.; Bahnck, C.; Felock, P. J.; Lu, M.; Lai, M.-T.; Touch, S.; Moyer, G.; DiStefano, D. J.; Flynn, J. A.; Liang, Y.; Sanchez, R.; Prasad, S.; Yan, Y.; Perlow-Poehnelt, R.; Torrent, M.; Miller, M.; Vacca, J. P.; Williams, T. M.; Anthony, N. J. Substituted Tetrahydroquinolines as Potent Allosteric Inhibitors of Reverse Transcriptase and Its Key Mutants. *Bioorg. Med. Chem. Lett.* **2009**, 19 (17), 5119–5123.
- (23) Zhang, J.; Zhan, P.; Wu, J.; Li, Z.; Jiang, Y.; Ge, W.; Pannecouque, C.; De Clercq, E.; Liu, X. Synthesis and Biological Evaluation of Novel 5-Alkyl-2-Arylthio-6-((3,4-Dihydroquinolin-1(2H)-Yl)methyl)Pyrimidin-4(3H)-Ones as Potent Non-Nucleoside HIV-1 Reverse Transcriptase Inhibitors. *Bioorg. Med. Chem.* **2011**, 19 (14), 4366–4376.
- (24) Chander, S.; Wang, P.; Ashok, P.; Yang, L.-M.; Zheng, Y.-T.; Murugesan, S. Rational Design, Synthesis, Anti-HIV-1 RT and Antimicrobial Activity of Novel 3-(6-Methoxy-3,4-Dihydroquinolin-1(2H)-Yl)-1-(Piperazin-1-Yl)Propan-1-One Derivatives. *Bioorg. Chem.* **2016**, 67, 75–83.
- (25) Ramesh, E.; Manian, R. D. R. S.; Raghunathan, R.; Sainath, S.; Raghunathan, M. Synthesis and Antibacterial Property of Quinolines with Potent DNA Gyrase Activity. *Bioorg. Med. Chem.* **2009**, 17 (2), 660–666.
- (26) Jarvest, R. L.; Berge, J. M.; Berry, V.; Boyd, H. F.; Brown, M. J.; Elder, J. S.; Forrest, A. K.; Fosberry, A. P.; Gentry, D. R.; Hibbs, M. J.; Jaworski, D. D.; O'Hanlon, P. J.; Pope, A. J.; Rittenhouse, S.; Sheppard, R. J.; Slater-Radosti, C.; Worby, A. Nanomolar Inhibitors of Staphylococcus Aureus Methionyl tRNA Synthetase with Potent Antibacterial Activity against Gram-Positive Pathogens. *J. Med. Chem.* **2002**, 45 (10), 1959–1962.
- (27) Muñoz, A.; Sojo, F.; Arenas, D. R. M.; Kouznetsov, V. V.; Arvelo, F. Cytotoxic Effects of New Trans-2,4-Diaryl-r-3-Methyl-1,2,3,4-Tetrahydroquinolines and Their Interaction with Antitumoral Drugs Gemcitabine and Paclitaxel on Cellular Lines of Human Breast Cancer. *Chem.-Biol. Interact.* **2011**, 189 (3), 215–221.
- (28) Kouznetsov, V. V.; Merchan-Arenas, D. R.; Tangarife-Castaño, V.; Correa-Royero, J.; Betancur-Galvis, L. Synthesis and Cytotoxic Evaluation of Novel 2-Aryl-4-(4-Hydroxy-3-Methoxyphenyl)-3-Methyl-6,7-Methylenedioxy-1,2,3,4-Tetrahydroquinolines, Podophyllotoxin-like Molecules. *Med. Chem. Res.* **2016**, 25 (3), 429–437.
- (29) Zhu, S.; Das, A.; Bui, L.; Zhou, H.; Curran, D. P.; Rueping, M. Oxygen Switch in Visible-Light Photoredox Catalysis: Radical Additions and Cyclizations and Unexpected C-C-Bond Cleavage Reactions. *J. Am. Chem. Soc.* **2013**, 135 (5), 1823–1829.
- (30) Yadav, A. K.; Yadav, L. D. S. Visible Light Photoredox Catalysis with N-Hydroxyphthalimide for [4 + 2] Cyclization between N-Methylanilines and Maleimides. *Tetrahedron Lett.* **2017**, 58 (6), 552–555.
- (31) Guo, J.-T.; Yang, D.-C.; Guan, Z.; He, Y.-H. Chlorophyll-Catalyzed Visible-Light-Mediated Synthesis of Tetrahydroquinolines from N,N-Dimethylanilines and Maleimides. *J. Org. Chem.* **2017**, 82 (4), 1888–1894.
- (32) Liang, Z.; Xu, S.; Tian, W.; Zhang, R. Eosin Y-Catalyzed Visible-Light-Mediated Aerobic Oxidative Cyclization of N,N-Dimethylanilines with Maleimides. *Beilstein J. Org. Chem.* **2015**, 11, 425–430.
- (33) Nicholls, T. P.; Constable, G. E.; Robertson, J. C.; Gardiner, M. G.; Bissember, A. C. Brønsted Acid Cocatalysis in Copper(I)-Photocatalyzed α -Amino C-H Bond Functionalization. *ACS Catal.* **2016**, 6 (1), 451–457.
- (34) Chen, L.; Chao, C. S.; Pan, Y.; Dong, S.; Teo, Y. C.; Wang, J.; Tan, C.-H. Amphiphilic Methyleneamino Synthons through Organic Dye Catalyzed-Decarboxylative Aminoalkylation. *Org. Biomol. Chem.* **2013**, 11 (35), 5922–5925.
- (35) Xin, J.-R.; Guo, J.-T.; Vigliaturo, D.; He, Y.-H.; Guan, Z. Metal-Free Visible Light Driven Synthesis of Tetrahydroquinoline Derivatives Utilizing Rose Bengal. *Tetrahedron* **2017**, 73 (31), 4627–4633.
- (36) Nicholls, T. P.; Constable, G. E.; Robertson, J. C.; Gardiner, M. G.; Bissember, A. C. Brønsted Acid Cocatalysis in Copper(I)-Photocatalyzed α -Amino C-H Bond Functionalization. *ACS Catal.* **2016**, 6 (1), 451–457.
- (37) Xu, G.-Q.; Li, C.-G.; Liu, M.-Q.; Cao, J.; Luo, Y.-C.; Xu, P.-F. Dual C-H Functionalization of N-Aryl Tetrahydroisoquinolines: A Highly Diastereoselective Synthesis of Dibenzo[a, f]Quinolizines via Visible-Light Induced Oxidation and Inverse Electron-Demand Aza-Diels-Alder Reaction. *Chem. Commun.* **2016**, 52 (6), 1190–1193.

- (38) Sun, D.; Hubig, S. M.; Kochi, J. K. Oxetanes from [2 + 2] Cycloaddition of Stilbenes to Quinone via Photoinduced Electron Transfer. *J. Org. Chem.* **1999**, *64* (7), 2250–2258.
- (39) Wu, G.; Li, Y.; Yu, X.; Gao, Y.; Chen, H. Acetic Acid Accelerated Visible-Light Photoredox Catalyzed N-Demethylation of N,N-Dimethylaminophenyl Derivatives. *Adv. Synth. Catal.* **2017**, *359* (4), 687–692.
- (40) Xu, K.; Fang, Y.; Yan, Z.; Zha, Z.; Wang, Z. A Highly Tunable Stereoselective Dimerization of Methyl Ketone: Efficient Synthesis of E- and Z-1,4-Enediones. *Org. Lett.* **2013**, *15* (9), 2148–2151.
- (41) Curran, D. P.; Keller, A. I. Radical Additions of Aryl Iodides to Arenes Are Facilitated by Oxidative Rearomatization with Dioxygen. *J. Am. Chem. Soc.* **2006**, *128* (42), 13706–13707.
- (42) Li, J.; Bao, W.; Zhang, Y.; Rao, Y. Cercosporin-Photocatalyzed Sp³ (C-H) Activation for the Synthesis of Pyrrolo[3,4-c]-Quinolones. *Org. Biomol. Chem.* **2019**, *17* (40), 8958–8962.
- (43) Ju, X.; Li, D.; Li, W.; Yu, W.; Bian, F. The Reaction of Tertiary Anilines with Maleimides under Visible Light Redox Catalysis. *Adv. Synth. Catal.* **2012**, *354* (18), 3561–3567.
- (44) Liu, Q.; Li, Y.-N.; Zhang, H.-H.; Chen, B.; Tung, C.-H.; Wu, L.-Z. Reactivity and Mechanistic Insight into Visible-Light-Induced Aerobic Cross-Dehydrogenative Coupling Reaction by Organo-photocatalysts. *Chem. - Eur. J.* **2012**, *18* (2), 620–627.
- (45) Yamaguchi, M.; Fujiwara, S.; Manabe, K. Synthesis of 2,2,5-Trisubstituted 2H-Pyrroles and 2,3,5-Trisubstituted 1H-Pyrroles by Ligand-Controlled Site-Selective Dearomative C2-Arylation and Direct C3-Arylation. *Org. Lett.* **2019**, *21* (17), 6972–6977.
- (46) Xuan, J.; Feng, Z.-J.; Chen, J.-R.; Lu, L.-Q.; Xiao, W.-J. Visible-Light-Induced C-S Bond Activation: Facile Access to 1,4-Diketones from β -Ketosulfones. *Chem. - Eur. J.* **2014**, *20* (11), 3045–3049.
- (47) Sheldrick, G. M. A Short History of SHELX. *Acta Crystallogr., Sect. A: Found. Crystallogr.* **2008**, *64* (1), 112–122.
- (48) Sheldrick, G. M. Crystal Structure Refinement with SHELXL. *Acta Crystallogr., Sect. C: Struct. Chem.* **2015**, *71* (1), 3–8.
- (49) Barbour, L. J. X-Seed — A Software Tool for Supramolecular Crystallography. *J. Supramol. Chem.* **2001**, *1* (4), 189–191.
- (50) Mandal, A. K.; Sreejith, S.; He, T.; Maji, S. K.; Wang, X.-J.; Ong, S. L.; Joseph, J.; Sun, H.; Zhao, Y. Three-Photon-Excited Luminescence from Unsymmetrical Cyanostilbene Aggregates: Morphology Tuning and Targeted Bioimaging. *ACS Nano* **2015**, *9* (5), 4796–4805.
- (51) Katritzky, A. R.; Yao, G.; Lan, X.; Zhao, X. The Conversion of Secondary into Tertiary Amides Using Benzotriazole Methodology. *J. Org. Chem.* **1993**, *58* (8), 2086–2093.
- (52) Barker, T. J.; Jarvo, E. R. Umpolung Amination: Nickel-Catalyzed Coupling Reactions of N,N-Dialkyl-N-Chloroamines with Diorganozinc Reagents. *J. Am. Chem. Soc.* **2009**, *131* (43), 15598–15599.